Strategic Information Exchange

Dinah Rosenberg[∗] , Eilon Solan† and Nicolas Vieille‡

November 18, 2018

Abstract

We study a class of two-player repeated games with incomplete information and informational externalities. In these games, two states are chosen at the outset, and players get private information on the pair, before engaging in repeated play. The payoff of each player only depends on his 'own' state and on his own action. We study to what extent, and how, information can be exchanged in equilibrium. We prove that provided the private information of each player is valuable for the other player, the set of sequential equilibrium payoffs converges to the set of feasible and individually rational payoffs as players become patient.

Whether and how to acquire information is a question faced by most decision makers. In statistical decision problems, the decision maker tries to learn the value of an unknown parameter, and he can sample from an exogenous population at a fixed cost per draw. In other contexts, information is held by strategic agents. In signalling games for instance, a player holding payoff-relevant private information tries to influence the action choice of an uninformed party. There, the rationale for disclosing/hiding information is that the uninformed party's action affects the payoff of the informed player. We here study a class of repeated games with private

[∗]Economics and Decision Sciences Department, HEC Paris, and GREG-HEC; . e-mail: rosenberg@hec.fr

[†]The School of Mathematical Sciences, Tel Aviv University, Tel Aviv 69978, Israel. e-mail: eilons@post.tau.ac.il

[‡]Economics and Decision Sciences Department, HEC Paris, and GREG-HEC. e-mail: vieille@hec.fr.

information, in which there is no such direct strategic interaction: as in setups of social learning, or of strategic experimentation, payoffs do not depend upon other players' actions. However, players hold private information that has value to other players. Our goal is to understand to what extent information can be exchanged at equilibrium along the play, assuming communication is costly. This assumption of pure informational externalities plays a dual role. Obviously, it simplifies the analysis of the model and allows to study the exchange of information in isolation from other strategic considerations. But it leads to a game in which we might least expect exchange of information, and any positive result in this setup might potentially open the way for the analysis of other setups.

As an illustration, consider the following game. There are two biased coins, C_1 and C_2 (say, with parameter $\frac{2}{3}$), which are tossed independently *once* at the outset of the game. Each of two players, $i = 1, 2$, has to repeatedly guess the outcome of $\text{coin } C_i$. A correct guess yields a payoff of one, while an incorrect one yields zero, and successive payoffs are discounted. If past payoffs are not observed, there is no role for direct inference, and it is natural to expect that player i will repeatedly 'guess' the most likely outcome, for an expected payoff of $\frac{2}{3}$. Assume that, once coins are drawn, each player *i* gets to observe the outcome of coin C_j , where $j \neq i$, but that cheap talk is excluded. This private information has no 'direct' value, but it might have a strategic value, because it is valuable to the other player. In this game, is there an equilibrium payoff that improves upon $(\frac{2}{3}, \frac{2}{3})$ $\frac{2}{3}$)? While stylized, this game is similar to the situation faced by executives of two different firms, who hold private information on their own firm. Trading the stock of one's own firm is illegal, at times where private information is most valuable, and the executives may be tempted to implement some implicit collusive scheme of information exchange through time.

We start with few simple observations. Since cheap talk is assumed away, exchange of information is to take place through actions, by 'encoding' privately held information into actions, and by conditioning the action choice of player i on the outcome of coin C_j . Plainly, there is no equilibrium in which each player *i fully* 'discloses' the outcome of C_j at stage 1. Indeed, once informed of the outcome of C_2 , player 2 can make the correct guess in all subsequent periods, and has no incentive whatsoever to incur the cost of disclosing information to player 1, including in stage 1. But then, player 1 would not be willing to play the myopically suboptimal action in stage 1. More generally, private information is here an asset to be exchanged for private information, at a cost. On the one hand, a player cannot disclose information without assigning positive probability to his myopically suboptimal action, and thereby incurring a cost.^{[1](#page-2-0)} On the other hand, a player is not willing to play a myopically suboptimal action, unless he expects to be rewarded with valuable information in return: no player wants to be the last one to disclose information. Not surprisingly, when the horizon is finite we prove that $(\frac{2}{3}, \frac{2}{3})$ $\frac{2}{3}$) is the unique equilibrium payoff. Having an infinite horizon raises the possibility of a gradual, open-ended exchange of information. Since however, the total 'amount' of information to be exchanged is bounded, the feasibility of such a process is not ensured.

Our model is a generalization of the stylized example. Two 'states', s and t, are drawn independently, and players get private information on both s and t^2 t^2 . Next, the players repeatedly choose actions, which are publicly disclosed. The crucial assumption we make is that a player's payoff only depends on 'his' state, and on his own action.

We prove that, provided that the information held by each player is valuable to the other player, the limit set (as $\delta \rightarrow 1$) of sequential equilibrium payoffs coincides with the set of all feasible payoffs, that are at least equal to the initial, myopic optimal payoffs. In the simple example discussed above, this limit set is thus equal to the set $\left[\frac{2}{3}, 1\right] \times \left[\frac{2}{3}\right]$ $\frac{2}{3}$, 1. Not only information can be shared, but the rate of information exchange can be arbitrarily high relative to the discount rate. Our equilibria share the following features. Players start by reporting truthfully whatever information they received on their own state. This leads to a continuation game in which no player holds private information on his own state. As a result, each player is able to compute how costly it is for the other player to play his suboptimal action, and is therefore able to adjust accordingly the amount of information he discloses as a 'reward'. Players next exchange information in an open-ended manner. The analysis presents two main and mostly independent difficulties. One is to design open-ended equilibrium processes, according to which information is exchanged. In our construction, the bulk of information exchange takes place early in the game. Later information disclosure only serves as a means to compensate for previously incurred costs. The second consists in adjusting this continuation play so as to provide the incentives for truthful reporting of one's information on one's own state.

¹Unless if indifferent, but this possibility will play no role.

²That is, each player receives information both on his state and on the other player's state.

Our motivation stems from repeated games with incomplete information. The literature on such games started with Aumann and Maschler (1966, 1995), and was extensively developed under the assumption of no discounting, see chapters 5 and 6 in Aumann and Hart (1992). When there is no discounting, communication through actions becomes costless, and our model, trivial. Besides the literature on reputation models, see Mailath and Samuelson (2006) for a survey, there is only limited work dealing with discounted repeated games with incomplete information. Recent con-tributions are Cripps and Thomas (200[3](#page-3-0)), Peski (2008) and Wiseman (2005).³ Both Cripps and Thomas (2003), and Peski (2008) look at games with one-sided information, in which each of the two players knows his own payoff function, and one of the two is unsure of the payoff function of the other player. Cripps and Thomas (2003) prove that a Folk Theorem type of result holds in the limit where the prior belief converges to the case of complete information. Peski (2008) essentially shows that all equilibria are payoff-equivalent to equilibria that involve finitely many rounds of information revelation. Wiseman (2005) looks at situations of common uncertainty. Players share the same information on the underlying state of nature, and refine this information by observing actual choices and payoffs.

Starting with Crawford and Sobel (1982), the huge literature on strategic information transmission and on cheap-talk games addresses issues related to ours. The paper that is closest to our work is Aumann and Hart (2003). There, prior to playing a game once, two players, one of which is informed of the true game to be played, exchange messages during countably many periods. Aumann and Hart (2003) characterize the set of equilibrium payoffs. Following an example of Forges (1990), they show that allowing for an unbounded communication length may increase the set of equilibrium payoffs. There are however significant differences with our setup. On the one hand, this literature allows the game to exhibit informational and strategic interaction as well. On the other hand, information is one-sided, and communication is costless.

Finally, the pattern of information disclosure in our setup is reminiscent of the pattern of contributions in dynamic models of public good contributions, see Admati and Perry (1991), Marx and Matthews (2000), or in the dynamic resolution of the hold-up problem, see Che and Sakovics (2004) . More generally, and as Compte and Jehiel (2004) argue, the existence of a history-dependent outside option forces equi-

³An older one is Mayberry (1967).

librium concessions to be gradual in many bargaining situations, just as information disclosure has to be gradual and open-ended here. There are however differences between the results on the two models. First, in dynamic models of public good contributions the evolution of contributions follows a deterministic trend, while in games with incomplete information beliefs follow a martingale. A more significant difference is the following. In the former models, there is a one-to-one relation between the contributed amount and the cost incurred when contributing: the more a player contributes, the higher his cost by doing so. Here, the cost of disclosing information is independent of the amount of information that is disclosed. The reason is that the cost of disclosure is incurred when playing a (myopically) suboptimal action, while the amount of information is a function of how revealing such an action is. The more precise the belief, the higher the cost of disclosing information.

The paper is organized as follows. Section [1](#page-4-0) contains the model and a statement of our main results. Section [2](#page-11-0) presents the main ideas of the proof through a version of the example discussed above (allowing for private signals). Section [3](#page-19-0) is devoted to concluding comments. Proofs are provided in the Appendix.

1 Model and Main Result

1.1 The Game

We study a class of two-player repeated games with incomplete information. At the outset of the game, a state of the world is realized, and the two players receive private signals, $l \in L$ and $m \in M$ respectively. At each stage $n \geq 1$, players choose actions from the action sets A and B. Actions, and only actions, are then publicly disclosed.

All sets are finite. Players share a common discount factor $\delta < 1$.

We make the following assumption:

- **A.1** The set of states of the world is a product set, $S \times T$, with elements denoted by (s, t) . Player is payoff depends only on his own action and on the i-th component of the state. That is, the payoff function of player 1 is a function $u : S \times A \to \mathbf{R}$, while player 2's payoff is given by a function $v : T \times B \to \mathbf{R}$.
- **A.2** Signal sets are also product sets, $L = L_S \times L_T$ and $M = M_S \times M_T$, with elements denoted by $l = (l_S, l_T)$ and (m_S, m_T) . The random triples (s, l_S, m_S) and

 $(\mathbf{t}, \mathbf{l}_T, \mathbf{m}_T)$ are drawn independently of each other according to the distributions $p \in \Delta(S \times L_S \times M_S)$ and $q \in \Delta(T \times L_T \times M_T)$ respectively.^{[4](#page-5-0)}

Assumption A.1 ensures that the game is one of pure informational externalities. Player 1 cares about player 2's behavior only to the extent that player 2's behavior conveys information about s.

The independence assumption $\mathbf{A.2}$ is often made in games with two-sided incomplete information, see, e.g., Zamir (1992). Not only does it imply that the two states s and t are independent, but also that the two private signals, say of player 1, \mathbf{l}_S and \mathbf{l}_T , are independent. The first component \mathbf{l}_S of player 1's private signal should be thought of as the information received by player 1 on his own state s, while the second component \mathbf{l}_T is player 1's information on player 2's state, \mathbf{t} .^{[5](#page-5-1)} Besides allowing for tractability, assumption A.2 implies that behaving myopically is an equilibrium. That is, assume that player 1 repeatedly plays an action a_{\star} that maximizes the expectation of $u(\mathbf{s}, a)$, given \mathbf{l}_S . Then, by **A.2**, the belief held by player 2 over his own state does not change along the play, and it is a best reply for player 2 to repeatedly play an action b_{\star} that maximizes the expectation of $v(\mathbf{t}, b)$, given \mathbf{m}_T . And vice-versa. If instead the two states s and t were correlated, then the choice of a_{\star} might be informative about t, and may lead to a change in player 2's action, which would in turn be informative about s. But then, to manipulate this informational feedback, player 1 might be tempted to 'mis-represent' his myopically action a_{\star} . Such an example is provided in Section [3.2.](#page-21-0) Assumption A.2 assumes away these effects.

In a sense, assumptions $\mathbf{A.1}$ and $\mathbf{A.2}$ imply that the only motive for disclosing information on the other player's state is to get back information in exchange: no information will ever be exchanged, unless out of purely strategic reasons. We stress that, although the triples (s, l_S, m_S) and (t, l_T, m_T) are independent, we make no assumption on the distributions p and q . Thus, the two signals relative to a *given* state may be correlated in an arbitrary way among themselves, or with the state itself.

The main question that we ask is whether and to what extent valuable information can be exchanged at equilibrium, and how to organize the exchange of information

⁴Here, the state of the world is (s, t) , the signal to player 1 is (l_S, l_T) , and the signal to player 2 is (m_S, m_T) . We will refer to s (resp., to t) as to player 1's state (resp., player 2's state).

⁵The subscripts in l_S , l_T serve a mnemonic purpose. We use boldface letters to denote random variables, when we fear confusion might be at stake.

along the play, while meeting equilibrium requirements. Our main result consists of a characterization of the limit set of sequential equilibrium payoffs, as players become patient.

Strategies will be denoted by σ and τ for players 1 and 2 respectively. A behavior strategy of player i maps his private information and the public history of past moves, into a mixed action. Accordingly, behavior strategies are maps σ : $L \times H \to \Delta(A)$ and $\tau : M \times H \to \Delta(B)$, where $H = \cup_{n \geq 0} (A \times B)^n$ is the set of finite sequences of moves. A strategy pair (σ, τ) induces a probability distribution over the set of (infinite) plays. Expectations under this distribution are denoted by $\mathbf{E}_{p,q,\sigma,\tau}$. Thus, the expected discounted payoff of player 1 is given by $\gamma_{\delta}^1(p,q,\sigma,\tau) = \mathbf{E}_{p,q,\sigma,\tau}[(1-\delta)\sum_{n=1}^{\infty}\delta^{n-1}u(\mathbf{s},\mathbf{a}_n)].$ We denote by $\mathbf{p}_n \in \Delta(S \times M_S)$ the belief held by player 1 at stage n: it is the conditional distribution of the pair $({\bf s}, {\bf m}_s)$, given I, and the (public) sequence of previous moves. Note that p_1 depends only on l, because there is no history of moves prior to stage 1. However, the computation of \mathbf{p}_n involves player 2's strategy for $n > 1$. We denote by $\mathbf{q}_n \in \Delta(T \times L_T)$ the belief of player 2 at stage n on his state of nature and on the signal that player 1 received on this state of nature.

1.2 Preliminaries

Loosely put, our main result is the following. Provided that each player holds information that is valuable to the other player, and that players are patient, any feasible and individually rational payoff is a sequential equilibrium payoff. Before we state formally this result, we list some preliminary remarks, and give few definitions.

1.2.1 Myopically optimal payoffs

Given a probability distribution $\pi \in \Delta(S)$ and an action $a \in A$, we write $u(\pi, a)$ for the expected payoff of player 1 when holding the belief π and playing action a:

$$
u(\pi, a) = \mathbf{E}_{\pi}[u(\cdot, a)] = \sum_{s \in S} \pi(s)u(s, a).
$$

We will often abuse notation and write $u(\pi, a)$ whenever π is a distribution over a product space of the form $S \times Q$, for some finite set Q. In that case, $u(\pi, a)$ =

 \sum s∈S $\pi({s} \times Q)u(s, a)$. Given a distribution $\pi \in \Delta(S)$, the myopically optimal payoff when holding the belief π is

$$
u_\star(\pi) := \max_{a \in A} u(\pi, a). \tag{1}
$$

For a given $a \in A$, the map $\pi \mapsto u(\pi, a)$ is affine. As a supremum of finitely many affine functions, the map u_* is convex and piecewise linear. An action $a \in A$ is *optimal* at π if it achieves the maximum in [\(1\)](#page-7-0).

Recall that $\mathbf{p}_1 \in \Delta(S \times M_S)$ is the (*interim*) belief of player 1 prior to stage 1. Thus, the probability $\mathbf{p}_1(s, m_S)$ assigned to (s, m_S) is equal to $p(s, m_S|\mathbf{l}_S)$. A myopic strategy of player 1 is a strategy σ_m that repeats the same, optimal action at \mathbf{p}_1 . The (ex ante) payoff induced by σ_m does not depend on player 2's strategy, and is equal to

$$
u_{\star} := \mathbf{E}_p \left[u_{\star}(\mathbf{p}_1) \right] = \sum_{l_S \in L_S} p(l_S) u_{\star}(p(\cdot | l_S)).
$$

If player 2's strategy does not depend on \mathbf{m}_S , the belief of player 1 does not change along the play: $\mathbf{p}_n = \mathbf{p}_1$ for each $n \geq 1$. In such a case, the expected payoff of player 1 does not exceed u_{\star} . Thus, u_{\star} is the minmax value for player 1 in the repeated game. For similar reasons,

$$
v_{\star} := \mathbf{E}_p \left[v_{\star}(\mathbf{q}_1) \right] = \sum_{m_T \in M_T} q(m_T) v_{\star}(q(\cdot|m_T))
$$

is the minmax value for player 2.

Player 1's payoff is highest when he knows all he may possibly know about s, given the rules of the game,^{[6](#page-7-1)} that is, when player 2's signal \mathbf{m}_S is made public. Player 1's belief is then denoted by $\tilde{\mathbf{p}} \in \Delta(S)$; thus, $\tilde{\mathbf{p}}(s) = p(s|\mathbf{l}_S, \mathbf{m}_S)$, for each state $s \in S$. Conditional on both signals \mathbf{l}_S and \mathbf{m}_S , player 1's optimal payoff is $u_\star(\tilde{\mathbf{p}})$. Therefore, player 1's ex ante expected payoff does not exceed

$$
u_{\star\star} := \mathbf{E}_p[u_{\star}(\tilde{\mathbf{p}})] = \sum_{l_S \in L_S, m_S \in M_S} p(l_S, m_S)u_{\star}(p(\cdot|l_S, m_S)).
$$

Since u_* is convex, one has $u_{**} \geq u_*$. This reflects the fact that the marginal value of the information held by player 2 is nonnegative.

⁶This can be formally deduced from Blackwell Theorem [\[6\]](#page-24-0).

Conversely, any payoff in $[u_{\star}, u_{\star\star})$ is a feasible payoff for player 1, provided players are patient. Therefore, the (limit) set of feasible and individually rational payoffs for player 1 is the interval $[u_\star, u_{\star\star}].$

Similarly, we define $\tilde{\mathbf{q}} = q(\cdot|\mathbf{l}_T, \mathbf{m}_T)$ and $v_{**} := \mathbf{E}_q[v_*(\tilde{\mathbf{q}})]$. As for player 1, the limit set of feasible and individually rational payoffs for player 2 is the interval $[v_\star, v_{\star\star}]$.

The example discussed in the introduction will serve as a leading example. Here, all four sets S, T, A and B are equal to $\{0,1\}$. Payoffs are given by $u(s,a) = 1$ if $s = a$, and $u(s, a) = 0$ if $s \neq a$ (resp., $v(t, b) = 1$ if $t = b$, and $v(t, b) = 0$ if $t \neq b$). We will refer to this setup as the **Binary Example**, and we will use it repeatedly, with various information structures. Note that the myopically optimal action is $a = 1$ if and only if the belief assigned by player 1 to $s = 1$ is at least $1/2$.

Binary Example 1 Assume here that L_S is a singleton, while player 2 observes s (that is, $M_S = S$, and $p(s, m_S) = 0$ if $s \neq m_S$). The belief of player 1 can be identified with the probability π assigned to state 1, and $u_{\star}(\pi) = \max{\pi, 1 - \pi}$. On the other hand, since player 2 observes **s**, \tilde{p} is either 0 or 1, and $u_{\star\star} = 1$.

1.2.2 Valuable information

A player will not be willing to play a myopically suboptimal action unless he expects to receive information in return, the marginal value of which offsets the cost incurred when playing the suboptimal action. In particular, a *necessary* condition for improving upon (u_*, v_*) is that each player holds information that is valuable to the other player.

We stress that it is not enough that each player holds information on the other player's state. Indeed, if, e.g., the two signals \mathbf{l}_s and \mathbf{m}_s coincide p-a.s., player 1 already knows all player 2 knows relative to s. Nor is it enough that players have private information, as the next example shows.

Binary Example 2 Assume that $p(s = 1) = \frac{3}{4}$ and that player 2 receives a binary signal m_S which (conditional on s) is correct with probability $\frac{2}{3}$. Assume moreover that L_S is a singleton, so that player 1 has no information and $p_1 = p = \frac{3}{4}$ $\frac{3}{4}$, and $u_{\star} = \frac{3}{4}$ $\frac{3}{4}$. By Bayes' rule, the posterior probability $\tilde{\boldsymbol{p}}$ assigned to state 1 is equal to $\tilde{\mathbf{p}} = \frac{6}{7}$ $\frac{6}{7}$ if $\mathbf{m}_S = 1$, and is equal to $\frac{3}{5}$ if $\mathbf{m}_S = 0$. In either case, the optimal action at \tilde{p} is a = 1, and, therefore, knowing the information of player 2 will not change the

optimal behavior of player 1: the information held by player 2 is valueless. Note that $u_{\star\star} = \mathbf{E}[u(\tilde{\boldsymbol{p}}, 1)] = u(p, 1) = u_{\star}$, where the middle equality holds by the law of iterated expectations.

In the previous example, the information held by player 2 is valueless because it does not affect player 1's optimal action. Indeed, the precision of player 2's signal is lower than the prior evidence that the state is 1, hence player 2's signal cannot provide decisive evidence against state 1.

This observation motivates the definition below.

Definition 1 The information of player 2 is valuable for player 1, if

$$
\mathbf{E}\left[u_{\star}\left(\tilde{\mathbf{p}}\right)|\mathbf{l}_{S}\right] > u_{\star}(\mathbf{p}_{1}), \quad \text{with } p-probability \, 1. \tag{2}
$$

Similarly, the information of player 1 is valuable for player 2 if $\mathbf{E}[v_{\star}(\tilde{\mathbf{q}})|\mathbf{m}_T] >$ $v_{\star}(\mathbf{q}_1)$, with q-probability 1.

Condition [\(2\)](#page-9-0) is an interim requirement: it is equivalent to requiring that, after learning \mathbf{I}_S , player 1 assigns positive probability to the event that his optimal action would change, if player 2's signal \mathbf{m}_S were made public.

This condition implies that $u_{**} > u_{*}$, but it is not implied by this inequality. Indeed, the latter condition is an ex ante requirement, that amounts to requiring that $\mathbf{E}[u_{\star}(\tilde{\mathbf{p}})|\mathbf{l}_S] > u_{\star}(\mathbf{p}_1)$ holds with positive *p*-probability.

Definition [1](#page-9-1) does not provide a measure of information value, but only a criterium for deciding whether the information held by a player has positive value or not. Note that this criterium involves player 1's utility function u , and is therefore gamedependent.

1.3 Main result

Our main result is the following.

Main Theorem 1 Assume that the information of each player is valuable to the other player. Then, as $\delta \to 1$, the set of sequential equilibrium payoffs converges to the set $[u_\star, u_{\star\star}] \times [v_\star, v_{\star\star}].$

The proof of Theorem [1](#page-9-2) is provided in the Appendix. Formally we prove that given any compact set $W \subset (u_\star, u_{\star\star}) \times (v_\star, v_{\star\star})$, there is $\delta_0 < 1$ such that each vector in W is a sequential equilibrium payoff of the δ -discounted game, as soon as $\delta \geq \delta_0$.

Note that player 1's expected payoff does not exceed $(1 - \delta)u_{\star} + \delta u_{\star\star}$, because player 1's action in stage 1 is based only upon l. It will follow from the proof of Theorem [1](#page-9-2) that equilibrium payoffs as high as $u_{**} - (1 - \delta)c$ can be implemented in the δ -discounted game, for some constant c. These two estimates give a rate of convergence of the set of equilibrium payoffs in the δ -discounted game.

The conclusion of the theorem is still valid when players hold different discount factors, which converge to 1.

A few comments are in order. The theorem gives a sharp answer to the question we posed. There exist equilibria, in which almost all information can be exchanged, with a negligible delay, in spite of the fact that information exchange must be gradual and open-ended. This stands in sharp contrast to conclusions obtained in dynamic models of public good contribution, see Compte and Jehiel (2004). The driving force that explains this contrast is the following. Here, the cost of disclosing information is the opportunity cost of playing a suboptimal action, while the amount of information depends on the evolution of beliefs and, therefore, on the extent to which actions are correlated with private information. As a result, cost and amount are disentangled. By contrast, in a public good model, there is a one-to-one relation between the cost of a given monetary contribution, and the amount contributed.

When the information, say, of player 2, has a positive ex ante value $(u_{**} > u_{*}),$ but is not valuable according to Definition [1,](#page-9-1) the conclusion of the theorem does not hold. Indeed, whenever the signal \mathbf{l}_S received by player 1 fails to satisfy inequality [\(2\)](#page-9-0), player 1 will infer (at the interim stage) that the information held by player 2 has no value. Then, player 1 will not be willing to disclose any information to player 2, although the information he holds may be valuable to player 2. In Section [3,](#page-19-0) we extend the characterization of the theorem to cover such cases.

The extension to an arbitrary number of players is outside of the scope of this paper. With more than two players, there exist cases where some player i holds no private information, yet receives information in equilibrium. The basic intuition is that player i may be 'rewarded' by some other player j , for information that has been disclosed to j by a third player k .

2 The Binary Example

We here explain the main ideas of the equilibrium construction within the binary setup. For simplicity, we assume here that as in the example in the Introduction, each player knows exactly the other player's state.[7](#page-11-1)

All equilibria share the feature that each player starts by reporting truthfully his private signal relative on his *own* state, \mathbf{l}_S and \mathbf{m}_T respectively. The rationale for this report is the following. In the continuation game, no player then holds *private* information on his *own* state. Consequently, player i will always know the posterior belief of player j on j's state. This in turn allows player i to compute the perceived cost incurred by player j when playing either action, and to adjust accordingly the amount of information he discloses. Of course, adequate incentives will have to be provided to ensure truthful reporting.

We first deal with games in which players get no information on their own state, which we call *self-ignorant games*. We next explain how to provide incentives for a truthful report in the general case. The discussion of how to combine these two logically independent steps is relegated to the Appendix.

2.1 The self-ignorant case

In this section, we analyze the game discussed in the introduction. The states s and t are drawn according to $p \in \Delta(S)$ and $q \in \Delta(T)$. Player 1 is told t, and player 2 is told s. Hence, p (and q) may be viewed as a distribution over $S = \{0, 1\}$. We identify any distribution over S (resp., over T) with the probability assigned to state $s = 1$ (resp., to state $t = 1$). For concreteness, we assume that p and q are such that $p, q > \frac{1}{2}.$

2.1.1 A first equilibrium profile

We here describe one specific equilibrium profile that will serve as a building block for the general construction. In this profile, the play is divided into two phases. In the first phase, at odd stages (resp. at even stages) player 1 (resp. player 2) randomizes, thereby transmitting information to player 2 (resp. to player 1), while player 2 (resp. player 1) plays his myopically optimal action. The second phase starts

⁷That is, $I_T = t$ with q-probability 1, and $m_S = s$ with p-probability 1.

when the player who is randomizing plays his myopically optimal action, and lasts ad infinitum. Along this phase both players play their myopically optimal action. Thus, the play path looks as follows: in the first few stages the players alternately play their myopically suboptimal action, and then the play switches to myopic play.

Randomizations are informative: in stage 1 for instance, the mixed action used by player 1 depends on t, so that the belief of player 2 in stage 2 depends on player 1's action in stage 1.

The play in the first phase has a cyclic pattern, with period 4. The evolution of beliefs along the play is illustrated in Figure 1. This figure involves two parameters, p^* and q^* , which will later be pinned down by equilibrium requirements.

Figure 1: The play as long as both players play suboptimally.

How to read this figure? Consider first stage 1. In stage 1, player 2 plays with probability 1 his optimal action (which is $b = 1$ since $q > \frac{1}{2}$). Consequently, the belief of player 1 in stage 2 is p , the same as in stage 1. Meanwhile, player 1 plays his suboptimal action $a = 0$ with probability x_t . The values of x_0 and x_1 are defined to be $x_1 := \bar{x} =$ $1 - q$ q $\times \frac{q^{\star} - q}{\cdot}$ $\frac{q}{q+q^*-1}$ and $x_0 := \underline{x} =$ q $1 - q$ $\times \frac{q^{\star} - q}{\cdot}$ $\frac{q}{q+q^*-1}$, where q^* will be defined below. By Bayesian updating, the posterior belief q_2 of player 2 in stage 2 is equal to $1-q < \frac{1}{2}$ following $a = 0$, and is equal to $q^* > q$ following $a = 1$. In player 2's eyes, the suboptimal action $a = 0$ is played with probability $x := q\bar{x} + (1-q)\underline{x} = \frac{1-\delta}{\delta^2}$ $\frac{-b}{\delta^2}$. Note that x and q^* solve $q = x(1-q) + (1-x)q^*$, which reflects the martingale property of beliefs. Because $q > 1/2$ it follows that $q^* > q$ and therefore $x_1 < x_0$.

Consider next stage 2. If player 1 played his optimal action $a = 1$ in stage 1, players stop exchanging information, and repeat forever their optimal actions, $a = 1$ and $b = 1$. If player 1 played his suboptimal action $a = 0$, player 2 reciprocates and assigns a probability y_s to his suboptimal action (which is now $b = 1$ because $1 - q < \frac{1}{2}$). The values of y_0 and y_1 , are set to $y_1 := \bar{y} =$ $1-p$ p $\times -\frac{p^{\star}-p}{1}$ $\frac{P}{p+p^*-1}$ and $y_0 := y =$ p $1-p$ $\times -\frac{p^{\star}-p}{\cdot}$ $\frac{p}{p+p^*-1}$ (where p^* is defined below) so that the belief of player 1 in stage 3 is either p^* or $1-p$. The overall probability assigned to the action $b=1$ is $y := p\bar{y} + (1 - p)y$, which solves $p = y(1 - p) + (1 - y)p^{*}$.

Consider now stage 3 (assuming $a = 0$ was played in stage 1). If the optimal action $b = 0$ was played in stage 2, players stop exchanging information and repeat their optimal actions, $a = 1$ and $b = 0$. If player 2 played his suboptimal action in stage 2, player 1 reciprocates and plays as in stage 1, only with the roles of the states/actions exchanged. To be precise, player 1 assigns positive probability to his suboptimal action (which is now $a = 1$ because $1 - p < \frac{1}{2}$). As in stage 1, this probability is set to \bar{x} , if the true state t happens to be the state which player 2 currently considers more likely (which is now state $t = 0$ because $\mathbf{q}_3 = 1 - q < \frac{1}{2}$), and it is set to \underline{x} otherwise. By Bayesian updating, the belief of player 2 in stage 4 is either q or $1 - q^*$. And so on.

Observable deviations^{[8](#page-13-0)} by player i are ignored by player j.

The value of p^* is dictated by equilibrium requirements. Observe first that, at equilibrium and at any node where player 1 is supposed to randomize, his expected continuation payoff must be equal to $u_{\star}(p)$. Indeed, consider any such node, say at stage $2n + 1$. If he plays his optimal action, player 1 gets an expected payoff of $u_{\star}(\mathbf{p}_{2n+1})$ in stage $2n+1$, and in all future stages as well, since the players then stop to exchange information. Note that $u_{\star}(\mathbf{p}_{2n+1}) = u_{\star}(p)$, because \mathbf{p}_{2n+1} is either equal to p or $1 - p$, depending wether n is even or odd. Because he is randomizing, player 1 should be indifferent between both actions, and the claim follows.

Let us now consider stage 1. If player 1 plays his suboptimal action in stage 1, his continuation payoff from stage 3 on, is equal to $u_{\star}(p)$ if player 2 plays $b = 1$ in stage 2, and to $u_{\star}(p^{\star})$ if player 2 plays $b = 0$. Since the probabilities of these two events

⁸Such deviations consist in playing the currently suboptimal action at a node at which the optimal action was expected.

are equal to y and $1 - y$ respectively, the overall payoff of player 1 is then equal to

$$
(1 - \delta)(1 - u_\star(p)) + \delta(1 - \delta)u_\star(p) + \delta^2(yu_\star(p) + (1 - y)u_\star(p^\star)),
$$

where the first two terms are the contributions of the first two stages to the overall payoff. When equating this last expression with $u_\star(p)$, and using $p > \frac{1}{2}$, one obtains

$$
p^* = p + (2p - 1)\frac{1 - \delta}{\delta^2 + \delta - 1}.
$$

The same argument, when applied to player 2, yields $q^* = q + (2q - 1) \frac{1 - \delta}{2q}$ $\frac{1}{\delta^2 + \delta - 1}.$

The parameter values $p^*, q^*, \bar{x}, \underline{x}, \bar{y}, y$ all lie in $(0, 1)$ as soon as $\varepsilon_\delta \leq p, q \leq 1 - \varepsilon_\delta$, where $\varepsilon_{\delta} = \frac{1-\delta}{\delta^2+\delta}$ $\frac{1-\delta}{\delta^2+\delta-1}$; that is, as soon as initial beliefs are not too precise. Note that ε_{δ} goes to 0 as δ goes to 1.

Conversely, and whenever $\varepsilon_{\delta} \leq p, q \leq 1-\varepsilon_{\delta}$, this profile is a Nash equilibrium of the repeated game. Observe that, while player 1's payoff is $u_{\star}(p)$, the payoff of player 2 is equal to

$$
f(q) := (1 - \delta)v_{\star}(q) + \delta (x v_{\star}(q) + (1 - x)v_{\star}(q^{\star}))
$$

= $v_{\star}(q) + \frac{1 - \delta}{\delta} (2v_{\star}(q) - 1) > v_{\star}(q).$

2.1.2 Further equilibrium payoffs

We here build upon the previous section, and introduce a class of simple equilibrium profiles, that implement all equilibrium payoffs (in the limit $\delta \to 1$).

In these equilibria, most of the information on player 2's state that player 1 will ever transmit is transmitted at stage 1 by randomizing in that stage, and similarly, most of the information on player 1's state that player 2 will ever transmit is transmitted in stage 2. From stage 3 on, the players either implement the equilibrium profile defined in the previous section, or switch to a myopic behavior. The equilibrium that is implemented from stage 3 on depends on the cost of information transmission in stages 1 and 2.

We will now be more precise. Let $\bar{q}, q, \bar{p}^0, p^0, \bar{p}^1$ and p^1 be arbitrary beliefs in (0, 1), such that $0 < \underline{q} \leq \frac{1}{2} < q < \overline{q} < 1$, and $0 < \underline{p}^{a} \leq \frac{1}{2} < p < \overline{p}^{a} < 1$ for $a \in \{0,1\}$. A further condition will be imposed later. We let the discount factor δ be sufficiently large such that all six beliefs, which are determined below, lie in the interval $(\varepsilon_{\delta}, 1-\varepsilon_{\delta}).$

Player 1 randomizes in stage 1, and plays his optimal action in stage 2. Player 2 plays his optimal action in stage 1, and randomizes in stage 2. We choose these state-dependent randomizations in such a way that the beliefs of the players evolve as indicated in Figure 2. Continuation payoffs from stage 3 also appear on this figure. Observe that the optimal action of player 2 in stage 2 depends on the action played by player 1 in stage 1, since $q \leq \frac{1}{2} < \bar{q}$.

p, q	b_1	p, \overline{q}	b_1	p, \overline{q}	p^1, \overline{q} : information exchange; continuation payoffs $u_\star(\overline{p}^0), f(\overline{q})$
b_0	p^1, \overline{q} : myopic play; continuation payoffs $u_\star(\underline{p}^0), v_\star(\overline{q})$				
a_1	b_0	p^2, \underline{q} : information exchange; continuation payoffs $u_\star(\overline{p}^1), f(\underline{q})$			
p, \overline{q}	b_1	p^2, \underline{q} : myopic play; continuation payoffs $u_\star(\underline{p}^1), v_\star(\underline{q})$			

Figure 2: The beliefs in the first three stages.

Since all beliefs lie in $(\varepsilon_{\delta}, 1 - \varepsilon_{\delta})$, this strategy profile is well-defined. As an example, consider the top final node in Figure 2. At that node, players' beliefs are \bar{p}^1 and \bar{q} , and players switch to the equilibrium profile which we designed in the previous section, taking \bar{p}^1 and \bar{q} as initial beliefs.

We claim that equilibrium conditions for player 2 are satisfied. Indeed, note that the function $f(\cdot)$ solves

$$
v_{\star}(\tilde{q}) = (1 - \delta)(1 - v_{\star}(\tilde{q})) + \delta f(\tilde{q}), \text{ for each } \tilde{q} \in [0, 1].
$$

Hence, whatever be the action played by player 1 in stage 1, player 2 is indifferent between his two actions at stage 2, as desired.

The overall payoff to player 2 is equal to $(1 - \delta)v_{\star}(q) + \delta (xv_{\star}(\bar{q}) + (1 - x)v_{\star}(q))$. As δ goes to 1, the weight of the first stage decreases to 0. Since q and \bar{q} were arbitrary, this payoff spans the whole interval $(q, 1) = (v_\star, v_{\star\star}).$

It remains to ensure that player 1 is indifferent between both actions in stage 1. That is, the difference in payoffs in stage 1 should be offset by a difference in the expected continuation payoffs. This is done by adjusting the amount of information disclosed by player 2 in stage 2 to the action played by player 1 in stage 1: more information is disclosed if player 1 plays his suboptimal action in stage 1.

Formally, the indifference condition translates to:

$$
(1-\delta)(u_{\star}(p)-(1-u_{\star}(p))) = \delta^{2}(y^{0}u_{\star}(\bar{p}^{0})+(1-y^{0})u_{\star}(\underline{p}^{0})) - \delta^{2}(y^{1}u_{\star}(\bar{p}^{1})+(1-y^{1})u_{\star}(\underline{p}^{1})).
$$
\n(3)

A necessary condition is that p^0, \bar{p}^0 be chosen such that the overall payoff when playing the suboptimal action in stage 1 is at least $u_\star(p)$. Conversely, one can check that for any such choice of p^0, \bar{p}^0 , there exist $p^1, \bar{p}^1 \in (p^0, \bar{p}^0)$ such that equation [\(3\)](#page-16-0) holds.

The overall payoff to player 1 is then equal to

$$
(1 - \delta)(1 - u_{\star}(p)) + \delta(1 - \delta)u_{\star}(p) + \delta^{2} (y^{0} u_{\star}(\bar{p}^{0}) + (1 - y^{0})u_{\star}(\underline{p}^{0}))
$$

A δ goes to 1, the weight of the first two stages decreases to zero. Since p^0, \bar{p}^0 are arbitrary (subject to the individual rationality condition), this payoff spans the whole interval $(p, 1) = (u_\star, v_{\star\star}).$

2.2 The self-informed case

While sticking to the binary setup, we now allow for private signals on one's own state, and we informally describe the main ideas of the construction. All details appear in the Appendix. All equilibria share a common structure. Equilibrium play is divided into four successive phases.

In phase 1, players 1 and 2 report their signals \mathbf{l}_S and \mathbf{m}_T , by means of encoding them into finite strings of actions in a one-to-one way.

The crucial phase is Phase 2. It is designed so as to provide incentives for truthfully reporting in phase 1. In expectation, very little information is exchanged in that phase. It is organized as follows. Each player i draws a random message, 9 and 'sends' it to player j (again, by encoding messages into sequences of actions). This message is slightly correlated with player j 's state, but is independent of the report of player j in phase 1^{10} 1^{10} 1^{10} Next, player j plays a long and deterministic sequence of actions. We refer to these two subphases as phases 2.1 and 2.2 respectively. The length of phase 2 is of the order of $\ln(1-\theta)/\ln \delta$ for some positive θ , hence phase 2 contributes a fraction of θ to the total discounted payoff.^{[11](#page-16-3)} The prescribed sequence of actions

⁹We distinguish between *signals*, drawn by nature, and *messages*, chosen by the players and possibly subject to strategic considerations.

¹⁰Even if drawn independently of player j's report, it is crucial that it is sent only after player j's report.

¹¹The value of θ is independent of δ , but may have to be adjusted to the equilibrium payoff.

depends both on player j's report in phase 1, and on player i's message in phase 2. The intuition behind this structure is discussed later.

In the continuation game that starts after phase 2, players hold no private information about their own state and they implement an equilibrium of the resulting self-ignorant game. The bulk of valuable information exchange takes place in phase 3. Each player i draws a random message that is correlated with player j 's state, and 'sends' it to player j. The degree of correlation is adjusted as a function of player j's equilibrium payoff. The (conditional) law of player i's message does not depend on player j 's report.

Phase 4 consists of the remaining stages. Prior to the first stage, N , of phase 4, each player i assesses the belief held by player j , assuming reports in phase 1 were truthful. He also assesses the total discounted cost incurred by player j in all previous stages – with the exception of the stages of phase 2.2. Players switch to the equilibrium profile, as designed in the previous section, which is associated with a payoff equal to the myopic payoff, $(u_*(p_N), v_*(q_N))$, plus a bonus that exactly offsets the cost incurred earlier. Because few stages are taken into account for the computation of the past cost, this bonus is at most of the order of $(1 - \delta)c$, for some positive c. Because it is small, hardly any valuable information is exchanged during phase 4. The role of the bonus is to ensure that each player, when randomizing in either phase 2 or phase 3, is indifferent between all messages.

Observable deviations of player i trigger a myopic play of player i .

We now provide a few insights into phase 2, by means of two examples. Phase 2 is designed so that the expected payoff of a player in phase 2.2 is strictly higher when reporting truthfully than when not.

Assume first that $L_S = \{\underline{l}_S, \overline{l}_S\}$, and that player 1 assigns to state $s = 1$ a posterior probability $\frac{1}{3}$ when observing l_S , and $\frac{2}{3}$ when observing \overline{l}_S . Note that the optimal action of player 1 is $a = 0$ in the former case, and $a = 1$ in the latter case. Here, incentives can be provided by simply requiring that player 1 repeats the action which is optimal given the signal he reported. Indeed, assume for concreteness that player 1 receives $I_S = I_S$. If player 1 reports truthfully I_S and plays as required the action $a = 0$, his expected payoff is $\frac{2}{3}$ in all stages of phase 2.2. If instead player 1 untruthfully claims that he received \bar{l}_S and next plays the action $a = 1$, which is optimal given his report but in fact suboptimal, his expected payoff is only $\frac{1}{3}$. Such

an incentive scheme is appropriate whenever the myopically optimal action of player 1 is in one-to-one relation with l_S .

In some cases, more complex schemes are needed. Assume that $L_S = \{ \underline{l}_S, l_S^*, \overline{l}_S \},\$ and that the posterior probability assigned to state $s = 1$ is respectively $\frac{1}{3}$ following $\underline{l}_S, \, \frac{1}{2}$ $\frac{1}{2}$ following l_S^* and $\frac{2}{3}$ following \bar{l}_S . Here, conditional on **s**, we let player 2 send a message μ in $\{*,0,1\}$. The probability of $\mu = *$ is close to 1, and does not depend on s, so that this message conveys no information. Conditional on $\mu \neq *,$ the message μ coincides with **s** with probability $1 - \varepsilon$, where $\varepsilon > 0$ is small. Next, player 1 is to repeat a sequence \vec{a} of actions of length 8, which depends on both report and disclosed state:

- If $\mu = *$, the sequence \vec{a} is independent of player 1's report;
- Following a report of l_{S}^{\star} , the sequence \vec{a} contains four 0's (and four 1's), irrespective of the message sent by player 2;
- Following a report of l_S , \vec{a} contains six 0's (and two 1's) if the player 2's message is $\mu = 0$, and seven 0's if $\mu = 1$;
- Following a report of \overline{l}_S , \overline{a} contains six 1's if $\mu = 1$, and seven 1's $\mu = 0$.

It is straightforward that strict incentive requirements are met for $\varepsilon > 0$ small enough.

We conclude by explaining the equilibrium logic. Assuming that player i chooses to truthfully report in phase 1, the computation of the continuation profile in phase 4 ensures that player i is indifferent between all messages in phases 2.1 and 3, and has no profitable deviation in phase 4. However, the prescribed sequence of actions in phase 2.2 may involve suboptimal actions,^{[12](#page-18-0)} and player i may consider deviating from this sequence. Such a deviation fails to be profitable as soon as the marginal value of the information received in phase 3 compensates the cost incurred along phase 2.2. This condition puts a (mild) constraint on the relation between the duration of phase 2 and the amount of information disclosed in phase 3.

What if player 1, say, chooses instead to mis-report his signal in phase 1, and claims that he received the signal l'_S when in fact receiving l_S ? If l_S and l'_S are

 12 The cost of which is *not* taken into account in the bonus.

equivalent, in the sense that the belief of player 1 is the same following either signal: $p(\cdot | l_S) = p(\cdot | l_S') \in \Delta(S)$, such a deviation is irrelevant.^{[13](#page-19-1)} Hence, assume that player 1 reports a signal $l'_{\mathcal{S}}$ that is not equivalent to $l_{\mathcal{S}}$, and plays consistently with his report thereafter.^{[14](#page-19-2)} Three properties combine to ensure that such a deviation fails to be profitable. First, observe that the joint distribution of the two messages received from player 2 (in phases 2 and 3) does not depend on player 1's report. Hence, misreporting does not affect the information received from player 2 in phases 2 and 3. Next, observe that the continuation strategy of player 2 in phase 4 entails almost no information disclosure. As we show in the Appendix, this implies that the best-reply continuation payoff of player 1 in phase 4, is at most $u_{\star}(\mathbf{p}_{N}) + (1 - \delta)C$, for some positive C, where \mathbf{p}_N is the 'true' belief of player 1. That is, in spite of the fact that player 2's continuation strategy is based on a *wrong* assessment of player 1's belief,^{[15](#page-19-3)} the maximal gain for player 1 is very small. Finally, by design, the expected payoff received by player 1 in phase 2 is *strictly* higher when reporting l_S than l_S' .

Because the weight of phase 2 in the total payoff is θ , the small gain obtained in phase 4 can not offset the loss incurred in phase 2.2 when reporting l'_{S} rather than l_{S} .

3 Extensions and concluding comments

3.1 Ex ante vs. interim valuable information

We here discuss how to adapt the statement of the Main Theorem to a situation in which information is ex ante valuable, but need not be *interim* valuable. That is, we assume that both inequalities $u_* < u_{**}$ and $v_* < v_{**}$ hold, but that the information held by player *i* need not be valuable in the sense of Definition [1.](#page-9-1)

We will assume that with p -probability 1, player 1 has a unique myopically optimal action at \mathbf{p}_1 , and that the symmetric property.^{[16](#page-19-4)} holds for player 2 Define $\bar{L}_S \subset L_S$

¹³We note that the definition of equivalent signals adopted here is quite specific to the binary case, and will be different in the general case.

¹⁴The case where player 1 fails to play consistently, and deviates from the prescribed sequence will pose no specific problem. It is left to the appendix.

¹⁵Because it is assuming that player 1 reported truthfully.

¹⁶If a player has two myopically optimal actions at \mathbf{p}_1 , he can costlessly reveal information to the other player.

to be the set of signals l_S for which the inequality [\(2\)](#page-9-0) does not hold, and define $\bar{M}_T \subset M_T$ in a symmetric way. We argue that the limit set of sequential equilibrium payoffs is equal to the set

$$
E := [u_\star, u_\star q(\bar{M}_T) + u_{\star\star} q(M_T \setminus \bar{M}_T)] \times [v_\star, v_\star p(\bar{L}_S) + v_{\star\star} p(L_S \setminus \bar{L}_S)].
$$

Observe first that at any equilibrium, if $m_T \in \bar{M}_T$, player 2 realizes (interim) that the information held by player 1 has no value, and player 2 will then repeat his unique, optimal action at \mathbf{q}_1 . Thus, conditional on \mathbf{m}_T , and using the independence assumption **A.2**, player 1's expected payoff is at most u_* if $m_T \in \overline{M}_T$, and at most u_{**} if $m_T \notin \bar{M}_T$. Using the same argument for player 2, this shows that any equilibrium payoff vector lies in the set E.

Conversely, consider the following class of strategy profiles. In the first two stages, each player i 'tells' player j whether the information held by j has positive value to i or not. This is done as follows. In stage 1, player i plays his myopically optimal action. In stage 2, player 1, say, repeats this action if $I_s \in \overline{L}_s$, and switches to a different (suboptimal) action if $I_s \notin \bar{L}_s$ to signal his willingness to disclose/acquire information. If both players switched in stage 2, they implement from stage 3 on an equilibrium such as we designed in the proof of the Main Theorem. Otherwise, players repeat their stage 1 action. The sole role of stage 1 is to instruct the other player how to interpret the action played in stage 2.

If $I_s \in \bar{L}_s$, it is strictly dominant for player 1 to repeat his optimal action throughout, as required. Indeed, playing a different action in stage 2 would only lower player 1's payoff, with no benefit since player 2's information is valueless.

If $\mathbf{l}_s \in L_S \setminus \overline{L}_S$, player 1's overall payoff is $u_\star(\mathbf{p}_1)$ if he pretends that the information held by player 2 is valueless. However, because there is a positive q-probability that $m_T \notin \bar{M}_T$, it is a best reply for player 1 to switch to a suboptimal action in stage 2 as soon as the value of the information disclosed by player 2 exceeds on average the cost incurred in stage 2.[17](#page-20-0)

In such an equilibrium, conditional on \mathbf{l}_S , player 1's payoff is $u_\star(\mathbf{p}_1)$, which is then also equal to $\mathbf{E}[u_{\star}(\tilde{\mathbf{p}})|\mathbf{l}_S]$, if $\mathbf{l}_S \in \bar{L}_S$. If instead $\mathbf{l}_S \notin \bar{L}_S$, then with probability $q(M_T \setminus \bar{M}_T)$, player 1's payoff may be as high as $\mathbf{E}[u_{\star}(\tilde{\mathbf{p}})|\mathbf{l}_S]$. Otherwise, player 1's payoff will be (approximately) $u_{\star}(\mathbf{p}_1)$. The *ex ante* expected payoff can therefore be

¹⁷It cannot be optimal for player 1 to pretend that $I_s \notin \bar{L}_s$, yet to lie about his optimal action.

as high as

$$
u_\star q(\bar{M}_T) + q(M_T \setminus \bar{M}_T)u_{\star\star}.
$$

3.2 The correlated case

The case where the independence assumption $\mathbf{A.2}$ does not hold raises significant challenges. For the sake of simplicity, we focus here on the binary setup.

A first difficulty is that myopic play need not be an equilibrium. Hence it is not clear what is the lowest equilibrium payoff.

Binary Example 3 Here, states and signals are functions of an auxiliary variable $ω$, which can assume four different possible values $ω_i$, $i ∈ {1, 2, 3, 4}$. The probabilities of the different values, and the way ω determines the states and signals is as follows:

When behaving in a myopically optimal way, players play as follows. In stage 1, player 2 plays $b = 0$. Indeed, the probability he assigns to state $t = 0$ is either 1 or $\frac{2}{3}$. Meanwhile, player 1 plays $a = 1$ if $l = l^1$, and $a = 0$ if $l = l^2$. Indeed, the probability assigned to state $s = 0$ is $\frac{1}{3}$ in the former case, and $\frac{2}{3}$ in the latter case. In stage 2, player 1 repeats his stage 1 action. On the other hand, player 2 can deduce ω from player 1's stage 1 choice. If player 1 played $a = 1$, player 2 repeats his stage 1 action. If instead player 1 played $a = 0$, player 2 will play either $b = 0$ or $b = 1$, depending on player 2's information. In the former case, players repeat forever their stage 2 action. In the latter, player 1 deduces ω from player 2 stage 2's action, and obtains a payoff of 1 in all later stages. This obviously creates an incentive for player 1 to deviate in stage 1, and to always play $a = 0$, in order to learn the value of ω in stage 3. Thus, it is not an equilibrium to behave myopically. In this example, information exchange is a consequence of equilibrium behavior.

A second difficulty lies in understanding the extent to which information can be exchanged. In Section [2.1.1](#page-11-2) we described specific periodic equilibria. In fact, there are many degrees of freedom in this construction, and the choice of using periodic equilibria was made to facilitate the computation of the equilibrium requirements. It turns out that when information is correlated there are no degrees of freedom in determining the beliefs, and the sequence of beliefs is not periodic: it satisfies uninspiring recursive equations, that do not seem to have closed-form solutions. Preliminary numerical evidence seems to suggest that some equilibria can be constructed along these lines.

3.3 The finite horizon case

We here comment on the assertion that the exchange of information relies on the horizon being unbounded. We argue that, when the horizon is finite, the myopic equilibrium is the unique Nash equilibrium of the binary example, as soon as $\mathbf{p}_1 \neq \frac{1}{2}$ 2 *p*-a.s., and $\mathbf{q}_1 \neq \frac{1}{2}$ $\frac{1}{2}$ q-a.s. This claim is an illustration that in many cases of interest, a bounded horizon prevents players from exchanging information.

Let T be the length of the game, and let a Nash equilibrium be given. We prove that, for any sequence h_n of length $n \geq 0$ of moves that occur with positive probability, the continuation payoff of player 1 is $u_\star(p_n)$ if $q_n \neq \frac{1}{2}$ $\frac{1}{2}$, and the continuation payoff of player 2 is $v_{\star}(q_n)$ if $p_n \neq \frac{1}{2}$ $\frac{1}{2}$. We argue by backward induction over *n*. The claim holds trivially for $n = T$. Assume that the claim holds for every history of length $n + 1$, and consider a history of length n such that $q_n \neq \frac{1}{2}$ $\frac{1}{2}$. We proceed in two steps.

Step 1: We claim that $\mathbf{E}[u_{\star}(\mathbf{p}_{n+1})|h_n] = u_{\star}(p_n)$.

Note first that the claim trivially holds if player 2 does not randomize following h_n , since one then has $\mathbf{p}_{n+1} = p_n$ with probability 1.

Assume now that player 2 randomizes at stage n following h_n . Observe that \mathbf{p}_{n+1} must then be equal to $\frac{1}{2}$ with positive probability. Otherwise, by the induction hypothesis the continuation payoff of player 2 would be $v_{\star}(q_{n+1})$, irrespective of the action played by player 2 in stage n . But then player 2 would not be indifferent in stage n between the two actions – a contradiction.

It follows that both possible values of p_{n+1} lie in the same half of the interval [0, 1]. The claim follows, because $\mathbf{E}[\mathbf{p}_{n+1}|h_n] = p_n$ and because u_* is linear in both $[0, \frac{1}{2}]$ $\frac{1}{2}$ and $\left[\frac{1}{2}, 1\right]$.

Step 2: Conclusion

Since $q_n \neq \frac{1}{2}$ $\frac{1}{2}$, the event $\mathbf{q}_{n+1} \neq \frac{1}{2}$ $\frac{1}{2}$ has a positive probability conditional on h_n . Let $a \in A$ be any action that is played with positive probability at h_n , and following which $\mathbf{q}_{n+1}\neq\frac{1}{2}$ $\frac{1}{2}$. When playing a and by the induction hypothesis, the continuation payoff of player 1 in stage $n+1$ is equal to $u_\star(\mathbf{p}_{n+1})$. As a consequence, and when playing a, the continuation payoff of player 1 in stage n is equal to $(1-\delta)u(p_n, a) + \mathbf{E}[u_\star(\mathbf{p}_{n+1})|h_n] \leq$ $u_{\star}(p_n)$, using **Step 1**. By the equilibrium property, it follows that the continuation payoff of player 1 in stage n (following h_n) is equal to $u_*(p_n)$.

We note, though, that there may be cases in which information exchange is possible, even when the game is finitely repeated.The following example illustrates this point.

Consider the binary example, and add to each action set one more action, that we denote by 2, and that yield payoff $\frac{2}{3}$ irrespective of the state. The optimal action of player 1, as a function of the belief assigned to state $s = 1$, is given by Figure 3 below, and the structure of player 2's best-response is similar.

Figure 3: The optimal action of player 1

Example 1 Assume that the two states are equally likely, and that player 1 learns t while player 2 learns **s**. Suppose that in stage 1 player 1 plays $\left[\frac{1}{3}\right]$ $\frac{1}{3}(a_0), \frac{2}{3}$ $\frac{2}{3}(a_1)$ if $t = 0$, and $\left[\frac{2}{3}\right]$ $\frac{2}{3}(a_0), \frac{1}{3}$ $\frac{1}{3}(a_1)$ if $t = 1$, and suppose that player 2 plays in an analog way. Player 1's belief in stage 2 is either $\left[\frac{1}{3}\right]$ $\frac{1}{3}(0), \frac{2}{3}$ $\frac{2}{3}(1)$] or $\left[\frac{2}{3}\right]$ $\frac{2}{3}(0), \frac{1}{3}$ $\frac{1}{3}(1)$], depending on player 2's action in stage 1. In the former case, we let player 1 play either a_0 or a_2 depending on **t**. In the latter case, we let player 1 play either a_1 or a_2 , depending on t^{18} t^{18} t^{18} Let the behavior of player 2 in stage 2 be analog. Provided δ is high enough, this strategy pair is an equilibrium, in which players exchange all information in two stages.

The type of construction presented in Example [3.3](#page-22-0) is valid only for intermediate values of initial beliefs, and for self-ignorant games. Moreover, it is not clear how to use such profiles as continuation profiles in general games.

¹⁸We let player 1 repeat a_2 forever if player 2 played b_2 .

3.4 Signals along the Play

How do our results change when players receive independent signals, both on their own state and on the other player's state, along the game? If player i receives signals on his own state along the play, then, because he is patient, he will wait until almost all the information that he can get about is own state is received. However, if in subsequent stages he keeps on receiving information on his own state, then his belief changes, even though the changes are small. These changes have the effect that the other player does not know how to compensate player i for playing suboptimally, and our construction fails.

If, on the other hand, player i received signals on the other player's state along the game, then again the players can wait until player i receives almost all the information that he can, and then the players can implement the equilibrium that we construct. Thus, our results remain valid in this case.

References

- [1] Admati, A.R. and M. Perry (1987) Strategic Delay in Bargaining, The Review of Economic Studies, 54, 345–364.
- [2] Aumann, R.J. and S. Hart (1992) Handbook of Game Theory with Economic Applications, vol. 1, Handbook of Game Theory with Economic Applications, Elsevier Science Publishers, North-Holland.
- [3] Aumann, R.J. and S. Hart (2003) Long Cheap Talk, Econometrica, 71, 1619– 1660.
- [4] Aumann, R.J. and M.B. Maschler (1966) Game Theoretic Aspects of Gradual Disarmament, Chapter V in Report to the US Arms Central and Disarmament Agency, Washington, DC.
- [5] Aumann, R.J. and M.B. Maschler (1995) Repeated Games with Incomplete Information, The MIT Press.
- [6] Blackwell, D. (1953), "Equivalent Comparison of Experiments," Ann. Math. Stat., 24, 265-272.
- [7] Che, Y.-K. and J. Sakovics (2004) A Dynamic Theory of Holdup, Econometrica, 72, 1063–1103.
- [8] Compte, O. and P. Jehiel (2004) Gradualism in Bargaining and Contribution Games, The Review of Economic Studies, 71, 975–1000.
- [9] Crawford, V.P. and J. Sobel (1982) Strategic Information Transmission, Econometrica, 50, 1431–1451.
- [10] Cripps, M.W. and J.P. Thomas (2003) Some Asymptotic Results in Discounted Repeated Games of One-Sided Incomplete Information, Mathematics of Operations Research, 28, 433–462.
- [11] Forges, F. (1990) Equilibria with Communication in a Job Market Example, The Quarterly Journal of Economics, 105, 375–398.
- [12] Forges, F. (1992) Repeated Games of Incomplete Information : Non-Zero-Sum, in Handbook of Game Theory with Economic Applications, vol. 1, R. Aumann et S. Hart , Handbook of Game Theory with Economic Applications, Elsevier Science Publishers, North-Holland, 155–177.
- [13] Mailath, G. and L. Samuelson (2006) Repeated Games and Reputations: Long-Run Relationships, Oxford University Press.
- [14] Marx, L.M. and S.A. Matthews (2000) Dynamic Voluntary Contribution to a Public Project, *The Review of Economic Studies*, **67**, 327–358.
- [15] Mayberry, J.P. (1967) Discounted Repeated Games with Incomplete Information, in Report for the U.S. Arms Control and Disarmament Agency, ST-116, Chapter V, 435–461.
- [16] Peski, M. (2008) Repeated Games with Incomplete Information on One Side, Theoretical Economics, 3, 29–84.
- [17] Wiseman, T. (2005) A Partial Folk Theorem for Games with Unknown Payoff Distributions, *Econometrica*, **73**, 629–645.
- [18] Zamir, S. (1992) Repeated Games with Incomplete Information: Zero-Sum, in Handbook of Game Theory, Vol. 1, (Chapter 5), Edited by R.J. Aumann and S. Hart, Elsevier Science Publishers B.V., 109–154.

A Self-Ignorant Games

We assume throughout the Appendix that all payoffs are in $[0, 1]$. In this section we assume that no player receives private information on his own state. Equivalently, both sets L_S and M_T are singletons. For simplicity, we here write L and M instead of L_T and M_S .

Let initial distributions $p \in \Delta(S \times M)$ and $q \in \Delta(T \times L)$ be given. We denote the corresponding game by $\Gamma(p,q)$. W.l.o.g., we assume that $p(m) > 0$ for each $m \in M$.^{[19](#page-26-0)}

We assume that the information of each player i is valuable to player j. Since $\Gamma(p,q)$ is a self-ignorant game, this is equivalent to assuming that there is no action $a \in A$ that is optimal at all distributions $p_m := p(\cdot|m)$, $m \in M$. Equivalently, one has

$$
u_{\star} = u_{\star}(p) < u_{\star \star} = \sum_{m \in M} p(m) u_{\star}(p_m). \tag{4}
$$

By Bayes' rule, the belief p_n of player 1 at stage n is the weighted average of ${p_m \otimes 1_m, m \in M}$, where the weight of $p_m \otimes 1_m$ is equal to the probability that the signal of player 2 is m, given player 1's information at stage n^{20} n^{20} n^{20} . Thus, the set of possible values of \mathbf{p}_n is

$$
\Delta^{\dagger}(S \times M) = \text{conv}\{p_m \otimes 1_m, m \in M\}.
$$
\n(5)

Because $p(m) > 0$, p lies in the (relative) interior of $\Delta^{\dagger}(S \times M)$, which we denote by ◦ ∆ [†] $(S \times M)$. We define $q_l = q(\cdot | l)$ for $l \in L$, and the set $\Delta^{\dagger}(T \times L)$ is defined in a symmetric way.

It is convenient to allow the initial distribution to vary, to account for the fact that beliefs may change along the play. Since all beliefs lie in $\Delta^{\dagger}(S \times M)$ and $\Delta^{\dagger}(T \times L)$, we will only consider initial distributions in these sets. We still denote arbitrary such distributions by p and q .

In this section, we prove the two propositions below.

¹⁹And that $q(l) > 0$ for each $l \in L$: to avoid useless repetitions, we sometimes state properties for player 1, with the implicit understanding that analog properties hold for player 2 as well.

 20 This is a way of stating that, as the play proceeds, the belief of player 1 on $\mathbf m$ evolves, but the distribution of s conditional on m remains fixed.

Proposition 1 Let $p \in \overset{\circ}{\Delta}$ † $(S \times M)$ and $q \in \overset{\circ}{\Delta}$ † $(T \times L)$ be given. There exists $\varepsilon > 0$ and δ < 1 such that the following holds. For every $\delta \geq \overline{\delta}$, every payoff vector in $[u_*(p), u_*(p) + \varepsilon] \times [v_*(q), v_*(q) + \varepsilon]$ is a sequential equilibrium payoff of $\Gamma(p, q)$.

Given such p and q, a payoff vector $\gamma \in [u_*(p), u_*(p) + \varepsilon] \times [v_*(q), v_*(q) + \varepsilon]$, and a discount factor δ , we will construct an equilibrium profile $(\sigma_{p,q,\gamma}, \tau_{p,q,\gamma})$ in $\Gamma(p,q)$, with payoff γ . Proposition [2](#page-27-0) bounds the possible gain of player 1 if player 2 has an incorrect belief on p, provided γ is close to the myopically optimal payoff.

Proposition 2 Let $p \in \overset{\circ}{\Delta}$ † $(S \times M), q \in \overset{\circ}{\Delta}$ † $(T \times L)$ and $c > 0$ be given. There exists a constant $C > 0$ with the following property. For every discount factor δ , every payoff vector γ such that $\|\gamma - (u_\star(p), v_\star(q))\|_\infty \le (1-\delta)c$, and every $p' \in \Delta^{\dagger}(S \times M)$, one has

$$
\gamma^1(p', q, \sigma, \tau_{p,q,\gamma}) \le u_\star(p') + (1 - \delta)C, \text{ for every strategy } \sigma.
$$

In Propositions [1](#page-26-2) and [2,](#page-27-0) ε , C and $\overline{\delta}$ may depend a priori on the choice of (p, q) . We will prove that they can be chosen in such a way that the conclusions hold uniformly throughout some neighborhoods of p and q .

A.1 Notations and Preliminaries

We here describe the main steps leading to the proof of Propositions [1](#page-26-2) and [2.](#page-27-0) Our goal is to mimic the recursive construction of the binary case. We let $[p^0, p^1]$ be any segment in the interior of $\Delta^{\dagger}(S \times M)$ such that u_{\star} is not affine on the segment $[p^0, p^1]$. The beliefs p^0 and p^1 take the role of p and $1 - p$ in the binary case.

An optimal action a at p^0 is not optimal at p^1 (and vice-versa). Otherwise, a would be optimal throughout the segment $[p^0, p^1]$, and then u_* would coincide with the affine map $u(\cdot, a)$ on that segment.

For $k = 0, 1$, we let $a^k \in A$ be an optimal action at p^k . We denote by D^1 the straight line spanned by p^0 and p^1 in $\mathbb{R}^{S \times M}$, and we denote by p and \bar{p} the endpoints of the segment $D^1 \cap \Delta^{\dagger}(S \times M)$, with the convention of Figure 4.

Figure 4

Let $\pi \in [p_0, \bar{p}]$, and assume that player 1 receives information that changes his belief from p^0 to either p^1 (with probability y) or π (with probability $1 - y$). So that the martingale property of beliefs holds we must have $p^0 = yp^1 + (1 - y)\pi$. Assume moreover that from the next stage on player 1 receives his myopically optimal payoff. The gain of player 1 from the information that is revealed to him relative to his myopically optimal payoff at p^0 , is then $h_{p^0}(\pi) = (yu_*(p^1) + (1-y)u_*(\pi)) - u_*(p^0)$. Since u_* is convex, $h_{p0}(\pi) \geq 0$ for each π . Since u_* is not affine on the interval $[p^0, p^1]$, one also has $h_{p0}(\pi) > 0$ for $\pi \in (p^0, \bar{p}]$, see Figure 5. In addition, h_{p0} is piecewise affine, and non-decreasing as π moves away from p^0 towards \bar{p} .

Similarly, define $h_{p^1} : [p^1, p] \to \mathbf{R}^+$ by $h_{p^1}(\pi) = (yu_*(p^0) + (1 - y)u_*(\pi)) - u_*(p^1)$, where y solves $p^1 = yp^0 + (1 - y)\pi$.

We proceed in a symmetric way with player 2. We let $[q^0, q^1]$ be an arbitrary segment in the interior of $\Delta^{\dagger}(T \times L)$ such that the restriction of v_{\star} to the segment $[q^0, q^1]$ is not an affine map. We denote by D^2 the straight line in $\mathbb{R}^{T\times L}$ spanned by q^0 and q^1 , and by q, \bar{q} the endpoints of the segment $D^2 \cap \Delta^{\dagger}(T \times L)$. Finally, we define $h_{q^0}: [q^0, \bar{q}] \to \mathbf{R}^+$ and $h_{q^1}: [q^1, q] \to \mathbf{R}^+$ by adapting the definitions of h_{p^0} and h_{p^1} .

Given a belief $\pi \in \Delta(S \times M)$, and an action $a \in A$, the cost of a at π is defined as the loss incurred when playing a instead of the optimal action at π :

$$
c(\pi, a) := u_{\star}(\pi) - u(\pi, a).
$$

The cost $c(\pi, b)$, for $\pi \in \Delta(T \times L)$ and $b \in B$ is defined analogously.

The proof of Propositions [1](#page-26-2) and [2](#page-27-0) relies on Lemmas [1](#page-29-0) and [2](#page-29-1) below.

Lemma 1 Let δ < 1 be such that $\frac{1-\delta}{s}$ δ $c(p^i,a^j)$ < max h_{p^i} and $\frac{1-\delta}{s}$ δ $c(q^i, b^j) <$ $\max h_{q^i}$, for $i, j = 0, 1$. Then the vector $(u_\star(p^0), v_\star(q^0) + \frac{1-\delta}{\delta}c(q^0, b^1))$ is a sequential equilibrium payoff of $\Gamma(p^0, q^0)$.

Lemma 2 Let $\varepsilon > 0$ be such that $\varepsilon < \max h_{p^i}$, and $\varepsilon < \max h_{q^j}$ for $i, j = 0, 1$. There is $\bar{\delta}$ < 1, such that for every discount factor $\delta \geq \bar{\delta}$, every payoff in $[u_*(p^0), v_*(q^0) +$ $\varepsilon] \times [v_{\star}(q^{0}), v_{\star}(q^{0}) + \varepsilon]$ is a sequential equilibrium payoff of $\Gamma(p^{0}, q^{0})$.

We will prove that the conclusion holds uniformly for all initial distributions \tilde{p}^i , \tilde{q}^j close to p^i and q^j . To be precise, there exist a neighborhood $V(p^i)$ of p^i , and a neighborhood $V(q^j)$ of q^j $(i, j \in \{0, 1\})$ such that, for every $\delta \geq \overline{\delta}$, and every $\tilde{p}^i \in V(p^i), \tilde{q}^j \in V(q^j)$, all vectors in $[u_\star(\tilde{p}^i), u_\star(\tilde{p}^i) + \varepsilon] \times [v_\star(\tilde{q}^j), v_\star(\tilde{q}^j) + \varepsilon]$ are sequential equilibrium payoffs of $\Gamma(\tilde{p}^i, \tilde{q}^j)$.

A.2 Proof of Lemma [1](#page-29-0)

In the construction of Section [2,](#page-11-0) the probabilities x and y assigned to suboptimal actions were pinned down by equilibrium requirements. The construction here is slightly more involved because the number of signals may be larger than 2.

We let δ be as stated. Define first $\bar{p}^1 \in [p^0, \bar{p})$ by the condition $h_{p^0}(\bar{p}^1) =$ $1-\delta$ δ $c(p^0, a^1)$, and $y^1 \in [0, 1)$ by the equality $p^0 = y^1 p^1 + (1 - y^1) \bar{p}^1$. The revelation of information just defined offsets the cost to player 1 of playing the suboptimal action a^1 when the belief if p^0 . \bar{p}^1 takes the role of p^* in the binary case. For $m \in M$, we set $y_m^1 =$ $p^1(m)$ $p^0(m)$ y¹. Because p^0 is in the (relative) interior of $\Delta^{\dagger}(S \times M)$, one has $p^{0}(m) > 0$ for each m, and $y_{m}^{1} \in (0, 1)$. Observe that $y^{1} = \sum_{m \in M} p^{0}(m)y_{m}^{1}$, and that the following Bayesian updating property holds. If player 1's belief is p^0 , and if player 2 plays two different actions b and b' with respective probabilities y_{m}^{1} and $1 - y_{m}^{1}$, then following b the posterior belief of player 1 is equal to p^1 , and it is equal to \bar{p}^1 following b' .

Similarly, we let $\bar{p}^0 \in (p^1, \underline{p})$ be defined by $h_{p^1}(\bar{p}^0) = \frac{1-\delta}{\delta}c(p^1, a^0)$, and we set $y_m^0 =$ $p^0(m)$ $p^1(m)$ y^{0} for $m \in M$, where y^{0} solves $p^{1} = y^{0}p^{0} + (1 - y^{0})\bar{p}^{0}$.

We next exchange the roles of the two players, and proceed in a slightly asymmetric way. We let $\bar{q}^1 \in (q^0, \bar{q})$ be defined by $h_{q^0}(\bar{q}^1) = \frac{1-\delta}{s}$ δ $c(q^0, b^1)$, we let x^0 be defined by $q^0 = x^0 q^1 + (1 - x^0) \bar{q}^1$, and we set $x_l^0 =$ $p^1(l)$ $p^0(l)$ x^0 for $l \in L$.

We finally define $\bar{q}^0 \in (q^1, \underline{q})$, $x^1 \in (0, 1)$, and x^1 = $q^0(l)$ $q^1(l)$ x^0 for $l \in L$ in a similar way.

We are now in a position to define strategies σ_{\star} and τ_{\star} . As long as players alternate in playing their suboptimal action, player 1 (resp., player 2) randomizes in each odd (resp., in each even) stage, and beliefs evolve cyclically:

$$
p^{0}, q^{0} \rightarrow p^{0}, q^{1} \rightarrow p^{1}, q^{1} \rightarrow p^{1}, q^{0} \rightarrow p^{0}, q^{0} \rightarrow \cdots
$$

Along this cycle, player 1 assigns a probability x_1^1 to his suboptimal action, a^1 , when player 2's belief is q^0 , and a probability x_1^0 to his suboptimal action, a^0 , when player 2's belief is $q¹$. Analog properties hold for player 2. This is summarized in Figure 6 below.

	Stage	player 1	player 2		belief Suboptimal action
		1 mod 4 $\ [x_1^1(a^1), (1-x_1^1)(a^0)] \ $		$ p^0,q^0 $	a^{\perp}
	2 mod 4		$\left[(y_{\textbf{m}}^{1}(b^{0}),(1-y_{\textbf{m}}^{1})(b^{1})]\ \right] p^{0},q^{1} \ \left \ \right.$		h^0
		3 mod 4 $\ [x_1^0(a^0), (1-x_1^0)(a^1)] \ $		p^1, q^1	a^0
$\overline{0}$	\mod 4		$[y_{\mathbf{m}}^0(b^1), (1-y_{\mathbf{m}}^0)(b^0)] p^1, q^0 $		

Figure 6: the first phase of play: information exchange.

As soon as either player 1 plays his optimal action in some odd stage, or player 2 plays his optimal action in some even stage, the players switch to myopic play forever, as described in columns 3 and 4 in Figure 7. Here and later, $o(p)$ (resp. $o(q)$) stands for an optimal action of player 1 at p (resp., of player 2 at q).

First stage in which			
myopically optimal action is played \parallel new belief \parallel player 1 \parallel player 2			
mod 4	p^0, \bar{q}^1	a^0	
mod 4	\bar{p}^1, q^1	$o(\bar{p}^1)$	b^1
mod 4	p^1, \bar{q}^0	a^1	$J(\vec{q}^{0})$
mod 4	\bar{p}^0, q^0	$o(\bar{v}^0)$	b^0

Figure 7: the second phase of play: myopic play.

We complete the definition of $(\sigma_{\star}, \tau_{\star})$ by specifying actions and beliefs at information sets that are ruled out by $(\sigma_{\star}, \tau_{\star})$. For concreteness, we focus on player 1. An information set of player 1 contains all histories of the form (l, h) , for a fixed signal $l \in L$, and a fixed sequence $h \in H$ of moves. Fix an information set that is reached with probability 0 under $(\sigma_{\star}, \tau_{\star})$. We denote it by $I_{l,h}^1$, with $h \in H$. Write $h = (h', (\bar{a}, \bar{b}))$, so that h' is the longest prefix of h, and assume that $I_{l,h'}^1$ is reached with positive probability.

We distinguish two cases. Assume first that the action b has probability zero conditional on h' . This is the case where player 2 deviates in an observable way at h' . We let the belief of player 1 at $I_{l,h}^1$ be equal to the belief held at $I_{l,h'}^1$ – the deviation by player 2 is interpreted as being non-informative about **. Assume now that** b **is** played with positive probability at h'. In that case, the belief of player 1 at $I_{l,h}^1$ can be computed by Bayes' rule, from the belief held at $I_{l,h'}^1$.

In both cases, we let the belief at all subsequent information sets be equal to the belief at $I_{l,h}^1$, and we let σ_\star repeat forever any action that is optimal at $I_{l,h}^1$.

Observe that, following any history in $I_{l,h}^1$, under τ_{\star} player 2 repeats forever the same action.^{[21](#page-31-0)} Indeed, either the sequence h of actions has probability 0, or it has positive probability. In the former case, the claim follows from the definition of τ_{\star} at zero probability information sets. In the latter case, this implies that the information

²¹To be precise, player 2 plays the same action at $I_{m,h}^1$ and in all subsequent information sets.

set $I_{l',h}^1$ has positive probability, for some $l' \neq l$. Since the support of player 1's mixed actions in the information phase does not depend on his signal, this implies that $I_{l,h}^1$ must belong to the myopic play phase. Using this observation, one can check that beliefs are consistent with the strategy profile $(\sigma_{\star}, \tau_{\star})$. We omit the proof.

Note that the strategy σ_{\star} is sequentially rational at any $I_{l,h}^1$ that is reached with probability 0. Indeed, since the belief of player 1 is the same at $I_{l,h}^1$ and at all subsequent information sets, it is a best reply to repeat any action that is optimal at $I^1_{l,h}.$

Lemma 3 The profile $(\sigma_{\star}, \tau_{\star})$ is a sequential equilibrium of $\Gamma(p^0, q^0)$, with payoff $(u_{\star}(p^{0}), v_{\star}(q^{0}) + \frac{1-\delta}{\delta}c(q^{0}, b^{1})).$

We will use this lemma for various distributions p^0, q^0 . To avoid confusion, we will then denote the profile $(\sigma_{\star}^{p^0,q^0}, \tau_{\star}^{p^0,q^0}).$

Proof. Each of the strategies σ_{\star} and τ_{\star} can be described by an automaton with 8 states: four states that implement the periodic play in Figure 6, and four states that implement the myopic play in Figure 7.

In addition, transitions between (automaton) states are deterministic and depend only on the public history of moves. Hence, player i can always compute the current state of player j's automaton. Moreover, as can be verified inductively, the belief of player i following any public history h of moves only depends on the current state of player j's automaton.

It follows that player i has a best response that can be implemented by an automaton that has the same (or smaller) number of states as the automaton of player j . The dynamic programming principle may be used to identify such a best response. Using this principle, it is routine to verify that τ_{\star} is a best response against σ_{\star} , and vice versa. Indeed, denoting the 8 states of the automata by $\Omega = \{(1, \text{periodic}), (2, \text{periodic}),\}$ $(3, \text{periodic})$, $(0, \text{periodic})$, $(1, \text{myopic})$, $(2, \text{myopic})$, $(3, \text{myopic})$, $(0, \text{myopic})$, the expected payoff to player 2 starting at any given $\omega \in \Omega$ is:

$$
V(1, \text{periodic}) = v_{\star}(q^{0}) + \frac{1-\delta}{\delta}c(q^{0}, b^{1}); \qquad V(1, \text{myopic}) = v_{\star}(q^{0});
$$

\n
$$
V(2, \text{periodic}) = v_{\star}(q^{1}); \qquad V(2, \text{myopic}) = v_{\star}(\bar{q}^{1});
$$

\n
$$
V(3, \text{periodic}) = v_{\star}(q^{*}) + \frac{1-\delta}{\delta}c(q^{1}, b^{0}); \qquad V(3, \text{myopic}) = v_{\star}(q^{1});
$$

\n
$$
V(0, \text{periodic}) = v_{\star}(q^{0}); \qquad V(0, \text{myopic}) = v_{\star}(\bar{q}^{0}).
$$

One may verify that for every $\omega \in \Omega$, V solves

$$
V(\omega) = \max_{b \in B} \left\{ (1 - \delta)r(\omega, b) + \delta \sum_{\omega' \in \Omega} V(\omega, b)[\omega'] \right\};
$$
\n(6)

here $r(\omega, b)$ stands for the expected payoff of player 2 when playing b in the (automaton) state ω , where the expectation is taken w.r.t. the belief held at state ω . П

A.3 Proof of Lemma [2](#page-29-1)

Let a payoff vector $\gamma \in [u_\star(p^0), u_\star(p^0) + \varepsilon] \times [v_\star(q^0), v_\star(q^0) + \varepsilon]$ be given. For δ high enough, we will define a sequential equilibrium profile in $\Gamma(p^0, q^0)$ with payoff γ , using the ideas in Section [2.1.2.](#page-14-0) We need some preparations.

Define γ_s^1 and γ_o^1 by the equations

$$
\gamma^1 = (1 - \delta)u_{\star}(p^0) + \delta(1 - \delta)u_{\star}(p^0) + \delta^2 \gamma_o^1,
$$

$$
\gamma^1 = (1 - \delta)u(p^0, a^1) + \delta(1 - \delta)u_{\star}(p^0) + \delta^2 \gamma_s^1.
$$

 γ_s^1 (resp. γ_o^1) are the continuation payoffs of player 1 at stage 2, which ensure that the expected payoff of player 1 is γ^1 , if player 1 plays the myopically suboptimal (resp. optimal) action at stage 1, and the myopically optimal action^{[22](#page-33-0)} at stage 2.

Define γ_o^2 be the equality

$$
\gamma^2 = (1 - \delta)v_\star(q^0) + \delta\gamma_o^2.
$$

Because $\gamma^1 > u_*(p^0) > u(p^0, a^1)$ and $\gamma^2 > v_*(q^0)$ it follows that $\gamma_s^1 > \gamma_o^1 \ge u_*(p^0)$ while $\gamma_o^2 \ge v_\star(q^0)$. For δ high enough, and by definition of ε , one has

 $\gamma_s^1 < u_\star(p^0) + \max h_{p_0} \text{ and } \gamma_o^2 < v_\star(q^0) + \max h_{q^0}.$

Hence, there exist $p_s, p_o \in [p^0, \bar{p})$, and $q_o \in [q^0, \bar{q})$ such that

$$
h_{p^{0}}(p_{o}) = \gamma_{o}^{1} - u_{\star}(p^{0}),
$$

\n
$$
h_{p^{0}}(p_{s}) = \gamma_{s}^{1} - u_{\star}(p^{0}),
$$

\n
$$
h_{q^{0}}(q_{o}) = \gamma_{o}^{1} - v_{\star}(q^{0}).
$$

Mimicking the previous section, we define

²²The letters s, o remind that γ_o^1 and γ_s^1 are continuation payoffs following an optimal and a suboptimal action respectively.

•
$$
y_{o,m} = \frac{p^1(m)}{p^0(m)} y_o
$$
, for $m \in M$, where y_o solves $p^0 = y_o p^1 + (1 - y_o) p_o$.

•
$$
y_{s,m} = \frac{p^1(m)}{p^0(m)} y_s \ (m \in M)
$$
, where y_s solves $p^0 = y_s p^1 + (1 - y_s) p_s$.

•
$$
x_l = \frac{q^1(m)}{q^0(m)} x
$$
 for $l \in L$, where x solves $q^0 = xq^1 + (1 - x)q_o$.

We are now in a position to define a profile as follows (see also Figure 8).

- **Stage 1:** Player 2 plays b^0 , while player 1 plays the two actions a^1 and a^0 with probabilities x_1 and $1 - x_1$. By Bayesian updating, following a^1 the belief of player 2 in stage 2 is equal to $q¹$, and it is equal to q_o following $a⁰$ (while the belief of player 1 is still p^0).
- **Stage 2:** Player 2 randomizes. Following a^1 , player 2 plays the two actions b^0 and b^1 with probabilities $y_{s,m}$ and $1 - y_{s,m}$ respectively. Following a^0 , he plays the two actions b^1 and $o(q_o)$ with probabilities $y_{o,\mathbf{m}}$ and $1-y_{o,\mathbf{m}}$ respectively. Meanwhile, player 1 plays a^0 . By Bayesian updating, the belief of player 1 is equal to (i) p^1 following either (a^0, b^1) or (a^1, b^0) , (ii) to p_s following $(a^0, o(q_o))$ and (iii) to p_o following (a^1, b^1) .
- Stage 3 and on: If player 2 played his optimal action in stage 2, players repeat their optimal action. The continuation payoff is then $(u_*(p^1), v_*(q^1))$ following (a^1, b^1) and is $(u_{\star}(p_s^1), v_{\star}(q_o))$ following $(a^0, o(q_o))$. Assume now that player 2 played b^1 in stage 2, following a^0 . Beliefs are then (p_o^1, q_c) and players switch to the equilibrium profile $(\sigma^{p_o^1,q_o}_{\star}, \tau^{p_o^1,q_o}_{\star})$ of $\Gamma(p_o^1, q_o)$, with payoff $(u_{\star}(p_o^1), v_{\star}(q_o) + \frac{1-\delta}{\delta}c(q_o, b^1)).$ Finally, assume that player 2 played b^0 in stage 2, following a^1 . Beliefs are then (p^1, q^1) , and players switch to the profile $(\sigma^{p^1,q^1}_{\star}, \tau^{p^1,q^1}_{\star}).$

$$
\begin{array}{ccccccccc}\n & & & & p^0, q^1 & & b^0 & & \text{belief } p^1, q^1, \text{ payoff } u_\star(p^1), v_\star(q^1) + \frac{1-\delta}{\delta}c(q^1, b^0)) \\
& & & & & b^1 & & \text{belief } p^1, q^1, \text{ payoff } u_\star(p^1), v_\star(q^1) \\
& & & & & b^1 & & \text{belief } p^1, q_0, \text{ payoff } u_\star(p^1), v_\star(q_0) + \frac{1-\delta}{\delta}c(q_0, b^1)) \\
& & & & & & p^0, q_0 & & o(q_c) & & \text{belief } p^1, q_0, \text{ payoff } u_\star(p^1), v_\star(q_0)\n\end{array}
$$

Figure 8: The evolution of beliefs and of continuation payoffs.

Beliefs and actions at information sets that are ruled by this description are defined as in the proof of Lemma [1.](#page-29-0) The equations defining γ_s^1, γ_o^1 (resp. γ_o^2) ensure that player 1 is indifferent in stage 1 (resp. player 2 in stage 2) between the two actions that are assigned positive probability. This implies the equilibrium property. Details are standard and omitted.

A.4 Proofs of Propositions [1](#page-26-2) and [2](#page-27-0)

We start with the proof of Proposition [1.](#page-26-2) The construction we provide here is more complex than needed for Proposition [1.](#page-26-2) However, it will facilitate the proof of Propo-sition [2.](#page-27-0) We let initial distributions p and q be given, in the interiors of $\Delta^{\dagger}(S \times M)$ and $\Delta^{\dagger}(T \times L)$. Choose a segment $[p^0, p^1]$ included in the interior of $\Delta^{\dagger}(S \times M)$, such that (i) u_{\star} is not affine on $[p⁰, p¹]$, and (ii) $p \in (p⁰, p¹)$.

By (i) and (ii), one has $u_\star(p) < y u_\star(p^0) + (1-y) u_\star(p^1)$, where y solves $y p^0 + (1 (y)p^1 = p$. Observe also that the quantity $\tilde{y}u_*(p^0) + (1-\tilde{y})u_*(\tilde{p}^1)$ (with $\tilde{y}p^0 + (1-\tilde{y})\tilde{p}^1 =$ p) is strictly decreasing in the neighborhood of p^1 , as $\tilde{p}^1 \in [p^0, p^1]$ moves away from $p¹$ and towards $p⁰$.

By Lemma [2,](#page-29-1) there exists $\varepsilon_0 > 0$, $\bar{\delta} < 1$, and neighborhoods $V(p^i)$ and $V(q^j)$ of p^i and q^j $(i, j \in \{0, 1\})$, such that any payoff in $[u_*(\tilde{p}^i), u_*(\tilde{p}^i) + \varepsilon_0] \times [v_*(\tilde{q}^j), v_*(\tilde{q}^i) + \varepsilon_0]$ is a sequential equilibrium payoff of the game $\Gamma(\tilde{p}^i, \tilde{q}^j)$, as soon as $\delta \geq \bar{\delta}$.

We now prove that the conclusion of Proposition [1](#page-26-2) holds with $\varepsilon = \varepsilon_0$. Let $\gamma \in$ $[u_*(p), u_*(p) + \varepsilon_0] \times [v_*(q), v_*(q) + \varepsilon_0]$ be given. We describe an equilibrium profile that implements γ .

One main feature of this profile is the following. As a result of information disclosure by player 2, player 1's belief will move in one stage from p to a belief \tilde{p}^i close to either p^0 or p^1 . Similarly, player 2's belief will change to a belief \tilde{q}^j close to either $q⁰$ or $q¹$ in exactly one stage. From that point on, players implement an equilibrium of $\Gamma(\tilde{p}^i, \tilde{q}^j)$ with the appropriate payoff. There is however one minor difference with previously defined equilibria. If $u_*(p) < \gamma^1 < y u_*(p^0) + (1 - y) u_*(p^1)$, then the expected payoff of player 1 if we follow the previous construction will be higher than γ^1 , which is the target payoff. There are two ways to overcome this difficulty. One way is to choose in this case p^0 and p^1 which are closer to p, thereby lowering the expected continuation payoff $yu_*(p^0) + (1 - y)u_*(p^1)$. A second way, which we adopt here, is to delay information revelation, so that the discounted payoff is lower than

 $yu_{\star}(p^{0}) + (1 - y)u_{\star}(p^{1}).$

Define $N_1 \geq 1$ to be the least integer^{[23](#page-36-0)} such that $\gamma_c^1 \geq y u_\star(p^0) + (1-y) u_\star(p^1)$, where γ_c^1 is defined by $\gamma^1 = (1 - \delta^{N_1}) u_\star(p) + \delta^{N_1} \gamma_c^1$. The inequality $\gamma_c^1 \geq y u_\star(p^0) + (1 - y) u_\star(p^1)$ ensures that if player 2 starts revealing information at stage N_1 , then one can support γ_c^1 as a continuation payoff of player 1 at that stage. Define N_2 in a similar way for player 2, and assume w.l.o.g. that $N_1 \leq N_2$. Information is first disclosed at stage N_1 . The choice of N_1 implies

$$
\gamma_c^1 - \left(y u_\star(p^0) + (1 - y) u_\star(p^1) \right) \leq \frac{1 - \delta}{\delta} \left(y u_\star(p^0) + (1 - y) u_\star(p^1) - u_\star(p) \right),
$$

provided δ is high enough.

This implies that for δ high enough, there is $\tilde{p}^1 \in V(p^1) \cap [p^0, p^1]$ such that $\gamma_c^1 =$ $\tilde{y}u_{\star}(p^{0}) + (1 - \tilde{y})u_{\star}(\tilde{p}^{1}), \text{ and } \tilde{y}p^{0} + (1 - \tilde{y})\tilde{p}^{1} = p.$

We first define a strategy pair (σ, τ) up to stage N_1+1 . Player 1 repeats an optimal action $o(p)$ at all stages $1, \ldots, N_1$. Player 2 plays $o(q)$ at all stages $1, \ldots, N_1 - 1$. In stage N_1 , player 2 plays both actions $o(q)$ and $b' \neq o(q)$ with probabilities such that beliefs in stage $N_1 + 1$ are (p^0, q) following b', and (\tilde{p}^1, q) following $o(q)$.

We now define the continuation of (σ, τ) following $o(q)$. Define γ_c^2 by the equality $\gamma^2 = (1 - \delta^{N_1}) v_\star(q) + \delta^{N_1} \gamma_c^2$. The continuation of (σ, τ) in the other case is defined in an analog way, except that γ_c^2 has to be replaced by γ_c^2 + $1-\delta$ δ $c(q, b')$, and the equations that describe equilibrium constraints have to be adjusted.

Let \tilde{N}_2 be the least integer (possibly infinite) such that $\tilde{\gamma}^2 \geq xv_*(q^0) + (1-x)v_*(q^1)$, where $\tilde{\gamma}^2$ is defined by $\gamma_c^2 = (1 - \delta^{\tilde{N}_2})v_\star(q) + \delta^{\tilde{N}_2} \tilde{\gamma}^2$. The choice of \tilde{N}_2 implies

$$
\tilde{\gamma}^2 - \left(xv_{\star}(q^0) + (1-x)v_{\star}(q^1)\right) \leq \frac{1-\delta}{\delta} \left(xu_{\star}(q^0) + (1-x)u_{\star}(q^1) - v_{\star}(q)\right),
$$

provided δ is high enough. This implies that for δ high enough, there is $\tilde{q}^1 \in V(q^1) \cap$ $[q^{0}, q^{1}]$ such that $\tilde{\gamma}^{2} = \tilde{x}v_{\star}(q^{0}) + (1 - \tilde{x})v_{\star}(\tilde{q}^{1}),$ and $\tilde{x}q^{0} + (1 - \tilde{x})\tilde{q}^{1} = q$.

The continuation profile is defined as follows. Player 2 repeats $o(q)$ in all stages $N_1+1,\ldots,N_1+\tilde{N}_2$. Player 1 repeats $o(\tilde{p}^1)$ in all stages $N_1,\ldots,N_1+\ldots\tilde{N}_2-1$. In stage $N_1 + \tilde{N}_2$, player 1 plays both actions $o(p^1)$ and $a \neq o(p^1)$ with probabilities such that the belief of player 2 is equal to \tilde{q}^1 following $o(\tilde{p}^1)$, and to q^0 following a.

²³ $N_1 = \infty$ if $\gamma^1 = u_\star(p)$.

Following $o(\tilde{p}^1)$, players switch to an equilibrium of the game $\Gamma(\tilde{p}^1, \tilde{q}^1)$ with payoff $(u_{\star}(\tilde{p}^1), v_{\star}(\tilde{q}^1)).$ Following a, players switch to an equilibrium of the game $\Gamma(\tilde{p}^1, q^0)$ with payoff $(u_\star(\tilde{p}^1) + \frac{1-\delta}{\delta}c(\tilde{p}^1, a), v_\star(q^0)).$

Beliefs and actions off-the-equilibrium-path are defined as in the proof of Lemma [1.](#page-29-0) The definition of beliefs and continuation payoffs ensure that players are indifferent whenever randomizing, and that the overall payoff is exactly γ .

Observe also that there exists a neighborhood $V(p)$ of p, with the following property. The two beliefs p^0 and p^1 associated with $p' \in V(p)$ can be chosen to be continuous in p' and $x'u_*(p'^0) + (1-x')u_*(p'^1)$ (with $x'p'^0 + (1-x')p'^1 = p'$) is bounded away from $u_{\star}(p')$ over $V(p)$. Together with the symmetric property for player 2, this ensures that the robustness result mentioned after Proposition [1](#page-26-2) holds. This concludes the proof of Proposition [1.](#page-26-2)

We next proceed to the proof of Proposition [2.](#page-27-0) Let $p \in \overset{\circ}{\Delta}$ † $(S \times M), q \in \overset{\circ}{\Delta}$ † $(T \times L),$ and $c > 0$ be given. Let γ be such that $|\gamma^1 - u_\star(p)| \leq (1-\delta)c$ and $|\gamma^2 - v_\star(q)| \leq (1-\delta)c$. Let $[p^0, p^1]$ be the segment associated with p in the proof of Proposition [1,](#page-26-2) and let y solve the equation $p = yp^{0} + (1 - y)p^{1}$. Set

$$
\eta := (yu_*(p^0) + (1 - y)u_*(p^1)) - u_*(p) > 0,
$$

and let N_1 be defined as in the proof of Proposition [1.](#page-26-2) By construction, one has

$$
(1 - \delta^{N_1 - 1})u_\star(p) + \delta^{N_1 - 1}(u_\star(p) + \eta) < \gamma^1 \le u_\star(p) + (1 - \delta)c,
$$

hence $\eta \delta^{N_1-1} \leq (1-\delta)c$. Similarly, one has $\eta \delta^{N_2-1} \leq (1-\delta)c$ (for a possibly lower value of η). In the construction of Proposition [1,](#page-26-2) players repeat the same action until stage min $\{N_1, N_2\}$. Therefore, for any $p' \in \Delta^{\dagger}(S \times M)$ and every strategy σ , one has

$$
\gamma^1(p', q, \sigma, \tau_{p,q,\gamma}) \le (1 - \delta^{\min\{N_1, N_2\}}) u_{\star}(p') + \delta^{\min\{N_1, N_2\}} \le u_{\star}(p') + (1 - \delta) \frac{\delta c}{\eta}.
$$

The result follows, with $C = c/\eta$.

B General games

We here complete the proof of the Main Theorem. We start with a few notations and remarks in the spirit of Section [A.](#page-26-3) We let initial distributions $p \in \Delta(S \times L_S \times M_S)$ and $q \in \Delta(T \times L_T \times M_T)$ be given. W.l.o.g., we also assume that $p(l_S) > 0$ and $p(l_T) > 0$ for each $l_S \in L_S$ and $l_T \in L_T$.^{[24](#page-38-0)} We assume that the information of each player i is valuable for the other player. This is equivalent to assuming that, for each $l_S \in L_S$, there is no action $a \in A$ that is optimal at all beliefs $p_{l_S,m_S} := p(\cdot | l_S, m_S)$, $m_S \in M_S$.

As the play proceeds, player 2 may disclose information relative to \mathbf{m}_{S} , and player 1's belief about \mathbf{m}_S may change. Analogously to the case of self-ignorant games (see Eq. [\(5\)](#page-26-4)), the belief \mathbf{p}_n of player 1 given $\mathbf{l}_S = l_s$ is always in the set

$$
\Delta_{l_S}^{\dagger}(S \times M_S) = \text{conv}\{p_{l_S,m_S} \otimes 1_{l_S} \otimes 1_{m_S}, m_S \in M_S\}.
$$

Note that $p(\cdot | l_S)$ lies in the relative interior of the set $\Delta^{\dagger}(S \times M_S)$.

For $m_T \in M_T$, we define $\Delta_{m_T}^{\dagger}(T \times L_T)$ in a symmetric way. The results of Section [A](#page-26-3) will be applied to the different sets $\Delta_{l_S}^{\dagger}(S \times M_S)$ and $\Delta_{m_T}^{\dagger}(T \times L_T)$ of initial distributions.

B.1 Providing Incentives

For simplicity, we focus here on player 1. Analog properties hold for player 2 as well. We first define an equivalence relation \sim over L_S . As we will see, two signals L_S and \bar{l}_S such that $\underline{l}_S \sim \bar{l}_S$ may be merged, and treated as a single signal. Given $l_S \in L_S$, we define a vector \vec{Z}^{l_S} of size $M_S \times A \times A$ by

$$
\vec{Z}_{m_S,a,a'}^{l_S} := p(m_S|l_S) \left(u(p_{l_S,m_S}, a) - u(p_{l_S,m_S}, a') \right), \text{ for } m_S \in M_S, \text{ and } a, a' \in A.
$$

Because the information held by player 2 is valuable for player 1, $\vec{Z}^{l_S} \neq \vec{0}$, for each $l_S \in L_S.^{25}$ $l_S \in L_S.^{25}$ $l_S \in L_S.^{25}$

Definition 2 Let $l_S, l_S \in L_S$ be given. The two signals l_S and \bar{l}_S are equivalent, written $\underline{l}_S \sim \overline{l}_S$, if the two vectors \overline{Z}^{l_S} and \overline{Z}^{l_S} are positively collinear, that is, if

$$
\exists \alpha > 0, \vec{Z}^{\bar{l}_S} = \alpha \vec{Z}^{l_S}.\tag{7}
$$

Plainly, if the two distributions $p(\cdot|\underline{l}_S)$ and $p(\cdot|\overline{l}_S)$ in $\Delta(S \times M_S)$ coincide, then $\underline{l}_S \sim \overline{l}_S$. However, the converse implication does not hold.

 24 And we make the symmetric assumption for player 2.

²⁵Indeed, if $\vec{Z}^{l_S} = \vec{0}$, then any action $a \in A$ is optimal at p_{l_S,m_S} , for each m_S .

Observe that, if $l_S \sim \bar{l}_S$ and $\vec{Z}^{\bar{l}_S} = \alpha \vec{Z}^{l_S}$, then for every two mixed actions $x, x' \in \Delta(A)$ we have:

$$
p(m_S|\bar{l}_S) \left(u(p_{\bar{l}_S,m_S}, x) - u\left(p_{\bar{l}_S,m_S}, x'\right)\right) = \alpha p(m_S|l_S) \left(u(p_{l_S,m_S}, x) - u\left(p_{l_S,m_S}, x'\right)\right). \tag{8}
$$

As a preparation for Lemma [4](#page-39-0) below, observe that a strategy σ may be viewed as a collection $(\sigma_{l_S})_{l_S \in L_S}$, with the interpretation that $\sigma_{l_S}: L_T \times H \to \Delta(A)$ is the 'interim' strategy used if $I_S = l_S$.^{[26](#page-39-1)}

Lemma 4 Let τ be any strategy of player 2. Then there exists a best reply σ of player 1 to τ such that $\sigma_{\bar{l}_S} = \sigma_{l_S}$ whenever $\bar{l}_S \sim \underline{l}_S$.

According to Lemma [4,](#page-39-0) player 1 has a best reply that depends only on the equivalence class of \mathbf{l}_S .

Proof. Let a strategy τ of player 2 be fixed throughout. Given $f : H \to \Delta(A)$, and $(l_S, l_T) \in L_S \times L_T$, we denote by $\gamma^1(f, \tau | l_S, l_T)$ the interim expected payoff of player 1, when getting $\mathbf{l}_S = l_S$, $\mathbf{l}_T = l_T$, and when playing according to f thereafter. Given $n \geq 1$, we also denote by $g_n^1(f, \tau | l_S, l_T)$ the corresponding payoff at stage n.

We let $\bar{l}_S \sim l_S$ be any two equivalent signals, so that $\vec{Z}^{\bar{l}_S} = \alpha \vec{Z}^{\bar{l}_S}$ for some $\alpha > 0$. We will prove that, for every two "interim strategies" $f : H \to \Delta(A)$ and $f': H \to \Delta(A)$, for every $l_T \in L_T$ and every stage $n \geq 1$, one has

$$
g_n^1(f, \tau | \bar{l}_S, l_T) - g_n^1(f', \tau | \bar{l}_S, l_T) = \alpha \left(g_n^1(f, \tau | \underline{l}_S, l_T) - g_n^1(f', \tau | \underline{l}_S, l_T) \right). \tag{9}
$$

Equation [\(9\)](#page-39-2) will imply that

$$
\gamma^1(f,\tau|\bar{l}_S,l_T) - \gamma^1(f',\tau|\bar{l}_S,l_T) = \alpha\left(\gamma^1(f,\tau|\underline{l}_S,l_T) - \gamma^1(f',\tau|\underline{l}_S,l_T)\right),
$$

from which the result follows. Indeed, if f is better than f' when the signal is \bar{l}_S , then it is also the case when the signal is l_S . Therefore if f is a best response when the signal is \bar{l}_S , then it is also a best response when the signal is \underline{l}_S .

We let a stage $n \geq 1$ be given. We fix $(l_S, l_T) \in L_S \times L_T$, and we decompose the payoff $g_n^1(f, \tau | l_S, l_T)$ as follows. For a given sequence of moves $h \in H_n := (A \times B)^{n-1}$, we denote by $\mathbf{P}_{f,\tau}(h|l_S, l_T)$ the probability that h occurs, when $(\mathbf{l}_S, \mathbf{l}_T) = (l_S, l_T)$ and

²⁶To be formal, $\sigma_{l_S}(l_T, h)$ is defined to be $\sigma(l_S, l_T, h)$.

players play according to f and τ . We denote by $\mathbf{P}_{f,\tau}(\cdot|h, l_S, l_T) \in \Delta(S \times M_S)$ the belief which is then held by player 1.

With these notations, one has

$$
g_n^1(f, \tau | l_S, l_T) = \sum_{h \in H_n} \mathbf{P}_{f, \tau}(h | l_S, l_T) u \left(\mathbf{P}_{f, \tau}(\cdot | h, l_S, l_T) \right), f(h)). \tag{10}
$$

The belief of player 1 following h is given by

$$
\mathbf{P}_{f,\tau}(s|h, l_S, l_T) = \frac{1}{\mathbf{P}_{f,\tau}(h, l_S, l_T)} \sum_{m_S \in M_S} \mathbf{P}_{f,\tau}(s, h, l_S, l_T, m_S), \ s \in S,
$$

where $\mathbf{P}_{f,\tau}(h, l_S, l_T) = \mathbf{P}_{f,\tau}(h|l_S, l_T) p(l_S) q(l_T)$. Because the state s and the history of moves until stage *n* are conditionally independent given (l_S, l_T, m_S) , this belief is equal to

$$
\mathbf{P}_{f,\tau}(s|h, l_S, l_T) = \frac{1}{\mathbf{P}_{f,\tau}(h, l_S, l_T)} \sum_{m_S \in M_S} \mathbf{P}_{f,\tau}(h|l_S, l_T, m_S) p_{l_S, m_S}(s) p(l_S, m_S) q(l_T). \tag{11}
$$

Plugging (11) into (10) , and using the linearity of u, one gets

$$
g_n^1(f, \tau | l_S, l_T) = \sum_{h \in H_n} \sum_{m_S \in M_S} p(m_S | l_S) \mathbf{P}_{f, \tau}(h | l_S, l_T, m_S) u(p_{l_S, m_S}, f(h)). \tag{12}
$$

Using [\(12\)](#page-40-2) for both f and f', and because $\sum_{h\in H_n} \mathbf{P}_{\sigma,\tau}(h \mid l_S, l_T, m_S) = \sum_{h'\in H_n} \mathbf{P}_{\sigma',\tau}(h' \mid l_S, l_T, m_S)$ l_S, l_T, m_S = 1, we obtain:

$$
g_n^1(f, \tau | l_S, l_T) - g_n^1(f', \tau | l_S, l_T)
$$
\n
$$
= \sum_{m_S \in M_S} p(m_S | l_S) \left(\sum_{h \in H_n} \mathbf{P}_{f, \tau}(h | l_S, l_T, m_S) u(p_{l_S, m_S}, f(h)) - \sum_{h' \in H_n} \mathbf{P}_{f', \tau}(h' | l_S, l_T, m_S) u(p_{l_S, m_S}, f'(h')) \right)
$$
\n
$$
= \sum_{m_S \in M_S} \sum_{h \in H_n} \sum_{h' \in H_n} \mathbf{P}_{f, \tau}(h | l_S, l_T, m_S) \mathbf{P}_{f', \tau}(h' | l_S, l_T, m_S) \times p(m_S | l_S) \left(u(p_{l_S, m_S}, f(h)) - u(p_{l_S, m_S}, f'(h')) \right).
$$
\n(13)

Because l_S and \bar{l}_S are equivalent, Eq. [\(9\)](#page-39-2) follows by [\(8\)](#page-39-3), and [\(13\)](#page-40-3) applied to both $l_S = \underline{l}_S$ and $l_S = \overline{l}_S$.

Lemma [4](#page-39-0) implies that any equilibrium of the modified game in which player 1 only observes the equivalence class of \mathbf{l}_S is an equilibrium of the original game. Put it differently, the set of equilibrium payoffs of the game in which players do not distinguish between equivalent signals is a subset of the set of equilibrium payoffs of the game we started with. Besides, the values of u_* and v_* (resp., of u_{**} and v_{**}) are the same for both games.

Therefore, it is sufficient to prove that the conclusion of the Main Theorem holds for the modified game. In particular, we may and will assume from here on that, for every two signals $\bar{l}_S \neq \underline{l}_S$, the vectors $\vec{Z}^{\bar{l}_S}$ and $\vec{Z}^{\bar{l}_S}$ are not positively collinear. We also make the symmetric assumption for player 2. A direct consequence of this assumption is Corollary [1](#page-41-0) below.

Corollary 1 Let $\bar{a} \in A$ be arbitrary. For $l_S \in L_S$, define the vector \vec{Y}^{l_S} of size $M_S \times A$ by

$$
\vec{Y}_{m_S,a}^{l_S} := p(m_S|l_S) \left(u(p_{l_S,m_S}, a) - u(p_{l_S,m_S}, \bar{a}) \right), \quad m_S \in M_S, a \in A.
$$

Then for every two signals $\bar{l}_S \neq \underline{l}_S$, the two vectors $\vec{Y}^{\bar{l}_S}$ and $\vec{Y}^{\underline{l}_S}$ are not positively collinear.

The vector \vec{Y}^{l_S} is equal to the projection of \vec{Z}^{l_S} on a lower-dimensional space. Hence, linear independence of $\vec{Y}^{\bar{l}_S}$ and of \vec{Y}^{l_S} does not follow in general from linear independence of $\vec{Z}^{\bar{l}_S}$ and $\vec{Z}^{\bar{l}_S}$, and an *ad hoc* proof is needed.

Proof. We argue by contradiction, and assume that $\vec{Y}^{\bar{l}s} = \alpha \vec{Y}^{l_s}$ for some $\alpha > 0$. Let $m_S \in M_S, a, a' \in A$ be arbitrary. Observe that $\vec{Z}_{m_S,a,a'}^{l_S} = \vec{Y}_{m_S,a}^{l_S} - \vec{Y}_{m_S,a'}^{l_S}$, for $l_S = \bar{l}_S, \underline{l}_S$. Hence $\vec{Z}^{\bar{l}_S} = \alpha \vec{Z}^{\underline{l}_S}$, a contradiction.

The next lemma is central to the provision of incentives (phase 2 of the equilibrium play). Given $x: L_S \times M_S \to \Delta(A)$, and for every $l_S, k \in L_S$, we define

$$
\mathbf{E}_x[l_S \to k] = \sum_{m_S \in M_S} p(m_S|l_S) u(p_{l_S, m_S}, x_{k, m_S}),
$$

with the following interpretation. The expression $\mathbf{E}_x[l_S \to k]$ is the expected stage payoff when player 1 gets $\mathbf{l}_S = l_S \in L_S$, 'reports' $k \in L_S$, is told \mathbf{m}_S , and plays the mixed action x_{k,\mathbf{m}_S} that depends on player 1's report, and on player 2's signal.^{[27](#page-41-1)} According to Lemma [5](#page-41-2) below, the map x can be chosen in a way that this expected payoff is highest when reporting truthfully.

²⁷We use the different letter k to distinguish between a signal and a report, although both belong to the same set L_S .

Lemma 5 There exists x^* : $L_S \times M_S \rightarrow \Delta(A)$, such that

$$
\mathbf{E}_{x^*}[l_S \to k] < \mathbf{E}_{x^*}[l_S \to l_S], \text{ for every } l_S, k \in L_S, l_S \neq k.
$$

Proof. Let $\bar{a} \in A$ be arbitrary, and let $x : L_S \times M_S \to \Delta(A)$ be given. For $k \in L_S$, we define a vector \vec{X}^k of size $M_S \times A$ by $\vec{X}_{m_S,a}^k := x_{k,m_S}(a)$, $m_S \in M_S$, $a \in A$. Observe that $x_{k,m_S}(\bar{a}) = 1 - \sum$ $a \neq \bar{a}$ $x_{k,m_S}(a)$. Hence, $\mathbf{E}_x[l_S \to k]$ may be rewritten as

$$
\mathbf{E}_{x}[l_{S} \rightarrow k] = \sum_{m_{S} \in M_{S}} p(m_{S}|l_{S}) u(p_{l_{S},m_{S}}, \bar{a})
$$

+
$$
\sum_{m_{S} \in M_{S}} \sum_{a \in A} p(m_{S}|l_{S}) x_{k,m_{S}}(a) (u(p_{l_{S},m_{S}}, a) - u(p_{l_{S},m_{S}}, \bar{a}))
$$

=
$$
\vec{Y}^{l_{S}} \cdot \vec{X}^{k} + \sum_{m_{S} \in M_{S}} p(m_{S}|l_{S}) u(p_{l_{S},m_{S}}, \bar{a}).
$$

Because the second term in the last displayed equation does not depend on k , it is sufficient to construct x such that

$$
\vec{Y}^{ls} \cdot \vec{X}^k < \vec{Y}^{ls} \cdot \vec{X}^{ls} \text{ for every } l_S, k \in L_S, l_S \neq k. \tag{14}
$$

For $l_S \in L_S$, define

an

$$
\tilde{X}^{l_S} := \frac{1}{\|\vec{Y}^{l_S}\|_2} \vec{Y}^{l_S},
$$

and let $l_S \neq k$ be arbitrary in L_S . Then by the Cauchy-Schwartz inequality,

$$
\tilde{X}^k \cdot \vec{Y}^{l_S} = \frac{\vec{Y}^k}{\|\vec{Y}^k\|_2} \cdot \vec{Y}^{l_S} < \|\vec{Y}^{l_S}\|_2 = \frac{\vec{Y}^{l_S}}{\|\vec{Y}^{l_S}\|_2} \cdot \vec{Y}^{l_S} = \tilde{X}^{l_S} \cdot \vec{Y}^{l_S},
$$

where the strict inequality holds since \vec{Y}^k and \vec{Y}^{l_S} are not positively collinear. There-fore, [\(14\)](#page-42-0) holds with $(\tilde{X}^{ls})_{l_S \in L_S}$. Note that (14) still holds when the same constant is added to all components, and/or when all components are multiplied by the same constant $\phi > 0$. Choose $\beta \in \mathbf{R}$ and $\phi > 0$ such that all components of $\phi \tilde{X}^{l_S} + \beta$ lie in $(0, \frac{1}{|M_{\odot}|})$ $\frac{1}{|M_S \times A|}$, for all l_S . Because $Y_{m_S,\bar{a}}^{l_S} = 0$, it suffices to set

$$
x_{l_S,m_S}^\star(a) = \phi \tilde{X}_{m_S,a}^{l_S} + \beta \text{ for } a \neq \bar{a},
$$
d
$$
x_{l_S,m_S}^\star(\bar{a}) = 1 - \sum_{a \neq \bar{a}} x_{l_S,m_S}(a). \quad \blacksquare
$$

Given $\varepsilon_2 : M_S \to \Delta(M_S)$, and $l_S, k \in L_S$, we define

$$
\mathbf{E}_{\varepsilon_2,x^\star}[l_S \to k] = \sum_{m_S,\mu \in M_S} p(m_S|l_S) \varepsilon_2(\mu|m_S) u(p_{l_S,\mu},x_{k,\mu}^\star).
$$

This is the expected stage payoff of player 1 when (i) player 1 gets $I_s = l_s$, and 'reports' k, (ii) player 2 draws $\mu \in M_S$ according to $\varepsilon_2(\cdot | \mathbf{m}_S)$ and (iii) player 1 plays $x_{k,\mu}^{\star}$. We here abuse notation and write $p_{l_S,\mu}$ for the belief of player 1, given l_S and $\mu^{.28}$ $\mu^{.28}$ $\mu^{.28}$

Observe that the expectation $\mathbf{E}_{\varepsilon_2,x^*}[l_S \to k]$ is continuous w.r.t. ε_2 , and that $\mathbf{E}_{\varepsilon_2,x^*}[l_S \to k]$ is equal to $\mathbf{E}_{x^*}[l_S \to k]$ when $\varepsilon_2(\cdot | m_S)$ assigns probability 1 to m_S , for each m_S . Corollary [2](#page-43-1) below therefore follows from Lemma [5](#page-41-2) by continuity.

Corollary 2 There exists ε_2 : $M_S \to \overset{\circ}{\Delta}(M_S)$, such that

$$
\mathbf{E}_{\varepsilon_2,x^{\star}}[l_S \to k] < \mathbf{E}_{\varepsilon_2,x^{\star}}[l_S \to l_S], \text{ for every } l_S, k \in L_S, l_S \neq k. \tag{15}
$$

We fix ε_2 and x^* for the rest of the paper. Because the distribution $\varepsilon_2(\cdot|m_S)$ has full support, the conditional distribution $p_{l_S,\mu}$ lies in the relative interior of $\Delta_{l_S}^{\dagger}(S \times M_S)$ (for each $\mu \in M_S$). Define ε_1 analogously.

B.2 Equilibrium strategies – Structure

We let a payoff vector $\gamma = (\gamma^1, \gamma^2)$ be given, with $u_* < \gamma^1 < u_{**}$ and $v_* < \gamma^2 < v_{**}$. We will construct a sequential equilibrium with payoff γ . We let the discount factor δ be given. In the construction we add one additional message, \Box , to each player.

Given $x \in \Delta(A)$, and given a number N of stages, we denote by $\vec{a}^N(x) \in A^N$, a sequence of actions of length N that provides the best approximation of the mixed action x in terms of discounted frequencies. That is, $\vec{a}^N(x) = (a_n)_{1 \leq n \leq N}$ is chosen to minimize $||x_{\delta}(\vec{a}^{N}) - x||_{\infty}$, where

$$
x_{\delta}(\vec{a}^{N})[a] := \frac{1-\delta}{1-\delta^{N}} \sum_{n=1}^{N} \delta^{n-1} 1_{\{a_{n}=a\}}, \ a \in A.
$$

The sequence $\vec{a}^N(x_{k,\mu_1}^*)$ will be the sequence of actions required from player 1 in phase 2.2, when player 1 reports $k \in L_S$ and player 2 sends the message $\mu_1 \in$

²⁸Note that, for fixed μ , the belief $p_{l_S,\mu}$ depends on ε_2 , although this is not emphasized in the notation.

 $M_S \cup \{\Box\}$. For $\mu_1 = \Box$, we let $\vec{a}^N(x_{k,\mu_1}^*)$ be an arbitrary sequence of actions, that does not depend on $k \in L_S$.

Similarly, $\vec{b}^{N}(y) \in B^{N}$ is a vector that approximates the mixed action y in terms of discounted frequencies.

We set $K_1 := \max\{|L_S|, |M_T|\}$, and we let $\alpha_1 : L_S \to A^{K_1}$ and $\beta_1 : M_T \to B^{K_1}$ be arbitrary one-to-one maps. Similarly, we set $K_2 := 1 + \max\{|L_T|, |M_S|\}$, and we let $\alpha_2 : L_T \cup \{\Box\} \to A^{K_2}$ and $\beta_2 : M_S \cup \{\Box\} \to B^{K_2}$ be arbitrary one-to-one maps. The maps α_1 and β_1 are used to encode reports on one's own state into sequences of actions, while the maps α_2 and β_2 are used to encode messages on the other player's state into sequences of actions.

We let $\pi^1 \in \overset{\circ}{\Delta}(L_T)$ and $\pi^2 \in \overset{\circ}{\Delta}(M_S)$ be arbitrary distributions with full support.

We now proceed to the definition of a strategy profile $(\sigma_{\delta}, \tau_{\delta})$. The definition involves additional parameters θ, ζ , and $\psi^i, \psi^i \mid (i = 1, 2)$, all in $(0, 1)$, which will be chosen later. We first define the profile only at information sets that are not ruled out by the definition of $(\sigma_{\delta}, \tau_{\delta})$ at earlier information sets. The definition of $(\sigma_{\delta}, \tau_{\delta})$ at information sets that are reached with probability zero will be provided after.

Phase 1 It lasts K_1 stages. Player 1 plays the sequence $\alpha_1(\mathbf{l}_S)$ of actions, and player 2 plays the sequence $\beta_1(\mathbf{m}_T)$ of actions.

Phase 2 It is divided into two subphases, Phase 2.1 and Phase 2.2.

Phase 2.1 It lasts K_2 stages. Player 1 first draws a message $\lambda_1 \in L_T \cup \{\Box\}.$ The probability assigned to \Box , (resp. to each $l'_T \in L_T$), is equal to $1 - \zeta$ (resp. $\zeta \times \varepsilon_1(l'_T | \mathbf{l}_T)$). Symmetrically, player 2 draws a message $\mu_1 \in M_S \cup$ $\{\Box\}$. The probability assigned to \Box , (resp. to each $m'_S \in M_S$) is equal to $1-\zeta$, (resp. $\zeta \times \varepsilon_2(m_S'|\mathbf{m}_S)$).

In that phase, the players play the sequences $\alpha_2(\lambda_1)$ and $\beta_2(\mu_1)$ of actions.

Phase 2.2 It lasts $\nu := \lfloor \frac{\ln(1-\theta)}{\ln \delta} \rfloor$ $\frac{(1-\theta)}{\ln \delta}$ stages. Player 1 infers μ_1 from the actions played by player 2 in Phase 2.1, and plays the sequence $\vec{a}^{\nu}(x_{1s,\mu_1}^{\star})$ of actions. Meanwhile, player 2 infers λ_1 from the actions played by player 1 in Phase 2.1, and plays the sequence $\vec{b}(y_{\mathbf{m}_T,\lambda_1})$ of actions.

Phase 3 It lasts K_2 stages. Player 1 draws a message $\lambda_2 \in L_T$. The distribution of λ_2 depends on λ_1 . If $\lambda_1 = \square$, the probability assigned to I_T (resp. to each $l'_T \neq l_T$), is equal $(1 - \psi_{\Box}^1) + \psi_{\Box}^1 \times \pi^1(l_T)$ (resp. $\psi_{\Box}^1 \times \pi^1(l'_T)$). If $\lambda_1 \neq \Box$, the probability assigned to λ_2 is equal to $(1 - \psi^1) + \psi^1 \times \pi^1(\lambda_2)$ if $\lambda_2 = \mathbf{I}_T$, and it is equal $\psi^1 \times \pi^1(\lambda_2)$ otherwise. Player 2 draws a message $\mu_2 \in M_S$. The distribution of μ_2 depends on μ_1 , and is obtained as for player 1.

In this phase, the players play the sequences $\alpha_2(\lambda_2)$ and $\beta_2(\mu_2)$ of actions.

Phase 4 It contains all remaining stages. We denote by $N = K_1 + 2K_2 + \nu + 1$ its first stage. Let $h = (a_n(h), b_n(h))_{n \le N} \in (A \times B)^{N-1}$ be the history of moves up to stage N. Player 2 infers from h the belief $p_n(h)$ held by player 1 in each stage $n < N$ along h. In this computation, the report of player 1 in **Phase 1** is assumed to be truthful. For $n < N$, the belief $q_n(h) \in \Delta(T \times L_T)$ is defined in a symmetric way. The players compute

$$
c_1(h) = \delta^{-N} \sum_n (1-\delta) \delta^{n-1} c(p_n(h), a_n(h)) \text{ and } c_2(h) = \delta^{-N} \sum_n (1-\delta) \delta^{n-1} c(q_n(h), b_n(h)),
$$

where the sum is taken over all stages n of **Phases 1, 2.1** and 3. Players then start playing according to the equilibrium profile of the semi-ignorant game $\Gamma(p_n(h), q_n(h))$, with payoff $(u_*(p_n(h)) + c_1(h), v_*(q_n(h)) + c_2(h)$.

Some interpretation may be helpful. In Phase 2.1, the message \Box is uninforma-tive,^{[29](#page-45-0)} and is sent with high probability. In Phase 3, the level noise in the message sent by player 1 depend on player 1's first message, and is either ψ^1 if the first message was informative, or ψ_{\square}^1 otherwise.

B.3 Equilibrium Strategies – Parameter values

We now fix the parameter values, starting with θ . As $\delta \rightarrow 1$, the discounted weight of the $\frac{\ln(1-\theta)}{\ln \delta}$ $\frac{(1-\theta)}{\ln \delta}$ stages of Phase 2.2 converges to θ . Thus, θ is a measure of the contribution of the checking phase 2.2 to the total payoff. We choose $\theta \in (0,1)$ to be small enough so that the following set of inequalities is satisfied:

$$
(1 - \theta)\mathbf{E}[u_{\star}(p_{l_S, \mathbf{m}_S})|\mathbf{l}_S = l_S, \mu_1 = m_S] > u_{\star}(p(\cdot|\mathbf{l}_S = l_S, \mu_1 = m_S)), \ \forall m_S \in M_3 \tag{17}
$$
\n
$$
(1 - \theta)u_{\star \star} > \gamma^1,
$$

 29 Since its probability does not depend on signals.

together with the symmetric conditions for player 2.

By construction, the conditional distribution of \mathbf{m}_S given $(l_S, \mu_1) = (l_S, m_S)$ is independent of ζ , and only depends on the fixed map ε_2 . Since $\varepsilon_2(\cdot|m_S')$ has full support for each m_S' , this conditional distribution has full support. Therefore, the residual information held by player 2 is still valuable to player 1, whatever be $\mu^1 \in$ $\{\Box\} \cup M_S$. In particular, [\(16\)](#page-45-1) holds with $\theta = 0$, and thus also for $\theta > 0$ small enough. Because γ^1 < $u_{\star\star}$, condition [\(17\)](#page-45-1) is also satisfied for small θ .

Condition [\(16\)](#page-45-1) ensures that, even if payoffs in phase 2.2 are very low, the weight θ of phase 2.2 is so small, that the residual value of the information held by player 2 can still offset the cost incurred when playing the prescribed sequence in phase 2.2. Condition [\(16\)](#page-45-1) is designed to make sure that, when in phase 2.2, player 1 will rather play the prescribed sequence of actions, than switch to an optimal action.

Observe that with probability $1-\zeta$, player 1 receives no information prior to phase 3. Hence, for $\zeta > 0$ small, the bulk of information exchange takes place in phase 3. Condition [\(17\)](#page-45-1) ensures that, even if all information exchange is postponed to phase 3, payoffs as high as γ^1 can be implemented.

Choose $\zeta \in (0,1)$ to be small enough so that the two inequalities

$$
(1 - \zeta)u_{\star} + \zeta u_{\star\star} < \gamma^1 < (1 - \zeta)(1 - \theta)u_{\star\star} \tag{18}
$$

hold, together with the analog inequalities for player 2.

In phase 2.2, the (conditional) optimal payoff of player 1 is u_{\star} if $\mu_1 = \Box$, and does not exceed $u_{\star\star}$ if $\mu_1 \neq \Box$. The first inequality ensures that the probability $1 - \zeta$ of not disclosing information in phase 2.1 ($\mu_1 = \Box$) is so high that the expectation of the optimal payoff given μ_1 does not exceed γ^1 . That is, additional information must be disclosed in phase 3 in order to implement γ . This inequality, together with [\(17\)](#page-45-1), will allow us to adjust other parameter values in a way that the overall payoff is γ . The second inequality in [\(18\)](#page-46-0) does not play a critical role.

We now choose the value of $\psi^2 \in (0,1)$ small enough so that, for every $l_S \in$ $L_S, m_S \in M_S$

$$
(1 - \theta)\mathbf{E}[u_{\star}(p(\cdot|\mathbf{l}_{S}, \mu_{1}, \mu_{2}))|\mathbf{l}_{S} = l_{S}, \mu_{1} = m_{S}] > u_{\star}(p(\cdot|\mathbf{l}_{S} = l_{S}, \mu_{1} = m_{S})). \tag{19}
$$

In this expression, $p(\cdot|I_s, \mu_1, \mu_2)$ is the belief held by player 1 at the beginning of phase 4, after having received the two messages μ_1, μ_2 of player 2. The left-hand side of [\(19\)](#page-46-1) is continuous w.r.t. ψ^2 . For $\psi^2 = 0$, μ_2 is equal to \mathbf{m}_S with probability 1, and [\(19\)](#page-46-1) therefore holds by [\(16\)](#page-45-1). Hence [\(19\)](#page-46-1) holds for $\psi^2 > 0$ small enough.

Observe that all parameters values ζ , θ , ψ^2 are independent of the discount factor. The last parameter, ψ_{\square}^2 is chosen such that the expected payoff of player 1 is γ^1 . We first argue that for a given ψ_{\square}^2 , the limit discounted payoff of player 1, as $\delta \to 1$, is equal \rm{to}^{30} \rm{to}^{30} \rm{to}^{30}

$$
\theta \mathbf{E}[u(p(\cdot|\mathbf{l}_S,\mu_1),x_{\mathbf{l}_S,\mu_1}^*)] + (1-\theta)\mathbf{E}[u_\star(\mathbf{p}_N)].
$$
\n(20)

Here is why. The contribution of Phases 1, 2.1 and 3 vanishes, as the length of these phases is fixed independently of δ . The expected payoff in phase 2.2 converges^{[31](#page-47-1)} to $\mathbf{E}[u(p(\cdot\vert {\bf I}_S,\mu_1),x^{\star}_{{\bf I}_S,\mu_1})]$. Finally, for a fixed δ , the expected continuation payoff from stage N is equal to $\mathbf{E}[u_{\star}(\mathbf{p}_N)+c_1(\mathbf{h}_N)]$. As Lemma [6](#page-47-2) will show, $\mathbf{E}[c_1(\mathbf{h}_N)]$ will converge to 0.

Observe that for $\psi_{\square}^2 = 0$, and following $\mu_1 = \square$, the message μ_2 of player 2 is noninformative. Thus, conditional on the event that $\mu_1 = \Box$, player 2 does not disclose information prior to phase 4. Thus, for $\psi_{\square}^2 = 0$, the left-hand side of [\(20\)](#page-47-3) does not exceed $(1 - \zeta)u_{\star} + \zeta u_{\star\star}$ which by [\(18\)](#page-46-0) is less than γ^1 . If $\psi_{\square}^2 = 1$, following $\mu_1 = \square$ the message μ_2 is fully informative, and the left-hand side of [\(20\)](#page-47-3) is at least equal to $\zeta u_{\star} + (1 - \zeta) (\theta u_{\star} + (1 - \theta) u_{\star \star}),$ which exceeds γ^1 by [\(18\)](#page-46-0). It follows that for δ high enough, say $\delta \geq \bar{\delta}_1$, there exists $\psi_{\square}^2(\delta) \in (0,1)$, such that the discounted payoff of player 1 is equal to γ^1 , and such that $\psi_{\square}^2(1) := \lim_{\delta \to 1} \psi_{\square}^2(\delta) \in (0, 1)$.

We conclude this section by discussing how high should δ be, for the profile $(\sigma_{\delta}, \tau_{\delta})$ to be well-defined, and by discussing beliefs and actions off-equilibrium.

We first argue that the costs $c_1(h)$ and $c_2(h)$ are small.

Lemma 6 There is $c > 0$ such that for every $\delta \geq \overline{\delta}_1$ and every $h \in H_N$, one has

$$
c_1(h) \le (1 - \delta)c.
$$

Proof. Because payoffs are bounded by 1, one has

$$
c_1(h) \le (K_1 + 2K_2)(1 - \delta)\delta^{-N} = (K_1 + 2K_2) \frac{(1 - \delta)}{\delta^{K_1 + 2K_2}} \delta^{-\lfloor \frac{\ln(1 - \theta)}{\ln \delta} \rfloor}
$$

$$
\le (K_1 + 2K_2) \frac{(1 - \delta)}{\delta^{K_1 + 2K_2 + 1}} \delta^{\frac{\ln(1 - \theta)}{\ln \delta}} = (K_1 + 2K_2) \frac{(1 - \delta)}{\delta^{K_1 + 2K_2 + 1}} \frac{1}{\ln(1 - \theta)},
$$

³⁰We here abuse notation, since $N \to +\infty$ as $\delta \to 1$. However, the limit of $\mathbf{E}[u_{\star}(\mathbf{p}_{N})]$ is welldefined.

³¹Because the approximation of x^* by $x_{\delta}(\vec{a}(x^*))$ becomes perfectly accurate as $\delta \to 1$.

and the result follows.

For $\delta \leq 1$ (including $\delta = 1$), denote by $\mathcal{P}(\delta)$ the support of \mathbf{p}_N when ψ^2_{\square} is set to $\psi_{\Box}^2(\delta)$, and define $\mathcal{Q}(\delta)$ in a symmetric way. Since π_2 and $\varepsilon_2(\cdot|m_S)$ have full support, and since $\psi^2, \psi^2_{\square}(1) \in (0, 1)$, one has $\mathbf{p}_N \in \overset{\circ}{\Delta}$ † $_{\mathbf{I}_S}(S \times M_S)$, with probability 1.

Because $\mathcal{P}(1)$ and $\mathcal{Q}(1)$ are finite sets, and by Proposition [1,](#page-26-2) there is $\delta_2 < 1$, $\varepsilon > 0$, and neighborhoods $V(p)$ of $p \in \mathcal{P}$, $V(q)$ of $q \in \mathcal{Q}$, such that any payoff in $[u_*(p'), u_*(p') + \varepsilon] \times [v_*(q'), v_*(q') + \varepsilon]$ is a sequential equilibrium of $\Gamma(p', q')$, for every $p \in \mathcal{P}, p' \in V(p)$, and $q \in \mathcal{Q}, q' \in V(q)$.

In addition, we choose the neighborhoods $V(p)$, $V(q)$ to be small enough, and $C > 0$ so that the conclusion of Proposition [2](#page-27-0) holds for every $p \in \mathcal{P}, p' \in V(p)$, and $q \in \mathcal{Q}, q' \in V(q).$

We choose $\bar{\delta}_3$ < 1 to be high enough so that the following conditions are met for each $\delta \geq \overline{\delta}_3$: (i) every $p' \in \mathcal{P}(\delta)$ belongs to $V(p)$ for some $p \in \mathcal{P}$; (ii) $(1 - \delta)c \leq \varepsilon$.

For $\delta \geq \bar{\delta}_3$, the profile $(\sigma_{\delta}, \tau_{\delta})$ is then well-defined, at any information set that is not ruled out by the definition of $(\sigma_{\delta}, \tau_{\delta})$ at earlier stages.

Consider now an information set $I_{l,h}^1$ that is reached with probability 0, and assume that the information set $I_{l,h'}^1$ is reached with positive probability, where h' is the longest prefix of h.

If the sequence h of actions has probability zero, then we let beliefs at $I_{l,h}^1$ and at all subsequent information sets coincide with the belief held at $I^1_{l,h'}$. Player 1 repeats the action that is optimal at $I^1_{l,h'}$.

Assume now that the sequence h has positive probability. This corresponds to the case where player 1 misreported in Phase 1, and played consistently with his report afterwards. Then the belief of player 1 at $I_{l,h}^1$ is well-defined by Bayes' rule (and is independent of player 1's strategy), and only assigns a positive probability to information sets $I_{m,h}^2$ that are reached with positive probability under τ_{\star} . We let σ_{\star} play at $I_{l,h}^1$ a best reply to τ_{\star} .

By construction, sequential rationality holds at any information set $I_{l,h}^1$ that is reached with probability zero. One can verify that beliefs are consistent with $(\sigma_{\star}, \tau_{\star})$. We omit the proof.

B.4 Equilibrium properties

We claim that the profile $(\sigma_{\delta}, \tau_{\delta})$ is a sequential equilibrium profile for $\delta < 1$ high enough.

Let $\eta > 0$ be small enough so that

$$
\mathbf{E}_{\varepsilon_2,x^{\star}}[l_S \to k] < \mathbf{E}_{\varepsilon_2,x^{\star}}[l_S \to l_S] - 2\eta \text{ for every } l_S, k \in L_S, l_S \neq k,
$$

and we choose δ_4 < 1 such that

 $\mathbf{E}_{\varepsilon_2,x_\delta(\vec{a}(x^*))}[l_S \to k] < \mathbf{E}_{\varepsilon_2,x_\delta(\vec{a}(x^*))}[l_S \to l_S] - \eta$ for every $l_S, k \in L_S, l_S \neq k$ and $\delta \geq \overline{\delta}_4$.

We finally choose $\delta_5 < 1$ to be such that $1 - \delta^{K_1+2K_2} + (1 - \delta)C < \eta \delta^{K_1+2K_2}$ for each $\delta \geq \delta_5$.

We now verify that $(\sigma_{\star}, \tau_{\star})$ is a sequential equilibrium, as soon as $\delta \geq \max{\{\bar{\delta}_4, \bar{\delta}_5\}}$. It is sufficient to check that sequential rationality holds at any information set that is reached with positive probability. Let such an information set $I_{l,h}$ be given, and let n be the stage to which $I_{l,h}$ belongs. If stage n belongs to phase 4, then sequential rationality at $I_{l,h}$ follows because continuation strategies in phase 4 form a sequential equilibrium of the associated self-ignorant game. Assume then that $n < N$.

We will make use of the following observation that holds because $\varepsilon_1(\cdot)$, $\varepsilon_2(\cdot)$, π^1 and π^2 have full support: if $I_{ls,lT,h}$ is reached with positive probability, then the set of actions that are played with positive probability at $I_{l_S,l_T,h}$ does not depend on l_T , and, therefore, the information set $I_{l_S,l'_T,h}$ is also reached with positive probability, for every $l'_{S} \in L_{S}$. We note that the compensation made in phase 4 implies that player 1 is indifferent at $I_{ls,lT,h}$ between all actions that are played with positive probability. One thus simply needs to check that player 1 cannot increase his continuation payoff by playing some other action, a.

Assume first that n belongs to either phase 2.1, 2.2 or to phase 3. In that case, the set of actions that are played at $I_{l,h}$ does not depend on l. Hence, when playing a, player 1 triggers a myopic play by player 2, and player 1's overall payoff in that case does not exceed

$$
(1 - \delta)u_{\star}(p_n) + \delta \mathbf{E}[u_{\star}(\mathbf{p}_{n+1})|l, h].
$$

On the other hand, the expected continuation payoff of player 1 at $I_{l,h}$ is at least $\delta^N \mathbf{E}[u_{\star}(\mathbf{p}_N)|l, h]$. Sequential rationality then follow from the choice of parameters.

Assume finally that stage n belongs to phase 1. Again, it is not profitable to switch to an action that triggers a myopic play from player 2. What if player 1, instead of reporting l_S , chooses to report $k \neq l_S$? Then, as above, the choice of parameters ensures that it is optimal for player 1 to play consistently with k , at least until phase 4. Such a deviation yields a payoff (discounted back to h) of at most

$$
\delta^{-n} \left(\delta^{K_1 + K_2} \mathbf{E}_{\varepsilon_2, x_{\delta}(\vec{a}(x^*))} [l_S \to k] + (1 - \delta) \left(1 + \cdots + \delta^{K_1 + 2K_2 - 1} \right) + \delta^N \mathbf{E} [u_\star(\mathbf{p}_N) + (1 - \delta)C] \right).
$$

On the other hand, player 1's continuation payoff when reporting truthfully is at least

$$
\delta^{-n} \left(\delta^{K_1 + K_2} \mathbf{E}_{\varepsilon_2, x_\delta(\vec{a}(x^*))} [l_S \to l_S] + \delta^N \mathbf{E} [u_\star(\mathbf{p}_N) + (1 - \delta)C] \right).
$$

We stress that the distribution of \mathbf{p}_N is the same in both expressions, because the distribution of (μ_1, μ_2) does not depend on player 1's report. The result follows, by the choice of $\bar{\delta}_4$ and $\bar{\delta}_5$.