

Tail behavior of stationary solutions of random difference equations: the case of regular matrices

Gerold Alsmeyer^{†*} and Sebastian Mentemeier[†]

Institut für Mathematische Statistik, Einsteinstr. 62, 48149 Münster, DE

September 5, 2018

Given a sequence $(M_n, Q_n)_{n \geq 1}$ of i.i.d. random variables with generic copy (M, Q) such that M is a regular $d \times d$ matrix and Q takes values in \mathbb{R}^d , we consider the random difference equation (RDE) $R_n = M_n R_{n-1} + Q_n$, $n \geq 1$. Under suitable assumptions stated below, this equation has a unique stationary solution R such that, for some $\kappa > 0$ and some finite positive and continuous function K on $S^{d-1} := \{x \in \mathbb{R}^d : |x| = 1\}$,

$$\lim_{t \rightarrow \infty} t^\kappa \mathbb{P}(xR > t) = K(x) \quad \text{for all } x \in S^{d-1}$$

holds true. A rather long proof of this result, originally stated by Kesten at the end of his famous article [12], was given by LePage [15]. The purpose of this article is to show how regeneration methods can be used to provide a much shorter argument (in particular for the positivity of K). It is based on a multidimensional extension of Goldie's implicit renewal theory developed in [9].

Keywords: Markov renewal theory; implicit renewal theory, Harris recurrence, regeneration, random operators and equations; stochastic difference equations; random dynamical systems

AMS 2010 Subject Classification 60K05, 60H25, 39A50, 37H10

1 Introduction

Let $(M_n, Q_n)_{n \geq 1}$ be a sequence of i.i.d. random variables with generic copy (M, Q) such that M is a real $d \times d$ matrix and Q takes values in \mathbb{R}^d . Suppose further that

$$\mathbb{E} \log^+ \|M\| < \infty \tag{A1}$$

*Corresponding author. Email: gerolda@math.uni-muenster.de

[†]Research supported by the Deutsche Forschungsgemeinschaft (SFB 878)

where $\|M\| := \sup_{|x|=1} |xM|$. Then, with $\Pi_n := M_1 \cdot \dots \cdot M_n$, there exists $\beta \in [-\infty, \infty)$ such that

$$\beta := \lim_{n \rightarrow \infty} n^{-1} \log \|\Pi_n\| \quad \mathbb{P}\text{-a.s.}$$

and defines the Liapunov exponent of the RDE

$$R_n = M_n R_{n-1} + Q_n, \quad n \geq 1. \quad (1)$$

If β is negative and

$$\mathbb{E} \log^+ |Q| < \infty, \quad (\text{A2})$$

then this recursive Markov chain has a unique stationary distribution which is given by the law of the almost surely convergent series

$$R := \sum_{n \geq 1} \Pi_{n-1} Q_n \quad (2)$$

and is also characterized as the unique solution to the stochastic fixed-point equation (SFPE)

$$Y \stackrel{d}{=} MY + Q \quad (3)$$

where $\stackrel{d}{=}$ means equality in law and where Y is understood to be independent of (M, Q) . This by now standard result may easily be deduced from a more general one for iterations of random Lipschitz maps, see e.g. [8] or [6]. Our concern here is the tail behavior of R in the case when M takes almost surely values in $GL(d, \mathbb{R})$, the group of regular $d \times d$ matrices with real entries.

For $x \in \mathbb{R}^d \setminus \{0\}$, we write x^\sim for its projection on the unit sphere $S := S^{d-1}$, thus $x^\sim := |x|^{-1}x$. Lebesgue measure on the space of real $d \times d$ -matrices, seen as \mathbb{R}^{d^2} , is denoted as \mathfrak{A} and the uniform distribution on S as \mathfrak{A}_S . Finally, the open δ -balls in S and $GL(d, \mathbb{R})$ with centers x and A are denoted as $B_\delta(x)$ and $B_\delta(A)$, respectively.

Theorem 1.1. *Consider the RDE (1) and suppose that, in addition to (A1), (A2) and $\beta < 0$, the following assumptions hold:*

$$\mathbb{P}(M \in GL(d, \mathbb{R})) = 1. \quad (\text{A3})$$

$$\max_{n \geq 1} \mathbb{P}((x\Pi_n)^\sim \in U) > 0 \text{ for any } x \in S \text{ and any open } \emptyset \neq U \subset S. \quad (\text{A4})$$

$$\mathbb{P}(\Pi_{n_0} \in \cdot) \geq \gamma_0 \mathbf{1}_{B_c(\Gamma_0)} \mathfrak{A} \text{ for some } \Gamma_0 \in GL(d, \mathbb{R}), n_0 \in \mathbb{N} \text{ and } c, \gamma_0 > 0. \quad (\text{A5})$$

$$\mathbb{P}(Mv + Q = v) < 1 \text{ for any column vector } v \in \mathbb{R}^d. \quad (\text{A6})$$

There exists $\kappa_0 > 0$ such that

$$\mathbb{E} \inf_{x \in S} |xM|^{\kappa_0} \geq 1, \quad \mathbb{E} \|M\|^{\kappa_0} \log^+ \|M\| < \infty \text{ and } 0 < \mathbb{E} |Q|^{\kappa_0} < \infty. \quad (\text{A7})$$

Then there exists a unique $\kappa \in (0, \kappa_0]$ such that

$$\lim_{n \rightarrow \infty} n^{-1} \log \mathbb{E} \|\Pi_n\|^\kappa = 0, \quad (4)$$

and

$$\lim_{t \rightarrow \infty} t^\kappa \mathbb{P}(xR > t) = K(x) \quad \text{for all } x \in S, \quad (5)$$

where K is a finite positive and continuous function on S .

Remark 1. This result (with one extra condition) was stated by Kesten at the end of his famous article [12] and later proved by Le Page [15] with the help of Kesten’s Markov renewal theorem [13] (and without assuming (A5)). Markov renewal theory also plays an essential role in our approach, but we make use of a different Markov renewal theorem taken from [1] to show how proofs can be shortened considerably using Harris recurrence, which is the primary intention of this article. Condition (A5) plays a crucial role in obtaining (5) for all $x \in S$. Not assumed by LePage, he instead imposes the extra condition

$$\mathbb{E}|Q|^{\kappa_0+\varepsilon} < \infty \quad \text{for some } \varepsilon > 0$$

to arrive at the same conclusion. A similar result was also derived by Klüppelberg and Pergamenchtchikov [14] for a more specialized model. As further references, we mention related work by de Saporta et al. [5], by Guivarc’h [10] and, most recently, by Buraczewski et al. [4] who obtain more precise information on the tails of R under the restriction that M is a similarity (product of a dilation and an orthogonal transformation).

Remark 2. The most interesting ingredient to our approach may be roughly described as a suitable combination of Goldie’s implicit renewal theory [9], lifted to the multidimensional situation, with the technique of sampling along ”nice” regeneration epochs for the considered RDE (see Sections 6, 7 and 8).

Remark 3. Assumption (A6) is a condition on the dependence of M and Q (there is no need for independence), and asserts particularly that no Dirac measure solves the RDE. In fact our assumptions assure a priori, that $\text{supp } R$ is unbounded in \mathbb{R}^d (see Lemma 8.1).

Remark 4. Note that condition $\mathbb{E} \inf_{x \in S} |xM|^{\kappa_0} \geq 1$ in (A7) may be restated as

$$\mathbb{E} \lambda_d(MM^\top)^{\kappa_0/2} \geq 1,$$

where $\lambda_d(MM^\top)$ denotes the smallest of the d eigenvalues of the symmetric matrix MM^\top . This follows because $|xM| = (xMM^\top x^\top)^{1/2}$.

The further organization is as follows: Section 2 discusses the two central assumptions (A4) and (A5) in terms of their implications for obtaining Harris recurrence of an intrinsic Markov chain $(X_n)_{n \geq 0}$ on the sphere (see Section 5). We then proceed in Section 3 with some useful results concerning a whole class of SFPE that are solved by R and obtained via the use of stopping times. In particular, we explain how geometric sampling allows us to simplify some assumptions in Theorem 1.1 before proving it. Section 4 collects some facts about Harris recurrence and Markov renewal theory which are used in Section 5 to show that $\lim_{t \rightarrow \infty} t^{-\kappa} \mathbb{P}(\sup_{n \in \mathbb{N}} |xM_1 \cdots M_n| > t)$ exists and is positive. This section further contains all necessary ingredients for the Markov renewal approach including the crucial measure change (harmonic transform) also used by Kesten. The proof of Theorem 1.1 is then provided in Sections 6, 7 and 8.

2 Minorization: Implications of (A4) and (A5)

It is useful to discuss at this early point the implications of the two conditions (A4) and (A5) in terms of the semigroup $(P^n)_{n \geq 1}$ of Markov transition kernels on S , defined by $P^n(x, A) :=$

$\mathbb{P}((x\Pi_n)^\sim \in A)$ for $x \in S$ and measurable $A \subset S$. The pertinent Markov chain being of interest here will be introduced in Section 5. For compact subsets C of $GL(d, \mathbb{R})$, we further define the substochastic kernels $P_C^n(x, \cdot) := \mathbb{P}((x\Pi_n)^\sim \in \cdot, \Pi_n \in C)$. Let I denote the identity matrix.

Lemma 2.1. *Suppose (A4) and (A5). Then there exists a compact subset C of $GL(d, \mathbb{R})$ such that, for each $x \in S$, there are $\delta, p > 0$, $m \in \mathbb{N}$, and a probability measure ϕ on $B_\delta(x)$ satisfying*

$$P^m(y, \cdot) \geq P_C^m(y, \cdot) \geq p\phi \quad (\text{MC})$$

for all $y \in S$.

Proof. Fix any $x \in S$. By (A4) and (A5), we can choose $n_1 \geq 1$, $\eta > 0$ and thereupon $0 < \delta < \eta$, $0 < \varsigma < c$ and a compact $B_1 \subset GL(d, \mathbb{R})$ in such a way that

- (i) $\zeta := \inf_{y \in B_\delta(x)} P_{B_1}^{n_1}(y, U) > 0$, where $U := B_\eta((x\Gamma_0^{-1})^\sim)$;
- (ii) $\Phi(A) := \int_{B_\varsigma(I)} \mathbf{1}_A((x\mathbf{m})^\sim) \mathfrak{L}(d\mathbf{m})$ defines nonzero measure on $B_\delta(x)$;
- (iii) for any $u \in U$, we have $u = (xF_u\Gamma_0^{-1})^\sim$ as well as $F_u\Gamma_0^{-1}B_c(\Gamma_0) \supset B_\varsigma(I)$ for some unitary matrix F_u .

Put $C := B_1 \cdot \overline{B_c(\Gamma_0)} := \{\Lambda_1\Lambda_2 : \Lambda_1 \in B_1, \Lambda_2 \in \overline{B_c(\Gamma_0)}\}$, which is a compact subset of $GL(d, \mathbb{R})$ (as the continuous image of the compact $B_1 \times \overline{B_c(\Gamma_0)}$). It then follows for any $y \in B_\delta(x)$ and measurable $A \subset S$ that

$$\begin{aligned} P_C^{n_0+n_1}(y, A) &\geq \int_U \int_{B_c(\Gamma_0)} \mathbf{1}_A((um)^\sim) \mathbb{P}(\Pi_{n_0} \in d\mathbf{m}) P_{B_1}^{n_1}(y, du) \\ &\geq \gamma_0 \int_U \int_{B_c(\Gamma_0)} \mathbf{1}_A((um)^\sim) \mathfrak{L}(d\mathbf{m}) P_{B_1}^{n_1}(y, du) \\ &= \gamma_0 \int_U \int_{B_c(\Gamma_0)} \mathbf{1}_A((xF_u\Gamma_0^{-1}\mathbf{m})^\sim) \mathfrak{L}(d\mathbf{m}) P_B^{n_1}(y, du) \\ &\geq \gamma_0 |\det(\Gamma_0)|^d P_{B_1}^{n_1}(y, U) \int_{B_\varsigma(I)} \mathbf{1}_A((x\mathbf{m})^\sim) \mathfrak{L}(d\mathbf{m}) \\ &\geq \gamma_0 \zeta |\det(\Gamma_0)|^d \int_{B_\varsigma(I)} \mathbf{1}_A((x\mathbf{m})^\sim) \mathfrak{L}(d\mathbf{m}) \end{aligned} \quad (6)$$

which proves (MC) for all $y \in B_\delta(x)$ with $m = n_0 + n_1$ and $\phi := \Phi(B_\delta(x))^{-1}\Phi$.

In order to extend (MC) to all $y \in S$, observe that for any y , we can pick $\varepsilon(y) > 0$, $n_2(y) \geq 1$ and compact $B^y \subset GL(d, \mathbb{R})$ such that

$$\inf_{z \in B_{\varepsilon(y)}(y)} P_{B^y}^{n_2(y)}(z, B_\delta(x)) > 0.$$

By compactness, $S = \bigcup_{i=1}^k B_{\varepsilon(y_i)}(y_i)$ for suitable y_1, \dots, y_k , and a straightforward argument then shows that $\inf_{y \in S} P_{B_2}^{n_2}(y, B_\delta(x)) > 0$ for a suitable $n_2 \geq \max_{i=1, \dots, k} n_2(y_i)$ and with $B_2 := \bigcup_{i=1}^k B^{y_i}$. It is now readily seen with the help of property (i) that

- (iv) $\xi := \inf_{y \in S} P_B^{n_1+n_2}(y, U) > 0$, where $B := B_1 \cdot B_2$.

By estimating $P_C^{n_0+n_1+n_2}$ with $C := B \cdot \overline{B_c(\Gamma_0)}$ following (6) and utilizing (iv) instead of (i), we finally obtain (MC) for all $y \in S$ (with the same ϕ and $m = n_0 + n_1 + n_2$). Further details can be omitted. \square

Remark 5. It is useful for later purposes (see Lemma 5.6) to point out that (MC) is "embedded" in a bivariate condition, obtained via consideration of the bivariate extensions $H^n(x, \cdot) := \mathbb{P}((x\Pi_n)^\sim, \Pi_n) \in \cdot$ and $H_C^n(x, \cdot) := \mathbb{P}((x\Pi_n)^\sim, \Pi_n) \in \cdot, \Pi_n \in C$ of $P^n(x, \cdot)$ and $P_C^n(x, \cdot)$, respectively.

Keeping notation and settings from above and with $\gamma_1 := |\det(\Gamma_0)|^{-d} \max_{\mathbf{m} \in B} |\det(\mathbf{m})|^d$ finite, a similar estimation as in (6) leads to

$$\begin{aligned} & H_C^{n_0+n_1+n_2}(y, D) \\ & \geq \gamma_0 \int_{U \times B} \int_{B_c(\Gamma_0)} \mathbf{1}_D((u\mathbf{m}_0)^\sim, \mathbf{m}_1\mathbf{m}_0) \boldsymbol{\lambda}(d\mathbf{m}_0) H_B^{n_1+n_2}(y, d(u, \mathbf{m}_1)) \\ & = \gamma_0 \int_{U \times B} \int_{B_c(\Gamma_0)} \mathbf{1}_D((xF_u\Gamma_0^{-1}\mathbf{m}_0)^\sim, \mathbf{m}_1\mathbf{m}_0) \boldsymbol{\lambda}(d\mathbf{m}_0) H_B^{n_1+n_2}(y, d(u, \mathbf{m}_1)) \\ & \geq \frac{\gamma_0}{\gamma_1} \int_{U \times B} \int_{B_\zeta(I)} \mathbf{1}_D((x\mathbf{m}_0)^\sim, \mathbf{m}_1\Gamma_0 F_u^{-1}\mathbf{m}_0) \boldsymbol{\lambda}(d\mathbf{m}_0) H_B^{n_1+n_2}(y, d(u, \mathbf{m}_1)) \end{aligned}$$

and thus to the bivariate minorization condition

$$H_C^{n_0+n_1+n_2}(y, \cdot) \geq q \psi(y, \cdot) \quad (\text{BMC})$$

for all $y \in S$, some $q > 0$ and a probability kernel $\psi(y, \cdot)$ on $S \times C$. It contains (MC) as a special case, for $P_C^{n_0+n_1+n_2}(y, \cdot) = H^{n_0+n_1+n_2}(y, \cdot \times C)$ and $\psi(\cdot \times C) = \phi$.

3 The stopped RDE and geometric sampling

Geometric sampling and, more generally, the use of stopping times for $(M_n, Q_n)_{n \geq 1}$ provides a useful technique in our subsequent analysis and is thus briefly discussed next.

3.1 R remains solution to the stopped equation

Let $(\mathcal{G}_n)_{n \geq 0}$ be a filtration such that $(M_n, Q_n)_{n \geq 1}$ is adapted to it and $(M_k, Q_k)_{k > n}$ is independent of \mathcal{G}_n for any $n \geq 0$. Consider any a.s. finite stopping time τ with respect to $(\mathcal{G}_n)_{n \geq 0}$ which, by suitable choice of the latter, includes the case that τ and $(M_n, Q_n)_{n \geq 1}$ are independent (pure randomization). Then it is readily checked that R defined in (2) satisfies

$$R = \Pi_\tau R^\tau + Q^\tau \quad (7)$$

where

$$Q^n := \sum_{k=1}^n \Pi_{k-1} Q_k \quad \text{and} \quad R^n := \sum_{k > n} \left(\prod_{j=n+1}^{k-1} M_j \right) Q_k$$

for $n \geq 1$. But since $(M_{\tau+n}, Q_{\tau+n})_{n \geq 1}$ is a copy of $(M_n, Q_n)_{n \geq 1}$ and also independent of $(M_n, Q_n)_{1 \leq n \leq \tau}$ and τ , it follows that R^τ is independent of (Π_τ, Q^τ) with $R^\tau \stackrel{d}{=} R$. In other words, (the law of) R also solves the stopped SFPE

$$Y \stackrel{d}{=} \Pi_\tau Y + Q^\tau \quad (8)$$

and provides a stationary distribution to the RDE

$$Y_n = M'_n Y_{n-1} + Q'_n, \quad n \geq 1, \quad (9)$$

where $(M'_n, Q'_n)_{n \geq 1}$ is a sequence of i.i.d. copies of (Π_τ, Q^τ) . Uniqueness follows if (A1), (A2) persist to hold for the "stopped pair" (Π_τ, Q^τ) together with

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \|\Pi_{\sigma_n}\| < 0 \quad \mathbb{P}\text{-a.s.}$$

where $(\sigma_n)_{n \geq 0}$ denotes a zero-delayed renewal process such that $\sigma_1 = \tau$ and

$$(\sigma_n - \sigma_{n-1}, (M_k, Q_k)_{\sigma_{n-1} < k \leq \sigma_n}), \quad n \geq 1$$

are i.i.d. For stopping times τ with finite mean this is indeed easily verified and we state the result (without proof) in the following lemma.

Lemma 3.1. *The law of R forms the unique solution to the SFPE (8) whenever $\mathbb{E} \tau < \infty$.*

In order for finding the tail behavior of R , we are now allowed to do so within the framework of any stopped SFPE (8) with finite mean τ . The idea is to pick τ in such a way that (Π_τ, Q^τ) has nice additional properties compared to (M, Q) . Geometric sampling provides a typical example that will be used hereafter and therefore discussed next. Another use of this technique appears in Section 8.

3.2 Geometric sampling

Suppose now that $(\sigma_n)_{n \geq 0}$ is independent of $(M_n, Q_n)_{n \geq 1}$ with geometric(1/2) increments, that is $\mathbb{P}(\tau = n) = 1/2^n$ for each $n \geq 1$. Then not only Lemma 3.1 holds true but also the following result:

Lemma 3.2. *If (M, Q) satisfies the assumption of Theorem 1.1 and thus (MC), then so does (Π_τ, Q^τ) with $n = n_0 = m = 1$ in (A4), (A5) and (MC). Also, $\lim_{n \rightarrow \infty} n^{-1} \log \mathbb{E} \|\Pi_{\sigma_n}\|^\kappa = 0$ implies $\lim_{n \rightarrow \infty} n^{-1} \log \mathbb{E} \|\Pi_n\|^\kappa = 0$, i.e. (4).*

Proof. That (A1), (A2) and $\lim_{n \rightarrow \infty} n^{-1} \log \|\Pi_{\sigma_n}\| < 0$ \mathbb{P} -a.s. persist to hold under any finite mean stopping time τ has already been pointed out before Lemma 3.1. As for (A3) to (A5), we just note that $\mathbb{P}(\Pi_\tau \in \cdot) = \sum_{k \geq 1} 2^{-k} \mathbb{P}(\Pi_k \in \cdot)$. Assumption (A6) ensures that the law of R is nondegenerate. But since R is also the unique solution to (8), (A6) must hold for (Π_τ, Q^τ) as well. Moreover,

$$\mathbb{E} \inf_{x \in S} |x \Pi_\tau| = \sum_{n \geq 1} 2^{-n} \mathbb{E} \inf_{x \in S} |x \Pi_n|$$

in combination with

$$\begin{aligned}
\mathbb{E} \inf_{x \in S} |x \Pi_n|^{\kappa_0} &= \mathbb{E} \left(\inf_{x \in S} |(x M_1 \cdot \dots \cdot M_{n-1})^\sim M_n|^{\kappa_0} \cdot |x M_1 \cdot \dots \cdot M_{n-1}|^{\kappa_0} \right) \\
&\geq \mathbb{E} \left(\inf_{x \in S} |x M_n|^{\kappa_0} \cdot \inf_{x \in S} |x \Pi_{n-1}|^{\kappa_0} \right) \\
&= \mathbb{E} \inf_{x \in S} |x M_n|^{\kappa_0} \mathbb{E} \inf_{x \in S} |x \Pi_{n-1}|^{\kappa_0} \\
&= \dots = \left(\mathbb{E} \inf_{x \in S} |x M|^{\kappa_0} \right)^n \geq 1
\end{aligned} \tag{10}$$

for each $n \geq 1$ shows the first assertion of (A7) for (Π_τ, Q^τ) . The remaining two moment assertions are again easily verified by standard estimates. We therefore omit further details. Finally, suppose that $\lim_{n \rightarrow \infty} n^{-1} \log \mathbb{E} \|\Pi_{\sigma_n}\|^\kappa = 0$. By subadditivity, $\xi := \lim_{n \rightarrow \infty} n^{-1} \log \mathbb{E} \|\Pi_n\|^\kappa = \inf_{n \geq 1} n^{-1} \log \mathbb{E} \|\Pi_n\|^\kappa$ exists in $[-\infty, \infty)$. Since

$$\frac{1}{n} \log \mathbb{E} \|\Pi_{\sigma_n}\|^\kappa = \frac{1}{n} \log \sum_{k \geq n} \mathbb{E} \|\Pi_k\|^\kappa \mathbb{P}(\sigma_n = k) \geq \frac{1}{n} \sum_{k \geq n} \mathbb{P}(\sigma_n = k) \log \mathbb{E} \|\Pi_k\|^\kappa$$

it is not difficult to see that $\xi > -\infty$. But then we further infer for any $\varepsilon > 0$ and all sufficiently large n that

$$\frac{1}{n} \log \mathbb{E} \|\Pi_{\sigma_n}\|^\kappa \geq \frac{1}{n} \log \sum_{k \geq n} e^{k(\xi - \varepsilon)} \mathbb{P}(\sigma_n = k) = \log \mathbb{E} e^{(\xi - \varepsilon)\tau}$$

and thus $\xi \leq 0$ upon taking $n \rightarrow \infty$ and then $\varepsilon \rightarrow 0$. By doing the same in the reverse inequality

$$\frac{1}{n} \log \mathbb{E} \|\Pi_{\sigma_n}\|^\kappa \leq \frac{1}{n} \log \sum_{k \geq n} e^{k(\xi + \varepsilon)} \mathbb{P}(\sigma_n = k) = \log \mathbb{E} e^{(\xi + \varepsilon)\tau}$$

finally shows $\xi = 0$ as claimed in (4). □

Remark 6. We note for later purposes that for all $\varepsilon, \delta > 0$, $x \in S$

$$\mathbb{P}(\|\Pi_\tau\| < \varepsilon, (x \Pi_\tau)^\sim \in B_\delta(x)) > 0,$$

because both, $\{\limsup_{n \rightarrow \infty} \|\Pi_n\| < \varepsilon\}$ and $\{(x \Pi_n)^\sim \in B_\delta(x) \text{ i.o.}\}$, are sets of probability one.

In view of the previous lemma we can now make the **standing assumption** that

$$\text{If (A4),(A5),(MC) and (BMC) hold, they hold with } n_0 = n = m = 1. \tag{SA}$$

4 Harris recurrence and Markov renewal theory

4.1 Strongly aperiodic Harris chains

Here and in the following subsection let S be a general separable metric space with Borel- σ -algebra \mathcal{S} . A Markov chain $(X_n)_{n \geq 0}$ on S is called *strongly aperiodic Harris chain*, if there

exists a set $\mathfrak{R} \in \mathcal{S}$, called *regeneration set*, such that $\mathbb{P}_x(X_n \in \mathfrak{R} \text{ infinitely often}) = 1$ for all $x \in S$ (recurrence) and, furthermore,

$$\inf_{x \in \mathfrak{R}} P(x, \cdot) \geq p \phi \quad (11)$$

for some $p > 0$, $r \in \mathbb{N}$ and a probability measure ϕ with $\phi(\mathfrak{R}) = 1$. Strong aperiodicity refers to the fact that P and not P^m for some $m \geq 2$ satisfies (11). If S itself is regenerative then $(X_n)_{n \geq 0}$ is called *Doebelin chain*. A strongly aperiodic Harris chain $(X_n)_{n \geq 0}$ possesses a nice regenerative structure as shown by the following regeneration lemma due to Athreya and Ney [2].

Lemma 4.1. *On a possibly enlarged probability space, one can redefine $(X_n)_{n \geq 0}$ together with an increasing sequence $(\sigma_n)_{n \geq 0}$ of random epochs such that the following conditions are fulfilled under any \mathbb{P}_x , $x \in S$:*

- (R1) *There is a filtration $\mathcal{G} = (\mathcal{G}_n)_{n \geq 0}$ such that $(X_n)_{n \geq 0}$ is Markov adapted and each σ_n a stopping time with respect to \mathcal{G} .*
- (R2) *$(\sigma_n - \sigma_1)_{n \geq 1}$ forms a zero-delayed renewal sequence with increment distribution $\mathbb{P}_\phi(\sigma_1 \in \cdot)$ and is independent of σ_1 .*
- (R3) *For each $k \geq 1$, the sequence $(X_{\sigma_k + n})_{n \geq 0}$ is independent of $(X_j)_{0 \leq j \leq \sigma_k - 1}$ with distribution $\mathbb{P}_\phi((X_n)_{n \geq 0} \in \cdot)$.*

The σ_n , called *regeneration epochs*, are obtained by the following coin-tossing procedure: If τ_n , $n \geq 1$, denote the successive return times of the chain to \mathfrak{R} , then at each such τ_n a p -coin is tossed. If head comes up, then $X_{\tau_n + 1}$ is generated according to ϕ , while it is generated according to $(1 - p)^{-1}(P(X_{\nu_n}, \cdot) - p\phi)$ otherwise. Hence, the $\sigma_n - 1$ are those return epochs at which the coin toss produces a head. More formally, this is realized by introducing i.i.d. Bernoulli(p) variables J_0, J_1, \dots with the following properties:

- (R4) *For each $n \geq 0$, J_n is independent of $\sigma((X_k)_{0 \leq k \leq n})$.*
- (R5) *$\sigma_0 := 0$ and $\sigma_n := \inf\{k > \sigma_{n-1} : X_{k-1} \in \mathfrak{R}, J_{k-1} = 1\}$ for $n \geq 1$.*

Note that $(X_n, J_n)_{n \geq 0}$, called *split chain* (see [16]), is also a strongly aperiodic Harris chain with state space $S \times \{0, 1\}$. Naturally, it depends on the choice of the regeneration set \mathfrak{R} .

4.2 Markov renewal theory

Let $(X_n, U_n)_{n \geq 0}$ be a temporally homogeneous Markov chain on $S \times \mathbb{R}$ such that

$$\mathbb{P}((X_{n+1}, U_{n+1}) \in A \times B | X_n, U_n) = P(X_n, A \times B) \quad \text{a.s.}$$

for all $n \geq 0$ and a transition kernel P . Then the associated sequence $(X_n, V_n)_{n \geq 0}$ with $V_n = V_{n-1} + U_n$ for $n \geq 1$ is also a Markov chain and called *Markov Random Walk (MRW) with driving chain* $(X_n)_{n \geq 0}$. This extends the notion of classical random walk with i.i.d. increments because, conditioned on $(X_n)_{n \geq 0}$, the U_n are independent, but no longer identically distributed. In fact, the conditional distribution of U_n given $(X_k)_{k \geq 0}$ is of the form $Q((X_{n-1}, X_n), \cdot)$ for each $n \geq 1$ and a suitable stochastic kernel Q . The MRW is called *d-arithmetic*, if there exists a minimal $d > 0$ and a measurable function $\gamma : S \rightarrow [0, d)$ such that

$$\mathbb{P}(U_1 - \gamma(x) + \gamma(y) \in d\mathbb{Z} | X_0 = x, X_1 = y) = 1$$

for $P_\pi((X_0, X_1) \in \cdot)$ almost all $(x, y) \in S^2$, and *nonarithmetic* otherwise. As usual, for any distribution λ on S , \mathbb{P}_λ means $\mathbb{P}_\lambda((X_0, V_0) \in \cdot) = \lambda \otimes \delta_0$. The Markov renewal measure $\sum_{n \geq 0} \mathbb{P}_\lambda((X_n, V_n) \in \cdot)$ associated with the given MRW under \mathbb{P}_λ is denoted as \mathbb{U}_λ .

Being enough for our purposes, we focus hereafter on the case when the driving chain is a strongly aperiodic Harris chain on compact state space and thus having a unique stationary distribution, denoted as π .

Defining the first exit time $N(t) := \inf\{n \geq 0 : V_n > t\}$ consider the residual lifetime process $R_t := (V_{N(t)} - t)\mathbf{1}_{\{N(t) < \infty\}}$ and the jump process $Z(t) := X_{N(t)}\mathbf{1}_{\{N(t) < \infty\}}$. A measurable function $g : S \times \mathbb{R} \rightarrow \mathbb{R}$ is called *π -directly Riemann integrable* if

$$g(x, \cdot) \text{ is } \lambda\text{-a.e. continuous for } \pi\text{-almost all } x \in S \quad (12)$$

$$\text{and } \int_S \sum_{n \in \mathbb{Z}} \sup_{t \in [n\delta, (n+1)\delta]} |g(x, t)| \pi(dx) < \infty \text{ for some } \delta > 0, \quad (13)$$

where λ denotes Lebesgue measure on \mathbb{R} . The following Markov renewal theorem (MRT) is the main result of [1]:

Theorem 4.2. *Let $(X_n, V_n)_{n \geq 0}$ be a nonarithmetic MRW with strongly aperiodic Harris driving chain $(X_n)_{n \geq 0}$ with stationary distribution π . Let $\alpha := \mathbb{E}_\pi V_1 > 0$. If $g : S \times \mathbb{R} \rightarrow \mathbb{R}$ is π -directly Riemann integrable then, for π -almost all $x \in S$,*

$$g * \mathbb{U}_x(t) := \mathbb{E}_x \left(\sum_{n \geq 0} g(X_n, t - V_n) \right) \rightarrow \frac{1}{\alpha} \int_S \int_{\mathbb{R}} g(u, v) dv \pi(du). \quad (14)$$

as $t \rightarrow \infty$. Moreover, if $f : S \times (0, \infty) \rightarrow (0, \infty)$ is bounded and continuous, then

$$\lim_{t \rightarrow \infty} \mathbb{E}_x \left(f(Z(t), R(t)) \mathbf{1}_{\{N(t) < \infty\}} \right) = L(f) \quad (15)$$

for π -almost all $x \in S$ and some constant $L(f) > 0$.

Remark 7. The following extension of the above result follows directly upon inspection of the coupling proof given in [1, Section 7]: If ϕ is any minorizing distribution for the transition kernel of the Harris driving chain $(X_n)_{n \geq 0}$, then $g * \mathbb{U}_\phi(t)$ is a bounded function and converges to the limit given in (14). This fact will be used in Section 7.

Remark 8. Note that $(V_n)_{n \geq 0}$ satisfies the strong law of large numbers, viz.

$$\lim_{n \rightarrow \infty} \frac{V_n}{n} = \alpha \quad \mathbb{P}\text{-a.s.} \quad (16)$$

The number $\alpha = \mathbb{E}_\pi V_1$ is called the *drift* of $(X_n, V_n)_{n \geq 0}$.

5 Measure change and tail behaviour of $\sup_{n \geq 1} |x \Pi_n|$

Returning to the model described in the Introduction, we proceed with a short account of the ideas in [12] and [15]. Recall that $S = S^{d-1}$ and define

$$X_n := (X_0 \Pi_n)^\sim \quad \text{and} \quad U_n := \log |X_{n-1} M_n|.$$

Since $U_n = \log |X_0 \Pi_n| - \log |X_0 \Pi_{n-1}|$, the sequence $(X_n, V_n)_{n \geq 0}$ forms a MRW on $S \times \mathbb{R}$ with initial values (X_0, V_0) , and

$$\{N(t) < \infty\} = \left\{ \sup_{n \geq 1} \log |X_0 \Pi_n| > t \right\}.$$

Under the assumptions of Theorem 1.1 (with $n_0 = n = 1$ in (A4) and (A5)), $(X_n)_{n \geq 0}$ is easily seen to be a strongly aperiodic Harris chain with transition kernel $P = P^1$ defined in Section 2. However, $(X_n, V_n)_{n \geq 0}$ does not satisfy the conditions of the MRT, since by (16),

$$\alpha = \lim_{n \rightarrow \infty} \frac{\log |X_0 \Pi_n|}{n} \leq \lim_{n \rightarrow \infty} \frac{\log \|\Pi_n\|}{n} = \beta < 0 \quad \mathbb{P}\text{-a.s.}$$

A MRW with positive drift is indeed obtained after a change of measure (harmonic transform) for which it is crucial that $\mathbb{P}(\log \|M\| > 0) > 0$ which in turn follows from assumption (A7).

Theorem 5.1. *Under the assumptions of Theorem 1.1, there exist $\kappa \in (0, \kappa_0]$ and a positive continuous function $r : S \rightarrow (0, \infty)$ such that*

$$\begin{aligned} & {}^\kappa \mathbb{E}_x f(X_0, V_0, X_1, V_1, \dots, X_n, V_n) \\ & := \frac{1}{r(x)} \mathbb{E}_x \left(r(X_n) e^{\kappa V_n} f(X_0, V_0, X_1, V_1, \dots, X_n, V_n) \right), \end{aligned} \quad (17)$$

for all bounded continuous functions f and all $n \geq 0$, defines a distribution ${}^\kappa \mathbb{P}_x$ for each $x \in S$. Under ${}^\kappa \mathbb{P}_x$, $(X_n, V_n)_{n \geq 0}$ is a MRW with positive drift and satisfies the assumptions of Theorem 4.2. The constant κ is the unique value such that (4) holds true, i.e.

$$\lim_{n \rightarrow \infty} n^{-1} \log \mathbb{E} \|\Pi_n\|^\kappa = 0.$$

Note that, for each $x \in S$, \mathbb{P}_x and ${}^\kappa \mathbb{P}_x$ are equivalent probability measures on any $\sigma((X_k, V_k) : k \leq n)$, $n \geq 0$. We often write \mathbb{P} -a.s. and ${}^\kappa \mathbb{P}$ -a.s. as shorthand for \mathbb{P}_x -a.s. and ${}^\kappa \mathbb{P}_x$ -a.s. for all $x \in S$, respectively. Moreover probabilities under \mathbb{P} without subscript are always understood as being independent of the initial state and thus the same under any \mathbb{P}_x .

5.1 Proof of Theorem 5.1: Choice of κ and r

Defining the positive operators

$$T_\varkappa : \mathcal{C}(S) \rightarrow \mathcal{C}(S), \quad f(x) \mapsto \mathbb{E} \left(e^{\varkappa \log |xM|} f((xM)^\sim) \right), \quad \varkappa \in (0, \kappa_0],$$

we must find κ such that T_κ has maximal eigenvalue 1 with positive eigenfunction r . Due to our standing assumption, T_\varkappa is even strictly positive in the sense that $T_\varkappa f$ is everywhere positive whenever $0 \neq f \geq 0$. Indeed, for any such f , the set $U_f = \{f > 0\}$ is nonempty and open by continuity whence, using (A4) with $n = 1$, we infer

$$T_\varkappa f(x) \geq \int |y|^\varkappa \mathbf{1}_{U_f}(y^\sim) f(y^\sim) \mathbb{P}(xM \in dy) > 0$$

for all $x \in S$. The strict positivity will enable us to provide an elegant proof of the important Lemma 5.4 below.

Lemma 5.2. *Let $\varrho(\varkappa)$ be the spectral radius of T_\varkappa , i.e. $\varrho(\varkappa) = \lim_{n \rightarrow \infty} \|T_\varkappa^n\|^{1/n}$. Then T_\varkappa has an eigenvalue of maximal modulus equal to $\varrho(\varkappa)$.*

Proof. The adjoint operator $T_\varkappa^* : \mathcal{C}(S)^* \rightarrow \mathcal{C}(S)^*$, $\mathcal{C}(S)^*$ being the space of regular bounded signed measures on S , is weakly compact, i.e. it maps bounded sets to weakly sequentially compact sets. This follows by Prokhorov's theorem because

$$\|T_\varkappa^*\| = \|T_\varkappa\| \leq \mathbb{E} \|M\|^\varkappa < \infty$$

and S is compact. By [7, Theorem VI.4.8], T_\varkappa is then weakly compact as well, and by [7, Corollary VI.7.5], T_\varkappa^2 is compact. Hence, by [7, Lemmata VII.4.5 & 6], the spectrum of T_\varkappa is pure point (maybe except for 0) and T_\varkappa possesses an eigenvalue λ_\varkappa that is maximal in modulus, i.e. $|\lambda_\varkappa| = \varrho(\varkappa)$. \square

The following argument shows the existence of $\kappa \in (0, \kappa_0]$ with $\varrho(\kappa) = 1$: As one can readily verify by induction, $T_\varkappa^n f(x) = \mathbb{E} \left(e^{\varkappa \log |x \Pi_n|} f((x \Pi_n)^\sim) \right)$, and we infer $\varrho(\kappa_0) \geq 1$ upon choosing $f = \mathbf{1}_S$ and using (10). If $\varrho(\kappa_0) = 1$ we are done, so suppose that $\varrho(\kappa_0) > 1$ and thus $\|T_{\kappa_0}^n\| > 1$ for all sufficiently large n .

Since $\varkappa \mapsto \|T_\varkappa^n f\|$ is log-convex and thus continuous on $(0, \kappa_0)$ and lower semicontinuous on $(0, \kappa_0]$ for each $f \in \mathcal{C}(S)$ and $n \geq 1$ (use Hölder's inequality), the same holds true for $\varkappa \mapsto \|T_\varkappa^n\|$ as its pointwise supremum. It follows that $\|T_{\kappa_1}^n\| > 1$ for some $\kappa_1 \in (0, \kappa_0)$ and all sufficiently large n and therefore $\varrho(\kappa_1) \geq 1$. But we also have $\varrho(\kappa_2) < 1$ for some $\kappa_2 \in (0, \kappa_0)$ because $\beta < 0$ (and by the Furstenberg-Kesten theorem).

Finally, again as pointwise limit of the log-convex functions $\varkappa \mapsto \|T_\varkappa^n\|^{1/n}$, $\varrho(\varkappa)$ is log-convex and thus continuous on $(0, \kappa_0)$. Hence, $\varrho(\kappa) = 1$ for some unique $\kappa \in (0, \kappa_0)$. That κ also satisfies (4) follows from the following more general lemma.

Lemma 5.3. *For each $\varkappa \in (0, \kappa_0]$,*

$$\varrho(\varkappa) = \lim_{n \rightarrow \infty} (\mathbb{E} \|\Pi_n\|^\varkappa)^{1/n}.$$

Proof. Obviously,

$$\varrho(\varkappa) = \lim_{n \rightarrow \infty} \sup_{x \in S} (\mathbb{E} |x \Pi_n|^\varkappa)^{1/n} \leq \lim_{n \rightarrow \infty} \inf (\mathbb{E} \|\Pi_n\|^\varkappa)^{1/n}.$$

For the converse note that, by [3, Prop. 3.2], $Z_{x_0} := \inf_{n \geq 0} \|\Pi_n\|^{-1} |x_0 \Pi_n| > 0$ a.s. for any $x_0 \in S$, whence

$$\sup_{x \in S} \mathbb{E} |x \Pi_n|^\varkappa \geq \mathbb{E} \|\Pi_n\|^\varkappa \frac{\mathbb{E} |x_0 \Pi_n|^\varkappa}{\mathbb{E} \|\Pi_n\|^\varkappa} \geq \mathbb{E} \|\Pi_n\|^\varkappa \frac{\mathbb{E} Z_{x_0} \|\Pi_n\|^\varkappa}{\mathbb{E} \|\Pi_n\|^\varkappa}$$

and therefore (using Jensen's inequality)

$$\varrho(\varkappa) \geq \lim_{n \rightarrow \infty} \sup (\mathbb{E} \|\Pi_n\|^\varkappa)^{1/n} \lim_{n \rightarrow \infty} \frac{\mathbb{E} Z_{x_0}^{1/n} \|\Pi_n\|^\varkappa}{\mathbb{E} \|\Pi_n\|^\varkappa} = \lim_{n \rightarrow \infty} \sup (\mathbb{E} \|\Pi_n\|^\varkappa)^{1/n}.$$

which completes the proof. \square

Lemma 5.4. *Let $\varrho(\kappa) = 1$. Then T_κ has maximal eigenvalue 1 with one-dimensional eigenspace containing a positive eigenfunction r which further is symmetric, i.e. $r(x) = r(-x)$ for all $x \in S$.*

Proof. The following argument goes back to Karlin [11, Section 5] and hinges on the strict positivity of T_κ . By Lemma 5.2, T_κ has eigenvalue λ with $|\lambda| = 1$. Let f be a corresponding eigenfunction. Obviously, $T_\kappa f = \lambda f$ implies

$$T_\kappa |f| \geq |f|.$$

Suppose that $T_\kappa |f| - |f| \neq 0$. Then, by the strict positivity of T_κ , we have that $T_\kappa(T_\kappa |f| - |f|)$ is positive and thus $> \eta$ for some $\eta > 0$ chosen small such that, furthermore, $T_\kappa |f| < 1/\eta$ (S compact). From this we further infer

$$T_\kappa^2 |f| - T_\kappa |f| > \eta > \eta^2 T_\kappa |f|, \quad \text{hence } T_\kappa^2 |f| > (1 + \eta^2) T_\kappa |f|$$

and thereby $T_\kappa^n T_\kappa |f| > (1 + \eta^2)^n T_\kappa |f|$ for all $n \geq 1$ upon iteration. Consequently, $\|T_\kappa^n\| > (1 + \eta^2)^n$ for all $n \geq 1$ and thus $\varrho(\kappa) > 1$, a contradiction that leads to the conclusion that $T_\kappa |f| = |f|$ and thus that $r := |f|$ is a positive eigenfunction for the eigenvalue 1.

Now, suppose there is another eigenfunction g , linearly independent of r and w.l.o.g. real-valued (for, if g is an eigenfunction, then so are its real and imaginary parts if nontrivial). Pick ε such that $h := r + \varepsilon g$ is nonnegative, but $h(x) = 0$ for some x . By linear independence, h does not vanish everywhere. Since it is again an eigenfunction, the strict positivity of T_κ implies that it must be positive everywhere which is a contradiction. Hence r must be the unique eigenfunction modulo scalars.

Finally, we must prove the asserted symmetry of r . To this end note first that T_κ maps symmetric functions to symmetric functions. Its weak compactness entails that T_κ^2 is a compact operator [7, Corollary VI.7.5] and thus maps bounded sequences to sequences with (strongly) convergent subsequences. As a consequence, any accumulation point g of the bounded sequence $n^{-1} T_\kappa \sum_{k=1}^n T_\kappa^k \mathbf{1}_S$, $n \geq 1$, is a *continuous* positive symmetric function with $Tg = g$ and thus a multiple of r . Hence, r must be symmetric. \square

Now ${}^\kappa \widehat{P}f(x, t) := r(x)^{-1} \mathbb{E}(|xM|^\kappa f((xM)^\sim, t + \log |xM|) r((xM)^\sim))$ defines a Markov transition kernel on $S \times \mathbb{R}$ corresponding to $({}^\kappa \mathbb{P}_x((X_n, V_n)_{n \geq 0} \in \cdot))_{x \in S}$ as defined by (17). Its associated "marginal"

$${}^\kappa P f(x) := \frac{1}{r(x)} \mathbb{E}(|xM|^\kappa f((xM)^\sim) r((xM)^\sim)) \quad (18)$$

is the transition kernel of $(X_n)_{n \geq 0}$ under $({}^\kappa \mathbb{P}_x)_{x \in S}$.

5.2 Proof of Theorem 5.1: Checking the assumptions of the MRT

This section corresponds to [12, Proposition 2], but provides a much shorter proof, even if technical details not mentioned here had been included. A random variable $T \geq 0$ is called geometrically bounded if it has exponentially decreasing tails.

Lemma 5.5. *Suppose (A4), (A5) and (SA). Then, for each $x \in S$, there exists some $\delta > 0$ such that $B_\delta(x)$ is a regeneration set with respect to P as well as ${}^\kappa P$, and the minorization condition holds with the same probability measure ϕ defined in Lemma 2.1. Moreover, $(X_n)_{n \geq 0}$ is a strongly aperiodic Doeblin chain under $(\mathbb{P}_y)_{y \in S}$ as well as under $({}^\kappa \mathbb{P}_y)_{y \in S}$.*

Proof. Let $x \in S$. By (MC) with $m = 1$, we know that $P(y, \cdot) \geq P_C(y, \cdot) \geq p\phi$ for suitable δ, p, C, ϕ and all $y \in S$ (see Lemma 2.1, especially (ii) in the proof for the definition of ϕ). In particular, $\inf_{y \in S} P(y, B_\delta(x)) \geq p\phi(B_\delta(x)) = p > 0$ which gives geometrically bounded times to visit $B_\delta(x)$ under any \mathbb{P}_y (uniformly in $y \in S$), thus $B_\delta(x)$ is regenerative with respect to P . In order to get the same with respect to ${}^\kappa P$, we first note that

$$\gamma_2 := \min_{z_1, z_2 \in S} \frac{r(z_1)}{r(z_2)} \quad \text{and} \quad \gamma_3 := \min_{z \in S, \mathbf{m} \in C} |z\mathbf{m}|^\kappa$$

are clearly both positive (here the compactness of the set $C \subset GL(d, \mathbb{R})$ enters in a crucial way). But then we infer

$$\begin{aligned} {}^\kappa P(y, A) &\geq {}^\kappa P_C(y, A) \geq \frac{1}{r(y)} \int_C |y\mathbf{m}|^\kappa \mathbf{1}_A((y\mathbf{m})^\sim) r((y\mathbf{m})^\sim) \mathbb{P}(M \in d\mathbf{m}) \\ &\geq \gamma_2 \gamma_3 \int_C \mathbf{1}_A((y\mathbf{m})^\sim) \mathbb{P}(M \in d\mathbf{m}) \\ &= \gamma_2 \gamma_3 P_C(y, A) \geq p \gamma_2 \gamma_3 \phi(A) \end{aligned}$$

for any measurable $A \subset B_\delta(x)$ and $y \in S$ and thereby that $B_\delta(x)$ is regenerative with respect to ${}^\kappa P$ as well. \square

The stationary distribution of this chain with respect to ${}^\kappa P$ is always denoted as π hereafter. Note that also $(X_n, U_n)_{n \geq 0}$ is a stationary sequence under $({}^\kappa \mathbb{P}_x)_{x \in S}$.

We will need further information on the behavior of $|xM_{\sigma_n}|$, which follows from the bivariate minorization condition stated in Remark 5 (with $n_0 + n_1 + n_2 = 1$ due to our standing assumption).

Lemma 5.6. *Let $B_\delta(x)$ be a regenerative ball with minorizing probability measure ϕ as in Lemma 2.1. Then we can choose a sequence of regeneration epochs $(\sigma_n)_{n \geq 0}$ such that, for suitable $\mathfrak{C} > \mathfrak{c} > 0$,*

$$\mathfrak{C} \geq \|M_{\sigma_n}\| \geq \inf_{y \in S} |yM_{\sigma_n}| \geq \mathfrak{c} \quad \mathbb{P}\text{-a.s.} \quad (\text{and thus } {}^\kappa \mathbb{P}\text{-a.s.}) \quad (19)$$

for all $n \geq 1$. As a particular consequence, $U_{\sigma_1} = \log |X_{\sigma_1-1}M_{\sigma_1}|$ is \mathbb{P} -a.s. bounded, that is taking values in some finite interval $[s_*, s^*]$.

Proof. We just note that, by (BMC), we may generate $(X_{\sigma_n}, M_{\sigma_n})$ given $X_{\sigma_n-1} = y$ at any regeneration epoch σ_n according to $\psi(y, \cdot)$ having first marginal ϕ , thus $X_{\sigma_n} \stackrel{d}{=} \phi$. Moreover, $M_{\sigma_n} \in C$ \mathbb{P} -a.s. for a compact $C \subset GL(d, \mathbb{R})$ which entails $\|M_{\sigma_n}\| \leq \mathfrak{C}$, $\|M_{\sigma_n}^{-1}\| \leq \mathfrak{c}^{-1}$ \mathbb{P} -a.s. for some constants. Since

$$\inf_{x \in S} |xM_{\sigma_n}| = \inf_{x \in S} \frac{|xM_{\sigma_n}|}{|xM_{\sigma_n}M_{\sigma_n}^{-1}|} = \frac{1}{\sup_{x \in S} |xM_{\sigma_n}^{-1}|} = \frac{1}{\|M_{\sigma_n}^{-1}\|},$$

we infer (19). The \mathbb{P} -almost sure boundedness of U_{σ_1} then follows directly from $\|M_{\sigma_1}\|^{-1} \leq e^{|U_{\sigma_1}|} \leq \|M_{\sigma_1}\|$. \square

From now on, we will always assume that (19) is in force when given sequence of regeneration epochs $(\sigma_n)_{n \geq 0}$. The regeneration set will always be some ball $B_\delta(x)$, the i.i.d. coin-tossing variables are denoted by J_n , so $J_n \stackrel{d}{=} \text{Bernoulli}(p)$ for $n \geq 0$, and the minorizing measure by ϕ as in Lemma 2.1 (which naturally depends on $B_\delta(x)$).

The following result will be needed in section 8:

Lemma 5.7. *For any sequence $(\sigma_n)_{n \geq 0}$ of regeneration epochs as described above, and all $y \in S$*

$$\mathbb{P}_y (\|\Pi_{\sigma_1}\| < 1) > 0.$$

Proof. Due to geometric sampling, in particular Remark 6,

$$\begin{aligned} \mathbb{P}_y (\|\Pi_{\sigma_1}\| < 1) &\geq \mathbb{P}_y (\|\Pi_2\| < 1, \sigma_1 \leq 2) \\ &\geq \mathbb{P} \left(\|\Pi_1\| < \frac{1}{\mathfrak{e}}, (x\Pi_1)^\sim \in B_\delta(x), J_1 = 1 \right) \\ &\geq p \mathbb{P} \left(\|\Pi_1\| < \frac{1}{\mathfrak{e}}, (x\Pi_1)^\sim \in B_\delta(x) \right) > 0, \end{aligned}$$

where $B_\delta(x)$ denotes the regenerative ball for $(\sigma_n)_{n \geq 0}$. \square

We now turn to the lattice-type of $(X_n, V_n)_{n \geq 0}$, which is the same under $(\mathbb{P}_x)_{x \in S}$ and $({}^\kappa \mathbb{P}_x)_{x \in S}$. Kesten [12] imposes an additional assumption involving so-called feasible matrices in order to ensure that $(X_n, V_n)_{n \geq 0}$ is nonarithmetic. But in view of assumption (A5) it should be no surprise that this is not needed here. The following lemma provides the confirmation in an even stronger form.

Lemma 5.8. *Suppose (A4), (A5) and (SA). Then $(X_n, V_n)_{n \geq 0}$ is nonarithmetic under $(\mathbb{P}_x)_{x \in S}$, in fact*

$$\mathbb{E}_x \left| \mathbb{E} \left(e^{itV_1} | X_0, X_1 \right) \right| < 1$$

for all $t \neq 0$ and π -almost all $x \in S$.

Proof. If the assertion fails to hold, there exists a distribution ν on S , absolutely continuous with respect to π , such that $\mathbb{E}_\nu \left| \mathbb{E} \left(e^{itV_1} | X_0, X_1 \right) \right| = 1$ for some $t \neq 0$. As a consequence,

$$\mathbb{E} \left(e^{itV_1} | X_0, X_1 \right) = e^{itf(X_0, X_1)} \mathbb{P}_x\text{-a.s.}$$

for some measurable function f and ν -almost all $x \in S$ or, equivalently,

$$\mathbb{P}_\nu \left(V_1 \in f(X_0, X_1) + t^{-1} \mathbb{Z} \right) = 1. \tag{20}$$

W.l.o.g. suppose $t = 1$ hereafter. Due to (A5) and (SA), a nonzero component of $\mathbb{P}_x ((X_1, V_1) \in \cdot)$ is given by

$$\Lambda_x(A \times B) := \gamma_0 \int_{B_c(\Gamma_0)} \mathbf{1}_A((x\mathbf{m})^\sim) \mathbf{1}_B(\log |x\mathbf{m}|) \mathfrak{L}(d\mathbf{m})$$

for measurable $A \subset S$, $B \subset \mathbb{R}$ and any $x \in S$. The mapping $\mathbf{m} \mapsto x\mathbf{m}$ induces an absolutely continuous measure on \mathbb{R}^d with some λ^d -density g , say. Switching to spherical coordinates, there are $\varepsilon_1, \varepsilon_2 > 0$ such that

$$\Lambda_x(A \times B) = \gamma_0 \int_{|x\Gamma_0|-\varepsilon_1}^{|x\Gamma_0|+\varepsilon_1} \int_{B_{\varepsilon_2}((x\Gamma_0)^\sim) \cap S} \mathbf{1}_A(\omega) \mathbf{1}_B(s) g(s\omega) \sigma(d\omega) \frac{1}{s^{1+d}} ds$$

where σ is a measure on the sphere S . Now, if (20) were true with $t = 1$, then

$$\Lambda_x(S \times \mathbb{R}) = \gamma_0 \int_{B_{\varepsilon_2}((x\Gamma_0)^\sim) \cap S} \int_{|x\Gamma_0|-\varepsilon_1}^{|x\Gamma_0|+\varepsilon_1} \mathbf{1}_{f(x,\omega)+\mathbb{Z}}(s) g(s\omega) \frac{1}{s^{1+d}} ds \sigma(d\omega) > 0$$

for all x which is impossible because the inner integral over a countable set is clearly zero for any fixed ω . \square

For the proof of Theorem 5.1, it finally remains to verify that $(X_n, V_n)_{n \geq 0}$ has positive drift under ${}^\kappa\mathbb{P}$. The subsequent argument simplifies the original one given by Kesten [12].

Lemma 5.9. *Under ${}^\kappa\mathbb{P}$, $(X_n, V_n)_{n \geq 0}$ has positive drift, given by*

$$\alpha := {}^\kappa\mathbb{E}_\pi \left(\frac{V_n}{n} \right) = \frac{1}{n} \int \frac{1}{r(x)} \mathbb{E} |x\Pi_n|^\kappa \log |x\Pi_n| r((x\Pi_n)^\sim) \pi(dx)$$

for any $n \geq 1$.

Proof. For each $n \geq 1$, the function

$$g_n(\varkappa) := \int \frac{1}{r(x)} \mathbb{E} |x\Pi_n|^\varkappa r((x\Pi_n)^\sim) \pi(dx)$$

is finite and thus convex for $\varkappa \in [0, \kappa_0]$. Moreover $g_n(\kappa) = 1$ and the left derivative at κ equals

$$\lim_{x \uparrow \kappa} \frac{g_n(x) - g_n(\kappa)}{x - \kappa} = \alpha n.$$

By convexity, α is positive if we can show that $g_n(\varkappa) < 1$ for some n and some $\varkappa < \kappa$. To this end pick any $\varkappa < \kappa$ and recall that $\varrho(\varkappa) < 1$. It follows that

$$g_n(\varkappa) = \int \frac{T_\varkappa^n r(x)}{r(x)} \pi(dx) = \int \frac{\|r\|_\infty}{r(x)} T_\varkappa^n \left(\frac{r(x)}{\|r\|_\infty} \right) \pi(dx) \leq C \|T_\varkappa^n\|$$

for some $C \in (0, \infty)$ and all $n \geq 1$. As $\|T_\varkappa^n\|^{1/n} \rightarrow \varrho(\varkappa)$, we infer $g_n(\varkappa) \rightarrow 0$ and thus the desired result. \square

5.3 Tail behavior of $\sup_{n \geq 1} |x \Pi_n|$

With the help of the MRT 4.2, we are now able to prove the following result on the tail behavior of $\sup_{n \geq 1} |x \Pi_n|$.

Proposition 5.10. *Under the conditions of Theorem 1.1 and with r as defined in Lemma 5.4,*

$$\lim_{t \rightarrow \infty} t^\kappa \mathbb{P} \left(\sup_{n \geq 1} |x \Pi_n| > t \right) = L r(x),$$

for π -almost all $x \in S$ and some $L > 0$.

Proof. The function $f : S \times (0, \infty) \rightarrow (0, \infty)$, $(y, s) \mapsto e^{-\kappa s}/r(y)$ is bounded and continuous whence, by an application of the MRT,

$$L(f) := \lim_{t \rightarrow \infty} {}^\kappa \mathbb{E}_x \left(f(Z(t), R(t)) \mathbf{1}_{\{N(t) < \infty\}} \right)$$

exists and is positive. On the other hand, we have

$$\begin{aligned} {}^\kappa \mathbb{E}_x \left(f(Z(t), R(t)) \mathbf{1}_{\{N(t) < \infty\}} \right) &= \sum_{n \geq 1} {}^\kappa \mathbb{E}_x \left(f(X_n, V_n - t) \mathbf{1}_{\{N(t) = n\}} \right) \\ &= \sum_{n \geq 1} {}^\kappa \mathbb{E}_x \left(\frac{1}{r(X_n)} e^{-\kappa V_n + \kappa t} \mathbf{1}_{\{N(t) = n\}} \right) \\ &= \frac{e^{\kappa t}}{r(x)} \sum_{n \geq 1} \mathbb{E}_x \left(\frac{1}{r(X_n)} e^{-\kappa V_n} r(X_n) e^{\kappa V_n} \mathbf{1}_{\{\tau(t) = n\}} \right) \\ &= \frac{e^{\kappa t}}{r(x)} \mathbb{P}_x(N(t) < \infty) \\ &= \frac{e^{\kappa t}}{r(x)} \mathbb{P} \left(\sup_{n \geq 1} \log |x \Pi_n| > t \right), \end{aligned} \tag{21}$$

which provides the asserted result upon substituting e^t by t . \square

5.4 Tail behavior of $\sup_{n \geq 1} |x \Pi_{\sigma_n - 1}|$

Let $B_\delta(x_0)$ be any regenerative ball with associated sequences $(\sigma_n)_{n \geq 0}$ and $(\tau_n)_{n \geq 1}$ of regeneration epochs and hitting times, respectively. In Section 8, we will need and therefore show below that

$$\limsup_{t \rightarrow \infty} t^\kappa \mathbb{P}_x \left(\sup_{n \geq 1} |x \Pi_{\sigma_n - 1}| > t \right) > 0$$

for π -almost all $x \in S$. The proof hinges on the following proposition similar to Proposition 5.10 above.

Proposition 5.11. *Under the conditions of Theorem 1.1 and with r defined in Lemma 5.4, there exists $L' > 0$ such that*

$$\lim_{t \rightarrow \infty} t^\kappa \mathbb{P}_x \left(\sup_{n \geq 1} |x \Pi_{\tau_n}| > t \right) = L' r(x),$$

for π -almost all $y \in B_\delta(x_0)$.

Since $V_{\tau_n} = \log |x\Pi_{\tau_n}|$ a.s. under \mathbb{P}_x and ${}^\kappa\mathbb{P}_x$, we can proceed exactly as in the proof of Proposition 5.10, provided that the assumptions of the MRT 4.2 hold for the sequence $(X_{\tau_n}, V_{\tau_n})_{n \geq 0}$ under $({}^\kappa\mathbb{P}_x)_{x \in S}$, which is verified by the subsequent lemma. We note that (17) extends to

$$\begin{aligned} & {}^\kappa\mathbb{E}_x f(X_0, V_0, X_1, V_1, \dots, X_{\tau_n}, V_{\tau_n}) \\ &= \frac{1}{r(x)} \mathbb{E}_x \left(r(X_{\tau_n}) e^{\kappa V_{\tau_n}} f(X_0, V_0, X_1, V_1, \dots, X_{\tau_n}, V_{\tau_n}) \right), \end{aligned} \quad (22)$$

as one can easily see by applying (17) to ${}^\kappa\mathbb{E}_x f(X_0, V_0, X_1, V_1, \dots, X_k, V_k) \mathbf{1}_{\{\tau_n=k\}}$ for each $k \geq 1$, which in turn is possible for the appearing indicator is a function of $(X_0, V_0, X_1, V_1, \dots, X_k, V_k)$.

Lemma 5.12. *The hit chain $(X_{\tau_n})_{n \geq 0}$ constitutes a strongly aperiodic Doeblin chain under $({}^\kappa\mathbb{P}_x)_{x \in S}$ with stationary distribution $\nu = \pi(\cdot \cap B_\delta(x_0))/\pi(B_\delta(x_0))$. Moreover, $(X_{\tau_n}, V_{\tau_n})_{n \geq 0}$ is a nonarithmetic MRW under $({}^\kappa\mathbb{P}_x)_{x \in S}$ with positive drift.*

Proof. For the first statement, we just note that $\{\sigma_n : n \geq 1\} \subset \{\tau_n : n \geq 1\}$. Next, due to Lemma 5.8 and the conditional independence of U_1, U_2, \dots given $(X_n)_{n \geq 0}$, we find that for $t \neq 0$

$$\begin{aligned} \mathbb{E}_x \left(\left| \mathbb{E} \left(e^{itV_{\tau_1}} | X_0, X_{\tau_1} \right) \right| \right) &\leq \mathbb{E}_x \left(\prod_{k=1}^{\tau_1} \left| \mathbb{E} \left(e^{itU_k} | X_{k-1}, X_k \right) \right| \right) \\ &\leq \mathbb{E}_x \left(\left| \mathbb{E} \left(e^{itV_1} | X_0, X_1 \right) \right| \right) < 1 \end{aligned}$$

for π -almost all and thus ν -almost all x . Consequently, $(X_{\tau_n}, V_{\tau_n})_{n \geq 0}$ is nonarithmetic under $(\mathbb{P}_x)_{x \in S}$ and $({}^\kappa\mathbb{P}_x)_{x \in S}$. Finally, we obtain for $\alpha' := {}^\kappa\mathbb{E}_\nu V_{\tau_1}$ that

$$\alpha' = \lim_{n \rightarrow \infty} \frac{V_{\tau_n}}{n} \geq \lim_{n \rightarrow \infty} \frac{V_{\tau_n}}{\sigma_n} \cdot \liminf_{n \rightarrow \infty} \frac{\tau_n}{n} \geq \alpha \cdot 1 > 0 \quad {}^\kappa\mathbb{P}\text{-a.s.}$$

where Remark 8 should be recalled. □

Proposition 5.13. *Let $x_0 \in S$ and $\delta > 0$ be such that $B_\delta(x_0)$ is regenerative with associated regeneration epochs σ_n , $n \geq 1$. Then*

$$\liminf_{t \rightarrow \infty} t^\kappa \mathbb{P}_x \left(\sup_{n \geq 1} |x\Pi_{\sigma_n-1}| > t \right) > 0 \quad (23)$$

for π -almost all $x \in B_\delta(x_0)$.

Proof. Let $(\tau_n)_{n \geq 1}$ denote the sequence of hitting times of $B_\delta(x_0)$ and observe that it contains $(\sigma_n - 1)_{n \geq 1}$ as a subsequence. By Proposition 5.11, we have

$$\lim_{t \rightarrow \infty} t^\kappa \mathbb{P}_x \left(\sup_{n \geq 1} |x\Pi_{\tau_n}| > t \right) = L' r(x) > 0,$$

for some $L' > 0$ and π -almost all $x \in B_\delta(x_0)$.

Fix any such x hereafter and put $\widehat{N}(t) := \inf\{n \geq 1 : |x\Pi_{\tau_n}| > t\}$, thus

$$\left\{ \sup_{n \geq 1} |x\Pi_{\tau_n}| > t \right\} = \{\widehat{N}(t) < \infty\}.$$

Since $\{\sup_{n \in \mathbb{N}} |x \Pi_{\sigma_n - 1}| > \varepsilon t\}$ contains

$$A(t) := \bigcup_{n \geq 1} \left\{ \widehat{N}(t) = n, |x \Pi_{\tau_n + 1}| > \varepsilon t, X_{\tau_n + 1} \in B_\delta(x_0), J_{\tau_n + 1} = 1 \right\}$$

as a subset, it suffices to show that $\liminf_{t \rightarrow \infty} t^\kappa \mathbb{P}_x(A(t)) > 0$ for suitably chosen $\varepsilon > 0$. To this end, we make the following estimation.

$$\begin{aligned} \mathbb{P}_x(A(t)) &= p \sum_{n \geq 1} \mathbb{P}_x \left(\widehat{N}(t) = n, |x \Pi_{\tau_n + 1}| > \varepsilon t, X_{\tau_n + 1} \in B_\delta(x_0) \right) \\ &= p \sum_{n \geq 1} \mathbb{P}_x \left(\widehat{N}(t) = n, |(x \Pi_{\tau_n})^\sim M_{\tau_n + 1}| > \frac{\varepsilon t}{|x \Pi_{\tau_n}|}, X_{\tau_n + 1} \in B_\delta(x_0) \right) \\ &\geq p \sum_{n \geq 1} \mathbb{P}_x \left(\widehat{N}(t) = n, \inf_{y \in S} |y M_{\tau_n + 1}| > \varepsilon, X_{\tau_n + 1} \in B_\delta(x_0) \right) \\ &\geq p \inf_{u \in B_\delta(x_0)} \mathbb{P}_u \left(\inf_{y \in S} |y M| > \varepsilon, (uM)^\sim \in B_\delta(x_0) \right) \mathbb{P}_x \left(\widehat{N}(t) < \infty \right) \\ &\geq p \inf_{u \in B_\delta(x_0)} \mathbb{P}_u \left(\inf_{y \in S} |y M_{\sigma_1}| > \varepsilon, \sigma_1 = 1 \right) \mathbb{P}_x \left(\widehat{N}(t) < \infty \right). \end{aligned}$$

Fixing any $\varepsilon \in (0, \mathbf{c})$, we now infer from (19) in Lemma 5.6 that

$$\mathbb{P}_u \left(\inf_{y \in S} |y M_{\sigma_1}| > \varepsilon, \sigma_1 = 1 \right) = \mathbb{P}_u(\sigma_1 = 1) = p > 0$$

for any $u \in B_\delta(x_0)$, whence we finally conclude

$$\mathbb{P}_x(A(t)) \geq p^2 \mathbb{P}_x \left(\sup_{n \geq 1} |y \Pi_{\tau_n}| > t \right)$$

for all $t > 0$ and thus $\liminf_{t \rightarrow \infty} t^\kappa \mathbb{P}_x(A(t)) > 0$. \square

6 Proof of Theorem 1.1: Implicit Markov renewal theory

We now turn to the proof of our main result Theorem 1.1, all assumptions of which will therefore be in force throughout, in fact in strengthened form given by our standing assumption.

Embarking on ideas by Goldie [9] and Le Page [15], a comparison of the distribution functions of xR and xMR will enable us to make use of a Markov modulated version of Goldie's implicit renewal theory. This will prove that $K(x) = \lim_{t \rightarrow \infty} t^\kappa \mathbb{P}(xR > t)$ exists for π -almost all $x \in S$.

We start with a simple lemma, stated without proof, which is just Lemma 9.3 in [9] adapted to our situation.

Lemma 6.1. *Let $x \in S$. If $K(x) := \lim_{t \rightarrow \infty} t^{-1} \int_0^t s^\kappa \mathbb{P}(xR > s) ds$ exists and is finite, then so does $\lim_{t \rightarrow \infty} t^\kappa \mathbb{P}(xR > t)$ and equals $K(x)$ as well.*

Substituting t' for e^t and a change of variables show that $t'^{-1} \int_0^{t'} s^\kappa \mathbb{P}(xR > s) ds$ equals $e^{-t} \int_{-\infty}^t e^{(\kappa+1)s} \mathbb{P}(xR > e^s) ds$ which is the form needed in the next result which provides us with the basic renewal theoretic identity.

Lemma 6.2. For all $t \in \mathbb{R}$,

$$\frac{e^{-t}}{r(x)} \int_{-\infty}^t e^{(\kappa+1)s} \mathbb{P}(xR > e^s) ds = \sum_{n \geq 0} \int \widehat{g}(y, t-u)^\kappa \mathbb{P}_x(X_n \in dy, V_n \in du), \quad (24)$$

where $\widehat{g}(y, t) := \int_{-\infty}^t e^{-(t-s)} g(y, s) ds$ is the exponential smoothing of

$$g(y, s) = \frac{e^{\kappa s}}{r(y)} [\mathbb{P}(yR > e^s) - \mathbb{P}(yMR > e^s)].$$

Proof. For arbitrary $n \in \mathbb{N}$, $x \in S$ and $s \in \mathbb{R}$, consider the following telescoping sum for $\mathbb{P}(xR > e^s)$ (recalling independence of R , M , and $(M_n)_{n \geq 1}$)

$$\begin{aligned} & \sum_{k=1}^n [\mathbb{P}(x\Pi_{k-1}R > e^s) - \mathbb{P}(x\Pi_kR > e^s)] + \mathbb{P}(x\Pi_nR > e^s) \\ &= \sum_{k=1}^n \left[\mathbb{P}_x(e^{V_{k-1}}X_{k-1}R > e^s) - \mathbb{P}_x(e^{V_{k-1}}X_{k-1}M_kR > e^s) \right] + \mathbb{P}_x(e^{V_n}X_nR > e^s) \\ &= \sum_{k=0}^{n-1} \left[\mathbb{P}_x(e^{V_k}X_kR > e^s) - \mathbb{P}_x(e^{V_k}X_kMR > e^s) \right] + \mathbb{P}_x(e^{V_n}X_nR > e^s) \\ &= \sum_{k=0}^{n-1} \int \mathbb{P}(yR > e^{s-u}) - \mathbb{P}(yMR > e^{s-u}) \mathbb{P}_x(X_k \in dy, V_k \in du) \\ & \qquad \qquad \qquad + \mathbb{P}_x(e^{V_n}X_nR > e^s) \end{aligned}$$

Multiply by $e^{\kappa s}/r(x) > 0$ to obtain

$$\begin{aligned} & \frac{e^{\kappa s}}{r(x)} \mathbb{P}(xR > e^s) \\ &= \sum_{k=0}^{n-1} \int \frac{e^{\kappa(s-u)}}{r(x)} [\mathbb{P}(yR > e^{s-u}) - \mathbb{P}(yMR > e^{s-u})] \\ & \quad \times \frac{r(y)}{r(y)} e^{\kappa u} \mathbb{P}_x(X_k \in dy, V_k \in du) + \frac{e^{\kappa s}}{r(x)} \mathbb{P}_x(e^{V_n}X_nR > e^s) \\ &= \sum_{k=0}^{n-1} \int g(y, s-u)^\kappa \mathbb{P}_x(X_k \in dy, V_k \in du) + \frac{e^{\kappa s}}{r(x)} \mathbb{P}_x(X_nR > e^{s-V_n}). \end{aligned}$$

Convolution with a standard exponential distribution then gives

$$\begin{aligned} \int_{-\infty}^t e^{-(t-s)} \frac{1}{r(x)} e^{\kappa s} \mathbb{P}(xR > e^s) ds &= \sum_{k=0}^{n-1} \int \widehat{g}(y, t-u)^\kappa \mathbb{P}_x(X_k \in dy, V_k \in du) \\ & \quad + \int_{-\infty}^t e^{-(t-s)} \frac{e^{\kappa s}}{r(x)} \mathbb{P}_x(X_nR > e^{s-V_n}) ds \end{aligned}$$

By the Cauchy-Schwarz inequality, $|X_nR| \leq |X_n||R| = |R|$ and thus

$$\mathbb{P}_x(X_nR > e^{s-V_n}) \leq \mathbb{P}_x(|X_nR| > e^{s-V_n}) \leq \mathbb{P}_x(|R| > e^{s-V_n}).$$

But the last term converges to 0 as $n \rightarrow \infty$ for any $s > 0$, because $\lim_{n \rightarrow \infty} V_n = -\infty$ \mathbb{P}_x -a.s. Hence assertion (24) follows by an appeal to the dominated convergence theorem. \square

Obviously, if ${}^\kappa\mathbb{U}_x := \sum_{n \geq 0} {}^\kappa\mathbb{P}_x((X_n, V_n) \in \cdot)$, then the right-hand side of (24) equals $\widehat{g} * {}^\kappa\mathbb{U}_x(t)$ for x outside a π -null set N provided that sum and integral may be interchanged for $x \notin N$. But the latter follows if we can prove hereafter that \widehat{g} is π -directly Riemann integrable which will also be the crucial condition that ensures applicability of the MRT 4.2. Indeed, if \widehat{g} has this property, then, by Equation (5.8) and Lemma A.5 in [1],

$$\begin{aligned} \widehat{g} * {}^\kappa\mathbb{U}_x(t) &= {}^\kappa\mathbb{E}_x \left(\sum_{k \geq 0} \widehat{g}(X_k, t - V_k) \right) \\ &= \int \sum_{k \geq 0} \widehat{g}(y, t - u) {}^\kappa\mathbb{P}_x(X_k \in dy, V_k \in du) < \infty \end{aligned}$$

for all $t \in \mathbb{R}$ and π -almost all $x \in S$. Split \widehat{g} in positive and negative part. This yields two π -null sets N_1, N_2 such that $\widehat{g}^+ * {}^\kappa\mathbb{U}_x(t)$ and $\widehat{g}^- * {}^\kappa\mathbb{U}_x(t)$ are finite for all $x \in (N_1 \cup N_2)^c$ and all $t \in \mathbb{R}$. By Fubini's theorem, sum and integral in (24) may be interchanged for all $x \in (N_1 \cup N_2)^c$. This is enough because the MRT asserts convergence of $\widehat{g} * {}^\kappa\mathbb{U}_x(t)$ only for x outside a π -null set.

Instead of π -direct Riemann integrability of \widehat{g} we will actually show the stronger property that

$$\sup_{y \in S} \sum_{n \in \mathbb{Z}} \sup_{t \in [n\delta, (n+1)\delta)} |\widehat{g}(y, t)| < \infty, \quad (25)$$

which can be done by resorting to the methods of Goldie [9, proof of Theorem 4.1] which are only summarized here. Let $L_1(\mathbb{R})$ as usual be the space of Lebesgue integrable functions.

Lemma 6.3. *If $f \in L_1(\mathbb{R})$ and $\widehat{f}(t) := \int_{-\infty}^t e^{-(t-u)} f(u) du$, then for any $\delta > 0$*

$$\sum_{n \in \mathbb{Z}} \sup_{t \in [n\delta, (n+1)\delta)} |\widehat{f}(t)| \leq \delta e^{2\delta} \int |f(t)| dt < \infty.$$

Proof. This is Lemma 9.2 in [9] \square

In view of the previous lemma, it suffices to show for (25) that $\int |g(y, s)| ds$ is uniformly bounded in y . First observe that (cf. [9, Corollary 2.4])

$$\begin{aligned} \int_{\mathbb{R}} |g(y, s)| ds &= \int_{\mathbb{R}} \frac{e^{\kappa s}}{r(y)} |\mathbb{P}(yR > e^s) - \mathbb{P}(yMR > e^s)| ds \\ &= \int_{\mathbb{R}} \frac{e^{\kappa s}}{r(y)} |\mathbb{P}(yMR + yQ > e^s) - \mathbb{P}(yMR > e^s)| ds \\ &= \frac{1}{\kappa r(y)} \mathbb{E} \left| ((yMR + yQ)^+)^{\kappa} - ((yMR)^+)^{\kappa} \right|. \end{aligned}$$

Then a case-by-case analysis with respect to the signs of yMR and yQ yields that

$$\sup_{y \in S} \frac{1}{\kappa r(y)} \mathbb{E} \left| ((yMR + yQ)^+)^{\kappa} - ((yMR)^+)^{\kappa} \right| < \infty,$$

see [9, Theorem 4.1]. Now we are ready to prove

Lemma 6.4. For π -almost all $x \in S$,

$$\lim_{t \rightarrow \infty} \frac{e^{-t}}{r(x)} \int_{-\infty}^t e^{(\kappa+1)s} \mathbb{P}(xR > e^s) ds = K_0,$$

where $K_0 := \frac{1}{\alpha\kappa} \int_S \frac{1}{r(y)} \mathbb{E} \left(((yR)^+)^{\kappa} - ((yMR)^+)^{\kappa} \right) \pi(dy) < \infty$ and α as before denotes the drift of $(X_n, V_n)_{n \geq 0}$.

Proof. Since \hat{g} is π -directly Riemann integrable, we may exchange sum and integral in (24) for π -almost all $x \in S$ and apply the MRT. This tells us that the right-hand side of (24) has the finite limit

$$\begin{aligned} & \frac{1}{\alpha} \int_S \int_{\mathbb{R}} e^{-t} \int_{-\infty}^t \frac{e^{(\kappa+1)s}}{r(y)} [\mathbb{P}(yR > e^s) - \mathbb{P}(yMR > e^s)] ds dt \pi(dy) \\ &= \frac{1}{\alpha} \int_S \int_{\mathbb{R}} g(y, t) dt \pi(dy) \\ &= \frac{1}{\alpha} \int_S \frac{1}{r(y)} \int_{\mathbb{R}} e^{\kappa t} [\mathbb{P}(yR > e^t) - \mathbb{P}(yMR > e^t)] dt \pi(dy) \\ &= \frac{1}{\alpha} \int_S \frac{1}{r(y)} \int_0^{\infty} u^{\kappa-1} [\mathbb{P}(yR > u) - \mathbb{P}(yMR > u)] du \pi(dy) \\ &= \frac{1}{\alpha\kappa} \int_S \frac{1}{r(y)} \mathbb{E} \left(((yR)^+)^{\kappa} - ((yMR)^+)^{\kappa} \right) \pi(dy) \end{aligned}$$

for π -almost all x . □

7 Proof of Theorem 1.1: Assertion (5) holds for all $x \in S$

So far we have proved our main assertion (5) (except for the positivity of $K(x)$) for π -almost all $x \in S$ and thus for all x from a dense subset of S (this is a direct consequence of (A4)). By employing a refined renewal argument, we will now remove this restriction. To this end, we fix an arbitrary $x \in S$ and $\delta > 0$ so small that $B_\delta(x)$ is regenerative for ${}^\kappa P$ with minorizing distribution ϕ and associated sequence $(\sigma_n)_{n \geq 1}$ of regeneration epochs such that Lemma 5.6 is in force. Put $\sigma := \sigma_1$. The task is to show that $\hat{g} * {}^\kappa \mathbb{U}_x(t)$ converges to K_0 , and we begin by pointing out that

$$\hat{g} * {}^\kappa \mathbb{U}_x(t) = {}^\kappa \mathbb{E}_x \left(\sum_{k \geq 0} \hat{g}(X_k, t - V_k) \right) = G(x, t) + \hat{g} * {}^\kappa \mathbb{U}_{\varphi(x, \cdot)}(t) \quad (26)$$

where $\varphi(x, \cdot) := {}^\kappa \mathbb{P}_x((X_\sigma, V_\sigma) \in \cdot)$ and

$$G(x, t) := {}^\kappa \mathbb{E}_x \left(\sum_{k=0}^{\sigma-1} \hat{g}(X_k, t - V_k) \right), \quad (x, t) \in S \times \mathbb{R}.$$

As for this last function, we now prove:

Lemma 7.1. *The function G is bounded and satisfies $\lim_{t \rightarrow \infty} G(y, t) = 0$ for all $y \in S$.*

Proof. By (25), $C := \sup\{|\hat{g}(y, t)| : y \in S, t \in \mathbb{R}\} < \infty$, and since $(X_n)_{n \geq 0}$ is a strongly aperiodic Doeblin chain, we infer

$$\sup_{y \in S, t \in \mathbb{R}} |G(y, t)| \leq C \sup_{y \in S} {}^\kappa \mathbb{E}_y \sigma < \infty.$$

Just note that the time it takes to hit the regenerative ball $B_\delta(x)$ pertaining to σ from any y is geometrically bounded (uniformly in y) and that a geometric number of coin tosses (the J_n) of such times determines σ . Turning to the convergence assertion, we point out that, again by property (25), $\lim_{t \rightarrow \infty} \hat{g}(y, t) = 0$ for all $y \in S$, which implies the desired result by an appeal to the dominated convergence theorem. \square

In view of (26), we are now left with a proof of $\hat{g} * {}^\kappa \mathbb{U}_{\varphi(x, \cdot)}(t) \rightarrow K_0$ defined in Lemma 6.4. This requires one more lemma.

Lemma 7.2. *The sequence $(X_\sigma, (X_n, U_n)_{n > \sigma})$ is independent of $(X_{\sigma-1}, V_{\sigma-1})$ under ${}^\kappa \mathbb{P}_x$ with distribution given by ${}^\kappa \mathbb{P}_\phi((X_0, (X_n, U_n)_{n \geq 1}) \in \cdot)$.*

Proof. The first assertion follows directly when observing that, by regeneration, $(X_{\sigma+n})_{n \geq 0}$ and $(X_{\sigma-1}, V_{\sigma-1})$ are independent under ${}^\kappa \mathbb{P}_x$, and the fact that the conditional distribution of U_k given $(X_n)_{n \geq 0}$ only depends on (X_k, X_{k-1}) . The proof is completed by the observation that ${}^\kappa \mathbb{P}_x((X_{\sigma+n})_{n \geq 0} \in \cdot) = {}^\kappa \mathbb{P}_\phi((X_n)_{n \geq 0} \in \cdot)$. \square

Define $V_{\sigma, n} := V_{\sigma+n} - V_\sigma$ for $n \geq 0$ and then

$$h(x, s, t) := {}^\kappa \mathbb{E}_x \left(\sum_{k \geq 0} \hat{g}(X_{\sigma+k}, t - s - V_{\sigma-1} - V_{\sigma, k}) \right)$$

for $s, t \in \mathbb{R}$. Lemma 7.2 implies

$$h(x, s, t) = \int_{\mathbb{R}} \hat{g} * {}^\kappa \mathbb{U}_\phi(t - s - r) {}^\kappa \mathbb{P}_x(V_{\sigma-1} \in dr).$$

As \hat{g} satisfies (25), we infer from the MRT 4.2 and the subsequent remark that $\hat{g} * {}^\kappa \mathbb{U}_\phi(t)$ is bounded and converges to K_0 . By the dominated convergence theorem, the same limit holds for $\lim_{t \rightarrow \infty} h(x, s, t)$ for all s .

Finally, the connection between $h(x, s, t)$ and $\hat{g} * {}^\kappa \mathbb{U}_{\varphi(x, \cdot)}(t)$ becomes apparent after the following observations: By Lemma 5.6, U_σ is taking values in some finite interval $[s_*, s^*]$. Hence we can estimate $\hat{g} * {}^\kappa \mathbb{U}_{\varphi(x, \cdot)}(t)$ by

$$\inf_{s \in [s_*, s^*]} h(x, s, t) \leq \hat{g} * {}^\kappa \mathbb{U}_{\varphi(x, \cdot)}(t) \leq \sup_{s \in [s_*, s^*]} h(x, s, t)$$

and thus arrive at the desired conclusion that $\lim_{t \rightarrow \infty} \hat{g} * {}^\kappa \mathbb{U}_{\varphi(x, \cdot)}(t) = K_0$.

8 Proof of Theorem 1.1: The limit $K(x)$ is positive

A combination of Lemma 6.1, Lemma 6.4 and the result of Section 7 renders convergence of $t^\kappa \mathbb{P}(xR > t)$ to the continuous function

$$K(x) := K_0 r(x) = \frac{r(x)}{\alpha\kappa} \int_S \frac{1}{r(y)} \mathbb{E} \left(((yR)^+)^{\kappa} - ((yMR)^+)^{\kappa} \right) \pi(dy)$$

for all $x \in S$. To complete the proof of Theorem 1.1, it remains to show that K or, equivalently, K_0 is positive, which is the topic of this final section.

Clearly, it suffices to show that $\limsup_{t \rightarrow \infty} t^\kappa \mathbb{P}(xR > t) > 0$ for some $x \in S$, or equivalently (since the limit exists) that the \liminf is positive. Notice that, as r is symmetric (Lemma 5.4), the same holds true for $K(x)$, hence

$$\lim_{t \rightarrow \infty} \mathbb{P}(xR > t) = \lim_{t \rightarrow \infty} \mathbb{P}(-xR > t) = \frac{1}{2} \lim_{t \rightarrow \infty} \mathbb{P}(|xR| > t).$$

So it is enough to show that $\liminf_{t \rightarrow \infty} t^\kappa \mathbb{P}(|xR| > t) > 0$ for some x . To this end we need the following lemma, originally due to Le Page [15, Lemma 3.11], which ensures that R and its "marginals" xR for any $x \in S$ have unbounded support. It is this result where the nondegeneracy assumption (A6), unused so far, enters in a crucial way. We postpone the proof until the end of this section.

Lemma 8.1. *For all $x \in S$ and $t \in \mathbb{R}$,*

$$\mathbb{P}(xR \leq t) < 1. \tag{27}$$

What this lemma shows is that, fixing any $x_0 \in S$, we can choose $\xi > 0$ and then sufficiently small $\zeta, \eta \in (0, 1)$ and $\delta \in (0, \zeta)$ such that

$$\mathbb{P}(zR > \xi) \geq \eta \quad \text{and} \quad \mathbb{P}(zR < (1 - \zeta)\xi) \geq \eta \tag{28}$$

for all $z \in B_\delta(x_0)$. Notice that

$$\inf_{z, y \in B_\delta(x_0)} zy > 1 - \delta. \tag{29}$$

We continue with a decomposition of xR with respect to entrances of $(x\Pi_k)^\sim$ into $B_\delta(x_0)$. In the following lemma, consider any (sub-)sequence $(\sigma_n)_{n \geq 1}$ of the hitting times (e.g. regeneration times). Note that (28) particularly holds for $z = X_{\sigma_n} = (x\Pi_{\sigma_n})^\sim$. Recall from Subsection 3.1 the definition of Q^n and R^n as well as $R^\tau \stackrel{d}{=} R$ for any a.s. finite stopping time τ with respect to $(\mathcal{F}_n)_{n \geq 0}$, the natural filtration of $(M_n, Q_n)_{n \geq 1}$.

Lemma 8.2. *Given any $x_0 \in S$ and sufficiently small $0 < \delta < 1$,*

$$\mathbb{P}(|xR| > t) \geq \eta \mathbb{P}_x \left(\sup_{n \geq 1} |xQ^{\sigma_n} + \xi x\Pi_{\sigma_n} y| > t \right)$$

holds true for all $x \in S$ and $y \in B_\delta(x_0)$.

Proof. This is an extension of Levy's inequality and inspired by [9, Prop. 4.2]. Since $R^{\sigma_k} \stackrel{d}{=} R$ for all $k \geq 1$ we see that (28) holds for R^{σ_k} as well. We show first that

$$\mathbb{P}(xR > t) \geq \eta \mathbb{P}_x \left(\sup_{n \geq 1} xQ^{\sigma_n} + \xi x\Pi_{\sigma_n} y > t \right)$$

and will consider $\mathbb{P}(-xR > t)$ in a second step. Define

$$C_k := \left\{ \max_{1 \leq j < k} (xQ^{\sigma_j} + \xi x\Pi_{\sigma_j} y) \leq t, xQ^{\sigma_k} + \xi x\Pi_{\sigma_k} y > t \right\}$$

and $D_k := \{x\Pi_{\sigma_k} R^{\sigma_k} > \xi x\Pi_{\sigma_k} y\}$.

By (29), $0 < (x\Pi_{\sigma_k})^\sim y \leq 1$ for all $y \in B_\delta(x_0)$, giving

$$D_k = \{(x\Pi_{\sigma_k})^\sim R^{\sigma_k} > \xi (x\Pi_{\sigma_k})^\sim y\} \supset \{(x\Pi_{\sigma_k})^\sim R^{\sigma_k} > \xi\}$$

and thus $\mathbb{P}_x(D_k | \mathcal{F}_{\sigma_k}) \geq \eta$ \mathbb{P}_x -a.s. In combination with $\bigcup_{k=1}^n (C_k \cap D_k) \subset \{xR > t\}$ and $C_k \in \mathcal{F}_{\sigma_k}$, this implies

$$\mathbb{P}(xR > t) \geq \sum_{k=1}^n \int_{C_k} \mathbb{P}_x(D_k | \mathcal{F}_{\sigma_k}) d\mathbb{P}_x \geq \eta \mathbb{P}_x \left(\bigcup_{k=1}^n C_k \right),$$

and thus $\mathbb{P}(xR > t) \geq \eta \mathbb{P}_x \left(\sup_{n \geq 1} (xQ^{\sigma_n} + \xi x\Pi_{\sigma_n} y) > t \right)$ by letting $n \rightarrow \infty$.

Turning to the respective inequality for $\mathbb{P}(-xR > t)$, define

$$C'_k := \left\{ \max_{1 \leq j < k} (-xQ^{\sigma_j} - \xi x\Pi_{\sigma_j} y) \leq t, -xQ^{\sigma_k} - \xi x\Pi_{\sigma_k} y > t \right\}$$

and $D'_k := \{-x\Pi_{\sigma_k} R^{\sigma_k} > -\xi x\Pi_{\sigma_k} y\} = \{(x\Pi_{\sigma_k})^\sim R^{\sigma_k} < \xi (x\Pi_{\sigma_k})^\sim y\}$.

Again by (29), $(x\Pi_{\sigma_k})^\sim y \geq 1 - \delta > 1 - \zeta$ for all $y \in B_\delta(x_0)$, giving

$$D'_k \supset \{(x\Pi_{\sigma_k})^\sim R^{\sigma_k} < (1 - \zeta)\xi\}$$

and thus $\mathbb{P}_x(D'_k | \mathcal{F}_{\sigma_k}) \geq \eta$ \mathbb{P}_x -a.s. Now reasoning as above,

$$\mathbb{P}(-xR > t) \geq \eta \lim_{n \rightarrow \infty} \mathbb{P}_x \left(\bigcup_{k=1}^n C'_k \right) = \eta \mathbb{P}_x \left(\sup_{n \geq 1} (-xQ^{\sigma_n} - \xi x\Pi_{\sigma_n} y) > t \right)$$

The desired result hence follows by a combination of this inequality with the one obtained for $\mathbb{P}(xR > t)$. \square

Proposition 8.3. *There exists $x \in S$ such that $\liminf_{t \rightarrow \infty} t^\kappa \mathbb{P}(|xR| > t)$ is positive.*

Proof. Pick any regenerative $B_\delta(x_0)$ with δ sufficiently small, such that Lemma 8.2 holds true, and let $\sigma_1, \sigma_2, \dots$ be the associated regeneration times, thus $X_{\sigma_n} \stackrel{d}{=} \phi$ for $n \geq 1$. By Proposition 5.13, $\liminf_{t \rightarrow \infty} t^\kappa \mathbb{P}_x \left(\sup_{n \geq 1} |x\Pi_{\sigma_{n-1}}| > t \right)$ is positive for π -almost all $x \in B_\delta(x_0)$. Fix any such x hereafter.

Define $\Pi_{j,k} := M_j \cdot \dots \cdot M_k$, $Q^{j,n} := \sum_{k=j}^n \Pi_{j,k-1} Q_k$ and (with $\sigma_0 := 0$ and any $y \in B_\delta(x_0)$, to be chosen in Lemma 8.4)

$$\begin{aligned} T_n &= xQ^{\sigma_n} + \xi x\Pi_{\sigma_n}y, \\ \Delta_n &:= Q^{\sigma_{n-1}+1, \sigma_n} - \xi(I - \Pi_{\sigma_{n-1}+1, \sigma_n})y, \\ U_n &= x\Pi_{\sigma_{n-1}}\Delta_n \end{aligned}$$

for $n \geq 1$. Then $T_n = T_{n-1} + U_n$ and $\{\sup_{n \geq 1} |T_n| > t\} \supset \{\sup_{n \geq 2} |U_n| > 2t\}$. Lemma 8.2 provides us with

$$\mathbb{P}(|xR| > t) \geq \eta \mathbb{P}_x \left(\sup_{n \geq 1} |T_n| > t \right)$$

for some $\eta > 0$.

By Lemma 5.6, $\inf_{y \in S} |yM_{\sigma_n}| \geq \mathbf{c}$ a.s. for all $n \geq 1$ and a suitable $\mathbf{c} > 0$. Hence, for all $t > 0$,

$$\begin{aligned} \mathbb{P}(|xR| > t) &\geq \eta \mathbb{P}_x \left(\sup_{n \geq 2} |U_n| \geq 2t \right) \\ &= \eta \mathbb{P}_x \left(\sup_{n \geq 1} |x\Pi_{\sigma_{n-1}}| |(x\Pi_{\sigma_{n-1}})^{\sim} M_{\sigma_n}| |X_{\sigma_n} \Delta_{n+1}| \geq 2t \right) \\ &\geq \eta \sum_{n \geq 1} \mathbb{P}_x \left(\bigcap_{k=1}^{n-1} A_k, |x\Pi_{\sigma_{n-1}}| > \frac{2t}{\mathbf{c}\varepsilon}, |X_{\sigma_n} \Delta_{n+1}| > \varepsilon \right) \\ &\geq \eta \sum_{n \geq 1} \mathbb{P}_x \left(\bigcap_{k=1}^{n-1} A_k, |x\Pi_{\sigma_{n-1}}| > \frac{2t}{\mathbf{c}\varepsilon} \right) \mathbb{P}_\phi(|X_0 \Delta_1| > \varepsilon) \quad (\text{use (R3)}) \\ &\geq \eta \mathbb{P}_\phi(|X_0 \Delta_1| > \varepsilon) \mathbb{P}_x \left(\sup_{n \geq 1} |x\Pi_{\sigma_{n-1}}| > \frac{2t}{\mathbf{c}\varepsilon} \right), \end{aligned}$$

where $A_k = \{|x\Pi_{\sigma_{k-1}}| \leq 2t/(\mathbf{c}\varepsilon)\}$ for $k \geq 1$ and some fixed $0 < \varepsilon < 1$. The proof is finished by the subsequent lemma where we show positivity of $\mathbb{P}_\phi(|X_0 \Delta_1| > \varepsilon)$. Together with (23) this clearly yields the desired conclusion. \square

Lemma 8.4. *In the situation of Proposition 8.3, there exist $\varepsilon > 0$ and $y \in B_\delta(x_0)$ such that (notice here the dependence of Δ_1 on y)*

$$\mathbb{P}_\phi(|X_0 \Delta_1| > \varepsilon) > 0.$$

Proof. Suppose that $X_0 \Delta_1 = X_0(Q^\sigma - \xi(I - \Pi_\sigma)y) = 0$ \mathbb{P}_ϕ -a.s. for all $y \in B_\delta(x_0)$, where $\sigma := \sigma_1$. Then the same holds true for all y in the convex hull of $B_\delta(x_0)$ (as a subset of \mathbb{R}^d) which contains a basis of \mathbb{R}^d . Consequently, the range of $Q^\sigma - \xi(I - \Pi_\sigma)$ and $\{tX_0 : t \in \mathbb{R}\}$ are orthogonal \mathbb{P}_ϕ -a.s. On the other hand, by Lemma 5.7, $\mathbb{P}_\phi(\|\Pi_\sigma\| < 1) > 0$ and thus $Q^\sigma - \xi(I - \Pi_\sigma)$ has full range \mathbb{R}^d on a set of positive probability under \mathbb{P}_ϕ . This contradicts our starting assumption and the lemma is proved. \square

We close this section with a proof of Lemma 8.1.

Proof of Lemma 8.1. We first show that $\text{supp } R$ is not a compact subset of \mathbb{R}^d . Use (7) to infer for each $n \geq 1$,

$$\Pi_n \text{supp } R + Q^n = \text{supp } R \quad \mathbb{P}\text{-a.s.}$$

and thus also ${}^\kappa\mathbb{P}$ -a.s., for \mathbb{P}_x and ${}^\kappa\mathbb{P}_x$ for any x are equivalent probability measures on each $\mathcal{F}_n = \sigma((M_j, Q_j)_{1 \leq j \leq n})$, $n \geq 1$. Now assume, that $\text{supp } R$ is bounded. By (A6), there exist at least two distinct $x_1, x_2 \in \text{supp } R$. Defining $v := x_1 - x_2$, it then follows that for all $n \geq 1$ and some $C \in (0, \infty)$

$$|\Pi_n v| \leq |\Pi_n x_1 + Q^n| + |\Pi_n x_2 + Q^n| < C \quad {}^\kappa\mathbb{P}\text{-a.s.}$$

and thereupon for all $x \in S$

$$C \geq |x \Pi_n v| = |x \Pi_n| |(x \Pi_n)^\sim v| \quad {}^\kappa\mathbb{P}\text{-a.s.}$$

The hitting times τ_n of $(x \Pi_n)^\sim$ in $B_\delta(v)$ are ${}^\kappa\mathbb{P}_x$ -a.s.-finite, yielding

$$\limsup_{n \rightarrow \infty} |x \Pi_{\tau_n}| \leq \frac{C}{X_{\tau_n} v} \leq \frac{C}{1 - \delta} \quad {}^\kappa\mathbb{P}_x\text{-a.s.}$$

for all $x \in S$, where (29) should be recalled for the final bound. Consequently,

$$\limsup_{n \rightarrow \infty} \frac{V_{\tau_n}}{\tau_n} = \limsup_{n \rightarrow \infty} \frac{1}{\tau_n} \log |X_0 \Pi_{\tau_n}| = 0 \quad {}^\kappa\mathbb{P}_\pi\text{-a.s.}$$

which contradicts Lemma 5.12.

Having thus shown that $\text{supp } R$ is not compact in \mathbb{R}^d , there exist sequences $(x_n)_{n \geq 1} \subset \text{supp } R$ with $\lim_{n \rightarrow \infty} |x_n| = \infty$ whence, by compactness of S , the following set is nonempty:

$$D := \left\{ y \in S : \exists (x_n)_{n \geq 1} \subset \text{supp } R, \lim_{n \rightarrow \infty} |x_n| = \infty, \lim_{n \rightarrow \infty} x_n^\sim = y \right\}.$$

Now suppose that $\mathbb{P}(x_0 R \leq t_0) = 1$ for some $(x_0, t_0) \in S \times \mathbb{R}$. For any $y_0 \in D$, choose a sequence $(x_n)_{n \geq 1} \subset \text{supp } R$ such that $x_n^\sim \rightarrow y_0$. It follows that $x_0 x_n < t_0$ for all n and thereby (since $|x_n| \rightarrow \infty$), that $x_0 y_0 \leq 0$ for all $y_0 \in D$. On the other hand, $(M x_n + Q)^\sim \rightarrow M y_0$ for the unbounded sequence $(M x_n + Q)_{n \geq 1}$ (which is \mathbb{P} -a.s. a subset of $\text{supp } R$) implies $M y_0 \in D$ \mathbb{P} -a.s. and therefore

$$\mathbb{P}(x_0 M y_0 \leq 0) = \mathbb{P}((x_0 M)^\sim y_0 \leq 0) = 1,$$

in particular $\mathbb{P}((x_0 M)^\sim \notin B_\delta(y_0)) = 0$ for sufficiently small $\delta > 0$ which is a contradiction to (A4) (with $n_0 = 1$). \square

References

- [1] G. Alsmeyer, *The Markov renewal theorem and related results*, Markov Proc. Rel. Fields 3 (1997), pp. 103–127.
- [2] K. Athreya and P. Ney, *A new approach to the limit theory of recurrent Markov chains*, Transactions of the American Mathematical Society 245 (1978), pp. 493–501.
- [3] P. Bougerol and J. Lacroix. *Products of random matrices with applications to Schrödinger operators*, volume 8 of *Progress in Probability and Statistics*. Birkhäuser Boston Inc., Boston, MA, 1985.

- [4] D. Buraczewski, E. Damek, Y. Guivarc'h, A. Hulanicki, and R. Urban, *Tail-homogeneity of stationary measures for some multidimensional stochastic recursions*, Probab. Theory Related Fields 145 (2009), pp. 385–420.
- [5] B. de Saporta, Y. Guivarc'h, and É. Le Page, *On the multidimensional stochastic equation $Y_{n+1} = A_n Y_n + B_n$* , Comptes Rendus Mathématique 339 (2004), pp. 499–502.
- [6] P. Diaconis and D. Freedman, *Iterated random functions*, SIAM Review 41 (1999), pp. 45–76.
- [7] N. Dunford and J.T. Schwartz, *Linear Operators, Part I, General Theory*, Wiley 1958.
- [8] J.H. Elton, *A multiplicative ergodic theorem for Lipschitz maps*, Stoch. Proc. Appl. 34 (1990), pp. 39–47.
- [9] C.M. Goldie, *Implicit renewal theory and tails of solutions of random equations*, Ann. Appl. Probab. 1 (1991), pp. 126–166.
- [10] Y. Guivarc'h, *Heavy tail properties of stationary solutions of multidimensional stochastic recursions*, in *Dynamics & stochastics*, IMS Lecture Notes Monogr. Ser., Vol. 48, Inst. Math. Statist., Beachwood, OH, 2006, pp. 85–99.
- [11] S. Karlin, *Positive operators*, J. Math. Mech. 8 (1959), pp. 907–937.
- [12] H. Kesten, *Random difference equations and renewal theory for products of random matrices*, Acta Math. 131 (1973), pp. 207–248.
- [13] H. Kesten, *Renewal Theory for Functionals of a Markov Chain with general state space*, The Annals of Probability 2 (1974), pp. 355–386.
- [14] C. Klüppelberg and S. Pergamenchtchikov, *The tail of the stationary distribution of a random coefficient AR(q) model*, Annals of Applied Probability 14 (2004), pp. 971–1005.
- [15] É. Le Page, *Théorèmes de renouvellement pour les produits de matrices aléatoires. Équations aux différences aléatoires*, in *Séminaires de probabilités Rennes 1983*, Univ. Rennes I, Rennes, 1983, p. 116.
- [16] E. Nummelin, *A splitting technique for Harris recurrent Markov chains*, Z. Wahrsch. Verw. Gebiete 43 (1978), pp. 309–318.