

A structure theorem for strong immersions

Zdeněk Dvořák*

Paul Wollan†

May 4, 2022

Abstract

A graph H is *strongly immersed* in G if H is obtained from G by a sequence of vertex splittings (i.e., lifting some pairs of incident edges and removing the vertex) and edge removals. Equivalently, vertices of H are mapped to distinct vertices of G (*branch vertices*) and edges of H are mapped to pairwise edge-disjoint paths in G , each of them joining the branch vertices corresponding to the ends of the edge and not containing any other branch vertices. We describe the structure of graphs avoiding a fixed graph as a strong immersion.

1 Introduction

In this paper, we consider graphs which can have parallel edges and loops, where each loop contributes 2 to the degree of the incident vertex. A graph without parallel edges and loops is called *simple*.

Various containment relations between graphs have been studied in structural graph theory. The best known ones are perhaps *minors* and *topological minors*. A graph H is a *minor* of G if it can be obtained from G by a sequence of edge and vertex removals and edge contractions. A graph H is a *topological minor* of G if a subdivision of H is a subgraph of G , or equivalently, if H can be obtained from G by a sequence of edge and vertex removals and by suppressions of vertices of degree two. In their fundamental series of papers, Robertson and Seymour developed the theory of graphs avoiding a fixed minor, giving a description of their structure [10] and proving that every proper minor-closed class of graphs is characterized by a finite set of forbidden minors [11]. The topological minor relation is somewhat harder to deal with (and in particular, there exist proper topological minor-closed classes that are not characterized by a finite set of forbidden topological minors), but a description of their structure is also available [6, 4].

In this paper, we consider the related notion of a graph immersion. Let H and G be graphs. An *immersion* of H in G is a function θ from vertices and edges of H such that

- $\theta(v)$ is a vertex of G for each $v \in V(H)$, and $\theta \upharpoonright V(H)$ is injective.
- $\theta(e)$ is a connected subgraph of G for each $e \in E(H)$, and if $f \in E(H)$ is distinct from e , then $\theta(e)$ and $\theta(f)$ are edge-disjoint.
- If $e \in E(H)$ is incident with $v \in V(H)$, then $\theta(v)$ is a vertex of $\theta(e)$, and if e is a loop, then $\theta(e)$ contains a cycle passing through $\theta(v)$.

An immersion θ is *strong* if it additionally satisfies the following condition:

*Computer Science Institute of Charles University, Prague, Czech Republic. E-mail: rakdver@iuuk.mff.cuni.cz. Supported the Center of Excellence – Inst. for Theor. Comp. Sci., Prague (project P202/12/G061 of Czech Science Foundation), and by project LH12095 (New combinatorial algorithms - decompositions, parameterization, efficient solutions) of Czech Ministry of Education.

†Department of Computer Science, University of Rome, “La Sapienza”, Rome, Italy wollan@di.uniroma1.it. Supported by the European Research Council under the European Union’s Seventh Framework Programme (FP7/2007-2013)/ERC Grant Agreement no. 279558.

- If $e \in E(H)$ is not incident with $v \in V(H)$, then $\theta(e)$ does not contain $\theta(v)$.

When we want to emphasize that an immersion does not have to be strong, we call it *weak*.

If H is a topological minor of G , then H is also strongly immersed in G . On the other hand, an appearance of H as a minor does not imply an immersion of H , and conversely, an appearance of H as a strong immersion does not imply the appearance as a minor or a topological minor. Nevertheless, many of the results for minors and topological minors have analogues for immersions and strong immersions. For example, any simple graph with minimum degree at least $200k$ contains a strong immersion of the complete graph K_k (DeVos et al. [2]), as compared to similar sufficient minimum degree conditions for minors ($\Omega(k\sqrt{\log k})$, Kostochka [8], Thomason [13]) and topological minors ($\Omega(k^2)$, Bollobás and Thomason [1], Komlós and Szemerédi [7]). Furthermore, every proper class of graphs closed on weak immersions is characterized by a finite set of forbidden immersions [12].

Let us restrict our attention for the moment to weak immersions. Fix t to be a positive integer. It is easy to show that if a graph G contains a set X of $t + 1$ vertices such that for every pair of vertices $x, y \in X$ there does not exist an edge cut of order less than t^2 separating x and y , then G contains K_t as a weak immersion. From this observation, we see that any graph which does not contain K_t as a weak immersion must either have a small number of large degree vertices, or alternatively, there exists a small edge cut separating two big degree vertices. This gives rise to an easy structure theorem for weak immersions as shown in DeVos et al. [3] and Wollan [14].

The same is not true for strong immersions. There exist graphs which are arbitrarily highly edge connected and still have no strong immersion of K_3 (although such graphs will necessarily not be simple by the extremal result of [2] mentioned above). As an example, let k be a positive integer and consider the graph P_k obtained from a path of length k by adding $k - 1$ parallel edges to each edge of the path. Then P_k is k edge connected but does not contain even K_3 as a strong immersion. The example can be made more complex as well. If we add edges to connect every pair of vertices at distance two on the path, the resulting graph will still have no strong immersion of K_4 . When the graph is assumed to be highly edge connected, this is essentially the only obstruction to a graph excluding a strong immersion of a fixed clique as shown by Marx and Wollan [9].

The main result of this article is a decomposition theorem for strong immersions. The basis is a theorem which says that if a graph avoids a strong immersion of a fixed clique and contains a large set X of vertices which are pairwise highly edge connected, then the graph must have a decomposition with respect to X yielding an obstruction to the existence of a strong clique immersion, similar in spirit to the construction of the highly edge connected graph avoiding a strong clique immersion described in the previous paragraph. From this decomposition theorem, it is straightforward to derive a structure theorem for graphs excluding a fixed graph as a strong immersion.

We first rigorously define the decomposition which arises. A *near-partition* of a set Z is a family of subsets Z_1, \dots, Z_k , possibly empty, such that $\bigcup_1^k Z_i = Z$ and $Z_i \cap Z_j = \emptyset$ for all $1 \leq i < j \leq k$. Let X be a subset of vertices of a graph G . A *path-like decomposition* \mathcal{P} of G with respect to X is an ordering x_1, x_2, \dots, x_t of the vertices of X and a near-partition B_0, \dots, B_t of $V(G) \setminus X$. The elements of the near-partition are called *bags* of the decomposition. For a vertex $x_i \in X$, the x_i -cut of the decomposition is the set of edges of G with one endpoint in $\{x_1, \dots, x_{i-1}\} \cup B_0 \cup \dots \cup B_{i-1}$ and the other endpoint in $\{x_{i+1}, \dots, x_t\} \cup B_i \cup \dots \cup B_t$. The *width* of the decomposition is the maximum size of an x -cut with $x \in X$. For a set $Z \subseteq V(G)$, let $b_{\mathcal{P}}(Z)$ denote the number of bags of the decomposition that intersect Z . For an integer $p \geq 0$, we say that Z is p -bounded in the decomposition \mathcal{P} when $|X \cap Z| + b_{\mathcal{P}}(Z) \leq p$. For an integer $k \geq 1$, we say that a set $W \subseteq V(G)$ is k -edge-connected if no two vertices of W are separated by an edge-cut of size less than k in G . A set $W \subseteq V(G)$ is α -linear if there exists a set $A \subseteq W$ of size at most α such that $G - A$ has a path-like decomposition \mathcal{P} with respect to $W \setminus A$ of width less than α and the neighborhood of every vertex of A is α -bounded in \mathcal{P} .

The following theorem is the main technical result of this article. It shows that every highly edge-connected set in a graph G is α -linear for some bounded value α if G does not contain a strong immersion of a fixed graph F .

Theorem 1. *For every graph F , there exists a value α such that if a graph G does not contain F as a strong immersion, then every α -edge-connected subset of $V(G)$ is α -linear.*

Conversely, we show as well that the property of being α -linear is a good approximate characterization of graphs excluding a fixed immersion in that every graph which satisfies this property for every highly edge connected set cannot contain a strong immersion of a big clique.

Assuming Theorem 1, we can derive a global structure theorem for graphs excluding a fixed graph as a strong immersion. The theorem and its proof are presented in Section 2. The proof of Theorem 1 is given in Section 3 and in the final section, we show that the converse statement to the decomposition and structure theorem are approximately true.

We establish some notation we will use going forward. Let G be a graph and $X \subseteq V(G)$. We use $G[X]$ to denote the subgraph of G induced by X . The graph $G - X$ refers to the subgraph of G induced on $V(G) \setminus X$. For a subset K of edges, $G - K$ is the subgraph with vertex set $V(G)$ and edge set $E(G) \setminus K$. We will use $\delta(X)$ to refer to the set of edges with one endpoint in X and one endpoint not in X (specifically, $\delta(X)$ does not contain any loops). A *separation* of G is a pair (X, Y) of non-empty subsets of $V(G)$ such that $X \cap Y = \emptyset$ and every edge has all its endpoints either contained in X or contained in Y .

2 A structure theorem

In this section, we show how Theorem 1 gives rise to a global structure theorem for graphs which exclude a fixed graph H as a strong immersion. We first present some further notation.

When studying graph minors, a natural decomposition is to break the graph on a small vertex cutset and look at the structure on each side of the cutset. This gives rise to the operation of clique sums on graphs. Given that graph immersions consist of a set of edge disjoint paths, it is natural to instead look at when the graph can be decomposed on a small edge cut. This motivates the definition of what we will refer to as edge sums in graphs.

Definition 1. *Let $G, G_1,$ and G_2 be graphs. Let $k \geq 1$ be a positive integer. The graph G is a k -edge sum of G_1 and G_2 if the following holds. There exist vertices $v_i \in V(G_i)$ such that $\deg(v_i) = k$ for $i = 1, 2$ and a bijection $\pi : \delta(v_1) \rightarrow \delta(v_2)$ such that G is obtained from $(G_1 - v_1) \cup (G_2 - v_2)$ by adding an edge from $x \in V(G_1) - v_1$ to $y \in V(G_2) - v_2$ for every pair e_1, e_2 of edges such that $e_1 \in \delta(v_1)$, $e_2 = \pi(e_1)$, the ends of e_1 are x and v_1 , and the ends of e_2 are y and v_2 .*

We will also refer to a k -edge sum as an edge sum of order k . The edge sum is grounded if there exist vertices v'_1 and v'_2 in G_1 and G_2 , respectively, such that for $i = 1, 2$, $v'_i \neq v_i$ and there exist k edge-disjoint paths linking v_i and v'_i . If G can be obtained by a k -edge sum of G_1 and G_2 , we write $G = G_1 \hat{\oplus}_k G_2$.

The following lemma appears in [14] and shows that edge sums preserve the presence of immersions.

Lemma 2 ([14]). *Let $G, G_1,$ and G_2 be graphs and let $k \geq 1$ be a positive integer. Assume $G = G_1 \hat{\oplus}_k G_2$, and assume that the edge sum is grounded. Let H be an arbitrary graph. If G_1 or G_2 admits an immersion of H , then G does as well. If the immersion in either G_1 or G_2 is strong, then the immersion in G is also strong.*

Just as clique sums give rise to tree decompositions, edge sums yield a natural tree-like decomposition of graphs.

Definition 2. *A tree-cut decomposition of a graph G is a pair (T, \mathcal{X}) such that T is a tree and $\mathcal{X} = \{X_t \subseteq V(G) : t \in V(T)\}$ is a near-partition of the vertices of G indexed by the vertices of T . For each edge $e = uv$ in T , $T - uv$ has exactly two components, namely T_v and T_u containing v and u respectively. The adhesion of the decomposition is*

$$\max_{uv \in E(T)} \left| \delta \left(\bigcup_{t \in V(T_v)} X_t \right) \right|$$

when T has at least one edge, and 0 otherwise. The sets $\{X_t : t \in V(T)\}$ are called the bags of the decomposition.

Note that the definition allows bags to be empty.

We will need to define one more operation on graphs. Let G be a graph and $X \subseteq V(G)$. The graph G' is obtained by *consolidating* X if we identify the vertices of X to a single vertex v_X and delete all loops incident to v_X .

Let G be a graph and (T, \mathcal{X}) a tree-cut decomposition of G . Fix a vertex $t \in V(T)$. The *torso of* (G, T, \mathcal{X}) at t is the graph H defined as follows. If $|V(T)| = 1$, then the torso H of (G, T, \mathcal{X}) at t is simply G itself. If $|V(T)| \geq 2$, let the components of $T - t$ be T_1, \dots, T_l for some positive integer l . Let $Z_i = \bigcup_{x \in V(T_i)} X_x$ for $1 \leq i \leq l$. Then H is made by consolidating each set Z_i to a single vertex z_i . The vertices X_t are called the *core vertices* of the torso. The vertices z_i are called the *peripheral vertices* of the torso. When there can be no confusion as to the graph G in question, we will also refer to the torso of (T, \mathcal{X}) at a vertex t .

The following lemma shows that tree cut decompositions can be combined in an edge sum of graphs.

Lemma 3 ([14]). *Let $G, G_1,$ and G_2 be graphs such that $G = G_1 \hat{\oplus}_k G_2$ for some $k \geq 0$. If G_i has a tree-cut decomposition (T_i, \mathcal{X}_i) for $i = 1, 2$, then G has a tree-cut decomposition (T, \mathcal{Y}) such that the adhesion of (T, \mathcal{Y}) is equal to*

$$\max\{k, \text{adhesion}(T_1, \mathcal{X}_1), \text{adhesion}(T_2, \mathcal{X}_2)\}.$$

Moreover, for every $t \in V(T)$, there exists $i \in \{1, 2\}$ and a vertex t' in $V(T_i)$ such that the torso H_t of (G, T, \mathcal{Y}) at t is isomorphic to the torso H' of $(G_i, T_i, \mathcal{X}_i)$ at t' . Finally, every core vertex of H_t is a core vertex of H' .

We can now state the structure theorem for graphs excluding a fixed clique immersion in terms of a tree-cut decomposition. The proof will follow easily assuming Theorem 1. We say that a graph is α -*basic* if the set of all its vertices of degree at least α is α -linear, i.e., it has a path-like decomposition such that all the vertices in its bags have degree less than α .

Theorem 4. *For every graph F , there exists an integer $\alpha = \alpha(F)$ such that if a graph G does not contain F as a strong immersion, then there exists a tree-cut decomposition (T, \mathcal{X}) of G of adhesion less than α such that each torso is α -basic.*

Proof (assuming Theorem 1). Fix the graph F and let α be the value given in Theorem 1. Assume the statement is false, and let G be a counterexample on a minimum number of edges. The set of vertices of degree at least α in G is not α -edge-connected, as otherwise Theorem 1 yields a contradiction. Thus, we may assume that there exists vertices x and y each of degree at least α such that there exists $X \subseteq V(G)$ with $x \in X, y \notin X$ and $|\delta(X)| < \alpha$.

Let G_X be the graph obtained by consolidating $V(G) \setminus X$ and G_Y the graph obtained by consolidating X . By construction, $G = G_X \hat{\oplus}_k G_Y$ for some positive integer $k < \alpha$, and if we assume we chose a minimum order edge cut separating x and y , the edge sum is grounded. Given that both x and y have degree at least α , we see that $|E(G_X)| < |E(G)|$ and $|E(G_Y)| < |E(G)|$. By Lemma 2, neither G_X nor G_Y contains F as a strong immersion. Both G_X and G_Y have the desired decomposition by minimality, and therefore, by Lemma 3, G has the desired decomposition as well. \square

3 Proof of Theorem 1

We prove a slightly stronger statement which gives a clearer picture on the relationship between the parameters involved. Let G be a graph and $a, w, p \geq 1$ be positive integers. A set $W \subseteq V(G)$ is (a, w, p) -*linear* if there exists a set $A \subseteq W$ of size at most a such that $G - A$ has a path-like decomposition \mathcal{P} with respect to $W \setminus A$ of width less than w and the neighborhood of every vertex of A is p -bounded in \mathcal{P} .

Theorem 5. *For every graph F , there exist integers a , w and p such that if a graph G does not contain F as a strong immersion, then every w -edge-connected subset of $V(G)$ is (a, w, p) -linear.*

To see that Theorem 5 is in fact a strengthening of Theorem 1, we observe the following. Assume that a subset W of vertices in a graph G is (a, w, p) -linear for positive integers a, w, p . Then a path-like decomposition of G with respect to W which certifies that W is (a, w, p) -linear trivially certifies as well that W is (a', w', p') -linear for all $a' \geq a$, $w' \geq w$, and $p' \geq p$. Thus, W is α -linear for $\alpha = \max\{a, w, p\}$, implying that Theorem 1 is an immediate consequence of Theorem 5.

For the remainder of this section, we define

$$d(k) = (2k + 1)^{8k+4} k^2 (k + 1).$$

We use the following result of Dvořák and Klímašová [5].

Theorem 6 ([5]). *Let G be a graph and $x \in V(G)$. Let $k \geq 3$ be an integer. Let $Y \subseteq V(G) \setminus \{x\}$ be a set of vertices such that G contains no edge cut of size less than k separating x from a vertex in Y . If a graph F of maximum degree at most k does not appear in a graph G as a strong immersion, then there exist sets $Y' \subseteq Y$ and $K \subseteq E(G)$ such that $k|Y'| + |K| < d(k)|V(F)|$ and the component of $(G - Y') - K$ that contains x does not contain any vertex of Y .*

For a graph G , a set $W \subseteq V(G)$ and an integer m , let $G(m, W)$ denote the graph with vertex set W such that two vertices x and y in W are adjacent in $G(m, W)$ iff $G - (W \setminus \{x, y\})$ contains at least m pairwise edge-disjoint paths joining x with y . As a corollary of Theorem 6, Dvořák and Klímašová [5] proved that if W is sufficiently edge-connected and sufficiently large and G avoids some fixed graph as a strong immersion, then $G(m, W)$ is connected. We need a strengthening of this claim.

Lemma 7. *Let F be a graph and let $k = \max\{\Delta(F), 3\}$. For all integers $a_0, m \geq 0$, there exists $w \geq k$ such that the following holds. Let G be a graph, let $W \subseteq V(G)$ be a w -edge-connected set and let A be a subset of W of size at most a_0 . Suppose that $G(m, W) - A$ is not connected and let (X_1, X_2) be a separation of $G(m, W) - A$. If G does not contain F as a strong immersion, then $G - A$ contains an edge-cut of size less than w separating X_1 from X_2 .*

Proof. Let $s = d(k)|V(F)|$, $w_0 = ms^3 + s^2$ and $w = \max\{2a_0sw_0, w_0 + a_0s\}$.

Claim 1. *For $i \in \{1, 2\}$, every vertex $x \in X_i$ is separated from X_{3-i} by an edge-cut of size less than w_0 in $G - A$.*

Proof. Suppose the claim is false. By symmetry, we can assume that $i = 1$. Apply Theorem 6 in G for x and X_2 , obtaining sets $Y \subseteq X_2$ and $K_0 \subseteq E(G)$, where $k|Y| + |K_0| < s$, such that the component of $(G - Y) - K_0$ that contains x does not contain any vertex of X_2 . For each $y \in Y$, apply Theorem 6 for y and X_1 , obtaining sets $Y_y \subseteq X_1$ and $K_y \subseteq E(G)$, where $k|Y_y| + |K_y| < s$, such that the component of $(G - Y_y) - K_y$ that contains y does not contain any vertex of X_1 . Let $K = K_0 \cup \bigcup_{y \in Y} K_y$ and let $Z = \bigcup_{y \in Y} Y_y$, and note that $|K| \leq s^2$ and $|Z| \leq s^2$.

By Menger's theorem, there exists a set S_0 of w_0 pairwise edge-disjoint paths from x to X_2 in $G - A$. Let $S \subseteq S_0$ consist of the paths that do not contain edges of K ; we have $|S| \geq w_0 - s^2$. Consider a path $P \in S$. Let v_0, v_1, \dots, v_t be the vertices of P in order, where $v_0 = x$ and $v_t \in X_2$. As the component of $(G - Y) - K_0$ that contains x does not contain any vertex of X_2 , the vertex v_t belongs to Y . Let j be the largest index such that v_j belongs to X_1 . As the component of $(G - Y_{v_t}) - K_{v_t}$ that contains v_t does not contain any vertex of X_1 , it follows that v_j belongs to $Y_{v_t} \subseteq Z$. Consequently, $G - A$ contains a set of $|S|$ pairwise edge-disjoint paths joining vertices of Y with vertices of Z and otherwise disjoint from W . By the pigeonhole principle, there exist vertices $u \in Z \setminus A \subseteq X_1$ and $v \in Y \setminus A \subseteq X_2$ contained in at least $\frac{|S|}{|Y||Z|} \geq m$ of these paths, and thus uv is an edge of $G(m, W)$. This contradicts the assumption that (X_1, X_2) is a separation of $G(m, W) - A$. \square

Consider any vertex $v \in X_1$. Since W is w -edge-connected, G contains at least w pairwise edge-disjoint paths from v to X_2 . By Claim 1, at least $w - w_0 + 1$ of these paths pass through a vertex of A . By pigeonhole principle, we have the following.

Claim 2. *For every $v \in X_1$, there exists a vertex $a_v \in A$ such that $G - (X_2 \cup (A \setminus \{a_v\}))$ contains at least s pairwise edge-disjoint paths from v to a_v .*

Suppose that the lemma is false and that $G - A$ does not contain an edge-cut of size less than w separating X_1 from X_2 . By Menger's theorem, G contains a set S_1 of w pairwise edge-disjoint paths from X_1 to X_2 and otherwise disjoint from W . By Claim 1, each vertex of X_2 is incident with less than w_0 of these paths, and thus we can select $S_2 \subseteq S_1$ such that $|S_2| \geq |S_1|/w_0 \geq 2a_0s$ and the paths in S_2 have pairwise distinct ends in X_2 . Furthermore, we can select $z \in A$ and $S_3 \subseteq S_2$ of size at least $|S_2|/a_0 \geq 2s$ such that every $v \in X_1$ incident with a path in S_3 satisfies $a_v = z$.

Let U be the set of endpoints of the paths of S_3 in X_2 and note that $|U| = |S_3| \geq 2s$. Consider any sets $U' \subseteq U$ and $K \subseteq E(G)$ such that $k|U'| + |K| < s$. We have $|U'| < s$, and thus $|U \setminus U'| \geq s$. Since $|K| < s$, there exists a path $P \in S_3$ ending in $U \setminus U'$ and disjoint with K . Let v be the endpoint of P in X_1 . By Claim 2, there exists a path from v to z disjoint with K and U' . Therefore, $(G - U') - K$ contains a path from z to U . Since this holds for every $U' \subseteq U$ and $K \subseteq E(G)$ with $k|U'| + |K| < s$, we obtain a contradiction with Theorem 6. \square

Furthermore, avoiding a strong immersion of a fixed graph restricts the structure of $G(m, W)$, as long as m is large enough.

Lemma 8. *Let F and G be graphs, let W be a subset of $V(G)$ and let m be an integer. If $m \geq 2|E(F)|$ and G does not contain F as a strong immersion, then $G(m, W)$ does not contain $K_{1,|V(F)|}$ as a minor.*

Proof. The claim is trivial if $|V(F)| \leq 1$. Suppose that $|V(F)| \geq 2$ and that $G(m, W)$ contains $K_{1,|V(F)|}$ as a minor, that is, $G(m, W)$ contains a subtree T with $|V(F)|$ leaves and at least one non-leaf vertex. Let c be a non-leaf vertex of T and let Z be the set of leaves of T . Let θ be an injective function mapping $V(F)$ to Z . By the definition of $G(m, W)$ and Menger's theorem, there exists a set S of $2|E(F)|$ pairwise edge-disjoint paths in G from Z to c , such that every vertex $z \in Z$ is contained in exactly $\deg_F(\theta^{-1}(z))$ of these paths. A *half-edge* of F is a pair (u, e) , where e is an edge of F and u is incident with e . Note that there exists a bijective function f from half-edges of F to S such that the path $f(u, e)$ contains the vertex $\theta(u)$ for every half-edge (u, e) . We extend θ to a strong immersion of F in G by defining $\theta(e) = f(u, e) + f(v, e)$ for every edge $e = uv$ of F . \square

The next lemma gives an approximate characterization of when a graph does not contain a large $K_{1,k}$ minor. A set X of vertices of a graph G is a *linearizing set* if $G - X$ is a vertex-disjoint union of paths.

Lemma 9 ([9]). *If a simple connected graph G does not contain $K_{1,k}$ as a minor, then G has a linearizing vertex set of size at most $4k$.*

We now give the proof of Theorem 5.

Proof of Theorem 5. Let $m = 2|E(F)|$, $a = 4|V(F)|$, $a_0 = a + 1$ and let w be the corresponding constant from Lemma 7. Let $k = \max(\Delta(F), 3)$ and $p = 3d(k)|V(F)| + 1$.

Let W be a w -edge-connected subset of $V(G)$, and let $H = G(m, W)$. If H is not connected, then there exists a separation (X_1, X_2) of H . By Lemma 7 applied with $A = \emptyset$, it follows that G contains an edge-cut of size less than w separating X_1 from X_2 . This is a contradiction, since W is w -edge-connected.

Therefore, H is a connected simple graph, and by Lemma 8, H does not contain $K_{1,|V(F)|}$ as a minor. Let A be the smallest linearizing set in H . By Lemma 9, we have $|A| \leq a$. Let x_1, x_2, \dots, x_t be an ordering of $W \setminus A$ such that for $2 \leq i \leq t - 1$, the neighbors of x_i in $H - A$ are contained in $\{x_{i-1}, x_{i+1}\}$.

If $t = 0$, we set $B_0 = V(G) \setminus A$. If $t = 1$, we set $B_0 = \emptyset$ and $B_1 = V(G) \setminus (A \cup \{x_1\})$. If $t = 2$, we set $B_0 = B_2 = \emptyset$ and $B_1 = V(G) \setminus (A \cup \{x_1, x_2\})$. In all the cases, we obtain a path-like decomposition of $G - A$ with respect to $W \setminus A$ of width 0, and since $p \geq 3$, the neighborhood of every vertex of A is p -bounded. Therefore, assume that $t \geq 3$.

Consider an index i such that $2 \leq i \leq t - 1$ and let $U_i = \{x_1, \dots, x_{i-1}\}$ and $V_i = \{x_{i+1}, \dots, x_t\}$. A set $Z \subset V(G) \setminus A$ is an i -separator if $x_i \notin Z$, $U_i \subseteq Z$ and $V_i \cap Z = \emptyset$. We set $s_i(Z)$ to be the number of edges of $G - A$ with one end in Z and the other end in $V(G - A) \setminus (Z \cup \{x_i\})$. Let $L_i \subset V(G) \setminus A$ be an i -separator with $s_i(L_i)$ as small as possible, and subject to that with $|L_i|$ minimal. By Lemma 7 applied with $A \cup \{x_i\}$, we have $s_i(L_i) < w$.

Claim 3. *If $2 \leq i < j \leq t - 1$, then $L_i \subset L_j$.*

Proof. Consider indices i and j such that $2 \leq i < j \leq t - 1$. Let $N_i = L_i \cap L_j$ and $N_j = L_i \cup L_j$. Note that N_i is an i -separator and N_j is a j -separator, and thus $s_i(N_i) \geq s_i(L_i)$ and $s_j(N_j) \geq s_j(L_j)$. Observe that $s_i(L_i) + s_j(L_j) - (s_i(N_i) + s_j(N_j)) = 2b_1 + b_2$, where b_1 is the number of edges with one end in $L_j \setminus (L_i \cup \{x_i\})$ and the other end in $L_i \setminus L_j$, and b_2 is the number of edges incident with x_i or x_j and with the other end in $L_i \setminus L_j$. Consequently, $s_i(L_i) + s_j(L_j) \geq s_i(N_i) + s_j(N_j)$. Putting the inequalities together, we have $s_i(N_i) = s_i(L_i)$ and $s_j(N_j) = s_j(L_j)$. Since L_i is chosen with $|L_i|$ minimal, it follows that $|N_i| \geq |L_i|$. Therefore, $L_i \subseteq L_j$. Furthermore, the inclusion is sharp, since $x_i \in L_j \setminus L_i$. \square

Let $L_1 = \emptyset$ and $L_t = V(G) \setminus (A \cup \{x_t\})$. Let $B_0 = B_t = \emptyset$, and for $1 \leq i \leq t - 1$, let us set $B_i = L_{i+1} \setminus (L_i \cup \{x_i\})$. By Claim 3, B_0, \dots, B_t is a near-partition of $V(G) \setminus W$, and thus the ordering x_1, \dots, x_t and the sets B_0, \dots, B_t form a path-like decomposition \mathcal{P} of $G - A$ with respect to $W \setminus A$. Since $s_i(L_i) < w$ for $2 \leq i \leq t - 1$, the width of \mathcal{P} is less than w .

Claim 4. *For $1 \leq i \leq t - 2$, each component of $G[B_i]$ contains a neighbor of x_i or x_{i+1} .*

Proof. Suppose that C is the vertex set of a component of $G[B_i]$ containing neighbors of neither x_i nor x_{i+1} . We say that an edge of $G - A$ with one end in C and the other end not in C is *backward* if its end not in C belongs to L_{i+1} , and it is *forward* otherwise. Note that neither forward nor backward edges are incident with x_{i+1} .

Let $L'_{i+1} = L_{i+1} \setminus C$ and note that L'_{i+1} is an $(i + 1)$ -separator. Since $|L'_{i+1}| < |L_{i+1}|$, the choice of L'_{i+1} implies that $s_{i+1}(L'_{i+1}) > s_{i+1}(L_{i+1})$, and thus there are more backward edges than forward ones. Since $B_i = L_{i+1} \setminus (L_i \cup \{x_i\})$ and C induces a component of $G[B_i]$ containing no neighbors of x_i , all backward edges are incident with vertices in L_i . Since $L_1 = \emptyset$, we have $i \geq 2$. However, then $L'_i = L_i \cup C$ is an i -separator with $s_i(L'_i) < s_i(L_i)$, which is a contradiction. \square

To complete the proof, we must show that every vertex of A is p -bounded in \mathcal{P} . Suppose that the neighborhood of a vertex $c \in A$ is not p -bounded. By Claim 4, G contains at least $(p - 1)/3 = d(k)|V(F)|$ paths from c to vertices of $W \setminus A$ whose vertex sets pairwise intersect only in c . Let X denote the set of their endpoints in $W \setminus A$. Suppose that sets $Y \subseteq X$ and $K \subseteq E(G)$ have the property that the component of $(G - Y) - K$ that contains c does not contain any vertex of X . Then each of the paths from c to X contains either a vertex of Y or an edge of K , and thus $k|Y| + |K| \geq d(k)|V(F)|$. This contradicts Theorem 6. \square

4 An approximate converse

We conclude by showing that the decomposition guaranteed by Theorem 5 does indeed give a good approximation of graphs excluding strong clique immersions.

Theorem 10. *For all integers d, a, w and p , there exists an integer n such that if every d -edge-connected subset of $V(G)$ is (a, w, p) -linear, then G does not contain K_n as a strong immersion.*

Proof. Let $n = \max\{2w + 2, (2p + 1)a + 1, d + 1\}$, and suppose that θ is a strong immersion of K_n in G . Let V be the vertex set of K_n and let $W = \theta(V)$. Since $n - 1 \geq d$, W is d -edge-connected in G , and thus it is (a, w, p) -linear. Let A be a subset of W of size at most a such that $G - A$ has a path-like decomposition \mathcal{P} with respect to $W \setminus A$ of width less than w and the neighborhood of every vertex of A is p -bounded in \mathcal{P} . Let x_1, \dots, x_m be the ordering of $W \setminus A$ according to \mathcal{P} and let B_0, \dots, B_m be the bags of \mathcal{P} . Since $m \geq n - a \geq 2|A|p + 1$ and the neighborhood of every vertex of A in $G - A$ is p -bounded in \mathcal{P} , there exists an index $1 \leq i \leq m$ such that the vertices of A have no neighbors in $\{x_i\} \cup B_{i-1} \cup B_i$. Let K be the union of the x_{i-1} -cut and the x_{i+1} -cut of \mathcal{P} . Consider an edge e of K_n incident with $\theta^{-1}(x_i)$ such that e is incident neither with $\theta^{-1}(x_{i-1})$ nor $\theta^{-1}(x_{i+1})$. Since the immersion is strong, $\theta(e)$ contains neither x_{i-1} nor x_{i+1} , and thus $\theta(e)$ contains an edge of K . Therefore, there are at most $|K| \leq 2(w - 1)$ such edges incident with x_i , which is a contradiction since $n \geq 2w + 2$. \square

Similarly, the structure described in Theorem 4 is sufficient to exclude a large clique as a strong immersion.

Theorem 11. *For every integer $\alpha \geq 1$ there exists an integer $n \geq 1$ such that if a graph G has a tree-cut decomposition (T, \mathcal{X}) of adhesion less than α such that each torso is α -basic, then G does not contain K_n as a strong immersion.*

Proof. Let $n = 2\alpha^2 + 2\alpha + 1$, and suppose that θ is a strong immersion of K_n in G . Let V be the vertex set of K_n and let $W = \theta(V)$. The set W is α -edge-connected in G , and since (T, \mathcal{X}) has adhesion less than α , we conclude that there exists $t \in V(T)$ such that $W \subseteq X_t$. Let H be the torso of (G, T, \mathcal{X}) at t , and observe that H also contains K_n as a strong immersion. However, we can now obtain a contradiction in the same way as in the proof of Theorem 10. \square

References

- [1] B. BOLLOBÁS AND A. THOMASON, *Proof of a conjecture of Mader, Erdős and Hajnal on topological complete subgraphs*, European J. Combin., 19 (1998), pp. 883–887.
- [2] M. DEVOS, Z. DVOŘÁK, J. FOX, J. McDONALD, B. MOHAR, AND D. SCHEIDE, *Minimum degree condition forcing complete graph immersion*, ArXiv e-prints, 1101.2630 (2011).
- [3] M. DEVOS, J. McDONALD, B. MOHAR, AND D. SCHEIDE, *A note on forbidding clique immersions*, ArXiv e-prints, 1207.2117 (2012).
- [4] Z. DVOŘÁK, *A stronger structure theorem for excluded topological minors*, ArXiv e-prints, 1209.0129 (2012).
- [5] Z. DVOŘÁK AND T. KLIMOŠOVÁ, *Strong immersions and maximum degree*, ArXiv e-prints, 1304.0728 (2013).
- [6] M. GROHE AND D. MARX, *Structure theorem and isomorphism test for graphs with excluded topological subgraphs*, CoRR, abs/1111.1109 (2011).
- [7] J. KOMLÓS AND E. SZEMERÉDI, *Topological cliques in graphs. II*, Combin. Probab. Comput., 5 (1996), pp. 79–90.
- [8] A. KOSTOCHKA, *Lower bound of the Hadwiger number of graphs by their average degree*, Combinatorica, 4 (1984), pp. 307–316.
- [9] D. MARX AND P. WOLLAN, *Immersions in highly edge connected graphs*, SIAM Journal on Discrete Mathematics, 28 (2014), pp. 503–520.

- [10] N. ROBERTSON AND P. D. SEYMOUR, *Graph Minors. XVI. Excluding a non-planar graph*, J. Combin. Theory, Ser. B, 89 (2003), pp. 43–76.
- [11] ———, *Graph Minors. XX. Wagner’s conjecture*, J. Combin. Theory, Ser. B, 92 (2004), pp. 325–357.
- [12] ———, *Graph Minors. XXIII. Nash-Williams’ immersion conjecture*, J. Combin. Theory, Ser. B, 100 (2010), pp. 181–205.
- [13] A. THOMASON, *The extremal function for complete minors*, J. Combin. Theory, Ser. B, 81 (2001), pp. 318–338.
- [14] P. WOLLAN, *The structure of graphs not admitting a fixed immersion*, Journal of Combinatorial Theory, Series B, (2014), pp. —.