

# MODIFICATIONS OF TUTTE-GROTHENDIECK INVARIANTS AND TUTTE POLYNOMIALS

MARTIN KOCHOL

ABSTRACT. We transform Tutte-Grothendieck invariants thus also Tutte polynomials on matroids so that the contraction-deletion rule for loops (isthmuses) coincides with the general case.

## 1. INTRODUCTION

A *Tutte-Grothendieck invariant* (shortly a *T-G invariant*)  $\Phi$  is a mapping from the class of finite matroids to a commutative ring  $(R, +, \cdot, 0, 1)$  such that  $\Phi(M) = \Phi(M')$  if  $M$  is isomorphic to  $M'$  and there are constants  $\alpha_1, \beta_1, \alpha_2, \beta_2 \in R$  such that

$$(1) \quad \begin{array}{ll} \Phi(M) = 1 & \text{if the ground set of } M \text{ is empty,} \\ \Phi(M) = \alpha_1 \cdot \Phi(M - e) & \text{if } e \text{ is an isthmus of } M, \\ \Phi(M) = \beta_1 \cdot \Phi(M - e) & \text{if } e \text{ is a loop of } M, \\ \Phi(M) = \alpha_2 \cdot \Phi(M/e) + \beta_2 \cdot \Phi(M - e) & \text{otherwise,} \end{array}$$

for every matroid  $M$  and every element  $e$  of  $M$ . We also say that  $\Phi$  is *determined* by the 4-tuple  $(\alpha_1, \beta_1, \alpha_2, \beta_2)$ . In certain sense (see [7, 2]), all T-G invariants can be reduced from the *Tutte polynomial* of  $M$

$$(2) \quad T(M; x, y) = \sum_{A \subseteq E} (x - 1)^{r(M) - r(A)} (y - 1)^{|A| - r(A)},$$

where  $E$  and  $r$  denote the ground set and rank function of  $M$ , respectively. This is very important invariant that encodes many properties of graphs and has applications in combinatorics, knot theory, statistical physics and coding theory (see cf. [1, 2, 9]).

$M - e = M/e$  if  $e$  is a loop or an isthmus of  $M$ . Thus the second (third) row of (1) is contained in the fourth row if  $\alpha_1 = \alpha_2 + \beta_2$  ( $\beta_1 = \alpha_2 + \beta_2$ ). In this case  $\Phi$  is called an *isthmus-smooth* (*loop-smooth*) T-G invariant.

We show that any T-G invariant can be transformed to an isthmus- and loop-smooth T-G invariants. The transformations are studied in framework of matroid duality. Furthermore, we discuss modifications of covariance and convolution formulas known for the Tutte polynomial. Notice that transformations into isthmus-smooth invariants are used by decomposition algorithms of T-G invariants in [5].

## 2. GENERAL MODIFICATIONS

**Lemma 1.** *Let  $\alpha_1, \beta_1, \alpha_2, \beta_2$  be arbitrary elements of a commutative ring  $(R, +, \cdot, 0, 1)$ . Then  $\tilde{T}(M; \alpha_1, \beta_1, \alpha_2, \beta_2) = \alpha_2^{r(M)} \beta_2^{r^*(M)} T(M; \alpha_1/\alpha_2, \beta_1/\beta_2)$  is the unique  $T$ - $G$  invariant determined by  $(\alpha_1, \beta_1, \alpha_2, \beta_2)$ .*

*Proof.* For any matroid  $M$ , denote  $\Phi(M) = \tilde{T}(M; \alpha_1, \beta_1, \alpha_2, \beta_2)$  (interpreting the formula as the substitution  $x_2 = \alpha_2, y_2 = \beta_2$  in the polynomial  $\tilde{T}(M; x_1, y_1, x_2, y_2)$ ). We use that  $(T; x, y)$  is determined by  $(x, y, 1, 1)$  and induction on  $|E|$ . The statement of lemma holds true if  $|E| = 0$ , otherwise choose  $e \in E$ . If  $e$  is an isthmus of  $M$ , then

$$\begin{aligned} \Phi(M) &= \alpha_2^{r(M)} \beta_2^{r^*(M)} T(M; \alpha_1/\alpha_2, \beta_1/\beta_2) = \\ &\alpha_2^{r(M-e)+1} \beta_2^{r^*(M-e)} \alpha_1/\alpha_2 T(M-e; \alpha_1/\alpha_2, \beta_1/\beta_2) = \alpha_1 \Phi(M-e) \end{aligned}$$

by induction hypothesis. If  $e$  is a loop of  $M$ , then

$$\Phi(M) = \alpha_2^{r(M-e)} \beta_2^{r^*(M-e)+1} \beta_1/\beta_2 T(M-e; \alpha_1/\alpha_2, \beta_1/\beta_2) = \beta_1 \Phi(M-e).$$

If  $e$  is neither a loop nor an isthmus of  $M$ , then

$$\begin{aligned} \Phi(M) &= \alpha_2^{r(M/e)+1} \beta_2^{r^*(M/e)} T(M/e; \alpha_1/\alpha_2, \beta_1/\beta_2) + \\ &\alpha_2^{r(M-e)} \beta_2^{r^*(M-e)+1} T(M-e; \alpha_1/\alpha_2, \beta_1/\beta_2) = \alpha_2 \Phi(M/e) + \beta_2 \Phi(M-e). \end{aligned}$$

This proves the statement.  $\square$

Lemma 1 also follows from results of Oxley and Welsh [7] (see [2, Corollary 6.2.6]).

**Theorem 1.** *Let  $\Phi$  be a  $T$ - $G$  invariant determined by  $(\alpha_1, \beta_1, \alpha_2, \beta_2)$ ,  $\beta_2 \neq 0$ , and  $\xi \in R$  be a multiple of  $\beta_2$ . Then  $\Phi_\xi^{\text{is}}(M) = \xi^{|E|} \left( \frac{\alpha_1 - \alpha_2}{\beta_2} \right)^{r^*(M)} \Phi(M)$  is an isthmus-smooth  $T$ - $G$  invariant such that for every matroid  $M$ ,*

$$\begin{aligned} \Phi_\xi^{\text{is}}(M) &= 1 && \text{if } E = \emptyset, \\ \Phi_\xi^{\text{is}}(M) &= \xi \beta_1 (\alpha_1 - \alpha_2) / \beta_2 \Phi_\xi^{\text{is}}(M-e) && \text{if } e \text{ is a loop of } M, \\ \Phi_\xi^{\text{is}}(M) &= \xi \alpha_2 \Phi_\xi^{\text{is}}(M/e) + \xi (\alpha_1 - \alpha_2) \Phi_\xi^{\text{is}}(M-e) && \text{otherwise.} \end{aligned}$$

*Proof.* By Lemma 1,  $\Phi(M) = \alpha_2^{r(M)} \beta_2^{r^*(M)} T(M; \alpha_1/\alpha_2, \beta_1/\beta_2)$  for each matroid  $M$ . Setting  $\zeta = \xi \left( \frac{\alpha_1 - \alpha_2}{\beta_2} \right)$  and using equality  $|E| = r(M) + r^*(M)$ , we get

$$\Phi_\xi^{\text{is}}(M) = \xi^{r(M)} \zeta^{r^*(M)} \Phi(M) = (\xi \alpha_2)^{r(M)} (\zeta \beta_2)^{r^*(M)} T(M; \frac{\xi \alpha_1}{\xi \alpha_2}, \frac{\zeta \beta_1}{\zeta \beta_2}),$$

whence by Lemma 1,  $\Phi_\xi^{\text{is}}$  is a  $T$ - $G$  invariant determined by  $(\xi \alpha_1, \zeta \beta_1, \xi \alpha_2, \zeta \beta_2)$ . Furthermore,  $\xi \alpha_1 = \xi \alpha_2 + \zeta \beta_2$ , i.e.,  $\Phi_\xi^{\text{is}}$  is an isthmus-smooth  $T$ - $G$  invariant.  $\square$

$\Phi_\xi^{\text{is}}$  is called the  $\xi$ -*isthmus-smooth modification* of  $\Phi$ . Notice that if  $\Phi$  is an isthmus-smooth invariant (i.e., if  $\alpha_1 = \alpha_2 + \beta_2$ ), then  $\Phi_\xi^{\text{is}}(M) = \xi^{|E|} \Phi(M)$  for every matroid  $M$ .

If  $R$  has zero divisors, then  $\xi/\beta_2$  does not need to be unique. In this case we should formally replace fraction  $\xi/\beta_2$  by  $\xi'$  where  $\xi = \xi' \beta_2$ . On the other hand if  $\alpha_1 - \alpha_2 = \xi'' \beta_2$ ,

it suffices to replace fraction  $(\alpha_1 - \alpha_2)/\beta_2$  by  $\xi''$  and allow  $\xi$  to be any element of  $R$ . If  $R$  contains no zero divisors, we can extend  $R$  into its quotient field and allow  $\xi$  to be any element of  $R$ , or any element of the quotient field.

If  $\Phi$  is a T-G invariant determined by  $(\alpha_1, \beta_1, \alpha_2, \beta_2)$ , then define  $\Phi^*$  as the T-G invariant determined by  $(\beta_1, \alpha_1, \beta_2, \alpha_2)$ . Clearly,  $\Phi = (\Phi^*)^*$ . By Lemma 1,  $\Phi(M) = \alpha_2^{r(M)} \beta_2^{r^*(M)} T(M; \alpha_1/\alpha_2, \beta_1/\beta_2)$  and  $\Phi^*(M^*) = \beta_2^{r^*(M)} \alpha_2^{r(M)} T(M^*; \beta_1/\beta_2, \alpha_1/\alpha_2)$  for each matroid  $M$ . The covariance formula (see [2]) is that  $T(M; x, y) = T(M^*; y, x)$ , whence

$$(3) \quad \Phi(M) = \Phi^*(M^*).$$

**Theorem 2.** *Let  $\Phi$  be a T-G invariant determined by  $(\alpha_1, \beta_1, \alpha_2, \beta_2)$ ,  $\alpha_2 \neq 0$ , and  $\xi \in R$  be a multiple of  $\alpha_2$ . Then  $\Phi_\xi^{\text{ls}}(M) = \xi^{|E|} \left( \frac{\beta_1 - \beta_2}{\alpha_2} \right)^{r(M)} \Phi(M)$  is a loop-smooth T-G invariant such that for every matroid  $M$ ,*

$$\begin{aligned} \Phi_\xi^{\text{ls}}(M) &= 1 && \text{if } E = \emptyset, \\ \Phi_\xi^{\text{ls}}(M) &= \xi \alpha_1 (\beta_1 - \beta_2) / \alpha_2 \Phi_\xi^{\text{ls}}(M - e) && \text{if } e \text{ is an isthmus of } M, \\ \Phi_\xi^{\text{ls}}(M) &= \xi (\beta_1 - \beta_2) \Phi_\xi^{\text{ls}}(M/e) + \beta_2 \Phi_\xi^{\text{ls}}(M - e) && \text{otherwise.} \end{aligned}$$

*Proof.* Set  $\Phi_\xi^{\text{ls}} = ((\Phi^*)_\xi^{\text{is}})^*$ . By (3) and Theorem 1,  $\Phi_\xi^{\text{ls}}(M) = ((\Phi^*)_\xi^{\text{is}})^*(M) = (\Phi^*)_\xi^{\text{is}}(M^*)$ . Applying Theorem 1 for  $\Phi^*$  and  $M^*$ , we get  $(\Phi^*)_\xi^{\text{is}}(M^*) = \xi^{|E|} \left( \frac{\beta_1 - \beta_2}{\alpha_2} \right)^{r(M)} \Phi^*(M^*) = \xi^{|E|} \left( \frac{\beta_1 - \beta_2}{\alpha_2} \right)^{r(M)} \Phi(M)$ . Furthermore by definition of  $\Phi^*$  and Theorem 1,  $((\Phi^*)_\xi^{\text{is}})^*$  is determined by  $(\xi \alpha_1 (\beta_1 - \beta_2) / \alpha_2, \xi \beta_1, \xi (\beta_1 - \beta_2), \xi \beta_2)$ .  $\square$

Notice that  $\Phi_\xi^{\text{ls}} = ((\Phi^*)_\xi^{\text{is}})^*$ , whence  $\Phi_\xi^{\text{is}} = (((\Phi^*)_\xi^{\text{is}})^*)^* = ((\Phi^*)_\xi^{\text{ls}})^*$ . Thus

$$(4) \quad \Phi_\xi^{\text{ls}} = ((\Phi^*)_\xi^{\text{is}})^* \text{ and } \Phi_\xi^{\text{is}} = ((\Phi^*)_\xi^{\text{ls}})^*.$$

$\Phi_\xi^{\text{ls}}$  is called the  $\xi$ -loop-smooth modification of  $\Phi$ . If  $\Phi$  is an isthmus invariant, then  $\Phi_\xi^{\text{ls}}(M) = \xi^{|E|} \Phi(M)$  for every matroid  $M$ .

In Theorems 1 and 2 we have assumed that  $\beta_2 \neq 0$  and  $\alpha_2 \neq 0$ , respectively. Let  $l_M$  ( $i_M$ ) denote the number of loops (isthmuses) in a matroid  $M$ . If  $\alpha_2 = 0$ , then by (1),  $\Phi(M) = \alpha_1^{r(M)} \beta_1^{l_M} \beta_2^{r^*(M) - l_M}$ , whence by (3),  $\Phi(M) = \beta_1^{r^*(M)} \alpha_1^{i_M} \alpha_2^{r(M) - i_M}$  if  $\beta_2 = 0$ . Thus  $\Phi(M)$  is easy to evaluate if  $\alpha_2 = 0$  or  $\beta_2 = 0$  (a contrast with the fact that the Tutte polynomial is difficult to evaluate, see [3, 4, 8]).

### 3. MODIFICATIONS OF THE TUTTE POLYNOMIAL

Let  $\xi \in \mathbb{Z}[x, y]$ . Then  $\xi$  is a multiple of 1 whence by Theorem 1, the  $\xi$ -isthmus-smooth modification of the Tutte polynomial of  $M$  is

$$(5) \quad T_\xi^{\text{is}}(M; x, y) = \xi^{|E|} (x - 1)^{r^*(M)} T(M; x, y)$$

and satisfies

$$(6) \quad \begin{aligned} T_\xi^{\text{is}}(M; x, y) &= 1 && \text{if } E = \emptyset, \\ T_\xi^{\text{is}}(M; x, y) &= \xi y(x-1)T_\xi^{\text{is}}(M-e; x, y) && \text{if } e \text{ is a loop of } M, \\ T_\xi^{\text{is}}(M; x, y) &= \xi T_\xi^{\text{is}}(M/e; x, y) + \xi(x-1)T_\xi^{\text{is}}(M-e; x, y) && \text{otherwise.} \end{aligned}$$

By Theorem 2, the  $\xi$ -loop-smooth modification of the Tutte polynomial of  $M$  is

$$(7) \quad T_\xi^{\text{ls}}(M; x, y) = \xi^{|E|}(y-1)^{r(M)}T(M; x, y)$$

and satisfies

$$(8) \quad \begin{aligned} T_\xi^{\text{ls}}(M; x, y) &= 1 && \text{if } E = \emptyset, \\ T_\xi^{\text{ls}}(M; x, y) &= \xi x(y-1)T_\xi^{\text{ls}}(M-e; x, y) && \text{if } e \text{ is an isthmus,} \\ T_\xi^{\text{ls}}(M; x, y) &= \xi(y-1)T_\xi^{\text{ls}}(M-e; x, y) + \xi T_\xi^{\text{ls}}(M/e; x, y) && \text{otherwise.} \end{aligned}$$

By (3),  $T_\xi^{\text{ls}}(M; x, y) = (T_\xi^{\text{ls}})^*(M^*; x, y)$ , and by (4),  $(T_\xi^{\text{ls}})^*(M^*; x, y) = (T^*)_\xi^{\text{is}}(M^*; x, y)$ . By (2) and (1), we have  $T^*(M^*; x, y) = T(M^*; y, x)$ , whence  $(T^*)_\xi^{\text{is}}(M^*; x, y) = T_\xi^{\text{is}}(M^*; y, x)$ , i.e., we have a variant of the covariance formula

$$(9) \quad T_\xi^{\text{ls}}(M; x, y) = T_\xi^{\text{is}}(M^*; y, x).$$

Kook, Reiner, and Stanton [6] introduced the convolution formula

$$T(M; x, y) = \sum_{A \subseteq E} T(M/A; x, 0) \cdot T(M|A; 0, y),$$

(where  $M|A$  and  $M/A$  denote the restriction of  $M$  to  $A$  and the contraction of  $A$  from  $M$ , respectively). Hence by (5) and (7),

$$T(M; x, y) = \sum_{A \subseteq E} \xi^{-|E|}(-1)^{-r(M/A)}T_\xi^{\text{ls}}(M/A; x, 0) \cdot (-1)^{-r^*(M|A)}T_\xi^{\text{is}}(M|A; 0, y).$$

Since  $r^*(M|A) = |A| - r(A)$ ,  $r(M/A) = r(M) - r(A)$ , and  $2r(A) - r(M) - |A|$  has the same parity as  $r(M) + |A|$ , we get a variant of the convolution formula

$$(10) \quad T(M; x, y) = \xi^{-|E|}(-1)^{r(M)} \sum_{A \subseteq E} (-1)^{|A|}T_\xi^{\text{ls}}(M/A; x, 0) \cdot T_\xi^{\text{is}}(M|A; 0, y).$$

The ring  $\mathbb{Z}[x, y]$  has no divisors of zero, therefore it has a quotient field  $\mathbb{F}[x, y]$ , consisting of all rational polynomials with integral coefficients. Thus, as pointed out in the remark after Theorem 1, for any  $\xi \in \mathbb{F}[x, y]$ , we can consider  $\xi$ -isthmus- and  $\xi$ -loop-smooth modifications of the Tutte polynomial, thus also formulas (9), (10).

If  $\Phi$  is a T-G invariant determined by  $(\alpha_1, \beta_1, \alpha_2, \beta_2)$  and  $\xi, \zeta \in R$ , then by Lemma 1, there exists a T-G invariant determined by  $(\xi\alpha_1, \zeta\beta_1, \xi\alpha_2, \zeta\beta_2)$  denoted by  $\Phi_{\xi, \zeta}$ . Clearly,  $\Phi_{\xi, \zeta}(M) = \xi^{r(M)}\zeta^{r^*(M)}\Phi(M)$  for each matroid  $M$ . Suppose that  $\Phi_{\xi, \zeta}$  is isthmus- and loop-smooth in the same time. Then  $\xi\alpha_1 = \zeta\beta_1 = \xi\alpha_2 + \zeta\beta_2$ , whence  $\xi/\zeta = \beta_1/\alpha_1 = \beta_2/(\alpha_1 - \alpha_2) = (\beta_1 - \beta_2)/\alpha_2$ , and thus

$$(11) \quad \beta_1 = \alpha_1\beta_2/(\alpha_1 - \alpha_2).$$

On the other hand (11) implies  $\beta_1/\alpha_1 = \beta_2/(\alpha_1 - \alpha_2)$  and  $(\beta_1 - \beta_2)/\alpha_2 = \beta_2/(\alpha_1 - \alpha_2)$ . Thus (11) is a necessary and sufficient condition for existence of  $\xi$  and  $\zeta$  such that  $\Phi_{\xi,\zeta}$  is an isthmus- and loop-smooth invariant. Therefore this kind of transformation cannot be applied for each  $\Phi$ . In particular, (11) is not valid for the Tutte polynomial because  $y \neq x/(x - 1)$ .

## REFERENCES

- [1] L. Beaudin, J. Ellis-Monaghan, G. Pangborn, and R. Shrock, *A little statistical mechanics for the graph theorist*, Discrete Math. **310** (2010) 2037–2053.
- [2] T. Brylawski and J. Oxley, *The Tutte polynomial and its applications*, in: Matroid Applications, (N. White, Editor), Cambridge University Press, Cambridge (1992), pp. 123-225.
- [3] L.A. Goldberg and M. Jerrum, *Inapproximability of the Tutte polynomial*, Inform. and Comput. **206** (2008) 908–929.
- [4] F. Jaeger, D.L. Vertigan, and D.J.A. Welsh, *On the computational complexity of the Jones and Tutte polynomials*, Math. Proc. Cambridge Philos. Soc. **108** (1990) 35–53.
- [5] M. Kochol, *Splitting formulas for Tutte-Grothendieck invariants*, manuscript (2014).
- [6] W. Kook, V. Reiner, and D. Stanton, *A convolution formula for the Tutte polynomial*, J. Combin. Theory Ser. B **76** (1999) 297–300.
- [7] J.G. Oxley and D.J.A. Welsh, *The Tutte polynomial and percolation*, in: Graph Theory and Related Topics, (J.A. Bondy and U.S.R. Murty, Editors), Academic Press, New York (1979), pp. 329-339.
- [8] D. Vertigan, *The computational complexity of Tutte invariants for planar graphs*, SIAM J. Comput. **135** (2005) 690–712.
- [9] D. J. A. Welsh, *Complexity: Knots, Colourings and Counting*, London Math. Soc. Lecture Notes Series 186, Cambridge University Press, Cambridge (1993).

MÚ SAV, ŠTEFÁNIKOVA 49, 814 73 BRATISLAVA 1, SLOVAKIA  
*E-mail address:* martin.kochol@mat.savba.sk