

Neutrino Lorentz invariance violation from the CPT –even SME coefficients through a tensor interaction with cosmological scalar fields

Rubén Cordero,^{1,*} Luis A. Delgadillo,^{1,†} O. G. Miranda,^{2,‡} and C. A. Moura^{3,§}

¹*Departamento de Física, Escuela Superior de Física
y Matemáticas del Instituto Politécnico Nacional,*

Unidad Adolfo López Mateos, Edificio 9, 07738 Ciudad de México, Mexico

²*Departamento de Física, Centro de Investigación y de Estudios
Avanzados del IPN Apdo. Postal 14-740 07000 Ciudad de México, Mexico*

³*Centro de Ciências Naturais e Humanas,
Universidade Federal do ABC - UFABC, Av. dos Estados,
5001, 09210-580, Santo André-SP, Brazil*

(Dated: July 29, 2024)

Abstract

Numerous non-standard interactions between neutrinos and scalar fields have been suggested in the literature. In this work, we have outlined the case of tensorial neutrino non-standard interactions with scalar fields, which can be related to the effective CPT –even dimension–4 operators of the Standard Model Extension (SME). Moreover, in this paper, we have mostly analyzed the projected sensitivities to the CPT –even isotropic $c_{\alpha\beta}$ and Z –spatial $c_{\alpha\beta}^{ZZ}$ SME coefficients, in the context of a long-baseline experimental configuration. We study the particular case of DUNE and show its sensitivity to this type of new physics. The current IceCube experiment and upcoming neutrino experiments such as KM3NeT, IceCube-Gen2, and GRAND proposals may clarify these classes of neutrino non-standard interactions.

* rcorderoe@ipn.mx

† ldelgadillof2100@alumno.ipn.mx

‡ omar.miranda@cinvestav.mx

§ celio.moura@ufabc.edu.br

I. INTRODUCTION

The discovery of the Universe's accelerating expansion [1, 2] is one of the most important, captivating, and puzzling open questions in cosmology [3, 4]. Several proposals to architecture Dark Matter (DM) and Dark Energy (DE) have been investigated in the literature. The Lambda Cold Dark Matter (Λ CDM) model is among the most popular explanations for cosmological observations, although recent results from the Dark Energy Spectroscopic Instrument (DESI) [5] show some tension with this model. Different candidates for DM particles have been considered, ranging in a very wide mass spectrum, making the Weakly Interacting Massive Particle (WIMP) paradigm one of the best motivated from a theoretical point of view [6]. However, despite thorough searches for WIMPs, they all have failed to find any signature.

Other DM and DE candidates have also been studied in an effort to find a plausible explanation to the cosmological observations. For instance, ultralight scalars are well-motivated proposals for cosmological DM [7–11]. Moreover, several ultralight scalars become apparent within the context of string theories [12–15].¹ Scalar fields minimally coupled to gravity are sufficiently justified models of DE [19], namely quintessence [20, 21] and k-essence [22–28].

In k-essence models, the scalar field plays a significant role in describing the DE puzzle. This field could have adequate behavior at early epochs and can reproduce the dynamical effects of the cosmological constant at late times. Classes of k-essence Lagrangians were introduced in several settings, for example, as a possible model for inflation [29, 30]. Subsequently, k-essence models were used as another possibility to describe the characteristics of DE and as an alternative mechanism for unifying DE and DM [31]. Purely kinetic k-essence models [32, 33] are, in a way, as simple as quintessence models, because they rely only on one function (F) through the expression of the Lagrangian density $\mathcal{L} = F(X)$, where X is the kinetic term.

Among other proposals, there are second-order derivative scalar field models, or generalized galileons [34–37], with the property that their equations of motion are second-order. Interesting examples of this class of models are the so-called kinetic gravity braiding models, that are formulated from a Lagrangian that includes a D'Alambertian operator and

¹ Some strategies to hunt for ultralight scalars as cosmological DM involve atomic clocks [16], resonant-mass detectors [17] and atomic gravitational wave detectors [18].

an arbitrary function of a non-canonical kinetic term giving rise to appealing cosmological effects [38–41].

At the cosmological level, all the aforementioned models of DM and DE can be described as a perfect fluid through their energy-momentum tensor $T_\varphi^{\mu\nu}$ [42]. At present, discrimination among them could be done at the level of the energy-momentum tensor perturbations. On the other hand, the search for signatures confirming the existence of scalars is an important challenge. Proposals considering the interaction of scalar fields with neutrinos have demonstrated that a possible signature might be detected in long-baseline neutrino experiments [43–53] or via modifications to the ultrahigh energy neutrino fluxes [54, 55]. Besides, it has also been argued that interactions of neutrinos with scalars, if they exist, could induce an apparent violation of Lorentz and *CPT* symmetries in the neutrino sector [56–67]. These scalars can be identified as either DM or DE candidates. We refer to [68–72] for comprehensive reviews concerning Lorentz and *CPT* symmetry violations in the neutrino sector, within the Standard Model Extension (SME) framework [73]. An initial proposal for *CPT* violation in the neutrino sector was discussed in Ref. [74].

Recently, there has been an increased interest in searches for Lorentz invariance and *CPT*–breakdowns in neutrino oscillation experiments [75–80]. For instance, comprehensive studies of the isotropic *CPT*–even SME coefficient, $(c_L)^{TT}$, at different long-baseline experiments can be found in Refs. [78, 79].² In a forthcoming publication [85], we examine the phenomenology of isotropic and anisotropic *CPT*–odd SME coefficients, $(a_L)^T$ and $(a_L)^Z$, considering the Deep Underground Neutrino Experiment (DUNE) [86] configuration.

In this paper, we study the sensitivity of the DUNE to quintessence and similar DE or DM models through their possible interaction with neutrinos. We consider the cases where an apparent Lorentz–violating signature manifests and show that, in these models, a tensorial interaction of neutrinos with scalar fields may arise and give a sizable signature at the DUNE.

The manuscript is organized as follows: In Section II, we introduce the theoretical foundations, where the basics of Lorentz invariance violations are presented together with the necessary Lagrangian formalism to describe the several models of scalar fields useful to model DM and DE. In Section III, we describe the phenomenology of a tensorial neutrino interac-

² As shown in [81], it is possible to relate the isotropic SME coefficients $(c_L)^{TT}$ with the effects of the violation of the equivalence principle (VEP) in the neutrino sector. Besides, there is a correspondence between quantum-decoherence effects in neutrinos and the effective coefficients of the SME [82–84].

tion with scalar fields as either DM or DE candidates and its connection with the CPT –even SME coefficients $(c_L)_{\alpha\beta}^{\mu\nu}$. In Section IV, we assess the sensitivities to the CPT –even SME coefficients, $(c_L)_{\alpha\beta}^{\mu\nu}$, considering the DUNE configuration. Finally, we give our conclusions.

II. THEORETICAL FRAMEWORK

Within the SME framework, violations of Lorentz invariance (LIV) in the fermion sector, are parameterized by the effective Lagrangian [72]

$$\mathcal{L}_{\text{eff}}^\psi = \mathcal{L}_{\text{SM}}^\psi + \mathcal{L}_{\text{LIV}} + \text{h.c.}, \quad (1)$$

where $\mathcal{L}_{\text{SM}}^\psi$ is the Standard Model (SM) fermion Lagrangian, \mathcal{L}_{LIV} being the Lorentz and CPT –violating Lagrangian term,

$$-\mathcal{L}_{\text{LIV}} = \frac{1}{2} \left\{ p_{\alpha\beta}^\mu \bar{\psi}_\alpha \gamma_\mu \psi_\beta + q_{\alpha\beta}^\mu \bar{\psi}_\alpha \gamma_5 \gamma_\mu \psi_\beta - ir_{\alpha\beta}^{\mu\nu} \bar{\psi}_\alpha \gamma_\mu \partial_\nu \psi_\beta - is_{\alpha\beta}^{\mu\nu} \bar{\psi}_\alpha \gamma_5 \gamma_\mu \partial_\nu \psi_\beta \right\}. \quad (2)$$

In the case of neutrinos, it is convenient to define

$$(a_L)_{\alpha\beta}^\mu = (p+q)_{\alpha\beta}^\mu \quad \text{and} \quad (c_L)_{\alpha\beta}^{\mu\nu} = (r+s)_{\alpha\beta}^{\mu\nu}, \quad (3)$$

where the $(a_L)_{\alpha\beta}^\mu$ coefficients are CPT –odd, while the $(c_L)_{\alpha\beta}^{\mu\nu}$ coefficients are CPT –even. These type of CPT –conserving terms, $(c_L)_{\alpha\beta}^{\mu\nu}$, were studied in the context of a Finslerian Geometrical model [87], and of neutrino interactions with a k-essence field [88]. For other scenarios of neutrino–dark energy interactions, we refer the reader to Refs. [89–91].

Lately, it has been suggested that violations of Lorentz invariance and CPT symmetry in the neutrino sector may arise from a neutrino–current coupled with a dynamical cosmic field, $\varphi(t)$ (see, e.g., Refs. [56–59, 64–66])

$$(a_L)_{\alpha\beta}^\mu \bar{\nu}_\alpha \gamma_\mu (1 - \gamma_5) \nu_\beta \rightarrow y_{\alpha\beta} \frac{\partial^\mu \varphi}{\Lambda} \bar{\nu}_\alpha \gamma_\mu (1 - \gamma_5) \nu_\beta, \quad (4)$$

being $y_{\alpha\beta}$ some coupling constants, Λ the energy scale of the interaction, and $\varphi = \varphi(t)$ a time–varying scalar field, which could be identified as either a dark matter or dark energy candidate.³ Here, violations of the Lorentz and CPT symmetries emerge via the CPT –odd SME coefficients $(a_L)_{\alpha\beta}^\mu \rightarrow y_{\alpha\beta} \partial^\mu \varphi \Lambda^{-1}$.

³ For ultra-relativistic neutrinos, the phenomenology with axion-like dark matter is the same as the ultralight dark matter scenario [62, 65, 92].

Motivated by the aforementioned proposals, let us consider the corresponding free scalar field Lagrangian and its energy-momentum tensor, which are relevant for describing both, scalar field dark matter and dark energy quintessence [8, 20, 42],

$$\begin{aligned}\mathcal{L}_\varphi &= \frac{1}{2}g^{\mu\nu}\partial_\mu\varphi\partial_\nu\varphi + V(\varphi), \\ T_\varphi^{\mu\nu} &= \frac{2}{\sqrt{-g}}\frac{\delta}{\delta g_{\mu\nu}}\left[\sqrt{-g}\left(\frac{1}{2}g^{\lambda\sigma}\partial_\lambda\varphi\partial_\sigma\varphi + V(\varphi)\right)\right].\end{aligned}\tag{5}$$

Therefore, the scalar field energy-momentum tensor $T_\varphi^{\mu\nu}$ can be expressed as

$$T_\varphi^{\mu\nu} = g^{\mu\nu}\left(\frac{1}{2}g^{\lambda\sigma}\partial_\lambda\varphi\partial_\sigma\varphi + V(\varphi)\right) - g^{\lambda\mu}g^{\sigma\nu}\partial_\lambda\varphi\partial_\sigma\varphi,\tag{6}$$

and we identify the density and pressure of the scalar field φ as

$$\begin{aligned}\rho_\varphi &= -\frac{1}{2}g^{\mu\nu}\partial_\mu\varphi\partial_\nu\varphi + V(\varphi) \quad \text{and} \\ p_\varphi &= -\frac{1}{2}g^{\mu\nu}\partial_\mu\varphi\partial_\nu\varphi - V(\varphi).\end{aligned}\tag{7}$$

The scalar field four-velocity, U^μ , is given as

$$U^\mu = \frac{-\partial_\mu\varphi}{\sqrt{-g^{\lambda\sigma}\partial_\lambda\varphi\partial_\sigma\varphi}}.\tag{8}$$

The isotropy and homogeneity of space-time require the stress-energy-momentum tensor for a free scalar field, φ , to be that of a perfect fluid,

$$\begin{aligned}T_\varphi^{ij} &= p_\varphi\delta_j^i; \quad T_\varphi^{i0} = T_{\varphi 0i} = 0; \quad T_\varphi^{00} = \rho_\varphi, \\ T_\varphi^{\mu\nu} &= p_\varphi g^{\mu\nu} + (p_\varphi + \rho_\varphi)U^\mu U^\nu,\end{aligned}\tag{9}$$

subject to the constraint $g_{\mu\nu}U^\mu U^\nu = -1$. The conservation of $T_\varphi^{\mu\nu}$ in a Friedmann-Lemaitre-Robertson-Walker (FLRW) background leads to the continuity equation,

$$\partial_0 T_\varphi^{00} = \dot{\rho}_\varphi + 3\frac{\dot{a}}{a}(p_\varphi + \rho_\varphi) = 0,\tag{10}$$

which gives the equation of motion of the scalar field,

$$\ddot{\varphi} + 3H\dot{\varphi} + \frac{dV(\varphi)}{d\varphi} = 0.\tag{11}$$

Nevertheless, dark energy can be modeled by a k-essence field φ which has very appealing properties and is described by the Lagrangian density $\mathcal{L} = K(\varphi, X)$ [23] which is associated with the following energy-momentum tensor,

$$T_{\mu\nu}^\varphi = K_{,X}\nabla_\mu\varphi\nabla_\nu\varphi - g_{\mu\nu}K,\tag{12}$$

where $X = \nabla_\alpha \varphi \nabla^\alpha \varphi / 2$ is the kinetic term and $K_{,X} = \partial K / \partial X$. The energy density and pressure are given by

$$\rho_\varphi = 2XK_{,X} - K \quad \text{and} \quad p_\varphi = K , \quad (13)$$

respectively, and the equation of motion can be obtained from the continuity equation,

$$(2XK_{,XX} + K_{,X})\ddot{\varphi} + K_{,X\varphi}\dot{\varphi}^2 + 3HK_{,X}\dot{\varphi} = K_{,\varphi} , \quad (14)$$

where $K_{,X\varphi} = \partial^2 K / \partial X \partial \varphi$ and $K_{,\varphi} = \partial K / \partial \varphi$.

Another possible interesting scalar field used to describe the dark energy evolution is the braided scalar field described by the Lagrangian:

$$\mathcal{L} = \square \varphi G(X, \varphi) , \quad (15)$$

where $\square \varphi = \nabla^\mu \nabla_\mu \varphi$ and the function G is arbitrary [39]. The energy-momentum tensor, $T_{\mu\nu}^\varphi$, in covariant form, is given by

$$T_{\mu\nu}^\varphi = \frac{2}{\sqrt{-g}} \frac{\delta S_\varphi}{\delta g^{\mu\nu}} = \mathcal{L}_X \nabla_\mu \varphi \nabla_\nu \varphi - g_{\mu\nu} P_\varphi - \nabla_\mu G \nabla_\nu \varphi - \nabla_\nu G \nabla_\mu \varphi , \quad (16)$$

where $P_\varphi = \nabla^\lambda \varphi \nabla_\lambda G_\varphi$, and $G_\varphi = \partial G / \partial \varphi$. This energy-momentum tensor can be described in terms of an imperfect fluid [39]. We start defining some quantities to describe relativistic fluids. A local rest frame is set defining the normalized four-velocity U_μ

$$U_\mu \equiv \frac{\nabla_\mu \varphi}{\sqrt{2X}}, \quad U_\mu U^\mu = 1 , \quad (17)$$

$$a_\mu \equiv U^\nu \nabla_\nu U_\mu , \quad (18)$$

with a_μ is the four-acceleration which is orthogonal to velocity, $U_\mu a^\mu = 0$. The expansion, ϑ , and the diffusivity, Ω , are written as

$$\vartheta = \nabla_\mu U^\mu, \quad \Omega = 2XG_X , \quad (19)$$

where $G_X = \partial G / \partial X$. The energy-momentum tensor also can be expressed in this way:

$$T_{\mu\nu}^\varphi = \rho_\varphi U_\mu U_\nu - \perp_{\mu\nu} p_\varphi + U_\mu q_\nu + U_\nu q_\mu , \quad (20)$$

where $m = \sqrt{2X} = \dot{\varphi}$ is the chemical potential, $q_\mu = -m\Omega a_\mu$ is the heat flux (purely spatial, $U_\mu q^\mu = 0$) and $\perp_{\mu\nu} = g_{\mu\nu} - U_\mu U_\nu$ is the transverse projector. The energy density

and isotropic pressure are:

$$\rho_\varphi \equiv T_{\mu\nu}^\varphi U^\mu U^\nu = -2XG_\varphi + \vartheta m\Omega, \quad (21)$$

$$p_\varphi \equiv -\frac{1}{3}T_\varphi^{\mu\nu} \perp_{\mu\nu} = -2XG_\varphi - \Omega\dot{m}. \quad (22)$$

The conservation of $T_{\mu\nu}^\varphi$ using these definitions can be expressed as:

$$U_\nu \nabla_\mu T_\varphi^{\mu\nu} = \dot{\rho}_\varphi + \vartheta(\rho_\varphi + p_\varphi) - \nabla_\lambda(m\Omega a^\lambda) + m\Omega a_\lambda a^\lambda = 0. \quad (23)$$

The equation of motion for φ can be obtained from the last equation

$$\nabla_\mu [2G_\varphi \nabla^\mu \varphi - \square \varphi G_X \nabla^\mu \varphi + G_X \nabla^\mu X] = \nabla^\lambda \varphi \nabla_\lambda G_\varphi. \quad (24)$$

In the cosmological background, the evolution of the scalar field reduces to the dynamics of a perfect fluid which is described only through its energy density and pressure.

III. TENSORIAL NEUTRINO LIV

Consider the effective Lagrangian (see Appendix A for further details)

$$-\mathcal{L}_{\text{eff}} = -i \frac{\lambda_{\alpha\beta}}{M_*^4} T_\varphi^{\mu\nu} \bar{\nu}_\alpha \gamma_\mu (1 - \gamma_5) \partial_\nu \nu_\beta, \quad (25)$$

being $\lambda_{\alpha\beta}$ a coupling constant matrix, M_* the energy scale of the interaction, and $T_\varphi^{\mu\nu}$ is the tensor associated to the scalar field φ , for instance those of Eqs. (6), (12), and (16). Hence, we can identify the *CPT*-even LIV coefficients of the SME $(c_L)_{\alpha\beta}^{\mu\nu} = c_{\alpha\beta}^{\mu\nu}$ as

$$c_{\alpha\beta}^{\mu\nu} \rightarrow \frac{\lambda_{\alpha\beta}}{M_*^4} T_\varphi^{\mu\nu}. \quad (26)$$

Here, the scalar field φ could be one of the DM or DE candidates described in Section II. Besides, the effective interaction (Eq. 25) may induce potential scattering among the neutrinos and the scalar particles. However, such interactions would be expected to be negligible (see Appendix B of Ref. [58]). Henceforth, we discuss the case of a tensorial neutrino-scalar field interaction.⁴

⁴ A study of the back-reaction effects from the tensorial neutrino-scalar field interaction is beyond the scope of this work.

A. Isotropic LIV coefficients $c_{\alpha\beta}$ from a neutrino-scalar field interaction

From the effective Lagrangian in Eq. (25), the corresponding isotropic *CPT*-even LIV coefficients $c_{\alpha\beta}$ are

$$c_{\alpha\beta} = c_{\alpha\beta}^{00} \rightarrow \frac{\lambda_{\alpha\beta}}{M_*^4} T_\varphi^{00} = \frac{\lambda_{\alpha\beta}}{M_*^4} \rho_\varphi, \quad (27)$$

considering the scalar field φ as ultralight dark matter (ULDM) [8, 9], with corresponding local DM density in the Milky Way, $\rho_{\varphi,\odot} \sim \rho_{\text{DM},\odot} \simeq 2 \times 10^{-6} \text{ eV}^4$ [93–95], an accelerator-based experiment similar to DUNE, with sensitivity $c_{\alpha\beta} \sim [1 - 10] \times 10^{-25}$ (left panel of Fig. 1 in Sec. IV), could potentially probe an energy scale of the interaction

$$M_* \sim [3 - 6] \times 10^4 \text{ eV} \left(\lambda_{\alpha\beta}/\mathcal{O}(1) \right) \left(\rho_{\varphi,\odot}/10^{-6} \text{ eV}^4 \right). \quad (28)$$

On the other hand, if the scalar field φ is considered to be a DE candidate, namely quintessence, the corresponding dark energy density $\rho_\varphi^{\text{DE}} \simeq 3\rho_{\text{DM}}^{\text{avg}} \sim 10^{-10} \text{ eV}^4$, in this scenario, the DUNE setup could potentially probe an energy scale

$$M_* \sim [5 - 10] \times 10^3 \text{ eV} \left(\lambda_{\alpha\beta}/\mathcal{O}(1) \right) \left(\rho_\varphi^{\text{DE}}/10^{-10} \text{ eV}^4 \right). \quad (29)$$

At the cosmological level, the aforementioned models of dark energy outlined in Section II, can be described by a dynamical scalar field $\varphi(t)$, thus, for all cases, they would predict $T_\varphi^{00} = \rho_\varphi^{\text{DE}}$. However, limits from astrophysical neutrinos ($60 \text{ TeV} \lesssim E_\nu \lesssim \text{PeV}$) employing the IceCube astrophysical neutrino flavour data-set constraint $c_{\alpha\beta} \lesssim 10^{-34}$ [76] (such limits are expected to be improved by the combination of a two-detector fit, namely IceCube-Gen2 and GRAND [80]), therefore ⁵

$$c_{\alpha\beta} \sim \frac{\lambda_{\alpha\beta}}{M_*^4} \rho_\varphi \lesssim 10^{-34}, \quad (30)$$

considering φ as DM, ultra-high energy (UHE) neutrino experiments could potentially be sensitive to an energy scale

$$M_* \gtrsim 10^7 \text{ eV} \left(\lambda_{\alpha\beta}/\mathcal{O}(1) \right) \left(\rho_{\varphi,\odot}/10^{-6} \text{ eV}^4 \right) \left(c_{\alpha\beta}/\lesssim 10^{-34} \right), \quad (31)$$

⁵ Besides, in the standard model, neutrinos are part of the $SU(2)_L$ doublet, hence, we could have potentially induced LIV effects for the charged leptons. However, for the case of electrons, limits from astrophysical observations constraint $c_{ee} \in (-80 \text{ to } 4) \times 10^{-20}$ [96], which are several orders of magnitude weaker than the constraints derived in the neutrino sector $c_{\alpha\beta} \lesssim 10^{-34}$ [76].

on the other hand, considering φ as DE, UHE neutrino experiments could potentially be sensitive to an energy scale

$$M_* \gtrsim 10^6 \text{ eV} \left(\lambda_{\alpha\beta}/\mathcal{O}(1) \right) \left(\rho_\varphi^{\text{DE}}/10^{-10} \text{ eV}^4 \right) \left(c_{\alpha\beta}/\lesssim 10^{-34} \right). \quad (32)$$

Besides, considering only the derivative coupling on the scalar fields (see Appendix A), unitarity bounds in collisions with quarks at the LHC set $M_* \gtrsim 30 \text{ GeV}$, while unitarity bounds from LEP when colliding electrons and positrons impose $M_* \gtrsim 3 \text{ GeV}$ [97].

B. Directional dependent LIV coefficients from a neutrino-scalar field interaction

Searches of directional dependent LIV effects can be performed at neutrino experiments such as the KM3NeT neutrino telescope and the IceCube neutrino observatory, with neutrino energies $E_\nu \gtrsim 10^5 \text{ GeV}$ [58, 98]. Besides, regarding directional dependent Lorentz violating effects in the Z -direction, the projected sensitivities of a neutrino long-baseline experiment similar to DUNE to the CPT -even Z -spatial LIV coefficients are, $c_{\alpha\beta}^{ZZ} \sim [1 - 10] \times 10^{-24}$ (right panel of Fig. 1 in Sec. IV) and $c_{\alpha\alpha}^{ZZ} - c_{\tau\tau}^{ZZ} \simeq [1 - 10] \times 10^{-24}$ (right panel of Fig. 2 in Sec. IV), both at 95% C.L., accordingly.

Furthermore, from the phenomenological Lagrangian in Eq. (25), the corresponding CPT -even Z -spatial LIV coefficients are

$$c_{\alpha\beta}^{ZZ} \rightarrow \frac{\lambda_{\alpha\beta}}{M_*^4} (T_\varphi^{ZZ} + \delta T_\varphi^{ZZ}), \quad (33)$$

where $\delta T_\varphi^{\mu\nu}$ accounts for the perturbations of the scalar field tensor $T_\varphi^{\mu\nu}$. For instance, perturbations for the several dark energy models outlined, quintessence, k-essence, or kinetic gravity braiding models are expected to be different. Subsequently, we refer to the dark energy quintessence model for simplicity. Still, a similar phenomenology applies to the other dark energy models.

The perturbed stress-energy-momentum tensor for the scalar field $\varphi(t)$, with perturbation $\delta\varphi(t, \mathbf{x})$ in the FLRW background with a Newtonian gauge is given as [7]

$$\delta T_\varphi^{ij} = \delta p_\varphi \delta_j^i; \quad \delta T_\varphi^{i0} = -\frac{1}{a} (\dot{\varphi}(t) \partial_i(\delta\varphi)); \quad \delta T_\varphi^{00} = \delta\rho_\varphi. \quad (34)$$

Considering φ as ULDM, from the equation of state of the scalar field $p_\varphi = \omega\rho_\varphi$, with $\omega = 0$, the leading contribution $T_\varphi^{ZZ} = 0$, hence, in this case the CPT -even Z -spatial LIV

coefficients are

$$|c_{\alpha\beta}^{ZZ}| \rightarrow \frac{\lambda_{\alpha\beta}}{M_*^4} |\delta T_\varphi^{ZZ}| \sim \frac{\lambda_{\alpha\beta}}{M_*^4} \left| \left(\dot{\varphi}(t) \delta \dot{\varphi}(t, \mathbf{x}) - \dot{\varphi}^2(t) \Phi(t, \mathbf{x}) - V(\varphi)_{,\varphi} \delta \varphi(t, \mathbf{x}) \right) \right|, \quad (35)$$

here $\Phi(t, \mathbf{x})$ plays the role of the gravitational potential, while $\delta \varphi(t, \mathbf{x})$ is the scalar field perturbation [7]. On the other hand, if φ is a DE candidate (quintessence) from the equation of state of the scalar field $p_\varphi = \omega \rho_\varphi^{\text{DE}}$, with $\omega = -1$,

$$|c_{\alpha\beta}^{ZZ}| \rightarrow \frac{\lambda_{\alpha\beta}}{M_*^4} |T_\varphi^{ZZ} + \delta T_\varphi^{ZZ}| \sim \frac{\lambda_{\alpha\beta}}{M_*^4} \rho_\varphi^{\text{DE}}. \quad (36)$$

In a similar fashion, considering φ as either ULDM or DE quintessence, the leading contribution $T_\varphi^{TZ} = 0$ (Eq. 9), therefore, the *CPT*-even coefficients from the *TZ*-sector can be described as

$$|c_{\alpha\beta}^{TZ}| \rightarrow \frac{\lambda_{\alpha\beta}}{M_*^4} |\delta T_\varphi^{TZ}| \sim \frac{\lambda_{\alpha\beta}}{M_*^4} \left| -\frac{1}{a} \left(\dot{\varphi}(t) \frac{\partial}{\partial z} \delta \varphi(t, x, y, z) \right) \right|. \quad (37)$$

IV. NUMERICAL ANALYSIS AND RESULTS

Long-baseline neutrino oscillation experiments play a significant role in both deciphering mysteries within the traditional three-neutrino oscillation picture and exploring other novel physics scenarios, including the potential breaking of the Lorentz and *CPT* symmetries. Hence, in this part of the paper, we focus on a long-baseline experimental configuration, the Deep Underground Neutrino Experiment (DUNE) as a case of study, which is a next-generation accelerator-based neutrino oscillation experiment that will consist of up to 40 kt of liquid argon (far) detector located at the Sanford underground research facility (SURF) in South Dakota [99]. Moreover, this configuration is expected to deliver a neutrino flux with a mean neutrino energy $E_\nu \sim 3$ GeV, which is going to be located at a distance of $L \sim 1300$ km from the beam source (on-axis) at Fermilab (we refer the reader to Refs. [78, 86], for a detailed discussion regarding the experimental configuration).

In order to obtain sensitivities to the LIV coefficients at DUNE, we use the GLOBES software [100, 101] and its additional NSI tool *snu.c* [102, 103] which was modified to implement the *CPT*-even coefficients of the SME at the Hamiltonian level. Moreover, to simulate the DUNE configuration, we employ the available GLOBES files [104] and specifications from the Technical Design Report (TDR) configuration [86]. Furthermore, in this work, we have contemplated a 10-year running time, evenly distributed among neutrino and anti-neutrino modes.

In this analysis, we have considered the following Hamiltonian

$$H = H_0 + H_{\text{MSW}} + H_{\text{LIV}}, \quad (38)$$

here, H_0 and H_{MSW} are the standard neutrino Hamiltonian in vacuum and matter, respectively. Besides, H_{LIV} is the contribution from the Lorentz invariance violation (LIV) sector, which can be parameterized as [105–107]

$$H_{\text{LIV}} = -\frac{E_\nu}{2} \left[(3 - \hat{N}_Z^2)(c_L)_{\alpha\beta}^{TT} + (3\hat{N}_Z^2 - 1)(c_L)_{\alpha\beta}^{ZZ} - 2\hat{N}_Z(c_L)_{\alpha\beta}^{TZ} \right], \quad (39)$$

where E_ν is the neutrino energy, $(c_L)_{\alpha\beta}^{\mu\nu}$ are the CPT –even LIV coefficients of the SME (being $\alpha, \beta = e, \mu, \tau$, and $\mu, \nu = T, X, Y, Z$), for non-diagonal $\alpha \neq \beta$, $c_{\alpha\beta}^{\mu\nu} = |c_{\alpha\beta}^{\mu\nu}|e^{i\phi_{\alpha\beta}^{\mu\nu}}$ (from now on, we will denote $(c_L)_{\alpha\beta}^{\mu\nu} = c_{\alpha\beta}^{\mu\nu}$, and $(c_L)_{\alpha\beta}^{TT} = c_{\alpha\beta}$), and

$$\hat{N}^Z = -\sin\chi \sin\theta \cos\phi + \cos\chi \cos\theta, \quad (40)$$

is the spatial Z –component factor, expressed in terms of local spherical coordinates at the detector, that represents the direction of neutrino propagation in the Sun-centered frame. Being χ the colatitude of the detector, θ the angle at the detector between the beam direction and vertical, and ϕ the angle between the beam and east of south [105]. In the particular case of the DUNE location, $\hat{N}^Z \simeq 0.16$ [85].

A. Sensitivity to CPT –even SME coefficients

In order to assess the statistical significance to CPT –even SME coefficients, we employ a chi-squared test, we have considered the muon neutrino disappearance channel $P(\nu_\mu \rightarrow \nu_\mu)$ as well as the electron neutrino appearance channel $P(\nu_\mu \rightarrow \nu_e)$, from a muon-neutrino beam with neutrino and antineutrino data sets. The total χ^2 –function is provided as in Ref. [52]

$$\chi^2 = \sum_k \tilde{\chi}_k^2 + \chi_{\text{prior}}^2, \quad (41)$$

where the corresponding $\tilde{\chi}_k^2$ –function for each channel (k), appearance or disappearance is given as in Ref. [108]

$$\begin{aligned} \tilde{\chi}_k^2 = \min_{\zeta_j} & \left[\sum_i^{n_{\text{bin}}} 2 \left\{ N_{i,\text{test}}^{3\nu+\text{LIV}}(\Pi, \Gamma, \{\zeta_j\}) - N_{i,\text{true}}^{3\nu} + N_{i,\text{true}}^{3\nu} \log \frac{N_{i,\text{true}}^{3\nu}}{N_{i,\text{test}}^{3\nu+\text{LIV}}(\Pi, \Gamma, \{\zeta_j\})} \right\} \right. \\ & \left. + \sum_j^{n_{\text{syst}}} \left(\frac{\zeta_j}{\sigma_j} \right)^2 \right], \end{aligned} \quad (42)$$

here, $N_{i,\text{true}}^{3\nu}$ refers to the simulated events at the i -th energy bin (considering the standard three neutrino oscillations picture), while $N_{i,\text{test}}^{3\nu+\text{LIV}}(\Pi, \Gamma, \{\zeta_j\})$ are the computed events at the i -th energy bin including CPT –even SME coefficients (one parameter at a time). In addition, $\Pi = \{\theta_{12}, \theta_{13}, \theta_{23}, \delta_{CP}, \Delta m_{21}^2, \Delta m_{31}^2\}$ is the set of neutrino oscillation parameters, while $\Gamma = \{|c_{\alpha\beta}|, \phi_{\alpha\beta}, c_{\alpha\alpha}, \dots, |c_{\alpha\beta}^{ZZ}|, \phi_{\alpha\beta}^{ZZ}, c_{\alpha\alpha}^{ZZ}, \dots\}$ is the set of either isotropic ($c_{\alpha\beta}$), Z –spatial ($c_{\alpha\beta}^{ZZ}$) or ($c_{\alpha\beta}^{TZ}$) SME coefficients, where $\{\zeta_j\}$ are the nuisance parameters to account for the systematic uncertainties. Furthermore, σ_j are the systematic uncertainties as reported in the DUNE TDR [86]. Besides, to obtain our simulated events; we have considered the corresponding neutrino oscillation parameters as *true* values, namely $\Delta m_{21}^2 = 7.5 \times 10^{-5} \text{ eV}^2$, $\Delta m_{31}^2 = 2.55 \times 10^{-3} \text{ eV}^2$, $\theta_{12} = 34.3^\circ$, $\theta_{13} = 8.53^\circ$, $\theta_{23} = 49.26^\circ$, and $\delta_{CP} = 1.08\pi$, corresponding to the best-fit values with normal mass ordering (NO) from Ref. [109] as displayed in Tab. I.

Oscillation parameter	best-fit NO
θ_{12}	34.3°
θ_{23}	49.26°
θ_{13}	8.53°
Δm_{21}^2 [10^{-5} eV^2]	7.5
$ \Delta m_{31}^2 $ [10^{-3} eV^2]	2.55
δ_{CP}/π	1.08

Table I: Standard oscillation parameters used in our analysis [109]. We consider the normal mass ordering (NO) throughout this study.

Furthermore, to include external input for the standard oscillation parameters in the total χ^2 –function, Gaussian priors [108] are utilized.

$$\chi_{\text{prior}}^2 = \sum_p^{n_{\text{priors}}} \frac{(\Pi_{p,\text{true}} - \Pi_{p,\text{test}})^2}{\sigma_p^2}, \quad (43)$$

the central values of the oscillation parameter priors Π_p were fixed to their best-fit [109], considering the normal mass ordering, σ_p is the uncertainty on the oscillation prior, which corresponds to a 1σ error (68.27% confidence level C.L.).




Figure 1: Projected 95% C.L. sensitivities to the SME coefficients $|c_{\alpha\beta}^{\mu\nu}|$ at the DUNE (TDR setup). Here, we have marginalized over the corresponding LIV phases $\phi_{\alpha\beta}$ and $\phi_{\alpha\beta}^{ZZ}$ from $[0-2\pi]$, as well as θ_{23} and δ_{CP} , considering a 1σ uncertainty of 10% and 15%, respectively. All the remaining oscillation parameters were fixed to their NO best fit values [109]. Refer to the text for details.

In Fig. 1, we display our results of the expected 95% C.L. sensitivities to the non-diagonal SME coefficients, namely the isotropic $|c_{\alpha\beta}| \lesssim 1.1 \times 10^{-24}$ (left panel) and Z -spatial $|c_{\alpha\beta}^{ZZ}| \lesssim 6.2 \times 10^{-24}$ (right panel), for the DUNE configuration. Here, we have marginalized over the corresponding LIV phases $\phi_{\alpha\beta}$ and $\phi_{\alpha\beta}^{ZZ}$ from $[0-2\pi]$, as well as the atmospheric mixing angle θ_{23} and the leptonic phase δ_{CP} , considering a 1σ uncertainty of 10% and 15%, respectively. All the remaining oscillation parameters were fixed to their best fit value with normal mass ordering [109]. Existing bounds from the Super-Kamiokande experiment constraint $|c_{e\mu}| < 8.0 \times 10^{-27}$ and $|c_{e\tau}| < 9.3 \times 10^{-25}$ at 95% C.L. [110], while IceCube sets $|c_{\mu\tau}| < 3.9 \times 10^{-28}$ at 99% C.L. [111]. Furthermore, a test for Lorentz and CPT violation with the MiniBooNE low-energy excess restrict $c_{e\mu}^{ZZ} < (2.6 \pm 0.8) \times 10^{-19}$ [96, 112], while limits from the Double Chooz experiment set $\text{Re}(c_{e\tau}^{ZZ}) < 4.9 \times 10^{-17}$ [113]. For projected sensitivities and some experimental constraints on the LIV coefficients, $c_{\alpha\beta}$ and $c_{\alpha\beta}^{ZZ}$, see Tab. II.

Regarding the diagonal SME coefficients $c_{\alpha\alpha} - c_{\tau\tau}$ and $c_{\alpha\alpha}^{ZZ} - c_{\tau\tau}^{ZZ}$, we have fixed $c_{\tau\tau} = 0$,

as well as $c_{\tau\tau}^{ZZ} = 0$, since we can redefine the diagonal elements up to a global constant.




Figure 2: Expected 95% C.L. sensitivities to the diagonal SME coefficients $c_{\alpha\alpha}^{\mu\nu} - c_{\tau\tau}^{\mu\nu}$ at the DUNE (TDR setup). Here, we have marginalized over the corresponding mixing angle θ_{23} and δ_{CP} , considering a 1σ uncertainty of 10% and 15%, respectively. All the remaining oscillation parameters were fixed to their NO best fit values [109]. Refer to the text for details.

Our results of the projected 95% C.L. sensitivities to the $c_{\alpha\alpha} - c_{\tau\tau}$ (left panel) and $c_{\alpha\alpha}^{ZZ} - c_{\tau\tau}^{ZZ}$ (right panel) coefficients at the DUNE experiment are shown in Fig. 2. We have marginalized over the corresponding atmospheric mixing angle θ_{23} and Dirac CP -violating phase δ_{CP} , considering a 1σ uncertainty of 10% and 15%, respectively. All the remaining oscillation parameters were fixed to their best fit value with normal mass ordering [109]. We observe that the DUNE will be able to set limits on the isotropic CPT -even SME coefficients $c_{ee} - c_{\tau\tau} < 1.4 \times 10^{-24}$ and $c_{\mu\mu} - c_{\tau\tau} < 2.3 \times 10^{-24}$, while for the Z -spatial LIV coefficients $c_{ee}^{ZZ} - c_{\tau\tau}^{ZZ} < 5.1 \times 10^{-24}$ and $c_{\mu\mu}^{ZZ} - c_{\tau\tau}^{ZZ} < 7.1 \times 10^{-24}$, all of them at 95% confidence level. For instance, the null observation of LIV at the IceCube neutrino observatory constraint $\text{Re}(c_{ee}) < 6 \times 10^{-33}$ and $\text{Re}(c_{\tau\tau}) < 8 \times 10^{-34}$ with a Bayes factor > 31.6 [76].




Figure 3: Projected 95% C.L. sensitivities at the DUNE (TDR setup), in the $|c_{\alpha\beta}^{ZZ}| - |c_{\alpha\beta}|$ and $(c_{\alpha\alpha}^{ZZ} - c_{\tau\tau}^{ZZ}) - (c_{\alpha\alpha} - c_{\tau\tau})$ planes, respectively. Here, we have marginalized over the corresponding LIV phases $\phi_{\alpha\beta}$ and $\phi_{\alpha\beta}^{ZZ}$ from $[0-2\pi]$, as well as θ_{23} and δ_{CP} , considering a 1σ uncertainty of 10% and 15%, respectively. All the remaining oscillation parameters were fixed to their NO best fit values [109]. Refer to the text for details.

In Fig. 3, we display the expected 95% C.L. sensitivity regions for the case of the DUNE setup in the $|c_{\alpha\beta}^{ZZ}| - |c_{\alpha\beta}|$ and $(c_{\alpha\alpha}^{ZZ} - c_{\tau\tau}^{ZZ}) - (c_{\alpha\alpha} - c_{\tau\tau})$ planes, left and right panel, respectively. We have marginalized over the corresponding LIV phases $\phi_{\alpha\beta}$ and $\phi_{\alpha\beta}^{ZZ}$ from $[0-2\pi]$, as well as the atmospheric mixing angle θ_{23} and the leptonic CP phase δ_{CP} , considering a 1σ uncertainty of 10% and 15%, accordingly. It is observed that flavor-changing LIV coefficients from the $(e - \mu)$ and $(\mu - \tau)$ sectors may be constrained approximately three times more effectively by the DUNE configuration than by those from the $(e - \tau)$ sector, namely $|c_{e\tau}| < 1.3 \times 10^{-24}$ ($|c_{e\tau}^{ZZ}| = 0$), and $|c_{e\tau}^{ZZ}| < 6.9 \times 10^{-24}$ ($|c_{e\tau}| = 0$), all limits at 95% C.L.




Figure 4: Expected 95% C.L. sensitivities $|c_{\alpha\beta}^{TZ}|$ and $c_{\alpha\alpha}^{TZ} - c_{\tau\tau}^{TZ}$ at the DUNE (TDR setup). Here, we have marginalized over the corresponding LIV phases $\phi_{\alpha\beta}^{TZ}$ from $[0-2\pi]$, as well as θ_{23} and δ_{CP} , considering a 1σ uncertainty of 10% and 15%, respectively. All the remaining oscillation parameters were fixed to their NO best fit values [109]. Refer to the text for details.

In Fig. 4, we show in left (right) panel, the projected 95% C.L. sensitivities to the SME coefficients $|c_{\alpha\beta}^{TZ}|$ and $c_{\alpha\alpha}^{TZ} - c_{\tau\tau}^{TZ}$, considering the DUNE configuration. We have marginalized over the corresponding LIV phases $\phi_{\alpha\beta}^{TZ}$ from $[0-2\pi]$, as well as the atmospheric mixing angle θ_{23} and the leptonic CP phase δ_{CP} , considering a 1σ uncertainty of 10% and 15%, accordingly. We observe that the sensitivities in the TZ -sector ($|c_{e\mu}^{TZ}|$, $|c_{\mu\tau}^{TZ}|$) $\lesssim 4.2 \times 10^{-24}$, are not as competitive as those from the isotropic and Z -spatial sectors ($|c_{e\mu}|$, $|c_{\mu\tau}|$) $\lesssim 0.35 \times 10^{-24}$, and ($|c_{e\mu}^{ZZ}|$, $|c_{\mu\tau}^{ZZ}|$) $\lesssim 1.2 \times 10^{-24}$, as illustrated in the left and right panels of Fig. 1. Which can be partially understood from their functional dependence in the Hamiltonian H_{LIV} of LIV (Eq. 39)

$$-H_{\text{LIV}} \simeq \frac{E_\nu}{2} [3c_{\alpha\beta} - c_{\alpha\beta}^{ZZ} - 2\hat{N}_Z c_{\alpha\beta}^{TZ} + \mathcal{O}(\hat{N}_Z^2)], \quad \hat{N}_Z \sim 0.1 . \quad (44)$$

Besides, the null observation of Lorentz violation employing the low-energy excess data set of the MiniBooNE experiment set $c_{e\mu}^{TZ} < (5.9 \pm 1.7) \times 10^{-20}$ [112], while bounds from the

Double Chooz experiment constraint $\text{Re}(c_{e\tau}^{TZ}) < 3.2 \times 10^{-17}$ [96, 113].

LIV sector	Limit (Sensitivity)
neutrino	$c_{\alpha\beta} \lesssim 10^{-34}$, IceCube [76]
neutrino	$ c_{e\mu} < 8.0 \times 10^{-27}$, Super-Kamiokande [110]
neutrino	$ c_{e\tau} < 9.3 \times 10^{-25}$, Super-Kamiokande [110]
neutrino	$ c_{\mu\tau} < 3.9 \times 10^{-28}$, IceCube [111]
neutrino	$c_{e\mu}^{ZZ} < (2.6 \pm 0.8) \times 10^{-19}$, MiniBooNE [96, 112]
neutrino	$\text{Re}(c_{e\tau}^{ZZ}) < 4.9 \times 10^{-17}$, Double Chooz [113]
neutrino	($ c_{\alpha\beta} \sim [1 - 10] \times 10^{-25}$), DUNE [78, 79]
neutrino	($ c_{\alpha\beta} \sim [1 - 10] \times 10^{-25}$), DUNE (this work)
neutrino	($ c_{\alpha\beta}^{ZZ} \sim [1 - 10] \times 10^{-24}$), DUNE (this work)
neutrino	$\text{Re}(c_{ee}) < 6 \times 10^{-33}$, IceCube [76]
neutrino	$\text{Re}(c_{\tau\tau}) < 8 \times 10^{-34}$, IceCube [76]
neutrino	$(c_{ee} - c_{\tau\tau} \simeq 1.3 \times 10^{-24})$, DUNE (this work)
neutrino	$(c_{\mu\mu} - c_{\tau\tau} \simeq 2.2 \times 10^{-24})$, DUNE (this work)
neutrino	$(c_{ee}^{ZZ} - c_{\tau\tau}^{ZZ} \simeq 5.0 \times 10^{-24})$, DUNE (this work)
neutrino	$(c_{\mu\mu}^{ZZ} - c_{\tau\tau}^{ZZ} \simeq 7.0 \times 10^{-24})$, DUNE (this work)
electron	$-80 \lesssim c_{ee}/10^{-20} \lesssim 4$ [96]

Table II: Current limits and projected sensitivities (shown in parenthesis) relevant to this study.

V. CONCLUSIONS

In contemporary cosmology, one of the most fascinating puzzles are the dark matter and dark energy conundrum. One of the most common fields employed to describe dark energy is the scalar field. Furthermore, a scalar field can also characterize ultralight dark matter, which is one of the most promising alternatives for dark matter.

On the other hand, if the *CPT* and Lorentz symmetries are broken at very high energies, well beyond the electroweak scale, oscillations of high energy neutrinos may probe energy scales where this possible violations arise. The aforementioned violation of Lorentz

invariance might arise from neutrino non-standard interactions with scalar fields.

In this study, we have outlined the case of an effective tensorial neutrino-scalar field interaction. In this case, the scalar field could be identified as either a dark energy or dark matter candidate. Moreover, in this paper, we have studied the sensitivities to the dimension-four *CPT*-even SME coefficients $c_{\alpha\beta}^{\mu\nu}$ of the SME, particularly the isotropic $c_{\alpha\beta}$ and Z -spatial $c_{\alpha\beta}^{ZZ}$. For an accelerator-based experiment similar to DUNE, the predicted sensitivities at 95% C.L. are $|c_{\alpha\beta}| \sim [1 - 10] \times 10^{-25}$ (left panel of Fig. 1) and $|c_{\alpha\beta}^{ZZ}| \sim [1 - 10] \times 10^{-24}$ (right panel of Fig. 1), while for the diagonal LIV coefficients $c_{ee} - c_{\tau\tau} \simeq 1.3 \times 10^{-24}$ and $c_{\mu\mu} - c_{\tau\tau} \simeq 2.2 \times 10^{-24}$ (left panel of Fig. 2); $c_{ee}^{ZZ} - c_{\tau\tau}^{ZZ} \simeq 5.0 \times 10^{-24}$ and $c_{\mu\mu}^{ZZ} - c_{\tau\tau}^{ZZ} \simeq 7.0 \times 10^{-24}$ (right panel of Fig. 2).

Upcoming and present neutrino experiments such as the KM3NeT, IceCube-Gen2 and GRAND proposals, as well as the IceCube neutrino observatory, could shed more light on these types of neutrino interactions.

This paper represents the views of the authors and should not be considered a DUNE collaboration paper.

ACKNOWLEDGMENTS

This work was partially supported by SNII-México and CONAHCyT research Grant No. A1-S-23238. Additionally the work of R. C. was partially supported by COFAA-IPN, Estímulos al Desempeño de los Investigadores (EDI)-IPN and SIP-IPN Grant No. 20241624.

Appendix A: Details of a tensorial neutrino model

A possible realization of the tensorial coupling discussed in this work could arise from conformal coupling and disformal transformations where the scalar field couples to different kinds of matter fields in the form [88, 97, 114–117]

$$S_M = \int d^4x \sqrt{-\tilde{g}} \mathcal{L}_m(\tilde{\Psi}_i, \tilde{g}_{\mu\nu}^i) , \quad (\text{A1})$$

where \tilde{g} is the determinant of the metric $\tilde{g}_{\mu\nu}^i$ and the disformal transformation can be written as [118–120]

$$\tilde{g}_{\mu\nu} = a(\varphi, X) g_{\mu\nu} + b(\varphi, X) \partial_\mu \varphi \partial_\nu \varphi . \quad (\text{A2})$$

Disformal transformations have been applied to a wide range of topics in cosmology, from inflation [121] through dark matter [122], dark energy [123] and its cosmological implications [124]. Besides, disformal transformations have been applied in generalized Palatini gravities [125], and it has been studied their possible non-trivial effects on radiation and signatures in laboratory tests [126]. Following the construction presented in [97, 114] we can decompose the transformation in the following form ⁶

$$\tilde{g}_{\mu\nu} = g_{\mu\nu} + \alpha(\varphi, X, M_*)g_{\mu\nu} + \frac{\beta(\varphi, X)}{M_*^4}\partial_\mu\varphi\partial_\nu\varphi = g_{\mu\nu} + h_{\mu\nu}, \quad (\text{A3})$$

where M_* is related to energy scale of the interaction. Considering the $h_{\mu\nu}$ term as a small correction to $g_{\mu\nu}$, we can expand the action at first order and obtain a derivative coupling of the scalar field with matter

$$S_M = \int d^4x \sqrt{-g} \mathcal{L}_m(\Psi_i, g_{\mu\nu}^i) + \sum_i \int d^4x \sqrt{-g} \frac{1}{2} T_i^{\mu\nu} \left(\alpha(\varphi, X, M_*) g_{\mu\nu} + \frac{\beta(\varphi, X)}{M_*^4} \partial_\mu\varphi\partial_\nu\varphi \right), \quad (\text{A4})$$

where the sum considers different matter components and $T^{\mu\nu} = \frac{2}{\sqrt{-g}} \frac{\delta S_M}{\delta g_{\mu\nu}}$ is the energy momentum tensor of the matter fields. In the case of fermions

$$S_M = \int d^4x \sqrt{-g} \frac{-i}{2} (\bar{\Psi} \gamma^\mu D_\mu \Psi - D_\mu \bar{\Psi} \gamma^\mu \Psi + 2im \bar{\Psi} \Psi), \quad (\text{A5})$$

and the energy-momentum tensor is

$$T_{\mu\nu}^\Psi = -\frac{i}{2} (\bar{\Psi} \gamma_{(\mu} D_{\nu)} \Psi - D_{(\mu} \bar{\Psi} \gamma_{\nu)} \Psi). \quad (\text{A6})$$

Therefore, the interaction term can be written as

$$S_I = - \int d^4x \sqrt{-g} \frac{i}{4} (\bar{\Psi} \gamma_{(\mu} D_{\nu)} \Psi - D_{(\mu} \bar{\Psi} \gamma_{\nu)} \Psi) \left(\alpha(\varphi, X, M_*) g^{\mu\nu} + \frac{\beta(\varphi, X)}{M_*^4} \partial^\mu\varphi\partial^\nu\varphi \right). \quad (\text{A7})$$

It is convenient to select a particular choice of the α and β functions to construct the energy-momentum tensor $T_\varphi^{\mu\nu}$ for the scalar field φ in the interaction

$$S_I = \int d^4x \sqrt{-g} \frac{ik}{2M_*^4} (\bar{\Psi} \gamma_{(\mu} D_{\nu)} \Psi + \text{h.c.}) T_\varphi^{\mu\nu}. \quad (\text{A8})$$

The effective Lagrangian has a similar structure that appears in modified gravity [129] where there is a coupling of the energy-momentum tensor with the Einstein tensor and a proposal of

⁶ In fact, there are more general disformal transformations akin [127], $\bar{g}_{\mu\nu} = F_0 g_{\mu\nu} + F_1 \varphi_\mu \varphi_\nu + 2F_2 \varphi_{(\mu} X_{\nu)} + F_3 X_\mu X_\nu$ where $\varphi_\mu := \nabla_\mu \varphi$, $X_\mu := \nabla_\mu X$. The functions F_i depend on the variables φ, X, Y, Z , where $Y := \varphi^\mu X_\mu$ and $Z := X^\mu X_\mu$. However, the consistency of the fermionic coupling requires that $F_3 = 0$ [128].

a coupling among the energy-momentum tensor and vector and scalar fields [130]. Moreover, a similar interaction was proposed in Refs. [131, 132] in the context of Galileon–neutrino couplings. For neutrinos, the interaction term can be written as

$$S_I = \int d^4x \sqrt{-g} \frac{i\lambda_{\alpha\beta}}{2M_*^4} (\bar{\nu}_\alpha \gamma_{(\mu}(1 - \gamma_5) D_\nu) \nu_\beta + \text{h.c.}) T_\varphi^{\mu\nu}. \quad (\text{A9})$$

In cases where the Minkowski metric ($\eta_{\mu\nu}$) is a good approximation, the former relation reduces to

$$S_I = \int d^4x \sqrt{-\eta} \frac{i\lambda_{\alpha\beta}}{2M_*^4} (\bar{\nu}_\alpha \gamma_\mu(1 - \gamma_5) \partial_\nu \nu_\beta + \text{h.c.}) T_\varphi^{\mu\nu}. \quad (\text{A10})$$

- [1] A. G. Riess *et al.* (Supernova Search Team), *Astron. J.* **116**, 1009 (1998), arXiv:astro-ph/9805201.
- [2] S. Perlmutter *et al.* (Supernova Cosmology Project), *Astrophys. J.* **517**, 565 (1999), arXiv:astro-ph/9812133.
- [3] V. Sahni, *Lect. Notes Phys.* **653**, 141 (2004), arXiv:astro-ph/0403324.
- [4] U. Alam, V. Sahni, and A. A. Starobinsky, *JCAP* **06**, 008 (2004), arXiv:astro-ph/0403687.
- [5] A. G. Adame *et al.* (DESI), (2024), arXiv:2404.03002 [astro-ph.CO].
- [6] A. Arbey and F. Mahmoudi, *Prog. Part. Nucl. Phys.* **119**, 103865 (2021), arXiv:2104.11488 [hep-ph].
- [7] J. Magana and T. Matos, *J. Phys. Conf. Ser.* **378**, 012012 (2012), arXiv:1201.6107 [astro-ph.CO].
- [8] E. G. M. Ferreira, *Astron. Astrophys. Rev.* **29**, 7 (2021), arXiv:2005.03254 [astro-ph.CO].
- [9] A. Suárez, V. H. Robles, and T. Matos, *Astrophys. Space Sci. Proc.* **38**, 107 (2014), arXiv:1302.0903 [astro-ph.CO].
- [10] L. Hui, J. P. Ostriker, S. Tremaine, and E. Witten, *Phys. Rev. D* **95**, 043541 (2017), arXiv:1610.08297 [astro-ph.CO].
- [11] T. Matos, L. A. Ureña López, and J.-W. Lee, *Front. Astron. Space Sci.* **11**, 1347518 (2024), arXiv:2312.00254 [astro-ph.CO].
- [12] A. Arvanitaki, S. Dimopoulos, S. Dubovsky, N. Kaloper, and J. March-Russell, *Phys. Rev. D* **81**, 123530 (2010), arXiv:0905.4720 [hep-th].

- [13] M. Cicoli, V. Guidetti, N. Righi, and A. Westphal, *JHEP* **05**, 107 (2022), arXiv:2110.02964 [hep-th].
- [14] B. S. Acharya, K. Bobkov, and P. Kumar, *JHEP* **11**, 105 (2010), arXiv:1004.5138 [hep-th].
- [15] D. J. E. Marsh, *Phys. Rept.* **643**, 1 (2016), arXiv:1510.07633 [astro-ph.CO].
- [16] A. Arvanitaki, J. Huang, and K. Van Tilburg, *Phys. Rev. D* **91**, 015015 (2015), arXiv:1405.2925 [hep-ph].
- [17] A. Arvanitaki, S. Dimopoulos, and K. Van Tilburg, *Phys. Rev. Lett.* **116**, 031102 (2016), arXiv:1508.01798 [hep-ph].
- [18] A. Arvanitaki, P. W. Graham, J. M. Hogan, S. Rajendran, and K. Van Tilburg, *Phys. Rev. D* **97**, 075020 (2018), arXiv:1606.04541 [hep-ph].
- [19] M. Li, X.-D. Li, S. Wang, and Y. Wang, *Commun. Theor. Phys.* **56**, 525 (2011), arXiv:1103.5870 [astro-ph.CO].
- [20] C. Wetterich, *Nucl. Phys. B* **302**, 668 (1988), arXiv:1711.03844 [hep-th].
- [21] I. Zlatev, L.-M. Wang, and P. J. Steinhardt, *Phys. Rev. Lett.* **82**, 896 (1999), arXiv:astro-ph/9807002.
- [22] T. Chiba, T. Okabe, and M. Yamaguchi, *Phys. Rev. D* **62**, 023511 (2000), arXiv:astro-ph/9912463.
- [23] C. Armendariz-Picon, V. F. Mukhanov, and P. J. Steinhardt, *Phys. Rev. Lett.* **85**, 4438 (2000), arXiv:astro-ph/0004134.
- [24] C. Armendariz-Picon, V. F. Mukhanov, and P. J. Steinhardt, *Phys. Rev. D* **63**, 103510 (2001), arXiv:astro-ph/0006373.
- [25] A. Melchiorri, L. Mersini-Houghton, C. J. Odman, and M. Trodden, *Phys. Rev. D* **68**, 043509 (2003), arXiv:astro-ph/0211522.
- [26] T. Chiba, *Phys. Rev. D* **66**, 063514 (2002), arXiv:astro-ph/0206298.
- [27] L. P. Chimento and A. Feinstein, *Mod. Phys. Lett. A* **19**, 761 (2004), arXiv:astro-ph/0305007.
- [28] L. P. Chimento, *Phys. Rev. D* **69**, 123517 (2004), arXiv:astro-ph/0311613.
- [29] C. Armendariz-Picon, T. Damour, and V. F. Mukhanov, *Phys. Lett. B* **458**, 209 (1999), arXiv:hep-th/9904075.
- [30] J. Garriga and V. F. Mukhanov, *Phys. Lett. B* **458**, 219 (1999), arXiv:hep-th/9904176.
- [31] R. J. Scherrer, *Phys. Rev. Lett.* **93**, 011301 (2004), arXiv:astro-ph/0402316.
- [32] R. de Putter and E. V. Linder, *Astropart. Phys.* **28**, 263 (2007), arXiv:0705.0400 [astro-ph].

- [33] X.-T. Gao and R.-J. Yang, *Phys. Lett. B* **687**, 99 (2010), arXiv:1003.2786 [gr-qc].
- [34] A. Nicolis, R. Rattazzi, and E. Trincherini, *Phys. Rev. D* **79**, 064036 (2009), arXiv:0811.2197 [hep-th].
- [35] C. de Rham and A. J. Tolley, *JCAP* **05**, 015 (2010), arXiv:1003.5917 [hep-th].
- [36] G. L. Goon, K. Hinterbichler, and M. Trodden, *Phys. Rev. D* **83**, 085015 (2011), arXiv:1008.4580 [hep-th].
- [37] G. Goon, K. Hinterbichler, and M. Trodden, *JCAP* **07**, 017 (2011), arXiv:1103.5745 [hep-th].
- [38] C. Deffayet, O. Pujolas, I. Sawicki, and A. Vikman, *JCAP* **10**, 026 (2010), arXiv:1008.0048 [hep-th].
- [39] O. Pujolas, I. Sawicki, and A. Vikman, *JHEP* **11**, 156 (2011), arXiv:1103.5360 [hep-th].
- [40] R. Kimura and K. Yamamoto, *JCAP* **04**, 025 (2011), arXiv:1011.2006 [astro-ph.CO].
- [41] D. Maity, *Phys. Lett. B* **720**, 389 (2013), arXiv:1209.6554 [hep-th].
- [42] S. Weinberg, *Cosmology* (Oxford University Press, 2008).
- [43] A. Berlin, *Phys. Rev. Lett.* **117**, 231801 (2016), arXiv:1608.01307 [hep-ph].
- [44] P. F. de Salas, R. A. Lineros, and M. Tórtola, *Phys. Rev. D* **94**, 123001 (2016), arXiv:1601.05798 [astro-ph.HE].
- [45] G. Krnjaic, P. A. N. Machado, and L. Necib, *Phys. Rev. D* **97**, 075017 (2018), arXiv:1705.06740 [hep-ph].
- [46] V. Brdar, J. Kopp, J. Liu, P. Prass, and X.-P. Wang, *Phys. Rev. D* **97**, 043001 (2018), arXiv:1705.09455 [hep-ph].
- [47] A. Y. Smirnov and X.-J. Xu, *JHEP* **12**, 046 (2019), arXiv:1909.07505 [hep-ph].
- [48] A. Dev, P. A. N. Machado, and P. Martínez-Miravé, *JHEP* **01**, 094 (2021), arXiv:2007.03590 [hep-ph].
- [49] M. Losada, Y. Nir, G. Perez, and Y. Shpilman, *JHEP* **04**, 030 (2022), arXiv:2107.10865 [hep-ph].
- [50] M. Losada, Y. Nir, G. Perez, I. Savoray, and Y. Shpilman, *JHEP* **03**, 032 (2023), arXiv:2205.09769 [hep-ph].
- [51] G.-y. Huang, M. Lindner, P. Martínez-Miravé, and M. Sen, *Phys. Rev. D* **106**, 033004 (2022), arXiv:2205.08431 [hep-ph].
- [52] R. Cordero, L. A. Delgadillo, and O. G. Miranda, *Phys. Rev. D* **107**, 075023 (2023), arXiv:2207.11308 [hep-ph].

- [53] M. Sen and A. Y. Smirnov, [JCAP](#) **01**, 040 (2024), arXiv:2306.15718 [hep-ph].
- [54] J. Barranco, O. G. Miranda, C. A. Moura, T. I. Rashba, and F. Rossi-Torres, [JCAP](#) **10**, 007 (2011), arXiv:1012.2476 [astro-ph.CO].
- [55] M. M. Reynoso and O. A. Sampayo, [Astropart. Phys.](#) **82**, 10 (2016), arXiv:1605.09671 [hep-ph].
- [56] P.-H. Gu, X.-J. Bi, and X.-m. Zhang, [Eur. Phys. J. C](#) **50**, 655 (2007), arXiv:hep-ph/0511027.
- [57] S. Ando, M. Kamionkowski, and I. Mocioiu, [Phys. Rev. D](#) **80**, 123522 (2009), arXiv:0910.4391 [hep-ph].
- [58] N. Klop and S. Ando, [Phys. Rev. D](#) **97**, 063006 (2018), arXiv:1712.05413 [hep-ph].
- [59] F. Capozzi, I. M. Shoemaker, and L. Vecchi, [JCAP](#) **07**, 004 (2018), arXiv:1804.05117 [hep-ph].
- [60] Y. Farzan and S. Palomares-Ruiz, [Phys. Rev. D](#) **99**, 051702 (2019), arXiv:1810.00892 [hep-ph].
- [61] S.-F. Ge and H. Murayama, (2019), arXiv:1904.02518 [hep-ph].
- [62] T. Gherghetta and A. Shkerin, [Phys. Rev. D](#) **108**, 095009 (2023), arXiv:2305.06441 [hep-ph].
- [63] C. A. Argüelles, K. Farrag, and T. Katori, [PoS ICRC2023](#), 1415 (2023), arXiv:2402.18126 [hep-ph].
- [64] C. A. Argüelles, K. Farrag, and T. Katori, in *9th Meeting on CPT and Lorentz Symmetry* (2023) arXiv:2401.15716 [hep-ph].
- [65] G. Lambiase and T. K. Poddar, [JCAP](#) **01**, 069 (2024), arXiv:2307.05229 [hep-ph].
- [66] R. Cordero and L. A. Delgadillo, [Phys. Lett. B](#) **853**, 138687 (2024), arXiv:2312.16320 [hep-ph].
- [67] C. A. Argüelles, K. Farrag, and T. Katori, (2024), arXiv:2404.10926 [hep-ph].
- [68] V. A. Kostelecky and M. Mewes, [Phys. Rev. D](#) **69**, 016005 (2004), arXiv:hep-ph/0309025.
- [69] J. S. Diaz, [Symmetry](#) **8**, 105 (2016), arXiv:1609.09474 [hep-ph].
- [70] M. D. C. Torri, [Universe](#) **6**, 37 (2020), arXiv:2110.09186 [hep-ph].
- [71] C. A. Moura and F. Rossi-Torres, [Universe](#) **8**, 42 (2022).
- [72] G. Barenboim, [Front. in Phys.](#) **10**, 813753 (2022).
- [73] D. Colladay and V. A. Kostelecky, [Phys. Rev. D](#) **58**, 116002 (1998), arXiv:hep-ph/9809521.
- [74] G. Barenboim and J. D. Lykken, [Phys. Lett. B](#) **554**, 73 (2003), arXiv:hep-ph/0210411.

- [75] G. Barenboim, M. Masud, C. A. Ternes, and M. Tórtola, *Phys. Lett. B* **788**, 308 (2019), arXiv:1805.11094 [hep-ph].
- [76] R. Abbasi *et al.* (IceCube), *Nature Phys.* **18**, 1287 (2022), arXiv:2111.04654 [hep-ex].
- [77] S. Sahoo, A. Kumar, and S. K. Agarwalla, *JHEP* **03**, 050 (2022), arXiv:2110.13207 [hep-ph].
- [78] S. K. Agarwalla, S. Das, S. Sahoo, and P. Swain, *JHEP* **07**, 216 (2023), arXiv:2302.12005 [hep-ph].
- [79] D. Raikwal, S. Choubey, and M. Ghosh, *Phys. Rev. D* **107**, 115032 (2023), arXiv:2303.10892 [hep-ph].
- [80] F. Testagrossa, D. F. G. Fiorillo, and M. Bustamante, (2023), arXiv:2310.12215 [astro-ph.HE].
- [81] F. N. Díaz, J. Hoefken, and A. M. Gago, *Phys. Rev. D* **102**, 055020 (2020), arXiv:2003.13712 [hep-ph].
- [82] G. Barenboim and N. E. Mavromatos, *JHEP* **01**, 034 (2005), arXiv:hep-ph/0404014.
- [83] V. De Romeri, C. Giunti, T. Stuttard, and C. A. Ternes, *JHEP* **09**, 097 (2023), arXiv:2306.14699 [hep-ph].
- [84] G. Barenboim and A. M. Gago, (2024), arXiv:2402.03438 [hep-ph].
- [85] L. A. Delgadillo, O. G. Miranda, G. Moreno-Granados, and C. A. Moura, In preparation.
- [86] B. Abi *et al.* (DUNE), (2020), arXiv:2002.03005 [hep-ex].
- [87] V. Antonelli, L. Miramonti, and M. D. C. Torri, *Eur. Phys. J. C* **78**, 667 (2018), arXiv:1803.08570 [hep-ph].
- [88] C. S. Gauthier, R. Saotome, and R. Akhoury, *JHEP* **07**, 062 (2010), arXiv:0911.3168 [hep-ph].
- [89] A. R. Khalifeh and R. Jimenez, *Phys. Dark Univ.* **31**, 100777 (2021), arXiv:2010.08181 [gr-qc].
- [90] A. R. Khalifeh and R. Jimenez, *Phys. Dark Univ.* **34**, 100897 (2021), arXiv:2105.07973 [astro-ph.CO].
- [91] A. R. Khalifeh and R. Jimenez, *Phys. Dark Univ.* **37**, 101063 (2022), arXiv:2111.15249 [astro-ph.CO].
- [92] G.-Y. Huang and N. Nath, *Eur. Phys. J. C* **78**, 922 (2018), arXiv:1809.01111 [hep-ph].
- [93] P. F. de Salas, K. Malhan, K. Freese, K. Hattori, and M. Valluri, *JCAP* **10**, 037 (2019), arXiv:1906.06133 [astro-ph.GA].

- [94] P. F. de Salas and A. Widmark, *Rept. Prog. Phys.* **84**, 104901 (2021), arXiv:2012.11477 [astro-ph.GA].
- [95] S. Sivertsson, J. I. Read, H. Silverwood, P. F. de Salas, K. Malhan, A. Widmark, C. F. P. Laporte, S. Garbari, and K. Freese, *Mon. Not. Roy. Astron. Soc.* **511**, 1977 (2022), arXiv:2201.01822 [astro-ph.GA].
- [96] V. A. Kostelecky and N. Russell, *Rev. Mod. Phys.* **83**, 11 (2011), arXiv:0801.0287 [hep-ph].
- [97] P. Brax and C. Burrage, *Phys. Rev. D* **90**, 104009 (2014), arXiv:1407.1861 [astro-ph.CO].
- [98] B. Telalovic and M. Bustamante, (2023), arXiv:2310.15224 [astro-ph.HE].
- [99] B. Abi *et al.* (DUNE), *Eur. Phys. J. C* **80**, 978 (2020), arXiv:2006.16043 [hep-ex].
- [100] P. Huber, M. Lindner, and W. Winter, *Comput. Phys. Commun.* **167**, 195 (2005), arXiv:hep-ph/0407333 [hep-ph].
- [101] P. Huber, J. Kopp, M. Lindner, M. Rolinec, and W. Winter, *Comput. Phys. Commun.* **177**, 432 (2007), arXiv:hep-ph/0701187 [hep-ph].
- [102] J. Kopp, *Int. J. Mod. Phys. C* **19**, 523 (2008), arXiv:physics/0610206.
- [103] J. Kopp, M. Lindner, T. Ota, and J. Sato, in *15th International Conference on Supersymmetry and the Unification of Fundamental Interactions (SUSY07)* (2007) pp. 756–759, arXiv:0710.1867 [hep-ph].
- [104] B. Abi *et al.* (DUNE), (2021), arXiv:2103.04797 [hep-ex].
- [105] V. A. Kostelecky and M. Mewes, *Phys. Rev. D* **70**, 076002 (2004), arXiv:hep-ph/0406255.
- [106] J. S. Diaz, V. A. Kostelecky, and M. Mewes, *Phys. Rev. D* **80**, 076007 (2009), arXiv:0908.1401 [hep-ph].
- [107] S. Mishra, S. Shukla, L. Singh, and V. Singh, (2023), arXiv:2309.01756 [hep-ph].
- [108] P. Huber, M. Lindner, and W. Winter, *Nucl. Phys. B* **645**, 3 (2002), arXiv:hep-ph/0204352.
- [109] P. F. de Salas, D. V. Forero, S. Gariazzo, P. Martínez-Miravé, O. Mena, C. A. Ternes, M. Tórtola, and J. W. F. Valle, *JHEP* **02**, 071 (2021), arXiv:2006.11237 [hep-ph].
- [110] K. Abe *et al.* (Super-Kamiokande), *Phys. Rev. D* **91**, 052003 (2015), arXiv:1410.4267 [hep-ex].
- [111] M. G. Aartsen *et al.* (IceCube), *Nature Phys.* **14**, 961 (2018), arXiv:1709.03434 [hep-ex].
- [112] T. Katori (MiniBooNE), in *5th Meeting on CPT and Lorentz Symmetry* (2010) pp. 70–74, arXiv:1008.0906 [hep-ph].

- [113] T. Katori and J. Spitz, in *6th Meeting on CPT and Lorentz Symmetry* (2014) pp. 9–12, arXiv:1307.5805 [hep-ph].
- [114] P. Brax, *Phys. Lett. B* **712**, 155 (2012), arXiv:1202.0740 [hep-ph].
- [115] M. Carrillo González, Q. Liang, J. Sakstein, and M. Trodden, *JCAP* **04**, 063 (2021), arXiv:2011.09895 [astro-ph.CO].
- [116] H. Yazdani Ahmadabadi and H. Mohseni Sadjadi, *Phys. Lett. B* **850**, 138493 (2024), arXiv:2201.02927 [hep-ph].
- [117] H. Yazdani Ahmadabadi and H. Mohseni Sadjadi, (2022), arXiv:2209.13367 [hep-ph].
- [118] J. D. Bekenstein, *Phys. Rev. D* **48**, 3641 (1993), arXiv:gr-qc/9211017.
- [119] C. Deffayet and S. Garcia-Saenz, *Phys. Rev. D* **102**, 064037 (2020), arXiv:2004.11619 [hep-th].
- [120] M. Zumalacárregui and J. García-Bellido, *Phys. Rev. D* **89**, 064046 (2014), arXiv:1308.4685 [gr-qc].
- [121] N. Kaloper, *Phys. Lett. B* **583**, 1 (2004), arXiv:hep-ph/0312002.
- [122] J. D. Bekenstein, *Phys. Rev. D* **70**, 083509 (2004), [Erratum: Phys.Rev.D 71, 069901 (2005)], arXiv:astro-ph/0403694.
- [123] M. Zumalacárregui, T. S. Koivisto, D. F. Mota, and P. Ruiz-Lapuente, *JCAP* **05**, 038 (2010), arXiv:1004.2684 [astro-ph.CO].
- [124] C. van de Bruck, J. Morrice, and S. Vu, *Phys. Rev. Lett.* **111**, 161302 (2013), arXiv:1303.1773 [astro-ph.CO].
- [125] G. J. Olmo, H. Sanchis-Alepuz, and S. Tripathi, *Phys. Rev. D* **80**, 024013 (2009), arXiv:0907.2787 [gr-qc].
- [126] P. Brax, C. Burrage, and A.-C. Davis, *Journal of Cosmology and Astroparticle Physics* **2012**, 016 (2012).
- [127] T. Ikeda, K. Takahashi, and T. Kobayashi, *Phys. Rev. D* **108**, 044006 (2023), arXiv:2302.03418 [gr-qc].
- [128] K. Takahashi, R. Kimura, and H. Motohashi, *Phys. Rev. D* **107**, 044018 (2023), arXiv:2212.13391 [gr-qc].
- [129] P. Asimakis, S. Basilakos, A. Lymperis, M. Petronikolou, and E. N. Saridakis, *Phys. Rev. D* **107**, 104006 (2023), arXiv:2212.03821 [gr-qc].

- [130] J. Beltrán Jiménez, J. A. R. Cembranos, and J. M. Sánchez Velázquez, *JHEP* **05**, 100 (2018), arXiv:1803.05832 [hep-th].
- [131] A. Kehagias, (2011), arXiv:1109.6312 [hep-ph].
- [132] E. Ciuffoli, J. Evslin, J. Liu, and X. Zhang, *ISRN High Energy Phys.* **2013**, 497071 (2013), arXiv:1109.6641 [hep-ph].