

NA 60 results on p_T spectra and the spectral function in In-In collisions

J. Seixas⁵ for the NA 60 Collaboration:

R. Amaldi¹⁰, R. Averbeck⁹, K. Banicz^{2,4}, J. Castor³,
 B. Chaurand⁷, C. Cicalo¹, A. Colla¹⁰, P. Cortese¹⁰,
 S. Damjanovic², A. David^{2,5}, A. De Falco¹, A. Devaux³, A.
 Dees⁹, L. Ducroux⁶, H. En'yo⁸, A. Ferretti¹⁰, M. Floris¹,
 A. Forster², P. Force³, N. Guettet³, A. Guichard⁶,
 H. Gulkanian¹¹, M. Keil^{2,5}, L. Kluber⁷, C. Lourenco²,
 J. Lozano⁵, F. Manso³, A. Masoni¹, P. Martinis^{2,5},
 A. Neves⁵, H. Ohnishi⁸, C. Oddo¹⁰, P. Parracho²,
 P. Pillot⁶, G. Puddu¹, E. Rademacher², P. Ramalho²,
 P. Rosinsky², E. Scomparin¹⁰, S. Serci¹, R. Shahoyan^{2,5},
 P. Sonderegger⁵, H. J. Specht⁴, R. Teulent⁶, G. U sai¹,
 R. Veenhof², H. K. W. Ohri¹

¹Univ. di Cagliari and INFN, Cagliari, Italy, ²CERN, Geneva, Switzerland,
³LPC, Univ. Blaise Pascal and CNRS-IN2P3, Clermont-Ferrand, France, ⁴Univ.
 Heidelberg, Heidelberg, Germany, ⁵LIP, Lisbon, Portugal, ⁶IPN-Lyon, Univ.
 Claude Bernard Lyon-I and CNRS-IN2P3, Lyon, France, ⁷LLR, Ecole
 Polytechnique and CNRS-IN2P3, Palaiseau, France, ⁸RIKEN, Wako, Saitama,
 Japan, ⁹SUNY Stony Brook, New York, USA, ¹⁰Univ. di Torino and INFN,
 Italy, ¹¹YerPhI, Yerevan, Armenia

Abstract. The NA 60 experiment at the CERN SPS has studied low-mass muon pairs in 158 A GeV In-In collisions. A strong excess of pairs is observed above the yield expected from neutral meson decays. The unprecedented sample size of close to 400K events and the good mass resolution of about 2% have made it possible to isolate the excess by subtraction of the decay sources (keeping the p_T). The shape of the resulting mass spectrum exhibits considerable broadening, but essentially no shift in mass. The acceptance-corrected transverse-momentum spectra have a shape atypical for radial flow and show a significant mass dependence, pointing to different sources in different mass regions.

1. Introduction

Thermal dilepton production in the mass region below $1 \text{ GeV}/c^2$ is largely mediated by the light vector mesons ρ , ω and ϕ . Among these, the ρ ($770 \text{ MeV}/c^2$) is the most important, due to its strong coupling to the $\pi\pi$ channel and its short lifetime of only $1.3 \text{ fm}/c$. Changes both in width and in mass were originally suggested as precursor signatures of the chiral transition [1] and subsequent models have tied these changes directly [2] or indirectly [3] to chiral symmetry restoration.

The first NA 60 results focused on the space-time averaged spectral function of the ρ [4]; more details were added recently [5]. The present paper concentrates mainly

z This work was supported in part by FCT under grant POCI/FP/63919/2005




Figure 1. Left: M mass spectra of the opposite-sign dimuons (upper histogram), combinatorial background (dashed), signal fake matches (dashed-dotted), and resulting signal (histogram with error bars). Right: Isolation of an excess above the hadron decay cocktail. Total data (open circles), individual cocktail sources (solid), difference data (thick triangles), sum of cocktail sources and difference data (dashed).

on new developments, in particular first results on acceptance-corrected p_T and m_T spectra for different mass windows

2. Experimental results

Details of the NA 60 apparatus can be found in [6, 7], while the different analysis steps (including the critical assessment of the combinatorial background from π and K decays through event mixing) are described in [8]. The results reported here were obtained from the analysis of data taken in 2003 for In-In at 158 A GeV. The left part of Fig. 1 shows the opposite-sign, background and signal dimuon mass spectra, integrated over all collision centralities. After subtracting the combinatorial background and the signal fake matches, the resulting net spectrum contains about 360 000 muon pairs in the mass range 0–2 GeV/c², roughly 50% of the total available statistics. The associated average charged-particle multiplicity density measured by the vertex tracker is $\langle dN_{ch}/d\eta \rangle = 120$. Vector mesons ρ and ω are completely resolved with a mass resolution at the level of 20 MeV/c². Most of the analysis is done in four classes of collision centrality (defined through the charged-particle multiplicity density): peripheral (4–30), semi-peripheral (30–110), semi-central (110–170) and central (170–240).

The peripheral data can essentially be described by the expected electromagnetic decays of the neutral mesons [4, 5]. This is not the case in the more central bins, due to the presence of a strong excess. To isolate this excess without any fits, a novel procedure has been devised [4, 5, 8]. The resulting spectrum for all centralities and all p_T shows a peaked structure residing on a broad continuum. The same feature can be seen for each centrality bin, with a yield strongly increasing with centrality, but remaining essentially centered around the position of the nominal ρ pole.

A more quantitative analysis of the shape of the excess mass spectra vs. centrality has been performed, using a finer subdivision of the data into 12 centrality bins [5, 6]. On the basis of this analysis it is possible to rule out that the excess shape can be accounted for by the cocktail residing on a broad continuum, independent of centrality.




Figure 2. Left: Acceptance relative to 4 for different mass windows. Right: Rapidity distribution of the excess data for the mass window $0.6 < M < 0.9$ GeV/c² and for three selected p_T bins. The measured data (full markers) are reflected around midrapidity (open markers).




Figure 3. Acceptance-corrected p_T spectra for three mass windows and for three centrality bins. For discussion of errors see text.

A differential analysis in p_T was originally performed and mass spectra associated with three different p_T windows, without acceptance correction, were obtained [5]. The NA 60 acceptance relative to 4 as a function of p_T as shown in Fig. 2 (left), implies that one should in principle perform the acceptance correction using a 3-dimensional grid in $(M; p_T; y)$ space. This can lead, however, to large errors once the correction is applied. Instead, the correction is performed in 2-dimensional $(M; p_T)$ space, using the measured rapidity distribution as an input. The latter was determined with an acceptance correction found, in an iterative way, from Monte Carlo simulations matched to the data in M and p_T . On the basis of this rapidity distribution, 0.1 GeV/c² bins in M and 0.2 GeV/c bins in p_T were used to determine the remaining 2-dimensional correction. After correction the results were integrated over the three extended mass windows $0.4 < M < 0.6$, $0.6 < M < 0.9$ and $1.0 < M < 1.4$ GeV/c². In Fig. 2 (right) the rapidity distribution of the central mass window is shown for three different p_T windows, exhibiting a close resemblance to the distribution of inclusive pion production, as measured by NA 49 for Pb-Pb and NA 60 for In-In. The results for the acceptance-corrected p_T spectra are summarized in Fig. 3. The errors shown are purely statistical. Systematic errors arise from the acceptance

corrections including the rapidity distribution used, the subtraction of the cocktail, and the subtraction of the combinatorial background plus fake matches. For $p_T < 0.5$ GeV/c, the combinatorial background contributes most, ranging from 10 to 25% for semiperipheral up to central. For $p_T > 1$ GeV/c, the statistical errors dominate. The data show a significant dependence on mass, but hardly any on centrality. To bear out the differences in mass more clearly, the data were summed over the three more central bins and plotted in Fig. 4 (left) as a function of m_T . The inverse slope parameter T_e as determined from differential fits of the m_T spectra with $\exp(-m_T/T_e)$, using a sliding window in p_T , is plotted on the right. Instead of flattening at very low m_T as should be expected from radial flow, strikingly all spectra steepen, which is equivalent to very small values of T_e . For comparison, the resonance, placed just in between the upper two mass windows, does flatten as expected. Moreover, depending on the m_T region, T_e covers an unusually large dynamic range. Finally, the largest masses have the steepest m_T spectrum, i.e. the smallest value of T_e everywhere, again contrary to radial flow and to what is usually observed for hadrons. All this suggests that different mass regions are coupled to basically different emission sources. Summarizing, the




Figure 4. Left: Acceptance-corrected m_T spectra for three mass windows, summed over centralities (excluding the peripheral bin). Right: Inverse slope parameter T_e from differential fits of the m_T spectra (see text).

previously measured excess mass spectra and the new acceptance-corrected p_T and m_T spectra present an unexpected behaviour. Beyond the spectral function this may lead to a better understanding of the continuum part of the spectra for $M > 1$ GeV/c², possibly disentangling parton-hadron duality.

References

- [1] R. D. Pisarski, Phys. Lett. 110B, 155 (1982)
- [2] G. E. Brown, M. Rho, Phys. Rept. 363, 85 (2002)
- [3] R. Rapp and J. Wambach, Adv. Nucl. Phys. 25, 1 (2000)
- [4] R. A. Makhiet al. (NA 60 Collaboration), Phys. Rev. Lett. 96 (2006) 162302
- [5] S. Damjanovic et al. (NA 60 Collaboration), nucl-ex/0609026; S. Damjanovic et al. (NA 60 Collaboration), nucl-ex/0701015
- [6] G. Usai et al. (NA 60 Collaboration), Eur. Phys. J. C 43, 415 (2005); G. Usai et al. (NA 60 Collaboration), these proceedings.
- [7] M. Keil et al., Nucl. Instrum. Meth. A 539, 137 (2005) and A 546, 448 (2005)
- [8] R. Shahoyan et al. (NA 60 Collaboration), Nucl. Phys. A 774, 677 (2006) R. Shahoyan et al. (NA 60 Collaboration), these proceedings