

Multi field DBI Inflation and Non-Gaussianities

Min-xin Huang

Department of Physics, CERN - Theory Division, CH-1211 Geneva 23, Switzerland

Gary Shiu and Bret Underwood

Department of Physics, University of Wisconsin, Madison, WI 53706, USA

We analyze the trajectories for multi field DBI inflation, which can arise in brane inflation models, and show that the trajectories are the same as in typical slow roll inflation. We calculate the power spectrum and find that the higher derivative terms of the DBI action lead to a suppression of the contribution from the isocurvature perturbations. We also calculate the bispectrum generated by the isocurvature perturbation, and find that it leads to distinctive features.

INTRODUCTION

Scalar field theories with non-canonical kinetic terms provide novel realizations of the inflationary paradigm [1]. One interesting class of such models which have been studied extensively in recent years is DBI inflation [2, 3], characterized by kinetic terms which arise from the Dirac-Born-Infeld (DBI) action. A particularly appealing phenomenological feature of DBI inflation is that it can lead to strong and unique non-Gaussian signatures in the Cosmic Microwave Background (CMB) [3, 4]. The DBI action is ubiquitous in string theory, e.g., as an effective theory for worldvolume degrees of freedom on branes. When embedded in brane inflation [5], the inflation field in DBI models can be given a natural geometrical interpretation as the position of a D-brane in extra dimensions. DBI inflation arises when the D-brane moves in a highly warped region of the internal space where the speed limit is small, and reduces to the usual slow roll brane inflation (with canonical kinetic term) when the brane is moving non-relativistically with respect to the local warp factor. Since the position of the brane in each compact direction is described by a scalar field, brane inflation is naturally a multi field inflationary model.

Multi field models are characterized by their trajectories in field space and can in general be decomposed into an adiabatic field, which parameterizes motion along the trajectory, and isocurvature fields, which describe the directions perpendicular to the trajectory. Features in the trajectory, such as a sharp turn, can convert isocurvature perturbations into adiabatic/curvature perturbations (even on superhorizon scales) and can give rise to interesting features in the primordial power spectrum and non-Gaussianity.

In this paper we study the effects of multiple fields in DBI inflation. In particular, we study the multi field DBI trajectories and show that they are identical to the usual slow-roll multi field case in which the trajectory is dominated by the field with the largest slope of the potential. We calculate the power spectrum for multiple DBI fields in the limit the trajectory makes a sharp turn and show that the contribution of the isocurvature perturbations

to the power spectrum is suppressed by the sound speed. Finally, we calculate the non-Gaussianity in the sharp turn limit and find that the non-Gaussianity is dominated at leading order in the sound speed by the usual single field DBI contribution, but has new multi field features at sub-leading order. We conclude by commenting on the impact of our results for DBI model building.

MULTIFIELD EQUATIONS OF MOTION

Consider a 10-dimensional warped throat background with the metric

$$ds^2 = f^{-1=2}(y) g_{\mu\nu} dx^\mu dx^\nu + f^{1=2}(y) g_{mn} dy^m dy^n \quad (1)$$

common to type IIB string compactifications [6, 7], where $f(y)$ is called the warp factor of the throat which can in principle depend on all of the coordinates of the internal space y .

A D3-brane in this background is described by (to lowest order in string coupling and to all orders in α'),

$$S_{DBI} = \int d^4x \sqrt{-g} \left[\frac{1}{f(\varphi_i)} - \frac{1}{2} \sum_i \left(\partial_\mu \varphi_i \right)^2 + V(\varphi_i) \right] \quad (2)$$

where the warp factor is rescaled $f(\varphi_i) = T_{D3}^{-1} f(y(\varphi_i))$, and the real canonical scalar fields associated with the motion of the brane are given by

$$\varphi_i = T_{D3}^{1=2} Y_i \quad (3)$$

where $i = 1, \dots, 6$. The potential $V(\varphi_i)$ can arise, for example, from interactions with $\overline{D}3$ -branes, $D7$ -branes, or from the breaking of the local isometries of the compact space, and we will leave it to be unspecified for the moment.

We will define the sound speed during inflation to be the inverse of the "Lorentz factor" of the DBI action for

arXiv:0709.3299v3 [hep-th] 15 Jan 2008

spatially homogeneous fields,

$$c_s = \frac{1}{\sqrt{1 - \sum_i \frac{X_i^2}{f_i}}} \quad (4)$$

We will be interested in the small sound speed limit $c_s \ll 1$ where the non-Gaussianity is observable. In the following we will use the convention that the y_i measure the distance from the "tip" of the throat and that the D3 brane moves towards the tip, so $\dot{y}_i < 0$.

Consider a FRW universe with four-dimensional metric,

$$ds^2 = dt^2 + a(t)^2 \sum_{i=1}^3 dx_i^2 \quad (5)$$

The Friedmann equation and equations of motion for the fields ϕ_i are (with $H = \frac{\dot{a}}{a}$)

$$3M_p^2 H^2 = \frac{1}{f(\phi_i)} \frac{1}{c_s^2} \sum_i \dot{\phi}_i^2 + V(\phi_i); \quad (6)$$

$$\frac{1}{a(t)^3} \frac{d}{dt} (a(t)^3 \frac{\dot{\phi}_i}{c_s}) = -\frac{\partial}{\partial \phi_i} V(\phi_i) + f(\phi_i)^{-1} (c_s^2 - 1) \quad (7)$$

Distributing the time derivative, the equation of motion (7) can also be written

$$\ddot{\phi}_i + 3H \dot{\phi}_i - \frac{c_s}{c_s} \ddot{\phi}_i + c_s \partial_i V + \frac{(c_s^2 - 1)}{f} \dot{\phi}_i = 0; \quad (8)$$

Clearly the equation of motion for a homogeneous scalar field with a canonical kinetic term is obtained from (7),(8) in the limit $c_s \rightarrow 1$.

We define the multi-field DBI inflationary parameters as [4]

$$\frac{H}{H^2} \quad (9)$$

$$M_{pl}^2 c_s \frac{\partial_i \partial_j V}{V}; \quad (10)$$

When these parameters are much smaller than one, the equations of motion (8) take a form similar to that of a slowly rolling field

$$3H \dot{\phi}_i \approx - \frac{\partial_i V}{c_s} + f^{-1} (c_s^2 - 1) \dot{\phi}_i; \quad (11)$$

where $\tan \alpha_{ij} = \frac{\partial_j V}{\partial_i V} = \frac{\partial_j V}{\partial_i V}$ to leading order in the DBI inflationary parameters (9-10).

As a specific example, for the "standard case" of an AdS warp factor that depends only on one of the fields




FIG. 1: The trajectory of a multiple field in a stationary system can be decomposed into an "adiabatic" field with components along the trajectory and an "entropy" field orthogonal to the trajectory.

$f(\phi_i) = \frac{1}{P} = 4$ and a separable potential of mass terms $V = \frac{1}{2} \sum_i m_i^2 \phi_i^2$, it can be shown that for small sound speed $c_s \ll 1$ and a large mass hierarchy ($m_i \gg m_j$ for some i, j) the multi-field DBI inflationary parameters are of order $O(\epsilon)$.

TRAJECTORIES IN MULTIFIELD BRANE INFLATION

To simplify our analysis we will restrict ourselves to a two-field model ($\phi_1; \phi_2$), but it is straightforward to generalize our analysis to any number of fields.

We will parameterize the classical trajectory by an "adiabatic field" that represents the component of the field motion along the trajectory [8] (see Figure 1),

$$\underline{\phi} = (\cos \alpha) \phi_1 + (\sin \alpha) \phi_2; \quad (12)$$

The angle α parameterizes the angle the classical trajectory makes with one of the field directions (here chosen to be ϕ_1) and should not be confused with the angular position of the D3 brane in the compact space, which is parameterized by the fields ϕ_i . The "entropy field" ϕ_\perp transverse to the classical trajectory gives rise to isocurvature fluctuations which are given by,

$$s = (\sin \alpha) \phi_1 + (\cos \alpha) \phi_2; \quad (13)$$

By definition, the entropy field is constant, $\dot{s} = 0$.

Using this parameterization, we can rewrite the exact equations of motion (8) as (in particular, the multi-field DBI parameters are not necessarily assumed small for

this expression),

$$-\frac{1}{c_s} + \frac{3H}{c_s} \frac{\partial \phi_2}{\partial \phi_1} + \partial_1 (V + f^{-1}(c_s - 1)) = 0 \quad (14)$$

$$\begin{aligned} -\frac{1}{c_s} &= \partial_1 (V + f^{-1}(c_s - 1)) \sin \theta \\ \partial_2 (V + f^{-1}(c_s - 1)) \cos \theta &: \end{aligned} \quad (15)$$

It is clear from (15) that the angle of the trajectory has a "fixed point" trajectory in field space — defined by

$$\tan \theta = \frac{\partial_2 (V + f^{-1}(c_s - 1))}{\partial_1 (V + f^{-1}(c_s - 1))} : \quad (16)$$

Furthermore this fixed point is stable to leading order when the slope of the potential is positive and dominates over the slope of the warp factor since small variations are driven to zero (notice that $\theta < 0$). This fixed point can also be seen from the equations of motion (11) in the "DBI slow roll regime" (e.g. when the multi field DBI parameters (9-10) are small),

$$\tan \theta = \frac{-2}{-1} \frac{\partial_2 (V + f^{-1}(c_s - 1))}{\partial_1 (V + f^{-1}(c_s - 1))} : \quad (17)$$

We see then that being in the DBI slow roll regime is equivalent to being at the stable "fixed point" of the trajectory, thus DBI slow roll is an attractor solution.

Let us examine (16) in more detail. First, we note that for an inflationary solution we require that the potential energy dominates over the kinetic energy, e.g. $V(\phi_i) \gg \dot{\phi}_i^2 = (c_s \dot{\phi}_i)^2$, so that for small c_s , $V(\phi_i) \gg c_s \dot{\phi}_i^2$ is automatically true. The trajectories (16) are now the same as in standard multi field inflation, in particular, the angle of the trajectory is controlled by the ratio of the curvatures of the potential: the trajectory follows the direction with the largest curvature. When the field with the largest curvature reaches its minimum the trajectory makes a sharp turn in field space, with the sharpness of the turn given by the ratio of the curvatures.

If we express the warped geometry as a cone over an angular base space X^5 ,

$$g_{mn} dy^m dy^n = dr^2 + ds_{X^5}^2 \quad (18)$$

we can identify one of the fields ϕ_1 as the radial coordinate and the other field ϕ_2 as one of the angular coordinates on the base X^5 of the D-brane. In this case we can roughly classify the trajectories based on which field dominates at early times, as shown in Figure 2: a radially dominated trajectory is when the slope of the potential in the angular direction ϕ_2 is much smaller than the radial direction ϕ_1 and so the trajectory is dominated at early times by motion in the radial direction; a diagonal trajectory is when the slopes of potentials of the two fields are approximately the same and so the trajectory is approximately a diagonal line composed of a linear combination of the radial and angular directions;




FIG. 2: Multi field models have a number of different trajectories, depending on the curvature of the potential for the fields. The sharpness of the turns is controlled by the ratio of the curvatures.

and an angularly dominated trajectory is when the slope of the potential in the radial direction ϕ_1 is the smallest so the trajectory is dominated at early times by motion in the angular ϕ_2 direction. Clearly the diagonal trajectory does not have a significant turn in field space, and so will not contribute to a generation of curvature perturbations as discussed above. In fact, the diagonal-type trajectory is just a linear combination of the fields and hence can be completely described by a single field, and so we will not consider this possibility.

THE POWER SPECTRUM OF MULTIFIELD DBI INFLATION

The equation of motion for the curvature perturbation R in a multi field model with fields ϕ_i can be solved exactly using the N formalism [9, 10], which states that the curvature perturbation is equal to the difference between the number of e-folds of the classical trajectory N_e and the perturbation to the classical trajectory N_e ,

$$\delta R = \delta N_e = N_e - \sum_i N_{\phi_i} \delta \phi_i ; \quad (19)$$

where $N_{\phi_i} = \frac{\partial N_e}{\partial \phi_i}$ is the derivative of the number of e-foldings with respect to the field evaluated when the mode exits the horizon ϕ_i . For two fields, the power spectrum is then given by,

$$\begin{aligned} 4^2 P_R &= h^2 = (N_{\phi_1})^2 h_{11}^2 \\ &+ 2(N_{\phi_1})(N_{\phi_2})h_{12} + (N_{\phi_2})^2 h_{22}^2 ; \end{aligned} \quad (20)$$

where we have allowed for cross correlation between the fields — for a canonical kinetic term, the cross coupling is zero. We see, then, that the power spectrum receives extra contributions in multi field inflation, both from cross

couplings and from the extra two point correlation functions of the additional fields.

We would like to analyze these extra contributions in more detail. Using the approximation that the DBI multi field parameters (10) are small and that the potential is separable $V(\tau_1; \tau_2) = V_1(\tau_1) + V_2(\tau_2)$ and dominates the energy density, the number of e-folds from the time of horizon crossing to the end of inflation is [10]

$$N_e = \frac{1}{M_p^2} \int_{Z^e}^Z \frac{V_1}{c_s \partial_{\tau_1} V_1} d\tau_1 + \frac{1}{M_p^2} \int_{Z^e}^Z \frac{V_2}{c_s \partial_{\tau_2} V_2} d\tau_2 : \quad (21)$$

As in [10], we can write the curvature perturbation (19) in a much simpler way by using a different set of multi field DBI parameters,

$$\tau_i = \frac{c_s M_p^2}{2} \frac{V_i^0}{V} \quad (22)$$

where a prime denotes a derivative with respect to the argument. Note that $\tau = H^{-2} = \tau_1 + \tau_2$. The curvature perturbation can now be written,

$$\zeta = \frac{1}{M_p^2} \frac{V_1 + Z^e}{2\tau_1} \delta\tau_1 + \frac{1}{M_p^2} \frac{V_2 + Z^e}{2\tau_2} \delta\tau_2 ; \quad (23)$$

with

$$Z^e = \frac{V_2^e \tau_1^e - V_1^e \tau_2^e}{e} \quad (24)$$

where the superscript 'e' denotes evaluation at the end of inflation. In the limit that the trajectory is in the radial or angular direction as shown above then at the end of inflation the one of the fields has settled into its minimum (say, τ_2 for concreteness) so $Z^e = V_2^e = \text{const}$. For mass-term dominated potentials (or equivalently when the vacuum energy from τ_2 is small $V_2^e \ll V_1$) then we can take $Z^e = 0$ and the expression for the curvature perturbation simplifies to include only the values of the potential and the slow roll parameters evaluated at horizon crossing,

$$\zeta = \frac{1}{M_p^2} \frac{V_1}{2\tau_1} \delta\tau_1 + \frac{1}{M_p^2} \frac{V_2}{2\tau_2} \delta\tau_2 : \quad (25)$$

This simple expression for the curvature perturbation will be useful later in evaluating the power spectrum.

The inflationary perturbation

In order to calculate the power spectrum we need to evaluate the two point correlation functions for the perturbations of the scalar fields in (20). In order to have

analytic control over our expressions we will assume that the trajectory is highly radial, e.g.,

$$\tan \theta = \frac{\dot{\tau}_2}{\dot{\tau}_1} = \frac{V_2^0}{V_1^0} c_s : \quad (26)$$

Under this assumption, the adiabatic perturbation simply the perturbation in the τ_1 direction and perturbations in the τ_2 direction are just isocurvature perturbations,

$$\delta\tau_2 = -\delta\tau_1 ; \quad \delta s = \delta\tau_2 : \quad (27)$$

Expanding the kinetic part of the Lagrangian in (2) to quadratic order in $\delta\tau$ and δs , we find

$$L_2 = \frac{a^3}{2c_s^3} [-\dot{\delta\tau}^2 - a^2 c_s^2 (r \delta\tau)^2] + \frac{a^3}{2c_s} [-\dot{\delta s}^2 - a^2 (r \delta s)^2] \quad (28)$$

Note that the isocurvature fluctuations (e.g. the fluctuations in the angular direction) scale differently with the sound speed; we will see soon that this has important consequences for the two point functions.

The quantization of the perturbations proceeds as usual with

$$(\tau; \mathbf{x}) = \frac{1}{(2\pi)^3} \int d^3k u(\tau; \mathbf{k}) a(\mathbf{k}) + u(\tau; -\mathbf{k}) \tilde{a}^\dagger(\mathbf{k}) e^{i\mathbf{k} \cdot \mathbf{x}} ;$$

where $u(\tau; \mathbf{k}) = \frac{H}{2k^3} (1 + ikc_s) e^{ikc_s \tau}$ (29)

for the adiabatic perturbation and

$$s(\tau; \mathbf{x}) = \frac{1}{(2\pi)^3} \int d^3k v(\tau; \mathbf{k}) b(\mathbf{k}) + v(\tau; -\mathbf{k}) \tilde{b}^\dagger(\mathbf{k}) e^{i\mathbf{k} \cdot \mathbf{x}} ;$$

where $v(\tau; \mathbf{k}) = H \frac{c_s}{2k^3} (1 + ik) e^{ik\tau}$ (30)

for the isocurvature modes. The creation and annihilation operators satisfy the usual commutation relation $[a(\mathbf{k}); a^\dagger(\mathbf{k}^0)] = [b(\mathbf{k}); b^\dagger(\mathbf{k}^0)] = (2\pi)^3 \delta^3(\mathbf{k} - \mathbf{k}^0)$, $u(\tau; \mathbf{k})$ and $v(\tau; \mathbf{k})$ are the solutions of the quadratic Lagrangian whose normalizations are fixed by the Wronskian conditions¹, and $\tau = \frac{1}{aH}$ is the conformal time.

It is now straightforward to calculate the two point functions,

$$\langle \zeta(\mathbf{k}_1) \zeta(\mathbf{k}_2) \rangle = (2\pi)^3 \delta^3(\mathbf{k}_1 + \mathbf{k}_2) \frac{H^2}{2k_1^3} ; \quad (31)$$

$$\langle \zeta(\mathbf{k}_1) s(\mathbf{k}_2) \rangle = (2\pi)^3 \delta^3(\mathbf{k}_1 + \mathbf{k}_2) \frac{c_s H^2}{2k_1^3} : \quad (32)$$

¹ This can be checked by computing the commutator $[\tau(\mathbf{x}_1); p(\mathbf{x}_2)] = i^3 \delta^3(\mathbf{x}_1 - \mathbf{x}_2)$, where $p = \frac{\partial L_2}{\partial \dot{\tau}}$ is the canonical momentum.

Here $\langle \delta^2(k) \rangle = \langle (\delta^2; k) \rangle_{j_i=0}$, $s(k) = s(\delta^2; k)_{j_i=0}$, and the Hubble parameter and sound speed are evaluated at the time of horizon crossing. Here we see that the isocurvature perturbations generated by δ^2 are suppressed by a factor of c_s^{-1} compared to the adiabatic perturbations generated by δ^1 . This suppression can be traced back to the different c_s dependence found in the quadratic Lagrangian (28). We note that the exact same calculation follows through if the trajectory is dominated by the angular direction after the replacement $\delta^1 \rightarrow \delta^2$. Finally, notice also that in the limit of a straight-line trajectory there is no cross correlation between the fields, so the cross term in the power spectrum (20) vanishes. Unfortunately, non-straight line trajectories are beyond our analytic control so it is not clear if the cross coupling will be significant, although we expect that in the diagonal limit the two point function should reduce to that of a simple single field model where the results are well known.

Using the two point functions (31,32) and the expression for the curvature perturbation (25), the power spectrum (20) becomes,

$$P = \frac{H^2}{4^2 M_p^2} \frac{1}{2_1} \frac{V_1}{V}^2 + \frac{c_s}{2_2} \frac{V_2}{V}^2 \quad (33)$$

$$\frac{H^2}{4^2 M_p^2} \frac{1}{2_1} \frac{V_1}{V}^2$$

where in the last line we assume that the second term is small compared to the first term in the small c_s limit.

Since the contribution of the angular modes to the power spectrum is highly suppressed by the sound speed we find that multi field DBI reduces essentially to the single field case, in contrast to multiple field slow roll inflation where additional fields may become important when the trajectory makes sharp turns in field space [10].

Multi field Non-Gaussianity

In the following we study the non-Gaussianities in more details, and as it turns out there are some potentially observable differences from the single field DBI inflation. To compute the non-Gaussianities, we expand the DBI Lagrangian to higher order. The leading order and subleading order cubic terms are

$$L_3 = \frac{a^3}{2c_s^3} [-^3 a^2 c_s^2 - (r \quad) a^2 c_s^2 - (r \quad s^2)] \quad (34)$$

The leading contribution to the non-Gaussianity comes from the first two terms, and their size is well known

$$f_{NL} = \frac{1}{c_s^2} \quad (35)$$

Since the angular modes is suppressed by a factor of $\frac{1}{c_s}$ comparing to the radial mode, we see that the third term in (34) contributes a non-Gaussianity of order

$$f_{NL} = \frac{1}{c_s} \quad (36)$$

Although it is sub-leading comparing to the first two terms, it is still potentially observable by future experiments if the sound speed c_s is small enough. In particular, this effect can be larger than the sub-leading effect of order $\frac{1}{c_s^2}$ computed in [4] when the sound speed satisfies $c_s > \frac{1}{2}$.

The calculation of the three-point function is standard, see e.g. [4] for details,

$$\langle \delta^i(k_1) \delta^j(k_2) \delta^k(k_3) \rangle = i(2\pi)^3 \delta^3(k_1 + k_2 + k_3) \int d^3x v(0; k_1) v(0; k_2) u(0; k_3) \frac{\partial N}{\partial \phi_1} \frac{\partial N}{\partial \phi_2} \frac{1}{c_s^2} \int_{-1}^0 du \frac{d u(\delta^i; k_3)}{d} [(k_1 \cdot k) v(\delta^i; k_1) v(\delta^j; k_2)] + c.c. + \text{cyclic} \quad (37)$$

here "cyclic" means two other terms by cyclically permuting $k_1; k_2; k_3$. We can see the non-Gaussianity vanishes in the squeezed limit when one of the momentum $k_i \rightarrow 0$, since $\frac{d u(\delta^i; k_3)}{d} \sim k_3^2$ and there is factor of $k_1 \cdot k$. This is the same as in single field DBI inflation.

Going away from the squeezed limit, we can compute the above three point function assuming k_1, k_2, k_3 are of the same order of magnitude. The non-Gaussianity has a very interesting shape as the following

$$A(k_1; k_2; k_3) = \frac{k_3^2 (k_1^2 + 3k_1 k_2 + k_2^2) (k_1 \cdot k)}{(k_1 + k_2)^3} + \text{cyclic} \quad (38)$$

We plot the non-Gaussianity as $A(k_1; k_2; 1) = (k_1 k_2)$ following the convention of [4] in Figure 3.

We can see from Figure 4 that the non-Gaussianity has a very interesting new feature, namely, the sign of non-Gaussianity is different in the middle of the "folded triangle" limit where $k_3 = k_1 + k_2$ from most of the other region of the configuration space. For example, one can directly check for the configuration of a folded triangle $k_1 = k_2 = \frac{k_3}{2}$, the shape is negative $A(k_1; k_2; k_3) = \frac{1:088}{k_3^3}$. This feature is not present in any other known inflationary models, so it can be used as a distinctive signature of multi field DBI inflation. The sign of non-Gaussianity determines the sense of skewness in the CMB temperature and matter density, and thus this change in sign can lead to potentially interesting observational effects². However, we should caution that since this is a

² The convention for the sign of f_{NL} (which characterizes the level




FIG. 3: The shape of non-Gaussianity in multi-eld DBI inflation is shown through a plot of $A(k_1, k_2; 1) = (k_1 k_2)$ in (38). Notice that in the folded triangle limit $k_1 = k_2 = k_3 = 2$ the bispectrum is negative, and constitutes a distinctive signature of multi-eld DBI inflation in the small c_s limit. The presence of opposite signs of the non-Gaussianity may give rise to interesting observational effects.




FIG. 4: The negative part of the non-gaussianity in the folded triangle limit is shown.

sub-leading effect, it might be difficult to disentangle the signature from experimental data. It would be interesting to extend this analysis to more than two-elds, which may lead to additional enhancements or suppressions of the multi-eld non-Gaussianities [12].

of non-Gaussianity) has been a source of confusion in the literature. See [11] for a discussion.

We have shown that the two-point function of the extra "angular" scalar-eld direction during inflation is suppressed by a factor of c_s compared to the usual single-eld contribution, thus the multi-eld DBI observables simply reduce to the single-eld case (in the limit where one of the-elds dominates the trajectory). There have been a lot of recent attempts in building single-eld DBI inflation models consistent with all known compactification constraints and precision cosmological observations [13]. In particular, these works find that combining observational constraints of the amplitudes and tilt of the scalar and tensor perturbations and the primordial non-gaussianity together with limitations on the-eld range coming from compactification puts severe constraints on the viable parameter space of single-eld DBI models, ruling out the most simple models.

In general, since we see that the small sound speed in multi-eld DBI inflation suppresses the multi-eld effects, multi-eld DBI inflationary observables can be well approximated by their single-eld values. Note that this also implies that the dramatic $O(1)$ effects expected at the end of multi-eld DBI inflation due to the inhomogeneous surface of tachyon condensation examined in [14] are now instead suppressed by $O(c_s)$ and are subdominant.

Variations of the basic DBI model, such as its IR version [15, 16], models involving wrapped branes [17] or different warped geometries, may be able to evade the strong constraints of [13]; since we have not made explicit use of any particular model we expect our results to hold in the small sound speed limit of these models as well.

Note: As this manuscript was being prepared, a preprint [18] appeared which contains some overlap with this work.

Acknowledgments:

We would like to thank Thorsten Battefeld, Damien Easson, Louis Leblond, Liam McAllister, Sarah Shandera and Gianmario Tasinato for helpful discussions and comments. The work of MH, GS and BU was supported in part by NSF CAREER Award No. PHY-0348093, DOE grant DE-FG-02-95ER 40896, a Research Innovation Award and a Cottrell Scholar Award from Research Corporation.

-
- [1] J. Garriga and V. F. Mukhanov, Phys. Lett. B 458, 219 (1999) [arXiv:hep-th/9904176]; C. Amendarias-Picon, T. Damour and V. F. Mukhanov, Phys. Lett. B 458, 209 (1999) [arXiv:hep-th/9904075].
 - [2] E. Silverstein and D. Tong, Phys. Rev. D 70, 103505 (2004) [arXiv:hep-th/0310221].

- [3] M. Alishahina, E. Silverstein and D. Tong, *Phys. Rev. D* 70, 123505 (2004) [arXiv:hep-th/0404084].
- [4] X. Chen, M. x. Huang, S. Kachru and G. Shiu, *JCAP* 0701, 002 (2007) [arXiv:hep-th/0605045]; M. x. Huang and G. Shiu, *Phys. Rev. D* 74, 121301 (2006) [arXiv:hep-th/0610235].
- [5] G. R. Dvali and S. H. H. Tye, *Phys. Lett. B* 450, 72 (1999) [arXiv:hep-ph/9812483].
- [6] S. B. Giddings, S. Kachru and J. Polchinski, *Phys. Rev. D* 66, 106006 (2002) [arXiv:hep-th/0105097].
- [7] I. R. Klebanov, M. J. Strassler, *JHEP* 0008, 052 (2000) [arXiv:hep-th/0007191].
- [8] C. Gordon, D. Wands, B. A. Bassett and R. Maartens, *Phys. Rev. D* 63, 023506 (2000), [arXiv:astro-ph/0009131].
- [9] M. Sasaki and E. D. Stewart, *Prog. Theor. Phys.* 95, 71 (1996) [arXiv:astro-ph/9507001]; A. A. Starobinsky, *JETP Lett.* 42, 152 (1985) [*Pis'ma Zh. Eksp. Teor. z.* 42, 124 (1985)]; D. Wands, K. A. Malik, D. H. Lyth, A. R. Liddle, *Phys. Rev. D* 62 (2000) 043527 [arXiv:astro-ph/0003278]; D. Seery and J. E. Lidsey, *JCAP* 0509, 011 (2005) [arXiv:astro-ph/0506056].
- [10] F. Vernizzi, D. Wands, *JCAP* 0605 (2006) 019 [astro-ph/0603799].
- [11] E. Komatsu, talk at "Cosmology and Strings" Workshop, ICTP, Trieste, July 2007.
- [12] T. Battefeld and R. Easther, *JCAP* 0703, 020 (2007) [arXiv:astro-ph/0610296]; D. Battefeld and T. Battefeld, *JCAP* 0705, 012 (2007) [arXiv:hep-th/0703012].
- [13] S. Kocemuti, J. Maiden, G. Shiu and B. Underwood, *JHEP* 0609, 076 (2006) [arXiv:hep-th/0605189]; G. Shiu and B. Underwood, *Phys. Rev. Lett.* 98, 051301 (2007) [arXiv:hep-th/0610151]; D. Baumann and L. McAllister, arXiv:0705.0240 [hep-th]; R. Bean, S. E. Shandera, S. H. H. Tye, J. Xu, *JCAP* 05 (2007) 004, arXiv:hep-th/0702107v1; H. V. Peiris, D. Baumann, B. Friedman, A. Cooray, arXiv:0706.1240v1 [astro-ph];
- [14] L. Leblond and S. Shandera, arXiv:hep-th/0610321.
- [15] X. Chen, *JHEP* 0508, 045 (2005) [arXiv:hep-th/0501184]; X. Chen, *Phys. Rev. D* 72, 123518 (2005) [arXiv:astro-ph/0507053].
- [16] S. Thomas and J. Ward, arXiv:hep-th/0702229.
- [17] T. Kobayashi, S. Mukohyama and S. Kinoshita, arXiv:0708.4285 [hep-th]; M. Becker, L. Leblond and S. E. Shandera, arXiv:0709.1170 [hep-th].
- [18] D. A. Easson, R. Gregory, D. F. Mota, G. Tasinato and I. Zavala, arXiv:0709.2666 [hep-th].