

Thermal Dimuon Emission in In-In at the CERN SPS

R. Arnaldi¹⁰, K. Banicz^{3,5}, J. Castor⁴, B. Chaurand⁷, C. Cicalò², A. Colla^{9,10}, P. Cortese^{9,10}, S. Damjanovic^{3,5}, A. David⁶, A. de Falco^{1,2}, A. Devaux⁴, L. Ducroux¹¹, H. En'yo⁸, J. Farjeix⁴, A. Ferretti^{9,10}, M. Floris^{1,2}, A. Förster³, P. Force⁴, N. Guettet^{3,4}, A. Guichard¹¹, H. Gulkanyan¹², J. Heuser⁸, M. Keil^{3,6}, L. Kluberg^{3,7}, C. Lourenço³, J. Lozano⁶, F. Manso⁴, P. Martins⁶, A. Masoni², A. Neves⁶, H. Ohnishi⁸, C. Oppedisano¹⁰, P. Parracho^{3,6}, P. Pillot¹¹, T. Poghosyan¹², G. Puoddu^{1,2}, E. Radermacher³, P. Ramalhoto^{3,6}, P. Rosinsky³, E. Scomparin¹⁰, J. Seixas⁶, S. Serçi^{1,2}, R. Shahoyan^{3,6}, P. Sonderegger⁶, H.J. Specht⁵, R. Tieulent¹¹, G. Usai^{1,2}, R. Venhof⁶ and H.K. Wöhri^{2,6}

(NA60 Collaboration)

¹Univ. di Cagliari, Italy; ²INFN Cagliari, Italy; ³CERN, Geneva, Switzerland; ⁴LPC, Univ. Blaise Pascal and CNRS-IN2P3, Clermont-Ferrand, France; ⁵Univ. Heidelberg, Heidelberg, Germany; ⁶IST-CFTP and LIP, Lisbon, Portugal; ⁷LLR, Ecole Polytechnique and CNRS-IN2P3, Palaiseau, France; ⁸RIKEN, Wako, Saitama, Japan; ⁹Univ. di Torino, Italy; ¹⁰INFN Torino, Italy; ¹¹IPN-Lyon, Univ. Claude Bernard Lyon-I and CNRS-IN2P3, Lyon, France; ¹²YerPhI, Yerevan, Armenia

The NA60 experiment at the CERN SPS has studied low-mass dimuon production in 158A GeV In-In collisions. A significant excess of pairs is observed above the yield expected from neutral meson decays, consistent with a dominant contribution from $\pi\pi \rightarrow \rho \rightarrow \mu\mu$. This paper presents precision results on the mass and transverse momentum spectra of the excess pairs. The space-time averaged rho spectral function associated to the measured mass distribution shows a significant broadening, but essentially no mass shift. The slope parameter T_{eff} extracted from the spectra rises with dimuon mass up to the ρ , followed by a sudden decline above. While the initial rise is consistent with the expectations for radial flow of a hadronic decay source, the decline indicates a transition to an emission source with much smaller flow, possibly of partonic origin.

The ultimate goal of heavy ion collisions is the detection of signatures of a phase transition from hadronic matter to a deconfined and/or chirally restored medium. Lepton pairs are a powerful probe of the hot and dense fireball formed in high-energy nuclear collisions. They are produced during the entire space-time evolution of the fireball and freely escape, undisturbed by final state interactions. In the invariant mass region <1 GeV, thermal dilepton production is mediated by the broad vector meson $\rho(770)$. In the mass region >1 GeV, thermal dileptons may be produced in either the early partonic or the late hadronic phase of the fireball.

In contrast to real photons, virtual photons decaying into lepton pairs can be characterized by two variables, mass M and transverse momentum p_T . The mass distribution can be directly connected to the space-time averaged spectral function of the intermediate vector meson. The measurement of p_T spectra of lepton pairs may offer access to their emission region, as p_T encodes the key properties of the expanding fireball (temperature and transverse flow). While hadrons always receive the full asymptotic flow reached at the moment of decoupling from the




Figure 1: Left: Isolation of an excess above the electromagnetic decays of neutral mesons (see text). Total data (open circles), individual cocktail sources (solid), difference data (thick triangles), sum of cocktail sources and difference data (dashed). Open charm not subtracted. Right: Transverse momentum spectra of the rho-like mass region for three centrality windows, arbitrarily normalized. The errors are purely statistical

flowing medium^{5,6}, lepton pairs are continuously emitted during the evolution of the system. They are thus produced with small flow and high temperature at early times, and larger flow and smaller temperatures at later times.

The NA60 experiment measured low-mass dimuon production in 158A GeV In-In collisions. The centrality-integrated net mass spectrum after subtraction of backgrounds¹⁴ is shown in Fig. 1. It contains about 430 000 dimuons in the mass range ≤ 1.4 GeV. The mass resolution at the ω is 20 MeV. The data are divided in centrality bins using the measured charged track density. While peripheral data can be described as a superposition of expected contribution from the electromagnetic decays of neutral mesons (“hadron cocktail”), this is not possible for the total data, due to the existence of a strong excess of pairs. To isolate this excess without any fits, the decay sources are subtracted from the data using local criteria which are only based on the measured mass distribution. The procedure is illustrated in Fig. 1. The narrow ω and ϕ peaks are fixed in order to get a smooth underlying continuum after subtraction. The yield of the η relative to the ω and ϕ is fixed from the data at $p_T > 1.0$ GeV. This contribution is relevant only for masses ≤ 0.4 GeV. The η two-body and ω Dalitz decays are then fixed as well. The ratio η'/η is assumed to be 0.12¹⁵. The ρ is not subtracted. Open charm, measured to be 0.30 ± 0.06 of the total yield in the mass interval $1.2 < M < 1.4$ GeV¹⁶, is subtracted throughout (but not yet in Fig. 1), with the spectral shape in M and p_T as described by Pythia¹⁶. For a detailed discussion of the subtraction procedure and associated systematic uncertainties see Ref.^{20,21}. After subtraction of the meson decays and charm, the remaining sample contains ~ 150 000 dimuons. The subtracted data for the η , ω and ϕ provide the p_T spectra for these mesons and are used later for comparison. A strong excess is found, with respect to the contribution expected from the “cocktail” ρ (bound to the ω with $\rho/\omega = 1$, as measured in elementary collisions). The excess is centered around the nominal ρ pole position and is compatible with a strong broadening of the ρ spectral function and no mass shift. It monotonically increases and broadens with centrality. More details can be found in Ref.^{4,8}

The data are corrected for the acceptance and for the centrality-dependent reconstruction efficiencies, estimated with an overlay Monte Carlo simulation (a Monte Carlo dimuon is reconstructed on top of a real event)¹⁰. Results on acceptance-corrected p_T -spectra for the mass window $0.6 \leq M \leq 0.9$ GeV and non-peripheral collisions are shown in Fig. 1; equivalent data for other mass windows can be found Ref.¹⁰. The data of the three centrality windows agree within errors; this also holds for the other mass windows¹⁰. Fig. 2 (left) displays the centrality-integrated data vs. transverse mass m_T for four mass windows; the ϕ is included for comparison. At very low m_T , a steepening is observed in all four mass windows, reminiscent of pion spectra and opposite to the expectation for radial flow at masses above the pion mass.




Figure 2: Left: Transverse mass spectra of the excess for four mass windows summed over centralities (excluding the peripheral bin), in comparison to the ϕ (absolute normalization). Right: Inverse slope parameter T_{eff} vs. dimuon mass M for $dN_{ch}/d\eta > 30$. The open squares correspond to the fit lines in the left pane. Open charm is subtracted throughout. For explanation of the inserted symbols and the errors see text.

This steepening is not observed for the hadron spectra. The rise only disappears for very peripheral collisions ($4 < dN_{ch}/d\eta < 10$). The lines in Fig. 2 (left) are fits with the exponential function $1/m_T dN/dm_T \propto \exp(-m_T/T_{\text{eff}})$. The fits are restricted to the range $0.4 < p_T < 1.8$ GeV (roughly $0.1 < (m_T - M) < 1.2$ GeV), to exclude the increased rise at low m_T . Fig. 2 (right) shows T_{eff} vs. pair mass, for the mass windows defined in the left pane (open squares) and for a finer mass binning (closed triangles). NA60 results from an independent analysis¹⁶ of the intermediate mass region (“IMR”) $1.16 < M < 2.56$ GeV, corrected for the contribution from Drell-Yan pairs, are shown for comparison. The hadron data as obtained from the subtraction procedure are also included; the value for the η has been obtained by tuning the GENESIS code¹⁵ to the η Dalitz decay and then referring back to the required T_{eff} of the mother. The errors shown for the low-mass data (“LMR”) are purely statistical. The systematic error is smaller than the statistical one (see Ref.²¹).

The results shown in Fig. 2 (right) can be summarized and interpreted as follows. The slope parameter T_{eff} rises nearly linearly with mass up to about 270 MeV at the pole position of the ρ , followed by a sudden decline to values of 190-200 MeV for masses > 1 GeV. The excess yield in the mass region $0.2 < M < 0.9$ GeV is generally attributed to thermal radiation from the fireball, dominated by pion annihilation $\pi^+\pi^- \rightarrow \rho \rightarrow \mu^+\mu^-$ via an in-medium modified ρ ⁴. The NA60 data in this region are now nearly quantitatively described by theoretical models^{17,18,19}. The linear rise of T_{eff} with M over the whole region up to the ρ peak is reminiscent of radial flow of a hadronic source. The hadron data show a similar linear rise, but their absolute values are surprisingly close to the excess values, contrary to the expectation for the temperature-flow folding. Hadrons should have larger flow, being emitted at freeze-out, while leptons are emitted throughout the whole lifetime of the fireball. To solve this seeming contradiction the vacuum ρ was studied. It can be isolated disentangling the peak from the broad continuum (Fig. 1, left)^{9,10,17,18,19}. It is found $T_{\text{eff}} = 300 \pm 17$ MeV for the peak and 231 ± 7 for the underlying continuum in the window $0.6 < M < 0.9$ MeV. The subtraction of a ρ modelled with the measured T_{eff} from the continuum data leads to differences of 4–20 MeV in T_{eff} depending on the closeness to the pole. The following interpretation emerges: the ρ is maximally coupled to pions and it is thus representative of the true freeze-out parameters of the fireball. The other hadrons freeze-out earlier, due to their smaller coupling to the pions. The large gap in T_{eff} between the vacuum ρ and the excess points is consistent with the expectations for the temperature-flow folding of the leptons. The linear rise of T_{eff} with M is consistent with the expectations for radial flow of a hadronic source ($\pi\pi \rightarrow \rho$) decaying into lepton pairs. Theoretical understanding of these results is underway^{17,18,19}, but does not yet describe the data in a satisfactory way. The large gap (> 50 MeV) in T_{eff} between the vacuum ρ and the ω only disappears for the lowest

peripheral “pp-like” selection $4 < dN_{ch}/d\eta < 10$, with $T_{\text{eff}} = 198 \pm 6$ MeV for the ρ and 201 ± 4 MeV for the ω . This implies that the “hot ρ ” is intimately connected to pions, disappearing as the $\pi\pi$ contribution to ρ production vanishes (with only the “cocktail” ρ left). The sudden decline of T_{eff} at masses > 1 GeV is the other most remarkable feature of the present data, which may indicate a transition to a source of a different nature. Extrapolating the lower-mass trend to beyond 1 GeV, a jump by about 50 MeV down to a low-flow situation is extremely hard to reconcile with emission sources of dominantly hadronic origin. If the rise is due to flow, then the sudden loss of flow may be explained as a transition to a qualitatively different source, suggesting dominantly early (partonic) processes like $q\bar{q} \rightarrow \mu^+\mu^-$ for which flow has not yet built up¹⁸. While still controversial¹⁷, this may well represent the first direct evidence for thermal radiation of partonic origin.

In conclusion, we have found strong evidence for radial flow in the region of thermal dilepton emission which has previously been associated with the ρ spectral function. The transition to a low-flow region above may signal a transition from a hadronic to a partonic source.

We are grateful to H. van Hees, R. Rapp, T. Renk and J. Ruppert for useful discussions. We acknowledge support from the BMBF (Heidelberg group), the C. Gulbenkian Foundation, Lisbon, and the Swiss Fund Kidagan (YerPhi group) and FCT (Lisbon group). M. Floris acknowledges support from the European Union “Marie Curie” Programme for the participation at the “XLIII Rencontres de Moriond” Conference.

References

1. R. D. Pisarski, Phys. Lett. **110B**, 155 (1982).
2. G. E. Brown and M. Rho, Phys. Rept. **363**, 85 (2002).
3. R. Rapp and J. Wambach, Adv. Nucl. Phys. **25**, 1 (2000).
4. R. Arnaldi *et al.*, Phys. Rev. Lett. **96**, 162302 (2006).
5. E. Schnedermann, J. Sollfrank and U. W. Heinz, Phys. Rev. C **48**, 2462 (1993)
6. U. W. Heinz, hep-ph/0407360.
7. K. Kajantie, M. Kataja, L. D. McLerran and P. V. Ruuskanen, Phys. Rev. D **34** (1986) 811.; M. Asakawa, C. M. Ko and P. Levai, Phys. Rev. Lett. **70** (1993) 398; T. Renk, private communication (2007).
8. G. Agakichiev *et al.*, Eur. Phys. J. **C41**, 475 (2005).
9. S. Damjanovic *et al.*, Eur. Phys. J. C **49**, 235 (2007)
10. S. Damjanovic *et al.*, Nucl. Phys. A **783**, 327 (2007).
11. G. Usai *et al.* and J. Seixas *et al.*, J. Phys. G **34** (2007) S233 and S1023.
12. K. Banicz *et al.*, Nucl. Instrum. Meth. A **546**, 51 (2005).
13. G. Usai *et al.*, Eur. Phys. J. **C43**, 415 (2005).
14. R. Shahoyan *et al.*, Eur. Phys. J. **C43** (2005) 209.
15. S. Damjanovic, A. De Falco and H. Wöhri, NA60 Internal Note 2005-1.
16. R. Shahoyan *et al.*, J. Phys. G **34** (2007) S1029.
17. H. van Hees and R. Rapp, Phys. Rev. Lett. **97**, 102301 (2006), R. Rapp, J. Phys. G **34** (2007) S405, H. van Hees and R. Rapp, J. Phys. G **34** (2007) S1051.
18. T. Renk and J. Ruppert, Phys. Rev. C **77** (2008) 024907 [arXiv:hep-ph/0612113]; J. Phys. G **34** (2007) S1047; J. Ruppert, C. Gale, T. Renk, P. Lichard and J. I. Kapusta, Phys. Rev. Lett. **100** (2008) 162301 [arXiv:0706.1934 [hep-ph]].z
19. K. Dusling, D. Teaney and I. Zahed, Phys. Rev. C **75** (2007) 024908; K. Dusling and I. Zahed, hep-ph/0701253.
20. R. Arnaldi *et al.* Phys. Rev. Lett. **100** (2008) 022302
21. S. Damjanovic *et al.* *Proceedings of the 20th International Conference on Ultra-Relativistic Nucleus Nucleus Collisions (Quark Matter 2008)* arXiv:0805.4153 [nucl-ex]