

Highlights from the NA 60 experiment

Alessandro De Falco for the NA 60 Collaboration

R Amaldi^a, K Banicz^{c e}, K Borer^d, J Castor^f, B Chaurand^g,
 W Chen^h, C Cicalo^j, A Colla^{b a}, P Cortese^b, S Damjanovic^{c e},
 A David^{e n}, A de Falco^{k j}, A Devaux^f, L Ducrouxⁱ, H En'yo^l,
 J Fargeix^g, A Ferretti^{b a}, M Floris^{k j}, P Force^f, A Forster^e,
 N Guettet^{e f}, A Guichardⁱ, H Gulkanian^m, J Heuser^l,
 M Keil^{e n}, L Kluberg^g, Z Li^h, C Lourenco^e, J Lozanoⁿ,
 F Manso^f, P Martins^{e n}, A Masoni^j, A Nevesⁿ, H Ohnishi^l,
 C Oppedisano^{b a}, P Parracho^{e n}, P Pillotⁱ, T. Poghosyan^m,
 G Puddu^{k j}, E Rademacher^e, P Ramalho^{e n}, P Rosinsky^e,
 E Scomparin^a, J Seixasⁿ, S Sercik^{k j}, R Shahoyanⁿ,
 P Sonderegger^{e n}, H J Specht^c, R Tieulentⁱ, G Usai^{k j},
 R Veenhof^e, H Wohri^{j n}

^a INFN Torino, Italy^b Università di Torino, Italy^c Physikalisches Institut der Universität Heidelberg, Germany^d Laboratory for High Energy Physics, University of Bern, Bern, Switzerland^e CERN, 1211 Geneva 23, Switzerland^f LPC, Université Blaise Pascal and CNRS-IN 2P3, Clermont-Ferrand, France^g LLR, Ecole Polytechnique and CNRS-IN 2P3, Palaiseau, France^h Brookhaven National Laboratory, Upton, New York, USAⁱ IPN-Lyon, Université Claude Bernard Lyon-I and CNRS-IN 2P3, Lyon, France^j INFN Cagliari, Italy^k Università di Cagliari, Italy^l RIKEN, Wako, Saitama, Japan^m Yerevan Physics Institute, Yerevan, Armeniaⁿ Instituto Superior Técnico, Lisbon, Portugal

E-mail: alessandro.de.falco@ca.infn.it

Abstract. NA 60 measured dimuon production in p-A and In-In collisions at the CERN SPS. This paper presents a high statistics measurement of $\mu^+\mu^-$ meson production in In-In collisions at 158 A GeV. Both the transverse momentum, rapidity, decay angular distributions and the absolute yield were measured as a function of centrality. The results are compared to previous measurements in order to shed light on the long standing puzzle. In addition, highlights on $\mu^+\mu^-$ meson production and on the dimuon excess below the J/ψ mass are presented.

1. Introduction

Lepton pairs production is particularly suitable as a probe of the hot and dense medium created in high-energy heavy ion collisions, because they can access several Quark-Gluon Plasma (QGP) signatures. Among them is strangeness enhancement [1] that can be addressed by measuring the ϕ and ω mesons production. The production of thermal dileptons [2], that can be detected in the intermediate mass region (IMR) between the ρ and the J/ψ peaks, is another key observable.

NA 60 is an SPS third generation experiment designed to study muon pairs production with high precision and statistics. In the following, after a short description of the experimental apparatus, key results on ϕ and ω production and on the excess in the intermediate mass region will be discussed.

2. NA 60 Detector and data selection

NA 60, a fixed target experiment at the CERN SPS, is mainly dedicated to the detection of muon pairs. The apparatus and data selection procedure are fully described in [3]. The key detectors are the muon spectrometer (MS) and the vertex tracker (VT). The MS detects the muon tracks in the rapidity range $3 < y < 4$, and is preceded by a 12 cm thick hadron absorber. The vertex tracker is a Si pixel detector placed between the target and the absorber in a 2.5 T dipole field, that tracks the muons, among the other tracks, before they suffer multiple scattering and energy loss. The muons detected in the MS are matched to the VT tracks both in momentum and angle, thus improving the mass resolution and making it possible to measure the distance between the primary vertex and the track impact point.

The sample used for the results presented in this paper was collected with a 158 A GeV In beam impinging on a segmented In target. 2.3 fb⁻¹ triggers were acquired. In order to avoid events with reinteractions of nuclear fragments in the subsequent targets, only events with one vertex in the target region were selected. The combinatorial background was evaluated by means of the event mixing technique, with an accuracy of 1% over the full dimuon mass spectrum. The contribution due to the incorrect match between the VT tracks and the muons detected in the MS was estimated either with an overlay Monte Carlo simulation or with a mixing technique. The former consists in generating a dimuon on top of a real event, while the latter mixes the MS muons of a given event with the VT tracks of another event. The two methods agree within 5%. The final sample consists of 440000 signal muon pairs, with an average signal to background ratio of about 1/7.

3. ϕ production in In-In collisions

NA 60 can detect the ϕ meson via its decay into both muon and kaon pairs, thus helping understanding the so-called ϕ puzzle. This takes its origin from the discrepancies

observed in the measurements performed by NA 50 [4] in the muon channel and by NA 49 [5] in the kaon channel, both in Pb-Pb collisions at 158 A GeV. The multiplicity measured by NA 50 is much higher than the one obtained by NA 49. Also the inverse slopes T_{eff} of the transverse mass distributions differ: NA 50 finds a value of about 230 MeV, almost independent on centrality, while NA 49 measures values increasing with the number of participants N_{part} from 250 MeV to 300 MeV. It has to be stressed that the acceptance of NA 50 is limited to $p_T > 1.1$ GeV/c, while NA 49 is dominated by low transverse momenta: in presence of radial flow, the measured T_{eff} will depend on the p_T range, being lower at high p_T .

The origin of this discrepancy has been longely discussed. It was suggested that kaons may suffer rescattering and absorption in the medium, resulting in a depletion of kaon pairs, especially at low p_T , that leads to a reduced yield and a hardening of the p_T distribution in the hadronic channel [6]. Recent results by CERES [7] both in the hadronic and leptonic channels confirm the NA 49 results, and thus suggests that there is no puzzle. However, the results in dielectrons, being affected by a large error bar, are not conclusive.

NA 60 measures the yield in the muon channel by counting the events in the mass window $0.98 < M < 1.06$ GeV/c² [8]. The signal/background ratio below the peak is 1/3, integrated over centrality. To account for the continuum under the peak, two side windows between $0.88 < M < 0.92$ GeV/c² and $1.12 < M < 1.16$ GeV/c² are subtracted to the window. By performing the measurement in several p_T , rapidity and decay angle intervals, raw differential distributions are extracted. They are then corrected for the acceptance using an overlay Monte Carlo tuned iteratively such that the resulting distributions reproduce the measured data. The systematic error is evaluated by varying the analysis parameters and selections. The measurement is performed for 5 centrality bins, corresponding to $\langle N_{\text{part}} \rangle = 15; 39; 75; 132; 183$.

The rapidity distribution, integrated over centrality, is fitted with a gaussian, giving a width $\Delta y = 1.13 \pm 0.06 \pm 0.05$. No centrality dependence is observed. The result is in agreement with the NA 49 results on several colliding systems [5],[9]. The decay angle distribution is obtained for several reference systems (Helicity, Collins-Soper, Gottfried-Jackson) resulting flat in all the cases, independent of centrality.

The transverse mass distributions, shown in fig. 1 (left) for 5 centrality bins, are fitted with the function $1 = m_T dN/dm_T = N_0 \exp(-m_T/T_{\text{eff}})$. The resulting inverse slopes are plotted as a function of N_{part} in fig. 1 (right), showing an increase from peripheral to mid-central collisions and a saturation for most central collisions. Since in presence of radial flow the T_{eff} value may depend on the p_T range, in order to compare our results with NA 49 and NA 50, the fit is also performed for $p_T < 1.6$ GeV/c and $p_T > 1.1$ GeV/c. The results are shown in fig. 2 for the two p_T ranges. At low p_T (fig. 2, left), the NA 60 In-In data show an increase with N_{part} , in agreement with the NA 49 results in the same range. At high p_T (fig. 2, right) a flattening of the inverse slope dependence on centrality is observed, analogously to the NA 50 results. However, while the difference between the low- p_T NA 49 results and the high- p_T NA 50 results




Figure 1. Left panel: transverse momentum distributions in indium-indium collisions as a function of centrality. From top to bottom: central to peripheral spectra. Right Panel: T_{eff} slope parameter as a function of the number of participants.

amounts to 70 MeV for central collisions, the same difference in the NA 60 In-In data is of 15 MeV. While the latter can presumably be due to radial flow, the magnitude of the effect in Pb-Pb can hardly be attributed entirely to the same cause. A further attenuating due to kaon rescattering and absorption may lead to larger T_{eff} values in the NA 49 data.




Figure 2. Centrality dependence of the T_{eff} slope parameter (full circles) compared to NA 49 (open circles) and to NA 50 (open squares). Left panel: t performed in the NA 49 range ($0 < p_T < 1.6 \text{ GeV}/c$). Right panel: t performed in the NA 50 range ($1.1 < p_T < 3 \text{ GeV}/c$).

The multiplicity was obtained with two different approaches, with largely independent systematic errors. In the first, the cross section is directly measured, counting the number of produced and the number of incident ions. The average multiplicity is then obtained dividing the cross section by the total In-In inelastic




Figure 3. $\langle \eta \rangle / N_{part}$ in fullphase space, compared to the NA49 measurement in several collision systems.




Figure 4. R_{CP} for the compared to the one for the and the expectation for primary production only.

cross section, estimated with the general formula for two nuclei of mass numbers A and B: $\sigma_{inel;AB} = r_0^2 A^{1/3} + B^{1/3} A^{1/3} + B^{1/3} A^{1/3}$ [11], with $r_0 = 1.3$ and $r_0 = 1.93$ which leads to $\sigma_{inel;In-In} = 4.32$ b. Alternatively, the multiplicity can be obtained using the J/ψ as a calibration process. The ratio between the J/ψ and the J/ψ multiplicities is given by: $\langle \eta \rangle = \langle J/\psi \rangle \frac{N_{rec; \eta}}{N_{rec; J/\psi}}$, where $N_{(J/\psi)}$, $A_{(J/\psi)}$ and $rec_{(J/\psi)}$ are the number of observed (J/ψ) resonances, the acceptance and the reconstruction efficiency respectively. Once corrected for anomalous and nuclear absorption, $\langle J/\psi \rangle$ scales with the number of binary collisions N_{coll} and can be written as $\langle J/\psi \rangle = \left(\frac{J/\psi}{NN} = \frac{NN}{NN} \right) N_{coll}$, where $\frac{J/\psi}{NN}$ and $\frac{NN}{NN}$ are the J/ψ and inelastic cross sections in nucleon-nucleon collisions, and N_{coll} is calculated from N_{part} using the Glauber model. By multiplying this value by the above ratio, one obtains $\langle \eta \rangle$. The two methods agree within 10%.

In Fig.3 the enhancement factor, expressed as the ratio $\langle \eta \rangle / N_{part}$ is plotted as a function of N_{part} , compared to the NA49 results in several collision systems [12], [9]. The value in central In-In collisions exceeds the corresponding NA49 one in Pb-Pb. A direct comparison with NA50 requires an extrapolation of the NA50 measurements to low p_T , and is difficult due to the discrepancies in the observed inverse slopes. Two extreme cases with $T_{eff} = 220$ MeV and $T_{eff} = 300$ MeV were considered. The corresponding results largely exceed the NA60 values even if the higher T_{eff} value is considered.

To conclude, the puzzle is not yet solved. The forthcoming NA60 measurement in the $\psi(2S) \rightarrow K^+ K^-$ channel may be decisive to clarify the picture.

4. Enhancement

The meson, carrying a strangeness content of 40%, is expected to be strangeness-enhanced in heavy ion collisions. NA60 detects the through its Dalitz decay $\psi(2S) \rightarrow \eta \pi^+ \pi^-$.

In Fig. 4 the R_{CP} of the Λ , defined as $R_{CP}(N_{part}) = \frac{h_{i=N_{part}}}{h_{i=N_{part}}^{peripheral}}$, is plotted as a function of N_{part} , compared to the Λ . An enhancement of the Λ by a factor 1.34 from most peripheral to most central collisions is observed. The values for the Λ and Λ mesons can be linked through the strangeness under-saturation parameter s . According to [13], the value of s is 0.7 in Si-Si and 0.9 in Pb-Pb central collisions. A value of 0.8 in central In-In collisions is assumed. Since Λ is produced only primarily, the dependence on centrality of s can be then extracted from the R_{CP} , according to the scaling: $h_{i=N_{part1}} = h_{i=N_{part2}} = \frac{2}{s}(N_{part1}) = \frac{2}{s}(N_{part2})$. The R_{CP} can then be calculated under the hypothesis that Λ is only primarily produced: in this case $\Lambda = N_{part} / (0.4 \frac{2}{s}(N_{part}) + 0.6)$. The result of the calculation is plotted in Fig. 4, assuming values for s that range from 0.7 to 0.9 in central collisions. It should be noted that the results agree with the expectations if only primary production is assumed, in contrast to Statistical Models where a large fraction of the Λ is produced from secondaries without strangeness [14]. For comparison, if the ratio between primary and secondary production (the latter coming mainly from the decays of $\Lambda(1400)$ and a_0 which are not s suppressed) would be 1:1, the expected enhancement from peripheral to central collisions would be 1:1.

5. Results in the intermediate mass region

The dimuon intermediate mass region (IMR), located between the Λ and the J/ψ peaks, is expected to be well suited for the search of thermal dimuons, due to its relative production yield with respect to the other expected contributions in this region. In p-A collisions, the IMR region is described as a superposition of the Drell-Yan and open charm sources [15]. The NA 38, NA 50 and HELIOS-3 experiments observed an excess in the IMR with respect to the extrapolation of these sources in S-U and Pb-Pb collisions [15], [16]. The mass spectrum of the excess has the same shape as the open charm. Possible explanations of this excess are the production of thermal dimuons or the enhancement of the charm cross section. In the former case, the excess would have a prompt origin, while in the latter the muons coming from charmed meson decays would be o -vertex. Due to limited vertex capabilities, it was not possible to distinguish between the two scenarios.

Thanks to its ability to measure the muon o set, NA 60 can distinguish, on a statistical basis, between prompt and o -vertex dimuons. The resolution of the o set distance between the muon tracks and the vertex, estimated using muon pairs from J/ψ decay, is 37 μ m in the x coordinate (bending plane) and 45 μ m in y. Since the o set resolution of the matched tracks is affected by multiple scattering and energy loss, a weighted o set is used in the analysis: $\chi^2 = x^2 V_{xx}^{-1} + y^2 V_{yy}^{-1} + 2xy V_{xy}^{-1}$ where V^{-1} is the inverse error matrix, accounting for the uncertainties of the vertex fit and of the muon kinematics, x and y are the differences between the vertex coordinates and those of the extrapolated muon track at $z = z_{vert}$. The dimuon o set is defined as $\chi^2 = (\chi^2_1 + \chi^2_2) = 2$.

NA60 observes an excess in the mass region $1.16 < M < 2.56 \text{ GeV}/c^2$ with respect to the expected sources, that increases from peripheral to central collision faster than N_{part} and Drell-Yan.

The dimuon onset distribution, shown in Fig. 5 for $1.16 < M < 2.56 \text{ GeV}/c^2$, is compared to the corresponding distributions for prompt dimuons and open charm, scaled with respect to extrapolations from p-A data. It can be noticed that, in order to describe the data, the prompt dimuon source must be scaled by a factor 2.43 with respect to the expected Drell-Yan, while no enhancement is observed for the open charm, within errors. Therefore, the excess can be ascribed to a prompt source.




Figure 5. Dimuon weighted onset compared to the superposition of Drell-Yan and open charm contributions.




Figure 6. Inverse slope parameter versus dimuon mass. For details, see [17].

In order to study the properties of the excess, the expected Drell-Yan and open charm contributions were subtracted to the data. The resulting excess mass distribution is not reproduced by the Drell-Yan process, being much flatter. The p_T spectrum of the excess is significantly softer than the Drell-Yan one. Moreover, the excess p_T spectra extracted from different mass windows in the IMR are clearly different, while for the Drell-Yan pairs mass and p_T spectra factorize, and are thus independent. More details on the analysis and results are reported in [3]. The properties of the observed prompt excess, being different from the Drell-Yan ones in many observables, suggest an interpretation as thermal radiation.

The inverse m_T slope extracted for three mass windows in the IMR, is plotted in Fig. 6 as a function of the mass, together with the corresponding values obtained in the low mass region [17]. The inverse slopes for the Drell-Yan and open charm contributions are shown for comparison. Below $1 \text{ GeV}/c^2$, the inverse slope parameters monotonically rise with mass from the dimuon threshold where $T \approx 180 \text{ MeV}$, up to the nominal pole of the ρ meson, where $T \approx 250 \text{ MeV}$. This is followed by a sudden decline to 190 MeV in the IMR. The initial rise is consistent with the expectation for radial flow

of a hadronic source (here ρ ! ρ !). The jump of ρ 50 MeV observed beyond $1 \text{ GeV}/c^2$, down to a low-ow situation, is extremely hard to reconcile with emission sources which continue to be of dominantly hadronic origin. Rather, the sudden loss of ow is most naturally explained as a transition to a qualitatively different source, implying dominantly early, i.e. partonic processes like $q\bar{q}$! , for which ow has not yet built up.

6. Summary

The NA 60 experiment measured dimuon production in In-In collisions at the CERN SPS. The yield observed in the leptonic channel is higher than the one observed by NA 49 in the hadronic channel in Pb-Pb collisions. The dependence of the inverse slope on p_T suggests that the difference between NA 49 and NA 50 is larger than expected from radial ow. The forthcoming NA 60 measurement in the ρ ! $K\bar{K}$ channel may be decisive to clarify the picture.

The meson shows an enhancement of a factor 1.34 from peripheral to central collisions. This enhancement, compared to the one, is consistent with the expectation in the hypothesis that the is only primarily produced.

In the intermediate mass region, an excess of prompt origin is observed. The Drell-Yan process can not reproduce the properties of the excess. The value of T_{eff} , significantly smaller than in the low mass region, suggests an emission from the partonic stage, when the transverse ow has not yet built up.

References

- [1] J. Rafelski and B. Müller, *Phys. Rev. Lett.* 48 (1982) 1066.
- [2] E. V. Shuryak, *Phys. Rep.* 61 (1980) 71.
- [3] R. A. Mankiet al. (NA 60 Coll.) *Eur. Phys. J. C* (2009) in press. arXiv:0810.3204.
- [4] B. Alessandro et. al. (NA 50 Coll.), *Phys. Lett. B* 555 (2003) 147.
- [5] S. V. Afanasev et. al. (NA 49 Coll.), *Phys. Lett. B* 491 (2000) 59.
- [6] E. V. Shuryak, *Nucl. Phys. A* 661 (1999) 119; Johnson S C, Jacak B V and Drees A, *Eur. Phys. J. C* 18 (2001) 645; Pal S, Ko C M and Lin Z, *Nucl. Phys. A* 707 (2002) 525
- [7] D. Adamova et. al. (CERES Coll.), *Phys. Rev. Lett.* 96 (2006) 152301.
- [8] M. Floris et. al. (NA 60 Coll.), *J. Phys. G* 35 (2008) 104054.
- [9] C. Alt et al. (NA 49 Coll.), *Phys. Rev. Lett.* 94 (2005) 052301;
- [10] C. Amsler et. al. (Particle Data Group), *Phys. Lett. B* 667 (2008) 1.
- [11] M. Kh. Anikina et. al., *Soviet Journal of Nuclear Physics, Yad. Fiz* 38 (1983) 88
- [12] V. Friese et al. (NA 49 Coll.), *Nucl. Phys. A* 698 (2002) 487; C. Alt et al. (NA 49 Coll.), *Phys. Rev. C* 73 (2006) 044910;
- [13] F. Becattini et al., *Phys. Rev. C* 73 (2006) 044905
- [14] F. Becattini, private communication.
- [15] M. C. Abreu et al. (NA 50 Coll.) *Eur. Phys. J. C* 14 (2000) 443
- [16] C. Lourenco et al., *Nucl. Phys. A* 566 (1994) 77; A. L. S. Angelis, *Eur. Phys. J. C* 13 (2000) 433.
- [17] R. A. Mankiet al. (NA 60 Coll.), *Phys. Rev. Lett.* 100 (2008) 022032