

Higgs boson decay into heavy quarks and heavy leptons: higher order corrections

Victor T. Kim^a

^aSt. Petersburg Nuclear Physics Institute of Russian Academy of Sciences 188300, Gatchina

Theoretical predictions for the decay width of Standard Model Higgs boson into bottom quarks and -leptons, in the case when $M_H = 2M_W$, are briefly reviewed. The effects of higher order perturbative QCD (up to $\frac{4}{s}$ -level) and QED corrections are considered. The uncertainties of the decay width of Higgs boson into $b\bar{b}$ and $H^+ + \bar{H}^-$ are discussed.

1. Introduction

Production cross-sections and decay widths of the Standard Electroweak Model Higgs boson are nowadays among the most extensively analyzed theoretical quantities (for a recent review, see, e.g., [1], [2]). Indeed, the main hope of scientific community is that this essential ingredient of the Standard Model may be discovered, if not at Fermilab Tevatron, then at the forthcoming LHC experiments at CERN. There is great interest in the "low-mass" region $114.5 \text{ GeV} < M_H < 2M_W$, because a "low-mass" Higgs boson is heavily favored by Standard Model analysis of the available precision data. The lower bound, 114.5 GeV , was obtained from the direct searches of Higgs boson at the LEP 2 $e^+ e^-$ -collider primarily through Higgs boson decay into a $b\bar{b}$ -pair.

It should be stressed, that the uncertainties in $(H^+ + b\bar{b})$, analytically calculated in QCD using the $\overline{\text{MS}}$ -scheme at the $\frac{4}{s}$ -level [3], dominate the theoretical uncertainty for the branching ratio of $H^+ + b\bar{b}$ decay, which is considered to be the most important process in searches for a "low mass" Higgs boson by CMS and ATLAS collaborations at the LHC.

Here we briefly discuss the uncertainties of the QCD predictions for $H^+ + b\bar{b} = (H^+ + b\bar{b})$, including those which come from the on-shell mass parameterizations of this quantity (previ-

ous discussions see in [4]-[9] and [10]) and from the resummations of the $\frac{2}{s}$ -terms, typical of the Mandelstam region (see [11]-[16]). We discuss also perturbative QED and QCD uncertainties for Higgs boson decay into heavy leptons, $H^+ + (H^- + \bar{H}^-)$.

2. QCD corrections for $H^+ + b\bar{b}$ in terms of pole and running b-quark mass

There are several approaches for Higgs boson decay $H^+ + b\bar{b}$ in perturbative QCD. One of them is based on pole (on-shell) mass consideration [4]-[4]:

$$H^+ + b\bar{b} = \frac{b_0}{m_b} \frac{1}{1 + \sum_{i=1}^X \frac{b_i a_s^i(M_H)}{m_b}} ; \quad (1)$$

where $b_0 = (3/2)^{1/2} = 8/G_F M_H m_b^2$, $a_s(M_H) = s(M_H)/m_b$ and M_H are the pole b-quark and Higgs boson masses, and b_i -coefficients are fifth-order polynomials of large logarithms $L_b = \ln(M_H^2/m_b^2)$. An another approach is based on $\overline{\text{MS}}$ -scheme framework:

$$H^+ + b\bar{b} = \frac{b_0 \overline{m}_b^2(M_H)}{m_b^2} \frac{1}{1 + \sum_{i=1}^X \frac{b_i a_s^i(M_H)}{\overline{m}_b}} ; \quad (2)$$

where $a_s(M_H) = s(M_H)/\overline{m}_b(M_H)$ and $\overline{m}_b(M_H)$ are the QCD running parameters, defined in the $\overline{\text{MS}}$ -scheme. The coefficients b_i can be expressed through the sum of the following contributions:

$$\frac{b_i}{m_b} = d_i^E + d_i^M : \quad (3)$$

delivered at the 3rd Joint International "Hadron Structure-2009" (HS'09) Workshop, Tatranska Strba, Slovakia, Aug. 30 (Sept. 3, 2009); e-mail: kim@pnpi.spb.ru.

Here the positive contributions d_i^E , calculated directly in the Euclidean region, and d_i^M are proportional to α_s^2 -factors, which are typical for the Minkowski-like region.

The corresponding expressions for \bar{m}_b^2 [17],[18] were derived at the α_s^4 -level in Ref. [3],[19]. Detailed analysis and results for Higgs decay width \bar{m}_{bb} at α_s^4 -level are presented in [9],[10]), where the β -function of QCD renormalization group (RG) and mass anomalous dimension function γ_m [20]-[25] were considered at the 5-loop level:

$$\frac{da_s}{d\ln^2} = (\alpha_s) = \alpha_0 \alpha_s^2 + \dots + \alpha_4 \alpha_s^6 + O(\alpha_s^7); \quad (4)$$

$$\frac{d\ln\bar{m}_b}{d\ln^2} = (\alpha_s) = \alpha_0 \alpha_s + \dots + \alpha_4 \alpha_s^5 + O(\alpha_s^6); \quad (5)$$

The 5-loop coefficients α_4 and α_5 are still unknown, but it can be estimated by Pade approximation procedure, developed in [26] (see discussion in Refs. [9],[10]).

It should be stressed, however, that the uncertainties of the estimated 5-loop contributions to the QCD β -function and mass anomalous dimension function γ_m are not so important in the definition of the running of the b-quark mass from the pole mass m_b to the pole mass of Higgs boson M_H . This effect of running is described by the solution of the following RG equation:

$$\begin{aligned} \bar{m}_b^2(M_H) &= \bar{m}_b^2(m_b) \exp \left(-2 \int_{\alpha_s(m_b)}^{\alpha_s(M_H)} \frac{m(x)}{x} dx \right) \quad (6) \\ &= \bar{m}_b^2(m_b) \left(\frac{\alpha_s(M_H)}{\alpha_s(m_b)} \right)^2 = \left(\frac{AD(\alpha_s(M_H))}{AD(\alpha_s(m_b))} \right)^2; \end{aligned}$$

where $AD(\alpha_s)$ is a polynomial of 4-th order in the QCD expansion parameter α_s [9],[10].

In the Higgs boson mass region of interest, Eq.(2) may be expressed in numerical form as

$$\begin{aligned} \frac{\bar{m}_{bb}}{m_b^2} &= \frac{\bar{m}_b^2(M_H)}{m_b^2} \quad (7) \\ &= 1 + 5.667\alpha_s(M_H) + 29.15\alpha_s(M_H)^2 \\ &\quad + 41.76\alpha_s(M_H)^3 - 825.7\alpha_s(M_H)^4 \end{aligned}$$

Substituting the value $\alpha_s(M_H) = 0.0366$ (which corresponds to $\alpha_s(M_H = 120 \text{ GeV}) = 0.115$) into Eq.(7), and decomposing the coefficients in the Minkowski series into Euclidean contributions and Minkowski-type α_s^2 -effects, one can get from Ref.[3] the following numbers

$$\begin{aligned} \frac{\bar{m}_{bb}}{m_b^2} &= \frac{\bar{m}_b^2(M_H)}{m_b^2} \quad 1 + 0.207 + 0.039 \quad (8) \\ &\quad + 0.0020 \quad 0.0015 \\ &= \frac{\bar{m}_b^2(M_H)}{m_b^2} \quad 1 + 0.207 + (0.056 \quad 0.017) \\ &\quad + (0.017 \quad 0.015) + (0.0063 \quad 0.0078); \end{aligned}$$

where the negative numbers in the round brackets come from the effects of analytical continuation. Having a look at Eq. (8) we may conclude that in the Euclidean region the perturbative series is well-behaved and the α_s^2 -contributions typical of the Minkowski region are also decreasing from order to order. However, due to the strong interplay between these two effects in the third and fourth terms, the latter ones are becoming numerically comparable. This feature spoils the convergence of the perturbation series in the Euclidean region. Therefore, to improve the accuracy of the perturbative prediction in the Minkowski region it seems natural to sum up these α_s^2 -terms using the ideas, developed in the 80s (see, e.g., [27]-[30]). These ideas now have a more solid theoretical background, see, e.g., Ref. [31].

Also, we stress that the truncated perturbative expansions of Eq.(7) have some additional uncertainties. These include M_H and t-quark mass dependent QCD [32], [33] and QED [34] contributions:

$$\bar{m}_{bb} = \frac{3}{8} \bar{m}_b^2 G_F M_H \bar{m}_b^2(M_H) \quad t + QED \quad (9)$$

where t and QED are defined following Refs. [33], [34] as

$$\begin{aligned} t &= \alpha_s^2 (3.111 \quad 0.667 L_t) \quad (10) \\ &+ \frac{\bar{m}_b^2}{M_H^2} \left(-10 + 4L_t + \frac{4}{3} \ln(\bar{m}_b^2/M_H^2) \right) \end{aligned}$$

$$\begin{aligned}
& + a_s^3 50.474 \quad 8.167 L_t \quad 1.278 L_t^2 \\
& + a_s^2 \frac{M_H^2}{m_t^2} 0.241 \quad 0.070 L_t \\
& + X_t \quad 1 \quad 4.913 a_s + a_s^2 (-72.117 \quad 20.945 L_t)
\end{aligned}$$

$L_t = \ln(M_H^2/m_t^2)$, $X_t = G_F m_t^2/(8^2 p_T^2)$, m_t is the t-quark pole mass, $\bar{m}_b = \bar{m}_b(M_H)$ and

$$QED = 0.472 \quad 3.336 \frac{\bar{m}_b^2}{M_H^2} a \quad 1.455 a^2 + 1.301 a a_s \quad (11)$$

Using $a = (M_H)/(m_t) = 0.0027$, $(M_H)^1 = 129$, $m_t = 175 \text{ GeV}$, $M_H = 120 \text{ GeV}$, $\bar{m}_b = 2.8 \text{ GeV}$, $G_F = 1.1667 \cdot 10^{-5} \text{ GeV}^{-2}$ we get

$$t = 4.84 \cdot 10^3 \quad 1.7 \cdot 10^5 \quad (12)$$

$$\begin{aligned}
& + 2.27 \cdot 10^3 + 1.85 \cdot 10^4 \\
& + 3.2 \cdot 10^3 \quad 5.75 \cdot 10^4 \quad 2.42 \cdot 10^4
\end{aligned}$$

$$QED = 1.1 \cdot 10^3 \quad 4.5 \cdot 10^6 \quad (13)$$

$$9 \cdot 10^6 \quad 1.2 \cdot 10^4 :$$

Comparing the numbers presented in Eq.(8) and Eq.(13)-Eq.(13), we conclude that $\frac{4}{s}$ -term s can be neglected at the current level of the experimental precision of "Higgs-hunting" at Fermilab and LHC. Indeed, one can see, that even for the light Higgs boson the numerical values of the order $\frac{4}{s}$ -contributions to Eq.(8) are comparable with the leading M_H - and m_t -dependent terms in Eqs. (12)-(13).

An another approach for $H \rightarrow b\bar{b}$, where the RG-controllable terms are summed up, may be written down as

$$\begin{aligned}
H \bar{b} \bar{b} &= \frac{b_0}{a_s(m_b)} \frac{a_s(M_H)}{a_s(m_b)}^{(24=23)} \quad (14) \\
& \frac{AD(a_s(M_H))^2}{AD(a_s(m_b))^2} \left[1 + \sum_{i=1}^X b_i a_s^i(M_H) \right] \\
& 1 - \frac{8}{3} a_s(m_b) \quad 18.556 a_s(m_b)^2 \\
& 175.76 a_s(m_b)^3 \quad 1892 a_s(m_b)^4 ;
\end{aligned}$$

where

$$\begin{aligned}
AD(a_s)^2 &= 1 + 2.351 a_s \quad (15) \\
& + 4.383 a_s^2 + 3.873 a_s^3 \quad 15.15 a_s^4 :
\end{aligned}$$

Here, an important relation between pole and running masses of Refs. [35],[36],[10] has been used. Detailed comparison of $H \bar{b} \bar{b}$ in RG-improved (Eq. (14)) and in pole mass truncated (Eq. (2)) approaches was presented in Refs. [9], [10].

The behavior of the RG-resummed expressions for $H \bar{b} \bar{b}$ and $R_{H \bar{b} \bar{b}}$ are more stable than in the case, when RG-summation of the mass-dependent terms is not used [4]-[9], [10] (Figs. 1,2). Difference of $H \bar{b} \bar{b}$ calculated the truncated pole-mass approach and the RG-improved parametrization of $H \bar{b} \bar{b}$ is becoming smaller in each successive order of perturbation theory.

Indeed, for the phenomenologically interesting value of Higgs boson mass $M_H = 120 \text{ GeV}$ we find that at the $\frac{2}{s}$ -level $H \bar{b} \bar{b} = 0.7 \text{ MeV}$, while for the $\frac{3}{s}$ -level it becomes smaller, namely $H \bar{b} \bar{b} = 0.3 \text{ MeV}$. At the $\frac{3}{s}$ -level of the RG-improved




Figure 1. Higgs boson width in the pole (on-shell) mass approach.




Figure 2. Higgs boson width in the approach with explicit RG-resummation.

$\overline{\text{MS}}$ -scheme series one has $\overline{\text{H}}_{\text{bb}} = 1.85 \text{ MeV}$ for $M_H = 120 \text{ GeV}$. For this scale the value of $\overline{\text{H}}_{\text{bb}}$ with the explicit dependence from the pole-mass is 16 % higher, than its RG-improved estimate.

There are different approaches to the treatment of the typical Minkowskiian γ^2 -contributions in the perturbative expressions for physical quantities, which demonstrated remarkable convergence properties [13]-[16]. At the moment, these approaches are developing for different phenomenological applications, which will allow a comparison with the existing methods.

3. Higgs boson decay into $^{+ -}$

Width of Higgs boson decay into $^{+ -}$ leptons in the $\overline{\text{MS}}$ -scheme can be read as [37]:

$$\begin{aligned} \overline{\text{H}} &= \overline{0} \frac{\overline{m}^2(M_H)}{m^2} \left(1 + a(M_H) \right)_1 \\ &+ a(M_H)^2 \left(2 + a(M_H) \right)_2^3 \\ &+ a(M_H)^2 a_s(M_H) \left(\text{QED} \times \text{QCD} \right)_3 ; \end{aligned} \quad (16)$$

where $\overline{0} = (\overline{2}=8) G_F M_H m^2$, $a(M_H)$ $\overline{\text{MS}}(M_H) = \overline{m}(M_H)$ are QED running parameters and $a_s(M_H)$ $\overline{\text{MS}}(M_H) = \overline{\text{QED}} \times \overline{\text{QCD}}$ is QCD parameter, and $\overline{\text{QED}} \times \overline{\text{QCD}}$ is a mixed QED-QCD correction to the coefficient function. Evolution of running γ -lepton mass in QED is similar to Eq. (6), but with $\overline{\text{QED}}$, $\overline{\text{QED}}$, $\overline{\text{2}}$ and $\overline{\text{QED}} \times \overline{\text{QCD}}$, complicated by quark fractional electric charge dependence [37]. $\overline{\text{3}}$ is known since [38], and $\overline{\text{3}}$ [23] is consistent with QED limit of Ref. [21]. At present for H to get accuracy of $\overline{\text{H}}_{\text{bb}}$ at s^3 -level it is enough to keep 2-loop running γ -lepton mass and 1-loop coefficient function $\overline{1}$ [37].

4. Summary

Different approaches based on the running and pole b -quark masses for the decay width of the $H \rightarrow b\bar{b}$ process become consistent in higher orders of perturbative QCD. However, different convergence in different approaches demonstrates an existence of additional theoretical QCD uncertainties, which are not usually considered in phenomenological studies.

Currently, for width of Higgs boson decay into heavy leptons H to have accuracy of $\overline{\text{H}}_{\text{bb}}$ at s^3 -level it is enough to take into account 2-loop running γ -lepton mass and 1-loop coefficient function $\overline{1}$.

The author thanks A. L. Kataev for fruitful collaboration on the results presented here. The author is also grateful to the Local Organizing Committee of the "Hadron Structure - 2009" (HS09) Workshop for enjoyable atmosphere and CERN Theory Unit for their warm hospitality. The work is supported in parts by Russian Foundation for Basic Research, grants 08-02-01184 and 06-02-72041-MNTIa, and the RF President grant NSh-378.2008.2.

REFERENCES

1. A. D. Djouadi, Phys. Rept. 457 (2008) 1.
2. K. A. Assamagan et al. [Higgs Working Group Coll.], The Higgs working group: Summary report 2003 [hep-ph/0406152].

3. P. A. Baikov, K. G. Chetyrkin and J. H. Kuhn, Phys. Rev. Lett. 96 (2006) 012003.
4. A. L. Kataev and V. T. Kim, Report ENSLAPP-A-407-92 [hep-ph/9304282].
5. A. L. Kataev and V. T. Kim, Proc. A IHEP-1993, Obereramergau, Germany, 4-8 October 1993, World Scientific 1994, pp.623-637.
6. A. L. Kataev and V. T. Kim, Mod. Phys. Lett. A 9 (1994) 1309.
7. L.R. Surguladze, Phys. Lett. B 341 (1994) 60.
8. A. L. Kataev and V. T. Kim, Proc. 13th Lomonosov Conf., Moscow, 23-29 August 2007, World Scientific 2008, pp.334-337 [arXiv:0712.2514 [hep-ph]].
9. A. L. Kataev and V. T. Kim, Proc. Int. Seminar on Contemp. Probl. Elem. Part. Phys., dedicated to I. L. Solovtsov, Dubna, January 17-18, 2008, JINR, 2008, pp.167-182 [arXiv:0804.3992 [hep-ph]].
10. A. L. Kataev and V. T. Kim, Proc. XI-Ith Adv. Comp. Anal. Techniq. Phys. Res., Erice, November 3-7, 2008, PoS(ACAT08) 004 [arXiv:0902.1442 [hep-ph]].
11. S.G. Gorishny, A. L. Kataev and S.A. Larin, Sov. J. Nucl. Phys. 40 (1984) 329.
12. D. J. Broadhurst, A. L. Kataev and C. J. Maxwell, Nucl. Phys. B 592 (2001) 247.
13. A. P. Bakulev, S. V. Mikhailov and N. G. Stefanis, Phys. Rev. D 75 (2007) 056005 [Erratum -ibid. D 77 (2008) 079901].
14. N. G. Stefanis and A. I. Karanikas, Proc. Int. Seminar on Contemp. Probl. Elem. Part. Phys., dedicated to I.L. Solovtsov, Dubna, January 17-18, 2008, JINR, 2008 [arXiv:0802.3176 [hep-ph]].
15. A. P. Bakulev, Phys. Part. Nucl. 40 (2009) [arXiv:0805.0829 [hep-ph]].
16. A. P. Bakulev, Proc. 15th Int. Seminar Quarks-2008, Sergiev Posad, 23-29 May, 2008, INR, Moscow [arXiv:0810.3315 [hep-ph]].
17. S.G. Gorishny, A. L. Kataev, S.A. Larin and L.R. Surguladze, Mod. Phys. Lett. A 5 (1990) 2703; Phys. Rev. D 43 (1991) 1633.
18. K. G. Chetyrkin, Phys. Lett. B 390 (1997) 309.
19. K. G. Chetyrkin, B. A. Kniehl and A. Sirlin, Phys. Lett. B 402 (1997) 359.
20. O. V. Tarasov, A. A. Vladimirov and A.Y. Zharkov, Phys. Lett. B 93 (1980) 429.
21. T. van Ritbergen, J. A. M. Vermaasen and S.A. Larin, Phys. Lett. B 400 (1997) 379.
22. S.A. Larin and J. A. M. Vermaasen, Phys. Lett. B 303 (1993) 334.
23. K. G. Chetyrkin, Phys. Lett. B 404 (1997) 161.
24. J. A. Vermaasen, S.A. Larin and T. van Ritbergen, Phys. Lett. B 405 (1997) 327.
25. M. Czakon, Nucl. Phys. B 710 (2005) 485.
26. J. R. Ellis, I. Jack, D. R. T. Jones, M. Karliner and M. A. Samuel, Phys. Rev. D 57 (1998) 2665.
27. F. J. Yndurain, Proc. 3rd Warsaw Symp. Elem. Part. Phys., May 1980, Nukleonika 26 (1981) 213.
28. M. R. Pennington and G. G. Ross, Phys. Lett. B 102 (1981) 167.
29. N. V. Krasnikov and A. A. Pivovarov, Phys. Lett. B 116 (1982) 168.
30. A. V. Radushkin, Report JINR-E2-159, 1982, JINR Rapid Commun. 78 (1996) 96.
31. D. V. Shirkov, Theor. Math. Phys. 127 (2001) 409.
32. S. A. Larin, T. van Ritbergen and J. A. M. Vermaasen, Phys. Lett. B 362 (1995) 134.
33. K. G. Chetyrkin and M. Steinhauser, Phys. Lett. B 408 (1997) 320.
34. A. L. Kataev, JETP Lett. 66 (1997) 327.
35. K. G. Chetyrkin and M. Steinhauser, Nucl. Phys. B 573 (2000) 617.
36. K. Melnikov and T. v. R. Ritbergen, Phys. Lett. B 482 (2000) 99.
37. A. L. Kataev and V. T. Kim, to appear in Proc. 14th Lomonosov Conf., Moscow, 19-25 August, 2009.
38. S.G. Gorishny, A. L. Kataev, S.A. Larin and L.R. Surguladze, Phys. Lett. B 256 (1991) 81.