

# Determination of $f_s/f_d$ for 7 TeV $pp$ collisions and a measurement of the branching fraction of the decay $B^0 \rightarrow D^- K^+$

R. Aaij et al., (The LHCb Collaboration)

The relative abundance of the three decay modes  $B^0 \rightarrow D^- K^+$ ,  $B^0 \rightarrow D^- \pi^+$  and  $B_s^0 \rightarrow D_s^- \pi^+$  produced in 7 TeV  $pp$  collisions at the LHC is determined from data corresponding to an integrated luminosity of  $35 \text{ pb}^{-1}$ . The branching fraction of  $B^0 \rightarrow D^- K^+$  is found to be  $\mathcal{B}(B^0 \rightarrow D^- K^+) = (2.01 \pm 0.18^{\text{stat}} \pm 0.14^{\text{systr}}) \times 10^{-4}$ . The ratio of fragmentation fractions  $f_s/f_d$  is determined through the relative abundance of  $B_s^0 \rightarrow D_s^- \pi^+$  to  $B^0 \rightarrow D^- K^+$  and  $B^0 \rightarrow D^- \pi^+$ , leading to  $f_s/f_d = 0.253 \pm 0.017 \pm 0.017 \pm 0.020$ , where the uncertainties are statistical, systematic, and theoretical respectively.

PACS numbers: 12.38.Qk, 13.60.Le, 13.87.Fh

Knowledge of the production rate of  $B_s^0$  mesons is required to determine any  $B_s^0$  branching fraction. This rate is determined by the  $b\bar{b}$  production cross-section and the fragmentation probability  $f_s$ , which is the fraction of  $B_s^0$  mesons amongst all weakly-decaying bottom hadrons. Similarly the production rate of  $B^0$  mesons is driven by the fragmentation probability  $f_d$ . The measurement of the branching fraction of the rare decay  $B_s^0 \rightarrow \mu^+ \mu^-$  is a prime example where improved knowledge of  $f_s/f_d$  is needed to reach the highest sensitivity in the search for physics beyond the Standard Model [1]. The ratio  $f_s/f_d$  is in principle dependent on collision energy and type as well as the acceptance region of the detector. This is the first measurement of this quantity at the LHC.

The ratio  $f_s/f_d$  can be extracted if the ratio of branching fractions of  $B^0$  and  $B_s^0$  mesons decaying to particular final states  $X_1$  and  $X_2$ , respectively, is known:

$$\frac{f_s}{f_d} = \frac{N_{X_2} \mathcal{B}(B^0 \rightarrow X_2) \epsilon(B^0 \rightarrow X_2)}{N_{X_1} \mathcal{B}(B_s^0 \rightarrow X_1) \epsilon(B_s^0 \rightarrow X_1)}. \quad (1)$$

The ratio of the branching fraction of the  $B_s^0 \rightarrow D_s^- \pi^+$  and  $B^0 \rightarrow D^- K^+$  decays is dominated by contributions from colour-allowed tree-diagram amplitudes and is therefore theoretically well understood. In contrast, the ratio of the branching ratios of the two decays  $B_s^0 \rightarrow D_s^- \pi^+$  and  $B^0 \rightarrow D^- \pi^+$  can be measured with a smaller statistical uncertainty due to the greater yield of the  $B^0$  mode, but suffers from an additional theoretical uncertainty due to the contribution from a  $W$ -exchange diagram. Both ratios are exploited here to measure  $f_s/f_d$  according to the equations [2, 3]

$$\frac{f_s}{f_d} = 0.971 \cdot \left| \frac{V_{us}}{V_{ud}} \right|^2 \left( \frac{f_K}{f_\pi} \right)^2 \frac{\tau_{B_d} \mathcal{N}_a \mathcal{N}_F \epsilon_{D^- \pi^+}}{\tau_{B_s} \mathcal{N}_a \mathcal{N}_F \epsilon_{D_s^- \pi^+}} \frac{N_{D_s^- \pi^+}}{N_{D^- K^+}}, \quad (2)$$

and

$$\frac{f_s}{f_d} = 0.982 \cdot \frac{\tau_{B_d} \mathcal{N}_a \mathcal{N}_F \mathcal{N}_E \epsilon_{D^- \pi^+}}{\tau_{B_s} \mathcal{N}_a \mathcal{N}_F \mathcal{N}_E \epsilon_{D_s^- \pi^+}} \frac{N_{D_s^- \pi^+}}{N_{D^- \pi^+}}. \quad (3)$$

Here  $\epsilon_X$  is the selection efficiency of decay  $X$  (including the branching fraction of the  $D$  decay mode used to re-

construct it),  $N_X$  is the observed number of decays of this type, the  $V_{ij}$  are elements of the CKM matrix,  $f_i$  are the meson decay constants and the numerical factors take into account the phase space difference for the ratio of the two decay modes. Inclusion of charge conjugate modes is implied throughout. The term  $\mathcal{N}_a$  parametrizes non-factorizable SU(3)-breaking effects;  $\mathcal{N}_F$  is the ratio of the form factors;  $\mathcal{N}_E$  is an additional correction term to account for the  $W$ -exchange diagram in the  $B^0 \rightarrow D^- \pi^+$  decay. Their values [2, 3] are  $\mathcal{N}_a = 1.00 \pm 0.02$ ,  $\mathcal{N}_F = 1.24 \pm 0.08$ , and  $\mathcal{N}_E = 0.966 \pm 0.075$ . The latest world average [4] is used for the  $B$  meson lifetime ratio  $\tau_{B_s}/\tau_{B_d} = 0.973 \pm 0.015$ . The numerical values used for the other factors are:  $|V_{us}| = 0.2252$ ,  $|V_{ud}| = 0.97425$ ,  $f_\pi = 130.41$  and  $f_K = 156.1$ , with negligible associated uncertainties [5].

The observed yields of these three decay modes in  $35 \text{ pb}^{-1}$  of data collected with the LHCb detector in the 2010 running period are used to measure  $f_s/f_d$  averaged over the LHCb acceptance and to improve the current measurement of the branching fraction of the  $B^0 \rightarrow D^- K^+$  decay mode [6].

The LHCb experiment [7] is a single-arm spectrometer, designed to study  $B$  decays at the LHC, with a pseudorapidity acceptance of  $2 < \eta < 5$  for charged tracks. The first trigger level allows the selection of events with  $B$  hadronic decays using the transverse energy of hadrons measured in the calorimeter system. The event information is subsequently sent to a software trigger, implemented in a dedicated processor farm, which performs a final online selection of events for later offline analysis. The tracking system determines the momenta of  $B$  decay products with a precision of  $\delta p/p = 0.35\text{--}0.5\%$ . Two Ring Imaging Cherenkov (RICH) detectors allow charged kaons and pions to be distinguished in the momentum range 2–100 GeV/ $c$  [8].

The three decay modes,  $B^0 \rightarrow D^-(K^+ \pi^- \pi^-) \pi^+$ ,  $B^0 \rightarrow D^-(K^+ \pi^- \pi^-) K^+$  and  $B_s^0 \rightarrow D_s^-(K^+ K^- \pi^-) \pi^+$ , are topologically identical and can therefore be selected using identical geometric and kinematic criteria, thus minimizing efficiency differences between them. Events

are selected at the first trigger stage by requiring a hadron with transverse energy greater than 3.6 GeV in the calorimeter. The second, software, stage [9, 10] requires a two, three, or four track secondary vertex with a high sum  $p_T$  of the tracks, significant displacement from the primary interaction, and at least one track with exceptionally high  $p_T$ , large displacement from the primary interaction, and small fit  $\chi^2$ .

The decays of  $B$  mesons can be distinguished from background using variables such as the  $p_T$  and impact parameter  $\chi^2$  of the  $B$ ,  $D$ , and the final state particles with respect to the primary interaction. In addition the vertex quality of the  $B$  and  $D$  candidates, the  $B$  lifetime, and the angle between the  $B$  momentum vector and the vector joining the  $B$  production and decay vertices are used in the selection. The  $D$  lifetime and flight distance are not used in the selection because the lifetimes of the  $D_s^-$  and  $D^-$  differ by about a factor of two.

The event sample is first selected using the gradient boosted decision tree technique [11], which combines the geometrical and kinematic variables listed above. The selection is trained on a mixture of simulated  $B^0 \rightarrow D^- \pi^+$  decays and combinatorial background selected from the sidebands of the data mass distributions. The distributions of the input variables for data and simulated signal events show excellent agreement, justifying the use of simulated events in the training procedure.

Subsequently,  $D^-$  ( $D_s^-$ ) candidates are identified by requiring the invariant mass under the  $K\pi\pi$  ( $KK\pi$ ) hypothesis to fall within the selection window  $1870_{-40}^{+24}$  ( $1969_{-40}^{+24}$ ) MeV/ $c^2$ , where the mass resolution is approximately 10 MeV/ $c^2$ . The final  $B^0 \rightarrow D^- \pi^+$  and  $B_s^0 \rightarrow D_s^- \pi^+$  subsamples consist of events that pass a particle identification (PID) criterion on the bachelor particle, based on the difference in log-likelihood between the charged pion and kaon hypotheses (DLL) of  $\text{DLL}(K-\pi) < 0$ , with an efficiency of 83.0%. The  $B^0 \rightarrow D^- K^+$  subsample consists of events with  $\text{DLL}(K-\pi) > 5$ , with an efficiency of 70.2%. Events not satisfying either condition are not used.

The relative efficiency of the selection procedure is evaluated for all decay modes using simulated events, where the appropriate resonances in the charm decays are taken into account. As the analysis is only sensitive to relative efficiencies, the impact of differences between data and simulation is small. The relative efficiencies are  $\epsilon_{D^- \pi^+} / \epsilon_{D^- K^+} = 1.221 \pm 0.021$ ,  $\epsilon_{D^- K^+} / \epsilon_{D_s^- \pi^+} = 0.917 \pm 0.020$ , and  $\epsilon_{D^- \pi^+} / \epsilon_{D_s^- \pi^+} = 1.120 \pm 0.025$ , where the errors are due to the limited size of the simulated event samples.

The relative yields of the three decay modes are extracted from unbinned extended maximum likelihood fits to the mass distributions shown in Fig. 2. The signal mass shape is described by an empirical model derived from simulated events. The mass distribution in the simulation exhibits non-Gaussian tails on either side of the

signal. The tail on the right-hand side is due to non-Gaussian detector effects and modeled with a Crystal Ball (CB) function [12]. A similar tail is present on the left-hand side of the peak. In addition, the low mass tail contains a second contribution due to events where hadrons have radiated photons that are not reconstructed. The sum of these contributions is modeled with a second CB function. The peak values of these two CB functions are constrained to be identical.

Various backgrounds have to be considered, in particular the crossfeed between the  $D^-$  and  $D_s^-$  channels, and the contamination in both samples from  $\Lambda_b \rightarrow \Lambda_c^+ \pi^-$  decays, where  $\Lambda_c^+ \rightarrow p K^- \pi^+$ . The  $D_s^-$  contamination in the  $D^-$  data sample is reduced by loose PID requirements,  $\text{DLL}(K-\pi) < 10$  (with an efficiency of 98.6%) and  $\text{DLL}(K-\pi) > 0$  (with an efficiency of 95.6%), for the pions and kaons from  $D$  decays, respectively. The resulting efficiency to reconstruct  $B_s^0 \rightarrow D_s^- \pi^+$  as background is evaluated, using simulated events, to be 30 times smaller than for  $B^0 \rightarrow D^- \pi^+$  and 150 times smaller than for  $B^0 \rightarrow D^- K^+$  within the  $B^0$  and  $D^-$  signal mass windows. Taking into account the lower production fraction of  $B_s^0$  mesons, this background is negligible.

The contamination from  $\Lambda_c$  decays is estimated in a similar way. However, different approaches are used for the  $B^0$  and  $B_s^0$  decays. A contamination of approximately 2% under the  $B^0 \rightarrow D^- \pi^+$  mass peak and below 1% under the  $B^0 \rightarrow D^- K^+$  peak is found, and therefore no explicit  $\text{DLL}(p-\pi)$  criterion is needed. The  $\Lambda_c$  background in the  $B_s^0$  sample is, on the other hand, large enough that it can be fitted for directly.

A prominent peaking background to  $B^0 \rightarrow D^- K^+$  is  $B^0 \rightarrow D^- \pi^+$ , with the pion misidentified as a kaon. The small  $\pi \rightarrow K$  misidentification rate (of about 4%) is compensated by the larger branching fraction, resulting in similar event yields. This background is modeled by obtaining a clean  $B^0 \rightarrow D^- \pi^+$  sample from the data and reconstructing it under the  $B^0 \rightarrow D^- K^+$  mass hypothesis. The resulting mass shape depends on the momentum distribution of the bachelor particle. The momentum distribution after the  $\text{DLL}(K-\pi) > 5$  requirement can be found by considering the PID performance as a function of momentum. This is obtained using a sample of  $D^{*+} \rightarrow D^0 \pi^+$  decays, and is illustrated in Fig. 1. The mass distribution is reweighted using this momentum distribution to reproduce the  $B^0 \rightarrow D^- \pi^+$  mass shape following the DLL cut.

The combinatorial background consists of events with random pions and kaons, forming a fake  $D^-$  or  $D_s^-$  candidate, as well as real,  $D^-$  or  $D_s^-$  mesons that combine with a random pion or kaon. The combinatorial background is modeled with an exponential shape.

Other background components originate from partially reconstructed  $B^0$  and  $B_s^0$  decays. In  $B^0 \rightarrow D^- \pi^+$  these originate from  $B^0 \rightarrow D^{*-} \pi^+$  and  $B^0 \rightarrow D^- \rho^+$  decays, which can also be backgrounds for  $B^0 \rightarrow$




FIG. 1: Probability, as a function of momentum, to correctly identify (full symbols) a kaon or a pion when requiring  $DLL(K - \pi) > 5$  or  $DLL(K - \pi) < 0$ , respectively. The correspondent probability to wrongly identify (open symbols) a pion as a kaon, or a kaon as a pion is also shown. The data are taken from a calibration sample of  $D^* \rightarrow D(K\pi)\pi$  decays; the statistical uncertainties are too small to display.

$D^-K^+$  in the case of a misidentified bachelor pion. In  $B^0 \rightarrow D^-K^+$  there is additionally background from  $B^0 \rightarrow D^{*-}K^+$  decays. The invariant mass distributions for the partially reconstructed and misidentified backgrounds are taken from large samples of simulated events, reweighted according to the mass hypothesis of the signal being fitted and the DLL cuts.

For  $B_s^0 \rightarrow D_s^- \pi^+$ , the  $B^0 \rightarrow D^- \pi^+$  background peaks under the signal with a similar shape. In order to suppress this peaking background, PID requirements are placed on both kaon tracks. The kaon which has the same sign in the  $B_s^0 \rightarrow D_s^- \pi^+$  and  $B^0 \rightarrow D^- \pi^+$  decays is required to satisfy  $DLL(K - \pi) > 0$ , while the other kaon in the  $D_s^+$  decay is required to satisfy  $DLL(K - \pi) > 5$ . Because of the similar shape, a Gaussian constraint is applied to the yield of this background. The central value of this constraint is computed from the  $\pi \rightarrow K$  misidentification rate. The  $\Lambda_b \rightarrow \Lambda_c^+ \pi^-$  background shape is obtained from simulated events, reweighted according to the PID efficiency, and the yield allowed to float in the fit. Finally, the relative size of the  $B_s^0 \rightarrow D_s^- \rho^+$  and  $B_s^0 \rightarrow D_s^{*-} \pi^+$  backgrounds is constrained to the ratio of the  $B^0 \rightarrow D^- \rho^+$  and  $B^0 \rightarrow D^{*-} \pi^+$  backgrounds in the  $B^0 \rightarrow D^- \pi^+$  fit, with an uncertainty of 20% to account for potential SU(3) symmetry breaking effects.

The free parameters in the likelihood fits to the mass distributions are the event yields for the different event types, *i.e.* the combinatorial background, partially reconstructed background, misidentified contributions, the signal, as well as the peak value of the signal shape. In addition the combinatoric background shape is left free in the  $B^0 \rightarrow D^- \pi^+$  and  $B_s^0 \rightarrow D_s^- \pi^+$  fits, and the signal width is left free in the  $B^0 \rightarrow D^- \pi^+$  fit. In the  $B_s^0 \rightarrow D_s^- \pi^+$  and  $B^0 \rightarrow D^- K^+$  fits the signal width is fixed to the value from the  $B^0 \rightarrow D^- \pi^+$  fit, corrected by

the ratio of the signal widths for these modes in simulated events.

The fits to the full  $B^0 \rightarrow D^- \pi^+$ ,  $B^0 \rightarrow D^- K^+$ , and  $B_s^0 \rightarrow D_s^- \pi^+$  data samples are shown in Fig. 2. The resulting  $B^0 \rightarrow D^- \pi^+$  and  $B^0 \rightarrow D^- K^+$  event yields are  $4103 \pm 75$  and  $252 \pm 21$ , respectively. The number of misidentified  $B^0 \rightarrow D^- \pi^+$  events under the  $B^0 \rightarrow D^- K^+$  signal as obtained from the fit is  $131 \pm 19$ . This agrees with the number expected from the total number of  $B^0 \rightarrow D^- \pi^+$  events, corrected for the misidentification rate determined from the PID calibration sample, of  $145 \pm 5$ . The  $B_s^0 \rightarrow D_s^- \pi^+$  event yield is  $670 \pm 34$ .




FIG. 2: Mass distributions of the  $B^0 \rightarrow D^- \pi^+$ ,  $B^0 \rightarrow D^- K^+$ , and  $B_s^0 \rightarrow D_s^- \pi^+$  candidates (top to bottom). The indicated components are described in the text.

The stability of the fit results has been investigated using different cut values for both the PID requirement on the bachelor particle and for the multivariate selection variable. In all cases variations are found to be small in comparison to the statistical uncertainty.

The relative branching fractions are obtained by correcting the event yields by the corresponding efficiency factors; the dominant correction comes from the PID efficiency. The dominant source of systematic uncertainty is the knowledge on the  $B^0 \rightarrow D^- \pi^+$  branching fraction (for the  $B^0 \rightarrow D^- K^+$  branching fraction measurement) and the knowledge of the  $D^-$  and  $D_s^-$  branching fractions (for the  $f_s/f_d$  measurement). An important source of systematic uncertainty is the knowledge of the PID efficiency as a function of momentum, which is needed to reweight the mass distribution of the  $B^0 \rightarrow D^- \pi^+$  decay under the kaon hypothesis for the bachelor track. This enters in two ways: firstly as an uncertainty on the correction factors, and secondly as part of the systematic uncertainty, since the shape for the misidentified backgrounds relies on correct knowledge of the PID efficiency as a function of momentum.

The performance of the PID calibration is evaluated by applying the same method from data to simulated events, and the maximum discrepancy found between the calibration method and the true mis-identification is attributed as a systematic uncertainty. The  $f_s/f_d$  measurement using  $B^0 \rightarrow D^- K^+$  and  $B_s^0 \rightarrow D_s^- \pi^+$  is more robust against PID uncertainties, since the final states have the same number of kaons and pions.

Other systematic uncertainties are due to limited simulated event samples (affecting the relative selection efficiencies), neglecting the  $\Lambda_b \rightarrow \Lambda_c^+ \pi^-$  and  $B_s^0 \rightarrow D_s^- \pi^+$  backgrounds in the  $B^0 \rightarrow D^- \pi^+$  fits, and the limited accuracy of the trigger simulation. Even though the ratio of efficiencies is statistically consistent with unity, the maximum deviation is conservatively assigned as a systematic uncertainty. The difference in interaction probability between kaons and pions is estimated using MC simulation. The systematic uncertainty due to possible discrepancies between data and simulation is expected to be negligible and it is not taken into account. The efficiency of the non-resonant  $D_s$  decays varies across the Dalitz plane, but has a negligible effect on the total  $B_s^0 \rightarrow D_s^- \pi^+$  efficiency. The sources of systematic uncertainty are summarized in Tab. I.

The efficiency corrected ratio of  $B^0 \rightarrow D^- \pi^+$  and  $B^0 \rightarrow D^- K^+$  yields is combined with the world average of the  $B^0 \rightarrow D^- \pi^+$  [5] branching ratio to give

$$\mathcal{B}(B^0 \rightarrow D^- K^+) = (2.01 \pm 0.18 \pm 0.14) \times 10^{-4}. \quad (4)$$

The first uncertainty is statistical and the second systematic.

The theoretically cleaner measurement of  $f_s/f_d$  uses  $B^0 \rightarrow D^- K^+$  and  $B_s^0 \rightarrow D_s^- \pi^+$  and is made according to Eq. 2. Accounting for the exclusive  $D$  branching fractions

TABLE I: Experimental systematic uncertainties for the  $\mathcal{B}(B^0 \rightarrow D^- K^+)$  and the two  $f_s/f_d$  measurements.

	$\mathcal{B}(B^0 \rightarrow D^- K^+)$	$f_s/f_d$
PID calibration	2.5%	1.0%/2.5%
Fit model	2.8%	2.8%
Trigger simulation	2.0%	2.0%
$\mathcal{B}(B^0 \rightarrow D^- \pi^+)$	4.9%	
$\mathcal{B}(D_s^+ \rightarrow K^+ K^- \pi^+)$		4.9%
$\mathcal{B}(D^+ \rightarrow K^- \pi^+ \pi^+)$		2.2%
$\tau_{B_s}/\tau_{B_d}$		1.5%

$\mathcal{B}(D^+ \rightarrow K^- \pi^+ \pi^+) = (9.14 \pm 0.20)\%$  [13] and  $\mathcal{B}(D_s^+ \rightarrow K^- K^+ \pi^+) = (5.50 \pm 0.27)\%$  [14], the value of  $f_s/f_d$  is found to be

$$f_s/f_d = (0.310 \pm 0.030^{\text{stat}} \pm 0.021^{\text{syst}}) \times \frac{1}{\mathcal{N}_a \mathcal{N}_F}, \quad (5)$$

where the first uncertainty is statistical and the second is systematic. The statistical uncertainty is dominated by the yield of the  $B^0 \rightarrow D^- K^+$  mode.

The statistically more precise but theoretically less clean measurement of  $f_s/f_d$  uses  $B^0 \rightarrow D^- \pi^+$  and  $B_s^0 \rightarrow D_s^- \pi^+$  and is, from Eq. 3,

$$f_s/f_d = (0.307 \pm 0.017^{\text{stat}} \pm 0.023^{\text{syst}}) \times \frac{1}{\mathcal{N}_a \mathcal{N}_F \mathcal{N}_E}. \quad (6)$$

The two values for  $f_s/f_d$  can be combined into a single value, taking all correlated uncertainties into account and using the theoretical inputs accounting for the  $SU(3)$  breaking part of the form factor ratio, the non-factorizable and W-exchange diagram:

$$f_s/f_d = 0.253 \pm 0.017^{\text{stat}} \pm 0.017^{\text{syst}} \pm 0.020^{\text{theor}}. \quad (7)$$

In summary, with  $35 \text{ pb}^{-1}$  of data collected using the LHCb detector during the 2010 LHC operation at a centre-of-mass energy of 7 TeV, the branching fraction of the Cabibbo-suppressed  $B^0$  decay mode  $B^0 \rightarrow D^- K^+$  has been measured with better precision than the current world average. Additionally, two measurements of the  $f_s/f_d$  production fraction are performed from the relative yields of  $B_s^0 \rightarrow D_s^- \pi^+$  with respect to  $B^0 \rightarrow D^- K^+$  and  $B^0 \rightarrow D^- \pi^+$ . These values of  $f_s/f_d$  are numerically close to the values determined at LEP and at the Tevatron [4].

#### Acknowledgments

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at CERN and at the LHCb institutes, and acknowledge support from the National Agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil);

CERN; NSFC (China); CNRS/IN2P3 (France); BMBF, DFG, HGF and MPG (Germany); SFI (Ireland); INFN (Italy); FOM and NWO (Netherlands); SCSR (Poland); ANCS (Romania); MinES of Russia and Rosatom (Russia); MICINN, XUNGAL and GENCAT (Spain); SNSF and SER (Switzerland); NAS Ukraine (Ukraine); STFC (United Kingdom); NSF (USA). We also acknowledge the support received from the ERC under FP7 and the Région Auvergne.

- 
- [1] Aaij, R. *et al.* (LHCb Collaboration). *Phys. Lett. B*, 699:330, 2011. arXiv:1103.2465[hep-ex].
  - [2] R. Fleischer, N. Serra, and N. Tuning. *Phys. Rev.*, D83:014017, 2011. arXiv:1012.2784[hep-ph].
  - [3] R. Fleischer, N. Serra, and N. Tuning. *Phys. Rev.*, D82:034038, 2010. arXiv:1004.3982[hep-ph].
  - [4] D. Asner *et al.* arXiv:1010.1589[hep-ex], 2010.
  - [5] K. Nakamura *et al.* *J. Phys. G*, 37(075021), 2010.
  - [6] Abe, K. *et al.* (BELLE Collaboration). *Phys. Rev. Lett.*, 87:111801, 2001. arXiv:0104051[hep-ex].
  - [7] A. A. Alves Jr. *et al.* (LHCb Collaboration). *JINST*, 3:S08005, 2008.
  - [8] A. A. Alves Jr. *et al.* The LHCb Detector at the LHC. *JINST*, 3:S08005, 2008.
  - [9] V. V. Gligorov. LHCb-PUB-2011-003.
  - [10] M. Williams *et al.* LHCb-PUB-2011-002.
  - [11] A. Hoecker *et al.* *PoS*, ACAT:040, 2007.
  - [12] T. Skwarnicki. DESY F31-86-02, 1986.
  - [13] Dobbs, S. *et al.* (CLEO Collaboration). *Phys. Rev.*, D76:112001, 2007. arXiv:0709.3783[hep-ex].
  - [14] Alexander, J. P. *et al.* (CLEO Collaboration). *Phys. Rev. Lett.*, 100:161804, 2008. arXiv:0801.0680[hep-ex].

## The LHCb Collaboration

R. Aaij<sup>23</sup>, B. Adeva<sup>36</sup>, M. Adinolfi<sup>42</sup>, C. Adrover<sup>6</sup>, A. Affolder<sup>48</sup>, Z. Ajaltouni<sup>5</sup>, J. Albrecht<sup>37</sup>, F. Alessio<sup>6,37</sup>, M. Alexander<sup>47</sup>, G. Alkhazov<sup>29</sup>, P. Alvarez Cartelle<sup>36</sup>, A.A. Alves Jr<sup>22</sup>, S. Amato<sup>2</sup>, Y. Amhis<sup>38</sup>, J. Amoraal<sup>23</sup>, J. Anderson<sup>39</sup>, R.B. Appleby<sup>50</sup>, O. Aquines Gutierrez<sup>10</sup>, L. Arrabito<sup>53</sup>, A. Artamonov<sup>34</sup>, M. Artuso<sup>52,37</sup>, E. Aslanides<sup>6</sup>, G. Auremma<sup>22,m</sup>, S. Bachmann<sup>11</sup>, J.J. Back<sup>44</sup>, D.S. Bailey<sup>50</sup>, V. Balagura<sup>30,37</sup>, W. Baldini<sup>16</sup>, R.J. Barlow<sup>50</sup>, C. Barschel<sup>37</sup>, S. Barsuk<sup>7</sup>, W. Barter<sup>43</sup>, A. Bates<sup>47</sup>, C. Bauer<sup>10</sup>, Th. Bauer<sup>23</sup>, A. Bay<sup>38</sup>, I. Bediaga<sup>1</sup>, K. Belous<sup>34</sup>, I. Belyaev<sup>30,37</sup>, E. Ben-Haim<sup>8</sup>, M. Benayoun<sup>8</sup>, G. Bencivenni<sup>18</sup>, S. Benson<sup>46</sup>, J. Benton<sup>42</sup>, R. Bernet<sup>39</sup>, M.-O. Bettler<sup>17,37</sup>, M. van Beuzekom<sup>23</sup>, A. Bien<sup>11</sup>, S. Bifani<sup>12</sup>, A. Bizzeti<sup>17,h</sup>, P.M. Bjørnstad<sup>50</sup>, T. Blake<sup>49</sup>, F. Blanc<sup>38</sup>, C. Blanks<sup>49</sup>, J. Blouw<sup>11</sup>, S. Blusk<sup>52</sup>, A. Bobrov<sup>33</sup>, V. Bocci<sup>22</sup>, A. Bondar<sup>33</sup>, N. Bondar<sup>29</sup>, W. Bonivento<sup>15</sup>, S. Borghi<sup>47</sup>, A. Borgia<sup>52</sup>, T.J.V. Bowcock<sup>48</sup>, C. Bozzi<sup>16</sup>, T. Brambach<sup>9</sup>, J. van den Brand<sup>24</sup>, J. Bressieux<sup>38</sup>, S. Brisbane<sup>51</sup>, M. Britsch<sup>10</sup>, T. Britton<sup>52</sup>, N.H. Brook<sup>42</sup>, A. Büchler-Germann<sup>39</sup>, A. Bursche<sup>39</sup>, J. Buytaert<sup>37</sup>, S. Cadeddu<sup>15</sup>, J.M. Caicedo Carvajal<sup>37</sup>, O. Callot<sup>7</sup>, M. Calvi<sup>20,j</sup>, M. Calvo Gomez<sup>35,n</sup>, A. Camboni<sup>35</sup>, P. Campana<sup>18,37</sup>, A. Carbone<sup>14</sup>, G. Carboni<sup>21,k</sup>, R. Cardinale<sup>19,i</sup>, A. Cardini<sup>15</sup>, L. Carson<sup>36</sup>, K. Carvalho Akiba<sup>23</sup>, G. Casse<sup>48</sup>, M. Cattaneo<sup>37</sup>, M. Charles<sup>51</sup>, Ph. Charpentier<sup>37</sup>, N. Chiapolini<sup>39</sup>, X. Cid Vidal<sup>36</sup>, P.E.L. Clarke<sup>46</sup>, M. Clemencic<sup>37</sup>, H.V. Cliff<sup>43</sup>, J. Closier<sup>37</sup>, C. Coca<sup>28</sup>, V. Coco<sup>23</sup>, J. Cogan<sup>6</sup>, P. Collins<sup>37</sup>, F. Constantin<sup>28</sup>, G. Conti<sup>38</sup>, A. Contu<sup>51</sup>, A. Cook<sup>42</sup>, M. Coombes<sup>42</sup>, G. Corti<sup>37</sup>, G.A. Cowan<sup>38</sup>, R. Currie<sup>46</sup>, B. D'Almeida<sup>7</sup>, C. D'Ambrosio<sup>37</sup>, P. David<sup>8</sup>, P.N.Y. David<sup>23</sup>, I. De Bonis<sup>4</sup>, S. De Capua<sup>21,k</sup>, M. De Cian<sup>39</sup>, F. De Lorenzi<sup>12</sup>, J.M. De Miranda<sup>1</sup>, L. De Paula<sup>2</sup>, P. De Simone<sup>18</sup>, D. Decamp<sup>4</sup>, M. Deckenhoff<sup>9</sup>, H. Degaudenzi<sup>38,37</sup>, M. Deissenroth<sup>11</sup>, L. Del Buono<sup>8</sup>, C. Deplano<sup>15</sup>, O. Deschamps<sup>5</sup>, F. Dettori<sup>15,d</sup>, J. Dickens<sup>43</sup>, H. Dijkstra<sup>37</sup>, P. Diniz Batista<sup>1</sup>, D. Dossett<sup>44</sup>, A. Dovbnya<sup>40</sup>, F. Dupertuis<sup>38</sup>, R. Dzhelyadin<sup>34</sup>, C. Eames<sup>49</sup>, S. Easo<sup>45</sup>, U. Egede<sup>49</sup>, V. Egorychev<sup>30</sup>, S. Eidelman<sup>33</sup>, D. van Eijk<sup>23</sup>, F. Eisele<sup>11</sup>, S. Eisenhardt<sup>46</sup>, R. Ekelhof<sup>9</sup>, L. Eklund<sup>47</sup>, Ch. Elsasser<sup>39</sup>, D.G. d'Enterria<sup>35,o</sup>, D. Esperante Pereira<sup>36</sup>, L. Esteve<sup>43</sup>, A. Falabella<sup>16,e</sup>, E. Fanchini<sup>20,j</sup>, C. Färber<sup>11</sup>, G. Fardell<sup>46</sup>, C. Farinelli<sup>23</sup>, S. Farry<sup>12</sup>, V. Fave<sup>38</sup>, V. Fernandez Albor<sup>36</sup>, M. Ferro-Luzzi<sup>37</sup>, S. Filippov<sup>32</sup>, C. Fitzpatrick<sup>46</sup>, M. Fontana<sup>10</sup>, F. Fontanelli<sup>19,i</sup>, R. Forty<sup>37</sup>, M. Frank<sup>37</sup>, C. Frei<sup>37</sup>, M. Frosini<sup>17,f,37</sup>, S. Furcas<sup>20</sup>, A. Gallas Torreira<sup>36</sup>, D. Galli<sup>14,c</sup>, M. Gandelman<sup>2</sup>, P. Gandini<sup>51</sup>, Y. Gao<sup>3</sup>, J.-C. Garnier<sup>37</sup>, J. Garofoli<sup>52</sup>, L. Garrido<sup>35</sup>, C. Gaspar<sup>37</sup>, N. Gauvin<sup>38</sup>, M. Gersabeck<sup>37</sup>, T. Gershon<sup>44</sup>, Ph. Ghez<sup>4</sup>, V. Gibson<sup>43</sup>, V.V. Gligorov<sup>37</sup>, C. Göbel<sup>54</sup>, D. Golubkov<sup>30</sup>, A. Golutvin<sup>49,30,37</sup>, A. Gomes<sup>2</sup>, H. Gordon<sup>51</sup>, M. Grabalosa Gándara<sup>35</sup>, R. Graciani Diaz<sup>35</sup>, L.A. Granado Cardoso<sup>37</sup>, E. Graugés<sup>35</sup>, G. Graziani<sup>17</sup>, A. Grecu<sup>28</sup>, S. Gregson<sup>43</sup>, B. Gui<sup>52</sup>, E. Gushchin<sup>32</sup>, Yu. Guz<sup>34</sup>, T. Gys<sup>37</sup>, G. Haefeli<sup>38</sup>, S.C. Haines<sup>43</sup>, T. Hampson<sup>42</sup>, S. Hansmann-Menzemer<sup>11</sup>, R. Harji<sup>49</sup>, N. Harnew<sup>51</sup>, J. Harrison<sup>50</sup>, P.F. Harrison<sup>44</sup>, J. He<sup>7</sup>, V. Heijne<sup>23</sup>, K. Hennessy<sup>48</sup>, P. Henrard<sup>5</sup>, J.A. Hernando Morata<sup>36</sup>, E. van Herwijnen<sup>37</sup>, W. Hofmann<sup>10</sup>, K. Holubyev<sup>11</sup>, P. Hopchev<sup>4</sup>, W. Hulsbergen<sup>23</sup>, P. Hunt<sup>51</sup>, T. Huse<sup>48</sup>, R.S. Huston<sup>12</sup>, D. Hutchcroft<sup>48</sup>, D. Hynds<sup>47</sup>, V. Iakovenko<sup>41</sup>, P. Ilten<sup>12</sup>, J. Imong<sup>42</sup>, R. Jacobsson<sup>37</sup>, M. Jahjah Hussein<sup>5</sup>, E. Jans<sup>23</sup>, F. Jansen<sup>23</sup>, P. Jaton<sup>38</sup>, B. Jean-Marie<sup>7</sup>, F. Jing<sup>3</sup>, M. John<sup>51</sup>, D. Johnson<sup>51</sup>, C.R. Jones<sup>43</sup>, B. Jost<sup>37</sup>, S. Kandybei<sup>40</sup>, T.M. Karbach<sup>9</sup>, J. Keaveney<sup>12</sup>, U. Kerzel<sup>37</sup>, T. Ketel<sup>24</sup>, A. Keune<sup>38</sup>, B. Khanji<sup>6</sup>, Y.M. Kim<sup>46</sup>, M. Knecht<sup>38</sup>, S. Koblitz<sup>37</sup>, P. Koppenburg<sup>23</sup>, A. Kozlinskiy<sup>23</sup>, L. Kravchuk<sup>32</sup>, K. Kreplin<sup>11</sup>, G. Krocker<sup>11</sup>, P. Krokovny<sup>11</sup>, F. Kruse<sup>9</sup>, K. Kruzelecki<sup>37</sup>, M. Kucharczyk<sup>20,25</sup>, S. Kukulak<sup>25</sup>, R. Kumar<sup>14,37</sup>, T. Kvaratskheliya<sup>30,37</sup>, V.N. La Thi<sup>38</sup>, D. Lacarrere<sup>37</sup>, G. Lafferty<sup>50</sup>, A. Lai<sup>15</sup>, D. Lambert<sup>46</sup>, R.W. Lambert<sup>37</sup>, E. Lanciotti<sup>37</sup>, G. Lanfranchi<sup>18</sup>, C. Langenbruch<sup>11</sup>, T. Latham<sup>44</sup>, R. Le Gac<sup>6</sup>, J. van Leerdam<sup>23</sup>, J.-P. Lees<sup>4</sup>, R. Lefèvre<sup>5</sup>, A. Leflat<sup>31,37</sup>, J. Lefrançois<sup>7</sup>, O. Leroy<sup>6</sup>, T. Lesiak<sup>25</sup>, L. Li<sup>3</sup>, Y.Y. Li<sup>43</sup>, L. Li Gioi<sup>5</sup>, M. Lieng<sup>9</sup>, R. Lindner<sup>37</sup>, C. Linn<sup>11</sup>, B. Liu<sup>3</sup>, G. Liu<sup>37</sup>, J.H. Lopes<sup>2</sup>, E. Lopez Asamar<sup>35</sup>, N. Lopez-March<sup>38</sup>, J. Luisier<sup>38</sup>, F. Machefert<sup>7</sup>, I.V. Machikhiliyan<sup>4,30</sup>, F. Maciuc<sup>10</sup>, O. Maev<sup>29,37</sup>, J. Magnin<sup>1</sup>, A. Maier<sup>37</sup>, S. Malde<sup>51</sup>, R.M.D. Mamunur<sup>37</sup>, G. Manca<sup>15,d</sup>, G. Mancinelli<sup>6</sup>, N. Mangiafave<sup>43</sup>, U. Marconi<sup>14</sup>, R. Märki<sup>38</sup>, J. Marks<sup>11</sup>, G. Martellotti<sup>22</sup>, A. Martens<sup>7</sup>, L. Martin<sup>51</sup>, A. Martín Sánchez<sup>7</sup>, D. Martinez Santos<sup>37</sup>, A. Massafferri<sup>1</sup>, Z. Mathe<sup>12</sup>, C. Matteuzzi<sup>20</sup>, M. Matveev<sup>29</sup>, E. Maurice<sup>6</sup>, B. Maynard<sup>52</sup>, A. Mazurov<sup>32,16,37</sup>, G. McGregor<sup>50</sup>, R. McNulty<sup>12</sup>, C. Mclean<sup>14</sup>, M. Meissner<sup>11</sup>, M. Merk<sup>23</sup>, J. Merkel<sup>9</sup>, R. Messi<sup>21,k</sup>, S. Miglioranzi<sup>37</sup>, D.A. Milanes<sup>13,37</sup>, M.-N. Minard<sup>4</sup>, S. Monteil<sup>5</sup>, D. Moran<sup>12</sup>, P. Morawski<sup>25</sup>, J.V. Morris<sup>45</sup>, R. Mountain<sup>52</sup>, I. Mous<sup>23</sup>, F. Muheim<sup>46</sup>, K. Müller<sup>39</sup>, R. Muresan<sup>28,38</sup>, B. Muryn<sup>26</sup>, M. Musy<sup>35</sup>, P. Naik<sup>42</sup>, T. Nakada<sup>38</sup>, R. Nandakumar<sup>45</sup>, J. Nardulli<sup>45</sup>, M. Nedos<sup>9</sup>, M. Needham<sup>46</sup>, N. Neufeld<sup>37</sup>, C. Nguyen-Mau<sup>38,p</sup>, P. M. Nicol<sup>7</sup>, S. Nies<sup>9</sup>, V. Niess<sup>5</sup>, N. Nikitin<sup>31</sup>, A. Oblakowska-Mucha<sup>26</sup>, V. Obraztsov<sup>34</sup>, S. Oggero<sup>23</sup>, S. Ogilvy<sup>47</sup>, O. Okhrimenko<sup>41</sup>, R. Oldeman<sup>15,d</sup>, M. Orlandea<sup>28</sup>, J.M. Otalora Goicochea<sup>2</sup>, B. Pal<sup>52</sup>, J. Palacios<sup>39</sup>, M. Palutan<sup>18</sup>, J. Panman<sup>37</sup>, A. Papanestis<sup>45</sup>, M. Pappagallo<sup>13,b</sup>, C. Parkes<sup>47,37</sup>, C.J. Parkinson<sup>49</sup>, G. Passaleva<sup>17</sup>, G.D. Patel<sup>48</sup>, M. Patel<sup>49</sup>, S.K. Paterson<sup>49</sup>, G.N. Patrick<sup>45</sup>, C. Patrignani<sup>19,i</sup>, C. Pavel-Nicorescu<sup>28</sup>, A. Pazos Alvarez<sup>36</sup>, A. Pellegrino<sup>23</sup>, G. Penso<sup>22,l</sup>, M. Pepe Altarelli<sup>37</sup>, S. Perazzini<sup>14,c</sup>, D.L. Perego<sup>20,j</sup>, E. Perez Trigo<sup>36</sup>, A. Pérez-Calero Yzquierdo<sup>35</sup>, P. Perret<sup>5</sup>, M. Perrin-Terrin<sup>6</sup>, G. Pessina<sup>20</sup>, A. Petrella<sup>16,37</sup>, A. Petrolini<sup>19,i</sup>, B. Pie Valls<sup>35</sup>, B. Pietrzyk<sup>4</sup>, T. Pilar<sup>44</sup>, D. Pinci<sup>22</sup>, R. Plackett<sup>47</sup>, S. Playfer<sup>46</sup>, M. Plo Casasus<sup>36</sup>, G. Polok<sup>25</sup>, A. Poluektov<sup>44,33</sup>, E. Polcarpo<sup>2</sup>, D. Popov<sup>10</sup>, B. Popovici<sup>28</sup>, C. Potterat<sup>35</sup>, A. Powell<sup>51</sup>, T. du Pree<sup>23</sup>, V. Pugatch<sup>41</sup>, A. Puig Navarro<sup>35</sup>, W. Qian<sup>52</sup>, J.H. Rademacker<sup>42</sup>, B. Rakotomiramanana<sup>38</sup>, I. Raniuk<sup>40</sup>, G. Raven<sup>24</sup>, S. Redford<sup>15</sup>, M.M. Reid<sup>44</sup>, A.C. dos Reis<sup>1</sup>, S. Ricciardi<sup>45</sup>, K. Rinnert<sup>48</sup>, D.A. Roa Romero<sup>5</sup>, P. Robbe<sup>7</sup>, E. Rodrigues<sup>47</sup>, F. Rodrigues<sup>2</sup>, C. Rodriguez Cobo<sup>36</sup>, P. Rodriguez Perez<sup>36</sup>, G.J. Rogers<sup>43</sup>, V. Romanovsky<sup>34</sup>, J. Rouvinet<sup>38</sup>, T. Ruf<sup>37</sup>, H. Ruiz<sup>35</sup>, G. Sabatino<sup>21,k</sup>, J.J. Saborido Silva<sup>36</sup>, N. Sagidova<sup>29</sup>, P. Sail<sup>47</sup>, B. Saitta<sup>15,d</sup>, C. Salzmann<sup>39</sup>, M. Sannino<sup>19,i</sup>, R. Santacesaria<sup>22</sup>, R. Santinelli<sup>37</sup>, E. Santovetti<sup>21,k</sup>, M. Sapunov<sup>6</sup>, A. Sarti<sup>18,l</sup>, C. Satriano<sup>22,m</sup>, A. Satta<sup>21</sup>, M. Savrie<sup>16,e</sup>, D. Savrina<sup>30</sup>, P. Schaack<sup>49</sup>, M. Schiller<sup>11</sup>, S. Schleich<sup>9</sup>, M. Schmelling<sup>10</sup>, B. Schmidt<sup>37</sup>, O. Schneider<sup>38</sup>, A. Schopper<sup>37</sup>, M.-H. Schune<sup>7</sup>, R. Schwemmer<sup>37</sup>, A. Sciubba<sup>18,l</sup>, M. Seco<sup>36</sup>, A. Semennikov<sup>30</sup>, K. Senderowska<sup>26</sup>, N. Serra<sup>39</sup>, J. Serrano<sup>6</sup>, P. Seyfert<sup>11</sup>, B. Shao<sup>3</sup>, M. Shapkin<sup>34</sup>, I. Shapoval<sup>40,37</sup>, P. Shatalov<sup>30</sup>, Y. Shcheglov<sup>29</sup>, T. Shears<sup>48</sup>, L. Shekhtman<sup>33</sup>, O. Shevchenko<sup>40</sup>, V. Shevchenko<sup>30</sup>, A. Shires<sup>49</sup>, R. Silva Coutinho<sup>54</sup>, H.P. Skottowe<sup>43</sup>, T. Skwarnicki<sup>52</sup>, A.C. Smith<sup>37</sup>, N.A. Smith<sup>48</sup>, K. Sobczak<sup>5</sup>, F.J.P. Soler<sup>47</sup>, A. Solomin<sup>42</sup>, F. Soomro<sup>49</sup>, B. Souza De Paula<sup>2</sup>, B. Spaan<sup>9</sup>, A. Sparkes<sup>46</sup>, P. Spradlin<sup>47</sup>, F. Stagni<sup>37</sup>, S. Stahl<sup>11</sup>, O. Steinkamp<sup>39</sup>, S. Stoica<sup>28</sup>, S. Stone<sup>52,37</sup>, B. Storaci<sup>23</sup>, U. Straumann<sup>39</sup>, N. Styles<sup>46</sup>, S. Swientek<sup>9</sup>, M. Szczekowski<sup>27</sup>, P. Szczypka<sup>38</sup>, T. Szumlak<sup>26</sup>, S. T'Jampens<sup>4</sup>, E. Teodorescu<sup>28</sup>, F. Teubert<sup>37</sup>, C. Thomas<sup>51,45</sup>,

E. Thomas<sup>37</sup>, J. van Tilburg<sup>11</sup>, V. Tisserand<sup>4</sup>, M. Tobin<sup>39</sup>, S. Topp-Joergensen<sup>51</sup>, M.T. Tran<sup>38</sup>, A. Tsaregorodtsev<sup>6</sup>, N. Tuning<sup>23</sup>, A. Ukleja<sup>27</sup>, P. Urquijo<sup>52</sup>, U. Uwer<sup>11</sup>, V. Vagnoni<sup>14</sup>, G. Valenti<sup>14</sup>, R. Vazquez Gomez<sup>35</sup>, P. Vazquez Regueiro<sup>36</sup>, S. Vecchi<sup>16</sup>, J.J. Velthuis<sup>42</sup>, M. Veltri<sup>17,g</sup>, K. Vervink<sup>37</sup>, B. Viaud<sup>7</sup>, I. Videau<sup>7</sup>, X. Vilasis-Cardona<sup>35,n</sup>, J. Visniakov<sup>36</sup>, A. Vollhardt<sup>39</sup>, D. Voong<sup>42</sup>, A. Vorobyev<sup>29</sup>, H. Voss<sup>10</sup>, K. Wacker<sup>9</sup>, S. Wandernoth<sup>11</sup>, J. Wang<sup>52</sup>, D.R. Ward<sup>43</sup>, A.D. Webber<sup>50</sup>, D. Websdale<sup>49</sup>, M. Whitehead<sup>44</sup>, D. Wiedner<sup>11</sup>, L. Wiggers<sup>23</sup>, G. Wilkinson<sup>51</sup>, M.P. Williams<sup>44,45</sup>, M. Williams<sup>49</sup>, F.F. Wilson<sup>45</sup>, J. Wishahi<sup>9</sup>, M. Witek<sup>25</sup>, W. Witzeling<sup>37</sup>, S.A. Wotton<sup>43</sup>, K. Wyllie<sup>37</sup>, Y. Xie<sup>46</sup>, F. Xing<sup>51</sup>, Z. Yang<sup>3</sup>, R. Young<sup>46</sup>, O. Yushchenko<sup>34</sup>, M. Zavertyaev<sup>10,a</sup>, L. Zhang<sup>52</sup>, W.C. Zhang<sup>12</sup>, Y. Zhang<sup>3</sup>, A. Zhelezov<sup>11</sup>, L. Zhong<sup>3</sup>, E. Zverev<sup>31</sup>, A. Zvyagin<sup>37</sup>.

<sup>1</sup>Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

<sup>2</sup>Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

<sup>3</sup>Center for High Energy Physics, Tsinghua University, Beijing, China

<sup>4</sup>LAPP, Université de Savoie, CNRS/IN2P3, Annecy-Le-Vieux, France

<sup>5</sup>Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, Clermont-Ferrand, France

<sup>6</sup>CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France

<sup>7</sup>LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France

<sup>8</sup>LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris, France

<sup>9</sup>Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany

<sup>10</sup>Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany

<sup>11</sup>Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

<sup>12</sup>School of Physics, University College Dublin, Dublin, Ireland

<sup>13</sup>Sezione INFN di Bari, Bari, Italy

<sup>14</sup>Sezione INFN di Bologna, Bologna, Italy

<sup>15</sup>Sezione INFN di Cagliari, Cagliari, Italy

<sup>16</sup>Sezione INFN di Ferrara, Ferrara, Italy

<sup>17</sup>Sezione INFN di Firenze, Firenze, Italy

<sup>18</sup>Laboratori Nazionali dell'INFN di Frascati, Frascati, Italy

<sup>19</sup>Sezione INFN di Genova, Genova, Italy

<sup>20</sup>Sezione INFN di Milano Bicocca, Milano, Italy

<sup>21</sup>Sezione INFN di Roma Tor Vergata, Roma, Italy

<sup>22</sup>Sezione INFN di Roma La Sapienza, Roma, Italy

<sup>23</sup>Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands

<sup>24</sup>Nikhef National Institute for Subatomic Physics and Vrije Universiteit, Amsterdam, Netherlands

<sup>25</sup>Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Cracow, Poland

<sup>26</sup>Faculty of Physics & Applied Computer Science, Cracow, Poland

<sup>27</sup>Soltan Institute for Nuclear Studies, Warsaw, Poland

<sup>28</sup>Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania

<sup>29</sup>Petersburg Nuclear Physics Institute (PNPI), Gatchina, Russia

<sup>30</sup>Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia

<sup>31</sup>Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia

<sup>32</sup>Institute for Nuclear Research of the Russian Academy of Sciences (INR RAN), Moscow, Russia

<sup>33</sup>Budker Institute of Nuclear Physics (SB RAS) and Novosibirsk State University, Novosibirsk, Russia

<sup>34</sup>Institute for High Energy Physics (IHEP), Protvino, Russia

<sup>35</sup>Universitat de Barcelona, Barcelona, Spain

<sup>36</sup>Universidad de Santiago de Compostela, Santiago de Compostela, Spain

<sup>37</sup>European Organization for Nuclear Research (CERN), Geneva, Switzerland

<sup>38</sup>Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland

<sup>39</sup>Physik-Institut, Universität Zürich, Zürich, Switzerland

<sup>40</sup>NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine

<sup>41</sup>Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine

<sup>42</sup>H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom

<sup>43</sup>Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom

<sup>44</sup>Department of Physics, University of Warwick, Coventry, United Kingdom

<sup>45</sup>STFC Rutherford Appleton Laboratory, Didcot, United Kingdom

<sup>46</sup>School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom

<sup>47</sup>School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom

<sup>48</sup>Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom

<sup>49</sup>Imperial College London, London, United Kingdom

<sup>50</sup>School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom

<sup>51</sup>Department of Physics, University of Oxford, Oxford, United Kingdom

<sup>52</sup>Syracuse University, Syracuse, NY, United States

<sup>53</sup>CC-IN2P3, CNRS/IN2P3, Lyon-Villeurbanne, France, associated member

<sup>54</sup>Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to <sup>2</sup>

<sup>a</sup>P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia

- <sup>b</sup> *Università di Bari, Bari, Italy*  
<sup>c</sup> *Università di Bologna, Bologna, Italy*  
<sup>d</sup> *Università di Cagliari, Cagliari, Italy*  
<sup>e</sup> *Università di Ferrara, Ferrara, Italy*  
<sup>f</sup> *Università di Firenze, Firenze, Italy*  
<sup>g</sup> *Università di Urbino, Urbino, Italy*  
<sup>h</sup> *Università di Modena e Reggio Emilia, Modena, Italy*  
<sup>i</sup> *Università di Genova, Genova, Italy*  
<sup>j</sup> *Università di Milano Bicocca, Milano, Italy*  
<sup>k</sup> *Università di Roma Tor Vergata, Roma, Italy*  
<sup>l</sup> *Università di Roma La Sapienza, Roma, Italy*  
<sup>m</sup> *Università della Basilicata, Potenza, Italy*  
<sup>n</sup> *LIFAELS, La Salle, Universitat Ramon Llull, Barcelona, Spain*  
<sup>o</sup> *Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain*  
<sup>p</sup> *Hanoi University of Science, Hanoi, Viet Nam*  
**(The LHCb Collaboration)**