



# Measurement of $CP$ observables in $B^0 \rightarrow DK^{*0}$ with $D \rightarrow K^+K^-$

The LHCb collaboration<sup>†</sup>

## Abstract

The decay  $B^0 \rightarrow DK^{*0}$  and the charge conjugate mode are studied using  $1.0 \text{ fb}^{-1}$  of  $pp$  collision data collected by the LHCb experiment at  $\sqrt{s} = 7 \text{ TeV}$  in 2011. The  $CP$  asymmetry between the  $B^0 \rightarrow DK^{*0}$  and the  $\bar{B}^0 \rightarrow D\bar{K}^{*0}$  decay rates, with the neutral  $D$  meson in the  $CP$ -even final state  $K^+K^-$ , is found to be

$$\mathcal{A}_d^{KK} = -0.45 \pm 0.23 \pm 0.02,$$

where the first uncertainty is statistical and the second is systematic. In addition, favoured  $B^0 \rightarrow DK^{*0}$  decays are reconstructed with the  $D$  meson in the non- $CP$  eigenstate  $K^+\pi^-$ . The ratio of the  $B$ -flavour averaged decay rates in  $D$  decays to  $CP$  and non- $CP$  eigenstates is measured to be

$$\mathcal{R}_d^{KK} = 1.36_{-0.32}^{+0.37} \pm 0.07,$$

where the ratio of the branching fractions of  $D^0 \rightarrow K^-\pi^+$  to  $D^0 \rightarrow K^+K^-$  decays is included as multiplicative factor. The  $CP$  asymmetries measured with two control channels, the favoured  $B^0 \rightarrow DK^{*0}$  decay with  $D \rightarrow K^+\pi^-$  and the  $\bar{B}_s^0 \rightarrow DK^{*0}$  decay with  $D \rightarrow K^+K^-$ , are also reported.

Submitted to JHEP

© CERN on behalf of the LHCb collaboration, license CC-BY-3.0.

<sup>†</sup>Authors are listed on the following pages.



## LHCb collaboration

R. Aaij<sup>38</sup>, C. Abellan Beteta<sup>33,n</sup>, A. Adametz<sup>11</sup>, B. Adeva<sup>34</sup>, M. Adinolfi<sup>43</sup>, C. Adrover<sup>6</sup>, A. Affolder<sup>49</sup>, Z. Ajaltouni<sup>5</sup>, J. Albrecht<sup>35</sup>, F. Alessio<sup>35</sup>, M. Alexander<sup>48</sup>, S. Ali<sup>38</sup>, G. Alkhazov<sup>27</sup>, P. Alvarez Cartelle<sup>34</sup>, A.A. Alves Jr<sup>22,35</sup>, S. Amato<sup>2</sup>, Y. Amhis<sup>7</sup>, L. Anderlini<sup>17,f</sup>, J. Anderson<sup>37</sup>, R. Andreassen<sup>57</sup>, R.B. Appleby<sup>51</sup>, O. Aquines Gutierrez<sup>10</sup>, F. Archilli<sup>18</sup>, A. Artamonov<sup>32</sup>, M. Artuso<sup>53</sup>, E. Aslanides<sup>6</sup>, G. Auriemma<sup>22,m</sup>, S. Bachmann<sup>11</sup>, J.J. Back<sup>45</sup>, C. Baesso<sup>54</sup>, V. Balagura<sup>28</sup>, W. Baldini<sup>16</sup>, R.J. Barlow<sup>51</sup>, C. Barschel<sup>35</sup>, S. Barsuk<sup>7</sup>, W. Barter<sup>44</sup>, A. Bates<sup>48</sup>, Th. Bauer<sup>38</sup>, A. Bay<sup>36</sup>, J. Beddow<sup>48</sup>, I. Bediaga<sup>1</sup>, S. Belogurov<sup>28</sup>, K. Belous<sup>32</sup>, I. Belyaev<sup>28</sup>, E. Ben-Haim<sup>8</sup>, M. Benayoun<sup>8</sup>, G. Bencivenni<sup>18</sup>, S. Benson<sup>47</sup>, J. Benton<sup>43</sup>, A. Berezhtoy<sup>29</sup>, R. Bernet<sup>37</sup>, M.-O. Bettler<sup>44</sup>, M. van Beuzekom<sup>38</sup>, A. Bien<sup>11</sup>, S. Bifani<sup>12</sup>, T. Bird<sup>51</sup>, A. Bizzeti<sup>17,h</sup>, P.M. Bjørnstad<sup>51</sup>, T. Blake<sup>35</sup>, F. Blanc<sup>36</sup>, C. Blanks<sup>50</sup>, J. Blouw<sup>11</sup>, S. Blusk<sup>53</sup>, A. Bobrov<sup>31</sup>, V. Bocci<sup>22</sup>, A. Bondar<sup>31</sup>, N. Bondar<sup>27</sup>, W. Bonivento<sup>15</sup>, S. Borghi<sup>51</sup>, A. Borgia<sup>53</sup>, T.J.V. Bowcock<sup>49</sup>, E. Bowen<sup>37</sup>, C. Bozzi<sup>16</sup>, T. Brambach<sup>9</sup>, J. van den Brand<sup>39</sup>, J. Bressieux<sup>36</sup>, D. Brett<sup>51</sup>, M. Britsch<sup>10</sup>, T. Britton<sup>53</sup>, N.H. Brook<sup>43</sup>, H. Brown<sup>49</sup>, A. Büchler-Germann<sup>37</sup>, I. Burducea<sup>26</sup>, A. Bursche<sup>37</sup>, J. Buytaert<sup>35</sup>, S. Cadeddu<sup>15</sup>, O. Callot<sup>7</sup>, M. Calvi<sup>20,j</sup>, M. Calvo Gomez<sup>33,n</sup>, A. Camboni<sup>33</sup>, P. Campana<sup>18,35</sup>, A. Carbone<sup>14,c</sup>, G. Carboni<sup>21,k</sup>, R. Cardinale<sup>19,i</sup>, A. Cardini<sup>15</sup>, H. Carranza-Mejia<sup>47</sup>, L. Carson<sup>50</sup>, K. Carvalho Akiba<sup>2</sup>, G. Casse<sup>49</sup>, M. Cattaneo<sup>35</sup>, Ch. Cauet<sup>9</sup>, M. Charles<sup>52</sup>, Ph. Charpentier<sup>35</sup>, P. Chen<sup>3,36</sup>, N. Chiapolini<sup>37</sup>, M. Chrzaszcz<sup>23</sup>, K. Ciba<sup>35</sup>, X. Cid Vidal<sup>34</sup>, G. Ciezarek<sup>50</sup>, P.E.L. Clarke<sup>47</sup>, M. Clemencic<sup>35</sup>, H.V. Cliff<sup>44</sup>, J. Closier<sup>35</sup>, C. Coca<sup>26</sup>, V. Coco<sup>38</sup>, J. Cogan<sup>6</sup>, E. Cogneras<sup>5</sup>, P. Collins<sup>35</sup>, A. Comerma-Montells<sup>33</sup>, A. Contu<sup>15</sup>, A. Cook<sup>43</sup>, M. Coombes<sup>43</sup>, G. Corti<sup>35</sup>, B. Couturier<sup>35</sup>, G.A. Cowan<sup>36</sup>, D. Craik<sup>45</sup>, S. Cunliffe<sup>50</sup>, R. Currie<sup>47</sup>, C. D'Ambrosio<sup>35</sup>, P. David<sup>8</sup>, P.N.Y. David<sup>38</sup>, I. De Bonis<sup>4</sup>, K. De Bruyn<sup>38</sup>, S. De Capua<sup>51</sup>, M. De Cian<sup>37</sup>, J.M. De Miranda<sup>1</sup>, L. De Paula<sup>2</sup>, W. De Silva<sup>57</sup>, P. De Simone<sup>18</sup>, D. Decamp<sup>4</sup>, M. Deckenhoff<sup>9</sup>, H. Degaudenzi<sup>36,35</sup>, L. Del Buono<sup>8</sup>, C. Deplano<sup>15</sup>, D. Derkach<sup>14</sup>, O. Deschamps<sup>5</sup>, F. Dettori<sup>39</sup>, A. Di Canto<sup>11</sup>, J. Dickens<sup>44</sup>, H. Dijkstra<sup>35</sup>, P. Diniz Batista<sup>1</sup>, M. Dogaru<sup>26</sup>, F. Domingo Bonal<sup>33,n</sup>, S. Donleavy<sup>49</sup>, F. Dordei<sup>11</sup>, A. Dosil Suárez<sup>34</sup>, D. Dossett<sup>45</sup>, A. Dovbnya<sup>40</sup>, F. Dupertuis<sup>36</sup>, R. Dzhelyadin<sup>32</sup>, A. Dziurda<sup>23</sup>, A. Dzyuba<sup>27</sup>, S. Easo<sup>46,35</sup>, U. Egede<sup>50</sup>, V. Egorychev<sup>28</sup>, S. Eidelman<sup>31</sup>, D. van Eijk<sup>38</sup>, S. Eisenhardt<sup>47</sup>, U. Eitschberger<sup>9</sup>, R. Ekelhof<sup>9</sup>, L. Eklund<sup>48</sup>, I. El Rifai<sup>5</sup>, Ch. Elsasser<sup>37</sup>, D. Elsby<sup>42</sup>, A. Falabella<sup>14,e</sup>, C. Färber<sup>11</sup>, G. Fardell<sup>47</sup>, C. Farinelli<sup>38</sup>, S. Farry<sup>12</sup>, V. Fave<sup>36</sup>, D. Ferguson<sup>47</sup>, V. Fernandez Albor<sup>34</sup>, F. Ferreira Rodrigues<sup>1</sup>, M. Ferro-Luzzi<sup>35</sup>, S. Filippov<sup>30</sup>, C. Fitzpatrick<sup>35</sup>, M. Fontana<sup>10</sup>, F. Fontanelli<sup>19,i</sup>, R. Forty<sup>35</sup>, O. Francisco<sup>2</sup>, M. Frank<sup>35</sup>, C. Frei<sup>35</sup>, M. Frosini<sup>17,f</sup>, S. Furcas<sup>20</sup>, E. Furfaro<sup>21</sup>, A. Gallas Torreira<sup>34</sup>, D. Galli<sup>14,c</sup>, M. Gandelman<sup>2</sup>, P. Gandini<sup>52</sup>, Y. Gao<sup>3</sup>, J. Garofoli<sup>53</sup>, P. Garosi<sup>51</sup>, J. Garra Tico<sup>44</sup>, L. Garrido<sup>33</sup>, C. Gaspar<sup>35</sup>, R. Gauld<sup>52</sup>, E. Gersabeck<sup>11</sup>, M. Gersabeck<sup>51</sup>, T. Gershon<sup>45,35</sup>, Ph. Ghez<sup>4</sup>, V. Gibson<sup>44</sup>, V.V. Gligorov<sup>35</sup>, C. Göbel<sup>54</sup>, D. Golubkov<sup>28</sup>, A. Golutvin<sup>50,28,35</sup>, A. Gomes<sup>2</sup>, H. Gordon<sup>52</sup>, M. Grabalosa Gándara<sup>5</sup>, R. Graciani Diaz<sup>33</sup>, L.A. Granado Cardoso<sup>35</sup>, E. Graugés<sup>33</sup>, G. Graziani<sup>17</sup>, A. Greco<sup>26</sup>, E. Greening<sup>52</sup>, S. Gregson<sup>44</sup>, O. Grünberg<sup>55</sup>, B. Gui<sup>53</sup>, E. Gushchin<sup>30</sup>, Yu. Guz<sup>32</sup>, T. Gys<sup>35</sup>, C. Hadjivasiliou<sup>53</sup>, G. Haefeli<sup>36</sup>, C. Haen<sup>35</sup>, S.C. Haines<sup>44</sup>, S. Hall<sup>50</sup>, T. Hampson<sup>43</sup>, S. Hansmann-Menzemer<sup>11</sup>, N. Harnew<sup>52</sup>, S.T. Harnew<sup>43</sup>, J. Harrison<sup>51</sup>, P.F. Harrison<sup>45</sup>, T. Hartmann<sup>55</sup>, J. He<sup>7</sup>, V. Heijne<sup>38</sup>, K. Hennessy<sup>49</sup>, P. Henrard<sup>5</sup>, J.A. Hernando Morata<sup>34</sup>, E. van Herwijnen<sup>35</sup>, E. Hicks<sup>49</sup>, D. Hill<sup>52</sup>, M. Hoballah<sup>5</sup>, C. Hombach<sup>51</sup>, P. Hopchev<sup>4</sup>, W. Hulsbergen<sup>38</sup>, P. Hunt<sup>52</sup>, T. Huse<sup>49</sup>, N. Hussain<sup>52</sup>, D. Hutchcroft<sup>49</sup>, D. Hynds<sup>48</sup>, V. Iakovenko<sup>41</sup>, P. Ilten<sup>12</sup>, J. Imong<sup>43</sup>, R. Jacobsson<sup>35</sup>,

A. Jaeger<sup>11</sup>, E. Jans<sup>38</sup>, F. Jansen<sup>38</sup>, P. Jatón<sup>36</sup>, F. Jing<sup>3</sup>, M. John<sup>52</sup>, D. Johnson<sup>52</sup>, C.R. Jones<sup>44</sup>,  
 B. Jost<sup>35</sup>, M. Kabbalo<sup>9</sup>, S. Kandybei<sup>40</sup>, M. Karacson<sup>35</sup>, T.M. Karbach<sup>35</sup>, I.R. Kenyon<sup>42</sup>,  
 U. Kerzel<sup>35</sup>, T. Ketel<sup>39</sup>, A. Keune<sup>36</sup>, B. Khanji<sup>20</sup>, O. Kochebina<sup>7</sup>, I. Komarov<sup>36,29</sup>,  
 R.F. Koopman<sup>39</sup>, P. Koppenburg<sup>38</sup>, M. Korolev<sup>29</sup>, A. Kozlinskiy<sup>38</sup>, L. Kravchuk<sup>30</sup>, K. Kreplin<sup>11</sup>,  
 M. Kreps<sup>45</sup>, G. Krocker<sup>11</sup>, P. Krokovny<sup>31</sup>, F. Kruse<sup>9</sup>, M. Kucharczyk<sup>20,23,j</sup>, V. Kudryavtsev<sup>31</sup>,  
 T. Kvaratskheliya<sup>28,35</sup>, V.N. La Thi<sup>36</sup>, D. Lacarrere<sup>35</sup>, G. Lafferty<sup>51</sup>, A. Lai<sup>15</sup>, D. Lambert<sup>47</sup>,  
 R.W. Lambert<sup>39</sup>, E. Lanciotti<sup>35</sup>, G. Lanfranchi<sup>18,35</sup>, C. Langenbruch<sup>35</sup>, T. Latham<sup>45</sup>,  
 C. Lazzeroni<sup>42</sup>, R. Le Gac<sup>6</sup>, J. van Leerdam<sup>38</sup>, J.-P. Lees<sup>4</sup>, R. Lefèvre<sup>5</sup>, A. Leflat<sup>29,35</sup>,  
 J. Lefrançois<sup>7</sup>, O. Leroy<sup>6</sup>, Y. Li<sup>3</sup>, L. Li Gioi<sup>5</sup>, M. Liles<sup>49</sup>, R. Lindner<sup>35</sup>, C. Linn<sup>11</sup>, B. Liu<sup>3</sup>,  
 G. Liu<sup>35</sup>, J. von Loeben<sup>20</sup>, J.H. Lopes<sup>2</sup>, E. Lopez Asamar<sup>33</sup>, N. Lopez-March<sup>36</sup>, H. Lu<sup>3</sup>,  
 J. Luisier<sup>36</sup>, H. Luo<sup>47</sup>, A. Mac Raighne<sup>48</sup>, F. Machefert<sup>7</sup>, I.V. Machikhiliyan<sup>4,28</sup>, F. Maciuc<sup>26</sup>,  
 O. Maev<sup>27,35</sup>, S. Malde<sup>52</sup>, G. Manca<sup>15,d</sup>, G. Mancinelli<sup>6</sup>, N. Mangiafave<sup>44</sup>, U. Marconi<sup>14</sup>,  
 R. Märki<sup>36</sup>, J. Marks<sup>11</sup>, G. Martellotti<sup>22</sup>, A. Martens<sup>8</sup>, L. Martin<sup>52</sup>, A. Martín Sánchez<sup>7</sup>,  
 M. Martinelli<sup>38</sup>, D. Martinez Santos<sup>39</sup>, D. Martins Tostes<sup>2</sup>, A. Massafferri<sup>1</sup>, R. Matev<sup>35</sup>,  
 Z. Mathe<sup>35</sup>, C. Matteuzzi<sup>20</sup>, M. Matveev<sup>27</sup>, E. Maurice<sup>6</sup>, A. Mazurov<sup>16,30,35,e</sup>, J. McCarthy<sup>42</sup>,  
 R. McNulty<sup>12</sup>, B. Meadows<sup>57,52</sup>, F. Meier<sup>9</sup>, M. Meissner<sup>11</sup>, M. Merk<sup>38</sup>, D.A. Milanes<sup>13</sup>,  
 M.-N. Minard<sup>4</sup>, J. Molina Rodriguez<sup>54</sup>, S. Monteil<sup>5</sup>, D. Moran<sup>51</sup>, P. Morawski<sup>23</sup>, R. Mountain<sup>53</sup>,  
 I. Mous<sup>38</sup>, F. Muheim<sup>47</sup>, K. Müller<sup>37</sup>, R. Muresan<sup>26</sup>, B. Muryn<sup>24</sup>, B. Muster<sup>36</sup>, P. Naik<sup>43</sup>,  
 T. Nakada<sup>36</sup>, R. Nandakumar<sup>46</sup>, I. Nasteva<sup>1</sup>, M. Needham<sup>47</sup>, N. Neufeld<sup>35</sup>, A.D. Nguyen<sup>36</sup>,  
 T.D. Nguyen<sup>36</sup>, C. Nguyen-Mau<sup>36,o</sup>, M. Nicol<sup>7</sup>, V. Niess<sup>5</sup>, R. Niet<sup>9</sup>, N. Nikitin<sup>29</sup>, T. Nikodem<sup>11</sup>,  
 S. Nisar<sup>56</sup>, A. Nomerotski<sup>52</sup>, A. Novoselov<sup>32</sup>, A. Oblakowska-Mucha<sup>24</sup>, V. Obraztsov<sup>32</sup>,  
 S. Oggero<sup>38</sup>, S. Ogilvy<sup>48</sup>, O. Okhrimenko<sup>41</sup>, R. Oldeman<sup>15,d,35</sup>, M. Orlandea<sup>26</sup>,  
 J.M. Otalora Goicochea<sup>2</sup>, P. Owen<sup>50</sup>, B.K. Pal<sup>53</sup>, A. Palano<sup>13,b</sup>, M. Palutan<sup>18</sup>, J. Panman<sup>35</sup>,  
 A. Papanestis<sup>46</sup>, M. Pappagallo<sup>48</sup>, C. Parkes<sup>51</sup>, C.J. Parkinson<sup>50</sup>, G. Passaleva<sup>17</sup>, G.D. Patel<sup>49</sup>,  
 M. Patel<sup>50</sup>, G.N. Patrick<sup>46</sup>, C. Patrignani<sup>19,i</sup>, C. Pavel-Nicorescu<sup>26</sup>, A. Pazos Alvarez<sup>34</sup>,  
 A. Pellegrino<sup>38</sup>, G. Penso<sup>22,l</sup>, M. Pepe Altarelli<sup>35</sup>, S. Perazzini<sup>14,c</sup>, D.L. Perego<sup>20,j</sup>,  
 E. Perez Trigo<sup>34</sup>, A. Pérez-Calero Yzquierdo<sup>33</sup>, P. Perret<sup>5</sup>, M. Perrin-Terrin<sup>6</sup>, G. Pessina<sup>20</sup>,  
 K. Petridis<sup>50</sup>, A. Petrolini<sup>19,i</sup>, A. Phan<sup>53</sup>, E. Picatoste Olloqui<sup>33</sup>, B. Pietrzyk<sup>4</sup>, T. Pilar<sup>45</sup>,  
 D. Pinci<sup>22</sup>, S. Playfer<sup>47</sup>, M. Plo Casasus<sup>34</sup>, F. Polci<sup>8</sup>, G. Polok<sup>23</sup>, A. Poluektov<sup>45,31</sup>,  
 E. Polcarpo<sup>2</sup>, D. Popov<sup>10</sup>, B. Popovici<sup>26</sup>, C. Potterat<sup>33</sup>, A. Powell<sup>52</sup>, J. Prisciandaro<sup>36</sup>,  
 V. Pugatch<sup>41</sup>, A. Puig Navarro<sup>36</sup>, W. Qian<sup>4</sup>, J.H. Rademacker<sup>43</sup>, B. Rakotomiaramananana<sup>36</sup>,  
 M.S. Rangel<sup>2</sup>, I. Raniuk<sup>40</sup>, N. Rauschmayr<sup>35</sup>, G. Raven<sup>39</sup>, S. Redford<sup>52</sup>, M.M. Reid<sup>45</sup>,  
 A.C. dos Reis<sup>1</sup>, S. Ricciardi<sup>46</sup>, A. Richards<sup>50</sup>, K. Rinnert<sup>49</sup>, V. Rives Molina<sup>33</sup>,  
 D.A. Roa Romero<sup>5</sup>, P. Robbe<sup>7</sup>, E. Rodrigues<sup>51</sup>, P. Rodriguez Perez<sup>34</sup>, G.J. Rogers<sup>44</sup>, S. Roiser<sup>35</sup>,  
 V. Romanovsky<sup>32</sup>, A. Romero Vidal<sup>34</sup>, J. Rouvinet<sup>36</sup>, T. Ruf<sup>35</sup>, H. Ruiz<sup>33</sup>, G. Sabatino<sup>22,k</sup>,  
 J.J. Saborido Silva<sup>34</sup>, N. Sagidova<sup>27</sup>, P. Sail<sup>48</sup>, B. Saitta<sup>15,d</sup>, C. Salzmann<sup>37</sup>,  
 B. Sanmartin Sedes<sup>34</sup>, M. Sannino<sup>19,i</sup>, R. Santacesaria<sup>22</sup>, C. Santamarina Rios<sup>34</sup>,  
 E. Santovetti<sup>21,k</sup>, M. Sapunov<sup>6</sup>, A. Sarti<sup>18,l</sup>, C. Satriano<sup>22,m</sup>, A. Satta<sup>21</sup>, M. Savrie<sup>16,e</sup>,  
 D. Savrina<sup>28,29</sup>, P. Schaack<sup>50</sup>, M. Schiller<sup>39</sup>, H. Schindler<sup>35</sup>, S. Schleich<sup>9</sup>, M. Schlupp<sup>9</sup>,  
 M. Schmelling<sup>10</sup>, B. Schmidt<sup>35</sup>, O. Schneider<sup>36</sup>, A. Schopper<sup>35</sup>, M.-H. Schune<sup>7</sup>, R. Schwemmer<sup>35</sup>,  
 B. Sciascia<sup>18</sup>, A. Sciubba<sup>18,l</sup>, M. Seco<sup>34</sup>, A. Semennikov<sup>28</sup>, K. Senderowska<sup>24</sup>, I. Sepp<sup>50</sup>,  
 N. Serra<sup>37</sup>, J. Serrano<sup>6</sup>, P. Seyfert<sup>11</sup>, M. Shapkin<sup>32</sup>, I. Shapoval<sup>40,35</sup>, P. Shatalov<sup>28</sup>,  
 Y. Shcheglov<sup>27</sup>, T. Shears<sup>49,35</sup>, L. Shekhtman<sup>31</sup>, O. Shevchenko<sup>40</sup>, V. Shevchenko<sup>28</sup>, A. Shires<sup>50</sup>,  
 R. Silva Coutinho<sup>45</sup>, T. Skwarnicki<sup>53</sup>, N.A. Smith<sup>49</sup>, E. Smith<sup>52,46</sup>, M. Smith<sup>51</sup>, K. Sobczak<sup>5</sup>,  
 M.D. Sokoloff<sup>57</sup>, F.J.P. Soler<sup>48</sup>, F. Soomro<sup>18,35</sup>, D. Souza<sup>43</sup>, B. Souza De Paula<sup>2</sup>, B. Spaan<sup>9</sup>,  
 A. Sparkes<sup>47</sup>, P. Spradlin<sup>48</sup>, F. Stagni<sup>35</sup>, S. Stahl<sup>11</sup>, O. Steinkamp<sup>37</sup>, S. Stoica<sup>26</sup>, S. Stone<sup>53</sup>,

B. Storaci<sup>37</sup>, M. Straticiu<sup>26</sup>, U. Straumann<sup>37</sup>, V.K. Subbiah<sup>35</sup>, S. Swientek<sup>9</sup>, V. Syropoulos<sup>39</sup>, M. Szczekowski<sup>25</sup>, P. Szczypka<sup>36,35</sup>, T. Szumlak<sup>24</sup>, S. T'Jampens<sup>4</sup>, M. Teklishyn<sup>7</sup>, E. Teodorescu<sup>26</sup>, F. Teubert<sup>35</sup>, C. Thomas<sup>52</sup>, E. Thomas<sup>35</sup>, J. van Tilburg<sup>11</sup>, V. Tisserand<sup>4</sup>, M. Tobin<sup>37</sup>, S. Tolk<sup>39</sup>, D. Tonelli<sup>35</sup>, S. Topp-Joergensen<sup>52</sup>, N. Torr<sup>52</sup>, E. Tournefier<sup>4,50</sup>, S. Tourneur<sup>36</sup>, M.T. Tran<sup>36</sup>, M. Tresch<sup>37</sup>, A. Tsaregorodtsev<sup>6</sup>, P. Tsopelas<sup>38</sup>, N. Tuning<sup>38</sup>, M. Ubeda Garcia<sup>35</sup>, A. Ukleja<sup>25</sup>, D. Urner<sup>51</sup>, U. Uwer<sup>11</sup>, V. Vagnoni<sup>14</sup>, G. Valenti<sup>14</sup>, R. Vazquez Gomez<sup>33</sup>, P. Vazquez Regueiro<sup>34</sup>, S. Vecchi<sup>16</sup>, J.J. Velthuis<sup>43</sup>, M. Veltri<sup>17,g</sup>, G. Veneziano<sup>36</sup>, M. Vesterinen<sup>35</sup>, B. Viaud<sup>7</sup>, D. Vieira<sup>2</sup>, X. Vilasis-Cardona<sup>33,n</sup>, A. Vollhardt<sup>37</sup>, D. Volyanskyy<sup>10</sup>, D. Voong<sup>43</sup>, A. Vorobyev<sup>27</sup>, V. Vorobyev<sup>31</sup>, C. Vob<sup>55</sup>, H. Voss<sup>10</sup>, R. Waldi<sup>55</sup>, R. Wallace<sup>12</sup>, S. Wandernoth<sup>11</sup>, J. Wang<sup>53</sup>, D.R. Ward<sup>44</sup>, N.K. Watson<sup>42</sup>, A.D. Webber<sup>51</sup>, D. Websdale<sup>50</sup>, M. Whitehead<sup>45</sup>, J. Wicht<sup>35</sup>, J. Wiechczynski<sup>23</sup>, D. Wiedner<sup>11</sup>, L. Wiggers<sup>38</sup>, G. Wilkinson<sup>52</sup>, M.P. Williams<sup>45,46</sup>, M. Williams<sup>50,p</sup>, F.F. Wilson<sup>46</sup>, J. Wishahi<sup>9</sup>, M. Witek<sup>23</sup>, W. Witzeling<sup>35</sup>, S.A. Wotton<sup>44</sup>, S. Wright<sup>44</sup>, S. Wu<sup>3</sup>, K. Wyllie<sup>35</sup>, Y. Xie<sup>47,35</sup>, F. Xing<sup>52</sup>, Z. Xing<sup>53</sup>, Z. Yang<sup>3</sup>, R. Young<sup>47</sup>, X. Yuan<sup>3</sup>, O. Yushchenko<sup>32</sup>, M. Zangoli<sup>14</sup>, M. Zavertyaev<sup>10,a</sup>, F. Zhang<sup>3</sup>, L. Zhang<sup>53</sup>, W.C. Zhang<sup>12</sup>, Y. Zhang<sup>3</sup>, A. Zhelezov<sup>11</sup>, A. Zhokhov<sup>28</sup>, L. Zhong<sup>3</sup>, A. Zvyagin<sup>35</sup>.

<sup>1</sup> *Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil*

<sup>2</sup> *Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil*

<sup>3</sup> *Center for High Energy Physics, Tsinghua University, Beijing, China*

<sup>4</sup> *LAPP, Université de Savoie, CNRS/IN2P3, Annecy-Le-Vieux, France*

<sup>5</sup> *Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, Clermont-Ferrand, France*

<sup>6</sup> *CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France*

<sup>7</sup> *LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France*

<sup>8</sup> *LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris, France*

<sup>9</sup> *Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany*

<sup>10</sup> *Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany*

<sup>11</sup> *Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany*

<sup>12</sup> *School of Physics, University College Dublin, Dublin, Ireland*

<sup>13</sup> *Sezione INFN di Bari, Bari, Italy*

<sup>14</sup> *Sezione INFN di Bologna, Bologna, Italy*

<sup>15</sup> *Sezione INFN di Cagliari, Cagliari, Italy*

<sup>16</sup> *Sezione INFN di Ferrara, Ferrara, Italy*

<sup>17</sup> *Sezione INFN di Firenze, Firenze, Italy*

<sup>18</sup> *Laboratori Nazionali dell'INFN di Frascati, Frascati, Italy*

<sup>19</sup> *Sezione INFN di Genova, Genova, Italy*

<sup>20</sup> *Sezione INFN di Milano Bicocca, Milano, Italy*

<sup>21</sup> *Sezione INFN di Roma Tor Vergata, Roma, Italy*

<sup>22</sup> *Sezione INFN di Roma La Sapienza, Roma, Italy*

<sup>23</sup> *Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*

<sup>24</sup> *AGH University of Science and Technology, Kraków, Poland*

<sup>25</sup> *National Center for Nuclear Research (NCBJ), Warsaw, Poland*

<sup>26</sup> *Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*

<sup>27</sup> *Petersburg Nuclear Physics Institute (PNPI), Gatchina, Russia*

<sup>28</sup> *Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia*

<sup>29</sup> *Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia*

<sup>30</sup> *Institute for Nuclear Research of the Russian Academy of Sciences (INR RAN), Moscow, Russia*

<sup>31</sup> *Budker Institute of Nuclear Physics (SB RAS) and Novosibirsk State University, Novosibirsk, Russia*

<sup>32</sup> *Institute for High Energy Physics (IHEP), Protvino, Russia*

<sup>33</sup> *Universitat de Barcelona, Barcelona, Spain*

- <sup>34</sup> *Universidad de Santiago de Compostela, Santiago de Compostela, Spain*
- <sup>35</sup> *European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- <sup>36</sup> *Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- <sup>37</sup> *Physik-Institut, Universität Zürich, Zürich, Switzerland*
- <sup>38</sup> *Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands*
- <sup>39</sup> *Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, The Netherlands*
- <sup>40</sup> *NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- <sup>41</sup> *Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- <sup>42</sup> *University of Birmingham, Birmingham, United Kingdom*
- <sup>43</sup> *H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- <sup>44</sup> *Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- <sup>45</sup> *Department of Physics, University of Warwick, Coventry, United Kingdom*
- <sup>46</sup> *STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- <sup>47</sup> *School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- <sup>48</sup> *School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- <sup>49</sup> *Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- <sup>50</sup> *Imperial College London, London, United Kingdom*
- <sup>51</sup> *School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- <sup>52</sup> *Department of Physics, University of Oxford, Oxford, United Kingdom*
- <sup>53</sup> *Syracuse University, Syracuse, NY, United States*
- <sup>54</sup> *Pontificia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to <sup>2</sup>*
- <sup>55</sup> *Institut für Physik, Universität Rostock, Rostock, Germany, associated to <sup>11</sup>*
- <sup>56</sup> *Institute of Information Technology, COMSATS, Lahore, Pakistan, associated to <sup>53</sup>*
- <sup>57</sup> *University of Cincinnati, Cincinnati, OH, United States, associated to <sup>53</sup>*
- <sup>a</sup> *P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia*
- <sup>b</sup> *Università di Bari, Bari, Italy*
- <sup>c</sup> *Università di Bologna, Bologna, Italy*
- <sup>d</sup> *Università di Cagliari, Cagliari, Italy*
- <sup>e</sup> *Università di Ferrara, Ferrara, Italy*
- <sup>f</sup> *Università di Firenze, Firenze, Italy*
- <sup>g</sup> *Università di Urbino, Urbino, Italy*
- <sup>h</sup> *Università di Modena e Reggio Emilia, Modena, Italy*
- <sup>i</sup> *Università di Genova, Genova, Italy*
- <sup>j</sup> *Università di Milano Bicocca, Milano, Italy*
- <sup>k</sup> *Università di Roma Tor Vergata, Roma, Italy*
- <sup>l</sup> *Università di Roma La Sapienza, Roma, Italy*
- <sup>m</sup> *Università della Basilicata, Potenza, Italy*
- <sup>n</sup> *LIFAEELS, La Salle, Universitat Ramon Llull, Barcelona, Spain*
- <sup>o</sup> *Hanoi University of Science, Hanoi, Viet Nam*
- <sup>p</sup> *Massachusetts Institute of Technology, Cambridge, MA, United States*

# 1 Introduction

Direct  $CP$  violation can arise in  $B^0 \rightarrow DK^{*0}$  decays<sup>1</sup> from the interference between two colour-suppressed transitions:  $\bar{b} \rightarrow \bar{c}$  (Cabibbo favoured) and  $\bar{b} \rightarrow \bar{u}$  (Cabibbo suppressed). The corresponding Feynman diagrams are shown in Fig. 1; interference occurs if the  $D^0$  and  $\bar{D}^0$  mesons decay to a common final state. The magnitude of the  $CP$ -violating asymmetry that arises from this interference is related to the value of the phase  $\gamma = \arg[-(V_{ud}V_{ub}^*)/(V_{cd}V_{cb}^*)]$ , the least-well determined angle of the Unitarity Triangle. A method to determine  $\gamma$  from hadronic  $B$ -decay rates was originally proposed by Gronau, London and Wyler (GLW) in Ref. [1] for various charged and neutral  $B \rightarrow DK$  decay modes and can be applied to the decay mode  $B^0 \rightarrow DK^{*0}$ . In this mode, the charge of the kaon from the  $K^{*0} \rightarrow K^+\pi^-$  decay unambiguously tags the flavour of the decaying  $B$  meson [2], hence no time-dependent tagged analysis is required.

The use of these neutral  $B$  decays is particularly interesting because the magnitude of the ratio of the suppressed over the favoured amplitude, which controls the size of the interference, is expected to be relatively large (naively a factor three larger than the analogous ratio for  $B^+ \rightarrow DK^+$  decays), hence the system can exhibit large  $CP$ -violating effects, depending on the  $D$  decay. Among the modes used in the GLW method, which are studied in this paper, large  $CP$  asymmetries can be expected when the  $D$  meson is reconstructed in a  $CP$  eigenstate. Contributions from  $B^0$  decays to the non-resonant  $DK^+\pi^-$  final state, which can pollute the  $DK^{*0}$  reconstructed signal candidates due to the large natural width of the  $K^{*0}$ , can be treated in a model-independent way, as shown in Ref. [3]. Studies with simulated events have shown that the  $B^0 \rightarrow DK^{*0}$  mode is one of the most promising channels to provide a precise measurement of  $\gamma$  at LHCb [4]. Results with this channel will therefore complement those from  $B^+ \rightarrow DK^+$ , which have recently




Figure 1: Feynman diagrams for (a)  $B^0 \rightarrow \bar{D}^0 K^{*0}$  and (b)  $B^0 \rightarrow D^0 K^{*0}$ .

<sup>1</sup>Here and in the following,  $D$  represents a neutral meson that is an admixture of  $D^0$  and  $\bar{D}^0$ . Inclusion of charge conjugate modes is implied unless specified otherwise.

been used by LHCb to constrain  $\gamma$  [5].

This paper presents the measurement of the  $B^0 - \bar{B}^0$  partial width asymmetry using  $D$  decays into the  $CP$  eigenstate  $K^+K^-$ ,

$$\mathcal{A}_d^{KK} = \frac{\Gamma(\bar{B}^0 \rightarrow D_{[K^+K^-]}\bar{K}^{*0}) - \Gamma(B^0 \rightarrow D_{[K^+K^-]}K^{*0})}{\Gamma(\bar{B}^0 \rightarrow D_{[K^+K^-]}\bar{K}^{*0}) + \Gamma(B^0 \rightarrow D_{[K^+K^-]}K^{*0})}, \quad (1)$$

together with the measurement of the ratio of the average of the  $B^0$  and  $\bar{B}^0$  partial widths with  $D \rightarrow K^+K^-$ , to the average partial width with  $D \rightarrow K^+\pi^-$  (where the sign of the kaon charge from the  $D$  decay is the same as that of the kaon from the  $K^{*0}$  decay),

$$\mathcal{R}_d^{KK} = \frac{\Gamma(\bar{B}^0 \rightarrow D_{[K^+K^-]}\bar{K}^{*0}) + \Gamma(B^0 \rightarrow D_{[K^+K^-]}K^{*0})}{\Gamma(\bar{B}^0 \rightarrow D_{[K^-\pi^+]}\bar{K}^{*0}) + \Gamma(B^0 \rightarrow D_{[K^+\pi^-]}K^{*0})}. \quad (2)$$

These quantities can be used together with other inputs to determine the value of  $\gamma$ . Note that the suppressed decay mode  $B^0 \rightarrow D_{[K^-\pi^+]}\bar{K}^{*0}$ , where the sign of the kaon charge from the  $D$  decay is opposite to that of the kaon from the  $K^{*0}$  decay, is not included in this analysis. This decay mode can exhibit large  $CP$ -violating effects and can be studied with a larger dataset. The measured asymmetry in the favoured decay  $B^0 \rightarrow D_{[K^+\pi^-]}K^{*0}$ ,

$$\mathcal{A}_d^{\text{fav}} = \frac{\Gamma(\bar{B}^0 \rightarrow D_{[K^-\pi^+]}\bar{K}^{*0}) - \Gamma(B^0 \rightarrow D_{[K^+\pi^-]}K^{*0})}{\Gamma(\bar{B}^0 \rightarrow D_{[K^-\pi^+]}\bar{K}^{*0}) + \Gamma(B^0 \rightarrow D_{[K^+\pi^-]}K^{*0})} \quad (3)$$

is a useful cross-check since it is expected to be compatible with zero given the size of the current dataset.

In  $pp$  collisions,  $B_s^0$  mesons are produced and can decay to the same final state,  $B_s^0 \rightarrow D\bar{K}^{*0}$  [6]. In these  $B_s^0$  decay modes, the interference between the two contributing amplitudes is expected to be small, since the relative magnitude of the suppressed to the favoured amplitude is small compared to the  $B^0$  case. Therefore, these modes are valuable control channels, and the asymmetry

$$\mathcal{A}_s^{KK} = \frac{\Gamma(\bar{B}_s^0 \rightarrow D_{[K^+K^-]}K^{*0}) - \Gamma(B_s^0 \rightarrow D_{[K^+K^-]}\bar{K}^{*0})}{\Gamma(\bar{B}_s^0 \rightarrow D_{[K^+K^-]}K^{*0}) + \Gamma(B_s^0 \rightarrow D_{[K^+K^-]}\bar{K}^{*0})}, \quad (4)$$

similar to that defined in Eq. (1), is also obtained in this analysis. Since the favoured (suppressed)  $\bar{B}_s^0$  ( $B_s^0$ ) decay gives kaons with opposite charges from  $D$  and  $K^{*0}$  decays,  $\mathcal{A}_s^{\text{fav}}$  is not used as a control measurement in the analysis, to avoid biasing a potential future measurement of  $\mathcal{A}_d^{\text{sup}}$ .

## 2 The LHCb detector, dataset and event selection

The study reported here is based on a data sample collected at the Large Hadron Collider (LHC) with the LHCb detector at a centre-of-mass energy of 7 TeV during the year 2011,



corresponding to an integrated luminosity of  $1.0 \text{ fb}^{-1}$ . The LHCb detector [7] is a single-arm forward spectrometer covering the pseudorapidity range  $2 < \eta < 5$ , designed for the study of particles containing  $b$  or  $c$  quarks. The detector includes a high precision tracking system consisting of a silicon-strip vertex detector surrounding the  $pp$  interaction region, a large-area silicon-strip detector located upstream of a dipole magnet with a bending power of about 4 Tm, and three stations of silicon-strip detectors and straw drift tubes placed downstream. The combined tracking system has a momentum resolution  $\Delta p/p$  that varies from 0.4% at 5 GeV/ $c$  to 0.6% at 100 GeV/ $c$ , and an impact parameter resolution of 20  $\mu\text{m}$  for tracks with high transverse momentum ( $p_{\text{T}}$ ). Charged hadrons are identified using two ring-imaging Cherenkov detectors. Photon, electron and hadron candidates are identified by a calorimeter system consisting of scintillating-pad and preshower detectors, an electromagnetic calorimeter and a hadronic calorimeter. Muons are identified by a system composed of alternating layers of iron and multiwire proportional chambers. The trigger [8] consists of a hardware stage, based on information from the calorimeter and muon systems, followed by a software stage which applies a full event reconstruction.

This analysis uses events selected by the hardware level trigger either when one of the charged particles of the signal decay gives a large enough energy deposit in the calorimeter system (hadron trigger), or when one of the particles in the event, not coming from the signal decay, fulfills the trigger requirements (*i.e.* mainly events triggered by one particle coming from the decay of the other  $B$  in the event). The software trigger requires a two-, three- or four-track secondary vertex with a high scalar sum of the  $p_{\text{T}}$  of the tracks and a significant displacement from the primary  $pp$  interaction vertices (PVs). At least one track should have  $p_{\text{T}} > 1.7 \text{ GeV}/c$  and an impact parameter (IP)  $\chi^2$  with respect to the PV greater than 16. The IP  $\chi^2$  is defined as the difference between the  $\chi^2$  of the PV reconstructed with and without the considered track. A multivariate algorithm is used for the identification of secondary vertices consistent with the decay of a  $b$  hadron.

Candidates are selected from combinations of charged particles.  $D$  mesons are reconstructed in the decay modes  $D \rightarrow K^+\pi^-$  and  $K^+K^-$ . The  $p_{\text{T}}$  of the daughters is required to be larger than 400 MeV/ $c$ . Particle identification (PID) is used to distinguish between charged pions and kaons. The difference between the log-likelihoods of the kaon and pion hypotheses ( $\text{DLL}_{K\pi}$ ) is required to be larger than 0 for kaons and smaller than 4 for pions. This aids the reduction of cross-feed between the signal  $D$  decay modes to a negligible level. A fit is applied to the two-track vertex, requiring that the corresponding  $\chi^2$  per degree of freedom is less than 5. In order to separate  $D$  mesons coming from a  $B$  decay from those produced at the PV, the  $D$  candidates are required to have an IP  $\chi^2$  greater than 4 with respect to any PV. To suppress background from  $B$  decays without an intermediate  $D$  meson ( $B^0 \rightarrow K^{*0}K^+K^-$  for example), for which all four charged hadrons are produced at the  $B$ -decay vertex, a condition on the  $D$  flight distance with respect to the  $B$  vertex is applied, requiring that it is larger than 0 by at least 2.5 standard deviations. Finally,  $D$  candidates with an invariant mass within  $\pm 20 \text{ MeV}/c^2$  of the nominal  $D^0$  mass are retained.

$K^{*0}$  mesons are reconstructed in the mode  $K^{*0} \rightarrow K^+\pi^-$ . The  $p_{\text{T}}$  of the  $K^+$  and  $\pi^-$  mesons must be larger than 300 MeV/ $c$ . PID is also used, requiring that  $\text{DLL}_{K\pi}$  is larger than 3 for the kaon and lower than 3 for the pion, reducing the cross-feed from

$B^0 \rightarrow D\rho^0$  to a manageable level and rejecting non-resonant  $B^0 \rightarrow DK^+K^-$  [9]. Possible contamination from protons in the kaon sample, *e.g.* from  $\Lambda_b^0 \rightarrow Dp\pi^-$  decays, is reduced by removing kaon candidates with a difference between the log-likelihoods of the proton and kaon hypotheses ( $\text{DLL}_{pK}$ ) of less than 10. The IP  $\chi^2$  of the  $K^{*0}$  mesons must be larger than 25, to select those coming from a  $B$  decay, and their invariant mass within  $\pm 50 \text{ MeV}/c^2$  of the nominal mass.

$B_{(s)}^{(0)}$  meson candidates are formed by combining  $D$  and  $K^{*0}$  candidates selected with the above requirements. A fit to a common vertex is performed, keeping only combinations with  $\chi^2$  per degree of freedom lower than 4, and a kinematic fit is performed to constrain the invariant mass of the reconstructed  $D$  to the nominal  $D^0$  mass [10]. Since  $B$  mesons are produced at the PV, only candidates with IP  $\chi^2$  lower than 9 are retained. In case several PVs are reconstructed, the one for which the  $B$ -candidate IP  $\chi^2$  is the smallest is taken as reference. Additionally, the momentum of the reconstructed  $B$  candidate is required to point back to the PV, by requiring that the angle between the  $B$  momentum direction and its direction of flight from the PV is smaller than 10 mrad. Furthermore, the sum of the square roots of the IP  $\chi^2$  of the four charged particles must be larger than 32. The absolute value of the cosine of the  $K^{*0}$  helicity angle is required to be larger than 0.4. This angle is defined as the angle between the kaon-daughter momentum direction in the  $K^{*0}$  rest frame, and the  $K^{*0}$  direction in the  $B$  rest frame.

Specific peaking backgrounds from  $B_{(s)}^0 \rightarrow D_{(s)}^\mp h^\pm$  decays, where  $h$  is a  $\pi$  or a  $K$  meson, are eliminated by vetoing candidates for which the invariant mass of  $K^+K^-\pi^+$  ( $K^-\pi^+\pi^+$  and  $K^+K^-\pi^+$ ) is within  $\pm 15 \text{ MeV}/c^2$  of the nominal mass of a  $D_s^+$  ( $D^+$ ) meson.

Where possible, data-driven methods are used to determine selection efficiencies and invariant mass distribution shapes. Otherwise, they are determined from fully simulated events. The  $pp$  collisions are generated using PYTHIA 6.4 [11] with a specific LHCb configuration [12] where, in particular, decays of hadronic particles are described by EVTGEN [13]. The interaction of the generated particles with the detector and its response are implemented using the GEANT4 toolkit [14] as described in Ref. [15].

### 3 Determination of signal yields

The numbers of reconstructed signal  $B^0$  and  $B_s^0$  candidates are determined from an unbinned maximum likelihood fit to their mass distributions. Candidates are split into four categories, which are fitted simultaneously:  $D(K^+K^-)K^{*0}$ ,  $D(K^+K^-)\bar{K}^{*0}$ ,  $D(K^+\pi^-)K^{*0}$ , and  $D(K^-\pi^+)\bar{K}^{*0}$ . The mass distribution of each category is fitted with a sum of probability density functions (PDF) modelling the different contributing components:

1. the  $B^0$  and  $B_s^0$  signals are described by double Gaussian functions;
2. the combinatorial background is described by an exponential function;
3. the cross-feed from  $B^0 \rightarrow D\rho^0$  decays, where one pion from the  $\rho^0 \rightarrow \pi^+\pi^-$  decay is misidentified as a kaon, is described by a non-parametric PDF [16] determined from

fully simulated and selected events;

4. the partially reconstructed  $B^0 \rightarrow D^* K^{*0}$  and  $\bar{B}_s^0 \rightarrow D^* K^{*0}$  decays, where the  $D^*$  is a  $D^{*0}$  or a  $\bar{D}^{*0}$  and the  $\pi^0$  or photon from the  $D^*$  decay is not reconstructed, are modelled by a non-parametric PDF determined from fully simulated and selected events.

There are 23 free parameters in the fit. These include the  $B^0$  PDF peak position, the core Gaussian resolution for the  $B^0$  and the  $B_s^0$  and the slope of the combinatorial background, all of which are common to the four fit categories. The remaining free parameters are yields for each fit component within each category. Yields for  $B_{(s)}^0$  and  $\bar{B}_{(s)}^0$  are constrained to be identical for the background components where  $CP$  violation effects can be excluded or are expected to be compatible with zero with the current data sample size.

A separate fit to  $B^0 \rightarrow D(K^+\pi^-)\rho^0$  candidates in the same data sample is performed. The yield of such candidates and the probability to reconstruct them as  $B^0 \rightarrow D(K^+\pi^-)K^{*0}$  is used to constrain the number of cross-feed events in the  $D(K^+\pi^-)K^{*0}$  category. The number of cross-feed candidates from  $B^0 \rightarrow D(K^+K^-)\rho^0$  in the  $D(K^+K^-)K^{*0}$  category is derived from the  $D(K^+\pi^-)K^{*0}$  category using the relative  $D$  branching fractions and  $B$  selection efficiencies. As no flavour asymmetry is expected for this background, the numbers of cross-feed events in the  $D\bar{K}^{*0}$  categories are constrained to be identical to those of the corresponding  $DK^{*0}$  categories.

The partially reconstructed background component accumulates at masses lower than the nominal  $B^0$  mass. Its shape depends on the unknown fraction of transverse polarisation in the  $\bar{B}_{(s)}^0 \rightarrow D^* K^{*0}$  decays. In order to model the  $\bar{B}_{(s)}^0 \rightarrow D^* K^{*0}$  contribution, a PDF is built from a linear combination of three non-parametric functions corresponding to the three orthogonal helicity eigenstates. The functions are derived from simulated  $\bar{B}_{(s)}^0 \rightarrow D^* K^{*0}$  events reconstructed as  $B^0 \rightarrow DK^{*0}$ . Each function corresponds to the weighted sum of the  $D^* \rightarrow D\gamma$  and  $D^* \rightarrow D\pi^0$  contributions for a defined helicity eigenstate, where the weights take into account the relative  $D^*$  decay branching fractions and the corresponding reconstruction efficiencies.

The invariant mass distributions together with the function resulting from the fit are shown in Fig. 2. Note that the decay  $\bar{B}_s^0 \rightarrow D(K^+\pi^-)K^{*0}$  is not observed since the charge combination of the kaons in the final state corresponds to the suppressed decay. The signal yield in each category is summarized in Table 1. The significance of the  $B^0 \rightarrow DK^{*0}$  signal for  $D \rightarrow K^+K^-$  decays, summing  $B^0$  and  $\bar{B}^0$  and including both statistical and systematic uncertainties, is found to be equal to  $5.1\sigma$ , by comparing the maximum of the likelihood of the nominal fit and the maximum with the yield of the  $B^0 \rightarrow D(K^+K^-)K^{*0}$  category set to zero.

The yields determined from the simultaneous mass fit are corrected for selection efficiencies in order to evaluate the asymmetries and ratios described in the introduction. The selection efficiencies account for the geometrical acceptance of the detector, the reconstruction, the PID, and the trigger efficiencies. All efficiencies are computed from

fully simulated events, except for the PID and trigger efficiencies, which are obtained directly from data using clean calibration samples of  $D^0 \rightarrow K^- \pi^+$  from  $D^{*+}$  decays.

Table 1: Signal yields with their statistical uncertainties.

Category	Signal yield	Category	Signal yield
$B^0 \rightarrow D_{[K^+K^-]} K^{*0}$	$21^{+6}_{-5}$	$\bar{B}^0 \rightarrow D_{[K^+K^-]} \bar{K}^{*0}$	$8 \pm 4$
$B_s^0 \rightarrow D_{[K^+K^-]} \bar{K}^{*0}$	$23^{+6}_{-5}$	$\bar{B}_s^0 \rightarrow D_{[K^+K^-]} K^{*0}$	$24^{+6}_{-5}$
$B^0 \rightarrow D_{[K^+\pi^-]} K^{*0}$	$108^{+12}_{-11}$	$\bar{B}^0 \rightarrow D_{[K^-\pi^+]} \bar{K}^{*0}$	$94 \pm 11$




Figure 2: Invariant mass distributions of (a)  $D_{[K^+K^-]} \bar{K}^{*0}$ , (b)  $D_{[K^+K^-]} K^{*0}$ , (c)  $D_{[K^-\pi^+]} \bar{K}^{*0}$  and (d)  $D_{[K^+\pi^-]} K^{*0}$  candidates. The  $D\bar{K}^{*0}$  distributions correspond to  $\bar{B}^0$  and  $B_s^0$  decays whereas the  $DK^{*0}$  distributions correspond to  $B^0$  and  $\bar{B}_s^0$  decays. The fit functions are superimposed; the different  $B$  decays and combinatorial background components are detailed in the legends.

## 4 Systematic uncertainties

Several sources of systematic uncertainty are considered, affecting either the determination of the signal yields or the computation of the efficiencies. They are summarized in Table 2. In order to take into account the measured difference in the production rate between  $\bar{B}^0$  and  $B^0$ , the  $\bar{B}^0$  yields are multiplied by a correction factor,

$$a_{\text{prod}}^d = \frac{1 - \kappa A_{\text{prod}}}{1 + \kappa A_{\text{prod}}}, \quad (5)$$

where  $A_{\text{prod}} = 0.010 \pm 0.013$  [17] is the asymmetry between  $\bar{B}^0$  and  $B^0$  at production in  $pp$  collisions, and  $\kappa$  is a decay-dependent factor,  $\kappa = \frac{\int_0^{+\infty} e^{-\Gamma t} \cos(\Delta m t) \epsilon(B^0 \rightarrow DK^{*0}, t) dt}{\int_0^{+\infty} e^{-\Gamma t} \epsilon(B^0 \rightarrow DK^{*0}, t) dt}$ , which takes into account dilution effects due to the  $B^0 - \bar{B}^0$  oscillation frequency,  $\Delta m$ , and includes the acceptance as a function of the decay time for the reconstructed decay,  $\epsilon(B^0 \rightarrow DK^{*0}, t)$ . The value of  $\kappa$  is found to be  $0.46 \pm 0.01$  using fully simulated events and PID efficiencies from calibration samples. The uncertainty on  $a_{\text{prod}}^d$  is propagated to the measured observables to estimate the systematic uncertainty from the production asymmetry. Owing to the large  $B_s^0$  oscillation frequency, the potential production asymmetry of  $B_s^0$  mesons does not significantly affect the measurement presented here and is neglected.

The PID calibration introduces a systematic uncertainty on the calculated PID efficiencies, which propagates to the final results. All PID correction factors are compatible with unity within their uncertainties which are of the order of 1%.

The systematic uncertainty associated to the trigger is estimated by varying in the simulation the fraction of events triggered by the hadron trigger with respect to the fraction of events triggered by the other  $b$ -hadron in the event. Other selection efficiencies cancel in the ratio of yields, except for the efficiencies of the  $p_T$  cuts on the  $D$  daughters, which are different between different  $D$  decay modes.  $\mathcal{R}_d^{KK}$  has to be corrected by a multiplicative factor  $0.94 \pm 0.04$ , where the statistical uncertainty on the correction, which arises from finite simulated sample size, is assigned as systematic uncertainty due to the relative selection efficiencies.

The fit procedure is validated with simulated experiments. A bias of statistical nature, owing to the small number of events in the  $B^0 \rightarrow D(K^+K^-)K^{*0}$  channel, is found to be 5% for  $B^0$  and 8% for  $\bar{B}^0$ . The signal yields are corrected for this bias before computing the asymmetries and ratios. A systematic uncertainty equal to half the size of the correction has been assigned.

Simulated experiments are also used to determine the systematic uncertainties due to the low-mass background, the  $B^0 \rightarrow D\rho^0$  cross-feed, and the signal shape. Samples are generated with different values of the polarisation parameters, the cross-feed fraction and the fixed signal parameters. The corresponding systematic uncertainty is estimated from the bias in the results obtained by performing the fit described in the previous section to these samples.

Table 2: Summary of the absolute systematic uncertainties on the measured observables.

Source	$\mathcal{A}_d^{KK}$	$\mathcal{A}_d^{\text{fav}}$	$\mathcal{A}_s^{KK}$	$\mathcal{R}_d^{KK}$
Production asymmetry	0.005	0.006	–	0.003
PID efficiency	0.004	0.008	0.005	0.014
Trigger efficiency	0.004	0.001	0.005	0.022
Selection efficiency	–	–	–	0.040
Bias correction	0.004	–	0.001	0.013
Low-mass background	0.017	0.001	0.004	0.042
$B^0 \rightarrow D\rho^0$ cross-feed	0.001	–	0.002	0.008
Signal description	0.001	0.001	0.001	0.005
$D$ branching fractions	–	–	–	0.022
Total	0.019	0.010	0.008	0.069

## 5 Results and summary

This paper reports the analysis of  $B^0 \rightarrow DK^{*0}$  decays using 1.0 fb<sup>-1</sup> of  $pp$  collision data. Potential contributions to the decay amplitudes from the non-resonant  $B^0 \rightarrow DK^+\pi^-$  mode are reduced by requiring that the  $K^{*0}$  reconstructed mass is within  $\pm 50$  MeV/ $c^2$  of the nominal mass and the absolute value of the cosine of the  $K^{*0}$  helicity angle is greater than 0.4. The results for the  $CP$ -violating observables are

$$\begin{aligned}
 \mathcal{A}_d^{KK} &= -0.45 \pm 0.23 \text{ (stat)} \pm 0.02 \text{ (syst)}, \\
 \mathcal{A}_d^{\text{fav}} &= -0.08 \pm 0.08 \text{ (stat)} \pm 0.01 \text{ (syst)}, \\
 \mathcal{A}_s^{KK} &= 0.04 \pm 0.16 \text{ (stat)} \pm 0.01 \text{ (syst)}, \\
 \mathcal{R}_d^{KK} &= 1.36 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.37 \\ 0.32 \end{smallmatrix} \text{ (stat)} \pm 0.07 \text{ (syst)}.
 \end{aligned}$$

The value of  $\mathcal{R}_d^{KK}$  takes into account the ratio of the branching fractions of  $D^0 \rightarrow K^+K^-$  to  $D^0 \rightarrow K^-\pi^+$  decays [10]. The correlation between  $\mathcal{A}_d^{KK}$  and  $\mathcal{R}_d^{KK}$  is equal to 0.16 and the correlations between the other observables are negligible.

These are the first measurements of  $CP$  asymmetries in  $B^0$  and  $\bar{B}_s^0$  to  $DK^{*0}$  decays with the neutral  $D$  meson decaying into a  $CP$ -even final state. Triggering, reconstructing and selecting a pure sample of these fully hadronic  $B$  decays is challenging in a high rate and high track-multiplicity environment, especially in the forward direction of LHCb. The present statistical limitations are due to a combination of several factors, the most important one being the trigger. In order to keep the output rate below its maximum of 1 MHz, the current hardware trigger imposes relatively restrictive criteria on the minimum

transverse momentum of hadrons, which affect the efficiency for fully-hadronic modes. This limitation is overcome in the proposed LHCb upgrade [18] by reading out the detector at the maximum LHC bunch-crossing frequency of 40 MHz. With more data, improved measurements of these and other quantities in  $B^0 \rightarrow DK^{*0}$  decays will result in important constraints on the angle  $\gamma$  of the Unitarity Triangle.

## Acknowledgements

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); NSFC (China); CNRS/IN2P3 and Region Auvergne (France); BMBF, DFG, HGF and MPG (Germany); SFI (Ireland); INFN (Italy); FOM and NWO (The Netherlands); SCSR (Poland); ANCS/IFA (Romania); MinES, Rosatom, RFBR and NRC ‘‘Kurchatov Institute’’ (Russia); MinECo, XuntaGal and GENCAT (Spain); SNSF and SER (Switzerland); NAS Ukraine (Ukraine); STFC (United Kingdom); NSF (USA). We also acknowledge the support received from the ERC under FP7. The Tier1 computing centres are supported by IN2P3 (France), KIT and BMBF (Germany), INFN (Italy), NWO and SURF (The Netherlands), PIC (Spain), GridPP (United Kingdom). We are thankful for the computing resources put at our disposal by Yandex LLC (Russia), as well as to the communities behind the multiple open source software packages that we depend on.

## References

- [1] M. Gronau and D. London, *How to determine all the angles of the unitarity triangle from  $B_d^0 \rightarrow DK_S$  and  $B_s^0 \rightarrow D\phi$* , Phys. Lett. **B253** (1991) 483; M. Gronau and D. Wyler, *On determining a weak phase from charged B decay asymmetries*, Phys. Lett. **B265** (1991) 172.
- [2] I. Dunietz, *CP violation with self-tagging  $B_d$  modes*, Phys. Lett. **B270** (1991) 75.
- [3] M. Gronau, *Improving bounds on  $\gamma$  in  $B^\pm \rightarrow DK^\pm$  and  $B^{\pm,0} \rightarrow DX_s^{\pm,0}$* , Phys. Lett. **B557** (2003) 198, [arXiv:hep-ph/0211282](#).
- [4] LHCb collaboration, B. Adeva *et al.*, *Roadmap for selected key measurements of LHCb*, [arXiv:0912.4179](#).
- [5] LHCb Collaboration, R. Aaij *et al.*, *Observation of CP violation in  $B^+$  to  $DK^+$  decays*, Phys. Lett. **B712** (2012) 203, [arXiv:1203.3662](#), erratum Phys. Lett. **B713** (2012) 351; R. Aaij *et al.*, *A model-independent Dalitz plot analysis of  $B^\pm \rightarrow DK^\pm$  with  $D \rightarrow K_S^0 h^+ h^-$  ( $h = \pi, K$ ) decays and constraints on the CKM angle  $\gamma$* , Phys. Lett. **B718** (2012) 43, [arXiv:1209.5869](#).

- [6] LHCb collaboration, R. Aaij *et al.*, *First observation of the decay  $\bar{B}_s^0 \rightarrow D^0 K^{*0}$  and a measurement of the ratio of branching fractions  $\frac{\mathcal{B}(\bar{B}_s^0 \rightarrow D^0 K^{*0})}{\mathcal{B}(\bar{B}^0 \rightarrow D^0 \rho^0)}$* , Phys. Lett. **B706** (2011) 32, [arXiv:1110.3676](#).
- [7] LHCb collaboration, A. A. Alves Jr. *et al.*, *The LHCb detector at the LHC*, JINST **3** (2008) S08005.
- [8] R. Aaij *et al.*, *The LHCb trigger and its performance*, [arXiv:1211.3055](#).
- [9] R. Aaij *et al.*, *Observation of  $B^0 \rightarrow \bar{D}^0 K^+ K^-$  and evidence of  $B_s^0 \rightarrow \bar{D}^0 K^+ K^-$* , Phys. Rev. Lett. **109** (2012) 131801, [arXiv:1207.5991](#).
- [10] Particle Data Group, J. Beringer *et al.*, *Review of particle physics*, Phys. Rev. **D86** (2012) 010001.
- [11] T. Sjöstrand, S. Mrenna, and P. Skands, *PYTHIA 6.4 physics and manual*, JHEP **05** (2006) 026, [arXiv:hep-ph/0603175](#).
- [12] I. Belyaev *et al.*, *Handling of the generation of primary events in GAUSS, the LHCb simulation framework*, Nuclear Science Symposium Conference Record (NSS/MIC) **IEEE** (2010) 1155.
- [13] D. J. Lange, *The EvtGen particle decay simulation package*, Nucl. Instrum. Meth. **A462** (2001) 152.
- [14] GEANT4 collaboration, J. Allison *et al.*, *Geant4 developments and applications*, IEEE Trans. Nucl. Sci. **53** (2006) 270; GEANT4 collaboration, S. Agostinelli *et al.*, *GEANT4: a simulation toolkit*, Nucl. Instrum. Meth. **A506** (2003) 250.
- [15] M. Clemencic *et al.*, *The LHCb simulation application, GAUSS: design, evolution and experience*, J. of Phys: Conf. Ser. **331** (2011) 032023.
- [16] K. S. Cranmer, *Kernel estimation in high-energy physics*, Comput. Phys. Commun. **136** (2001) 198, [arXiv:hep-ex/0011057](#).
- [17] LHCb collaboration, R. Aaij *et al.*, *First evidence of direct CP violation in charmless two-body decays of  $B_s$  mesons*, Phys. Rev. Lett. **108** (2012) 201601, [arXiv:1202.6251](#).
- [18] LHCb collaboration, R. Aaij *et al.*, *Letter of intent for the LHCb upgrade*, LHCb-PUB-2011-001; LHCb collaboration, I. Bediaga *et al.*, *Framework TDR for the LHCb upgrade*, CERN-LHCC-2012-007.