



First observation of the decay

$$B_c^+ \rightarrow J/\psi K^+$$

The LHCb collaboration[†]

Abstract

The decay $B_c^+ \rightarrow J/\psi K^+$ is observed for the first time using a data sample, corresponding to an integrated luminosity of 1.0 fb^{-1} , collected by the LHCb experiment in pp collisions at a centre-of-mass energy of 7 TeV. A yield of 46 ± 12 events is reported, with a significance of 5.0 standard deviations. The ratio of the branching fraction of $B_c^+ \rightarrow J/\psi K^+$ to that of $B_c^+ \rightarrow J/\psi \pi^+$ is measured to be $0.069 \pm 0.019 \pm 0.005$, where the first uncertainty is statistical and the second is systematic.

Submitted to JHEP

© CERN on behalf of the LHCb collaboration, license CC-BY-3.0.

[†]Authors are listed on the following pages.

LHCb collaboration

R. Aaij⁴⁰, C. Abellan Beteta^{35,n}, B. Adeva³⁶, M. Adinolfi⁴⁵, C. Adrover⁶, A. Affolder⁵¹, Z. Ajaltouni⁵, J. Albrecht⁹, F. Alessio³⁷, M. Alexander⁵⁰, S. Ali⁴⁰, G. Alkhazov²⁹, P. Alvarez Cartelle³⁶, A.A. Alves Jr^{24,37}, S. Amato², S. Amerio²¹, Y. Amhis⁷, L. Anderlini^{17,f}, J. Anderson³⁹, R. Andreassen⁵⁶, R.B. Appleby⁵³, O. Aquines Gutierrez¹⁰, F. Archilli¹⁸, A. Artamonov³⁴, M. Artuso⁵⁸, E. Aslanides⁶, G. Auriemma^{24,m}, S. Bachmann¹¹, J.J. Back⁴⁷, C. Baesso⁵⁹, V. Balagura³⁰, W. Baldini¹⁶, R.J. Barlow⁵³, C. Barschel³⁷, S. Barsuk⁷, W. Barter⁴⁶, Th. Bauer⁴⁰, A. Bay³⁸, J. Beddow⁵⁰, F. Bedeschi²², I. Bediaga¹, S. Belogurov³⁰, K. Belous³⁴, I. Belyaev³⁰, E. Ben-Haim⁸, G. Bencivenni¹⁸, S. Benson⁴⁹, J. Benton⁴⁵, A. Berezhnoy³¹, R. Bernet³⁹, M.-O. Bettler⁴⁶, M. van Beuzekom⁴⁰, A. Bien¹¹, S. Bifani⁴⁴, T. Bird⁵³, A. Bizzeti^{17,h}, P.M. Bjørnstad⁵³, T. Blake³⁷, F. Blanc³⁸, J. Blouw¹¹, S. Blusk⁵⁸, V. Bocci²⁴, A. Bondar³³, N. Bondar²⁹, W. Bonivento¹⁵, S. Borghi⁵³, A. Borgia⁵⁸, T.J.V. Bowcock⁵¹, E. Bowen³⁹, C. Bozzi¹⁶, T. Brambach⁹, J. van den Brand⁴¹, J. Bressieux³⁸, D. Brett⁵³, M. Britsch¹⁰, T. Britton⁵⁸, N.H. Brook⁴⁵, H. Brown⁵¹, I. Burducea²⁸, A. Bursche³⁹, G. Busetto^{21,p}, J. Buytaert³⁷, S. Cadeddu¹⁵, O. Callot⁷, M. Calvi^{20,j}, M. Calvo Gomez^{35,n}, A. Camboni³⁵, P. Campana^{18,37}, D. Campora Perez³⁷, A. Carbone^{14,c}, G. Carboni^{23,k}, R. Cardinale^{19,i}, A. Cardini¹⁵, H. Carranza-Mejia⁴⁹, L. Carson⁵², K. Carvalho Akiba², G. Casse⁵¹, L. Castillo Garcia³⁷, M. Cattaneo³⁷, Ch. Cauter⁹, M. Charles⁵⁴, Ph. Charpentier³⁷, P. Chen^{3,38}, N. Chiapolini³⁹, M. Chrzaszcz²⁵, K. Ciba³⁷, X. Cid Vidal³⁷, G. Ciezarek⁵², P.E.L. Clarke⁴⁹, M. Clemencic³⁷, H.V. Cliff⁴⁶, J. Closier³⁷, C. Coca²⁸, V. Coco⁴⁰, J. Cogan⁶, E. Cogneras⁵, P. Collins³⁷, A. Comerma-Montells³⁵, A. Contu¹⁵, A. Cook⁴⁵, M. Coombes⁴⁵, S. Coquereau⁸, G. Corti³⁷, B. Couturier³⁷, G.A. Cowan⁴⁹, D.C. Craik⁴⁷, S. Cunliffe⁵², R. Currie⁴⁹, C. D'Ambrosio³⁷, P. David⁸, P.N.Y. David⁴⁰, A. Davis⁵⁶, I. De Bonis⁴, K. De Bruyn⁴⁰, S. De Capua⁵³, M. De Cian³⁹, J.M. De Miranda¹, L. De Paula², W. De Silva⁵⁶, P. De Simone¹⁸, D. Decamp⁴, M. Deckenhoff⁹, L. Del Buono⁸, N. Déléage⁴, D. Derkach¹⁴, O. Deschamps⁵, F. Dettori⁴¹, A. Di Canto¹¹, H. Dijkstra³⁷, M. Dogaru²⁸, S. Donleavy⁵¹, F. Dordei¹¹, A. Dosil Suárez³⁶, D. Dossett⁴⁷, A. Dovbnya⁴², F. Dupertuis³⁸, R. Dzhelyadin³⁴, A. Dziurda²⁵, A. Dzyuba²⁹, S. Easo^{48,37}, U. Egede⁵², V. Egorychev³⁰, S. Eidelman³³, D. van Eijk⁴⁰, S. Eisenhardt⁴⁹, U. Eitschberger⁹, R. Ekelhof⁹, L. Eklund^{50,37}, I. El Rifai⁵, Ch. Elsasser³⁹, D. Elsby⁴⁴, A. Falabella^{14,e}, C. Färber¹¹, G. Fardell⁴⁹, C. Farinelli⁴⁰, S. Farry⁵¹, V. Fave³⁸, D. Ferguson⁴⁹, V. Fernandez Albor³⁶, F. Ferreira Rodrigues¹, M. Ferro-Luzzi³⁷, S. Filippov³², M. Fiore¹⁶, C. Fitzpatrick³⁷, M. Fontana¹⁰, F. Fontanelli^{19,i}, R. Forty³⁷, O. Francisco², M. Frank³⁷, C. Frei³⁷, M. Frosini^{17,f}, S. Furcas²⁰, E. Furfaro^{23,k}, A. Gallas Torreira³⁶, D. Galli^{14,c}, M. Gandelman², P. Gandini⁵⁸, Y. Gao³, J. Garofoli⁵⁸, P. Garosi⁵³, J. Garra Tico⁴⁶, L. Garrido³⁵, C. Gaspar³⁷, R. Gauld⁵⁴, E. Gersabeck¹¹, M. Gersabeck⁵³, T. Gershon^{47,37}, Ph. Ghez⁴, V. Gibson⁴⁶, V.V. Gligorov³⁷, C. Göbel⁵⁹, D. Golubkov³⁰, A. Golutvin^{52,30,37}, A. Gomes², H. Gordon⁵⁴, M. Grabalosa Gándara⁵, R. Graciani Diaz³⁵, L.A. Granado Cardoso³⁷, E. Graugés³⁵, G. Graziani¹⁷, A. Grecu²⁸, E. Greening⁵⁴, S. Gregson⁴⁶, P. Griffith⁴⁴, O. Grünberg⁶⁰, B. Gui⁵⁸, E. Gushchin³², Yu. Guz^{34,37}, T. Gys³⁷, C. Hadjivasiliou⁵⁸, G. Haefeli³⁸, C. Haen³⁷, S.C. Haines⁴⁶, S. Hall⁵², T. Hampson⁴⁵, S. Hansmann-Menzemer¹¹, N. Harnew⁵⁴, S.T. Harnew⁴⁵, J. Harrison⁵³, T. Hartmann⁶⁰, J. He³⁷, V. Heijne⁴⁰, K. Hennessy⁵¹, P. Henrard⁵, J.A. Hernando Morata³⁶, E. van Herwijnen³⁷, A. Hicheur¹, E. Hicks⁵¹, D. Hill⁵⁴, M. Hoballah⁵, C. Hombach⁵³, P. Hopchev⁴, W. Hulsbergen⁴⁰, P. Hunt⁵⁴, T. Huse⁵¹, N. Hussain⁵⁴, D. Hutchcroft⁵¹, D. Hynds⁵⁰, V. Iakovenko⁴³, M. Idzik²⁶, P. Ilten¹², R. Jacobsson³⁷, A. Jaeger¹¹, E. Jans⁴⁰, P. Jaton³⁸, A. Jawahery⁵⁷, F. Jing³,

M. John⁵⁴, D. Johnson⁵⁴, C.R. Jones⁴⁶, C. Joram³⁷, B. Jost³⁷, M. Kaballo⁹, S. Kandybei⁴²,
M. Karacson³⁷, T.M. Karbach³⁷, I.R. Kenyon⁴⁴, U. Kerzel³⁷, T. Ketel⁴¹, A. Keune³⁸,
B. Khanji²⁰, O. Kochebina⁷, I. Komarov³⁸, R.F. Koopman⁴¹, P. Koppenburg⁴⁰, M. Korolev³¹,
A. Kozlinskiy⁴⁰, L. Kravchuk³², K. Kreplin¹¹, M. Krepis⁴⁷, G. Krocker¹¹, P. Krokovny³³,
F. Kruse⁹, M. Kucharczyk^{20,25,j}, V. Kudryavtsev³³, T. Kvaratskheliya^{30,37}, V.N. La Thi³⁸,
D. Lacarrere³⁷, G. Lafferty⁵³, A. Lai¹⁵, D. Lambert⁴⁹, R.W. Lambert⁴¹, E. Lanciotti³⁷,
G. Lanfranchi^{18,37}, C. Langenbruch³⁷, T. Latham⁴⁷, C. Lazzeroni⁴⁴, R. Le Gac⁶,
J. van Leerdam⁴⁰, J.-P. Lees⁴, R. Lefèvre⁵, A. Leflat³¹, J. Lefrançois⁷, S. Leo²², O. Leroy⁶,
T. Lesiak²⁵, B. Leverington¹¹, Y. Li³, L. Li Gioi⁵, M. Liles⁵¹, R. Lindner³⁷, C. Linn¹¹, B. Liu³,
G. Liu³⁷, S. Lohn³⁷, I. Longstaff⁵⁰, J.H. Lopes², E. Lopez Asamar³⁵, N. Lopez-March³⁸, H. Lu³,
D. Lucchesi^{21,p}, J. Luisier³⁸, H. Luo⁴⁹, F. Machefert⁷, I.V. Machikhiliyan^{4,30}, F. Maciuc²⁸,
O. Maev^{29,37}, S. Malde⁵⁴, G. Manca^{15,d}, G. Mancinelli⁶, U. Marconi¹⁴, R. Märki³⁸, J. Marks¹¹,
G. Martellotti²⁴, A. Martens⁸, A. Martín Sánchez⁷, M. Martinelli⁴⁰, D. Martinez Santos⁴¹,
D. Martins Tostes², A. Massafferri¹, R. Matev³⁷, Z. Mathe³⁷, C. Matteuzzi²⁰, E. Maurice⁶,
A. Mazurov^{16,32,37,e}, B. Mc Skelly⁵¹, J. McCarthy⁴⁴, A. McNab⁵³, R. McNulty¹²,
B. Meadows^{56,54}, F. Meier⁹, M. Meissner¹¹, M. Merk⁴⁰, D.A. Milanes⁸, M.-N. Minard⁴,
J. Molina Rodriguez⁵⁹, S. Monteil⁵, D. Moran⁵³, P. Morawski²⁵, M.J. Morello^{22,r},
R. Mountain⁵⁸, I. Mous⁴⁰, F. Muheim⁴⁹, K. Müller³⁹, R. Muresan²⁸, B. Muryn²⁶, B. Muster³⁸,
P. Naik⁴⁵, T. Nakada³⁸, R. Nandakumar⁴⁸, I. Nasteva¹, M. Needham⁴⁹, N. Neufeld³⁷,
A.D. Nguyen³⁸, T.D. Nguyen³⁸, C. Nguyen-Mau^{38,o}, M. Nicol⁷, V. Niess⁵, R. Niet⁹, N. Nikitin³¹,
T. Nikodem¹¹, A. Nomerotski⁵⁴, A. Novoselov³⁴, A. Oblakowska-Mucha²⁶, V. Obraztsov³⁴,
S. Oggero⁴⁰, S. Ogilvy⁵⁰, O. Okhrimenko⁴³, R. Oldeman^{15,d}, M. Orlandea²⁸,
J.M. Otalora Goicochea², P. Owen⁵², A. Oyanguren³⁵, B.K. Pal⁵⁸, A. Palano^{13,b}, M. Palutan¹⁸,
J. Panman³⁷, A. Papanestis⁴⁸, M. Pappagallo⁵⁰, C. Parkes⁵³, C.J. Parkinson⁵², G. Passaleva¹⁷,
G.D. Patel⁵¹, M. Patel⁵², G.N. Patrick⁴⁸, C. Patrignani^{19,i}, C. Pavel-Nicorescu²⁸,
A. Pazos Alvarez³⁶, A. Pellegrino⁴⁰, G. Penso^{24,l}, M. Pepe Altarelli³⁷, S. Perazzini^{14,c},
D.L. Perego^{20,j}, E. Perez Trigo³⁶, A. Pérez-Calero Yzquierdo³⁵, P. Perret⁵, M. Perrin-Terrin⁶,
G. Pessina²⁰, K. Petridis⁵², A. Petrolini^{19,i}, A. Phan⁵⁸, E. Picatoste Olloqui³⁵, B. Pietrzyk⁴,
T. Pilar⁴⁷, D. Pinci²⁴, S. Playfer⁴⁹, M. Plo Casasus³⁶, F. Polci⁸, G. Polok²⁵, A. Poluektov^{47,33},
E. Polcarpo², A. Popov³⁴, D. Popov¹⁰, B. Popovici²⁸, C. Potterat³⁵, A. Powell⁵⁴,
J. Prisciandaro³⁸, A. Pritchard⁵¹, C. Prouve⁷, V. Pugatch⁴³, A. Puig Navarro³⁸, G. Punzi^{22,q},
W. Qian⁴, J.H. Rademacker⁴⁵, B. Rakotomiaramanana³⁸, M.S. Rangel², I. Raniuk⁴²,
N. Rauschmayr³⁷, G. Raven⁴¹, S. Redford⁵⁴, M.M. Reid⁴⁷, A.C. dos Reis¹, S. Ricciardi⁴⁸,
A. Richards⁵², K. Rinnert⁵¹, V. Rives Molina³⁵, D.A. Roa Romero⁵, P. Robbe⁷, E. Rodrigues⁵³,
P. Rodriguez Perez³⁶, S. Roiser³⁷, V. Romanovsky³⁴, A. Romero Vidal³⁶, J. Rouvinet³⁸,
T. Ruf³⁷, F. Ruffini²², H. Ruiz³⁵, P. Ruiz Valls³⁵, G. Sabatino^{24,k}, J.J. Saborido Silva³⁶,
N. Sagidova²⁹, P. Sail⁵⁰, B. Saitta^{15,d}, V. Salustino Guimaraes², C. Salzmann³⁹,
B. Sanmartin Sedes³⁶, M. Sannino^{19,i}, R. Santacesaria²⁴, C. Santamarina Rios³⁶,
E. Santovetti^{23,k}, M. Sapunov⁶, A. Sarti^{18,l}, C. Satriano^{24,m}, A. Satta²³, M. Savrie^{16,e},
D. Savrina^{30,31}, P. Schaack⁵², M. Schiller⁴¹, H. Schindler³⁷, M. Schlupp⁹, M. Schmelling¹⁰,
B. Schmidt³⁷, O. Schneider³⁸, A. Schopper³⁷, M.-H. Schune⁷, R. Schwemmer³⁷, B. Sciascia¹⁸,
A. Sciubba²⁴, M. Seco³⁶, A. Semennikov³⁰, K. Senderowska²⁶, I. Sepp⁵², N. Serra³⁹, J. Serrano⁶,
P. Seyfert¹¹, M. Shapkin³⁴, I. Shapoval^{16,42}, P. Shatalov³⁰, Y. Shcheglov²⁹, T. Shears^{51,37},
L. Shekhtman³³, O. Shevchenko⁴², V. Shevchenko³⁰, A. Shires⁵², R. Silva Coutinho⁴⁷,
T. Skwarnicki⁵⁸, N.A. Smith⁵¹, E. Smith^{54,48}, M. Smith⁵³, M.D. Sokoloff⁵⁶, F.J.P. Soler⁵⁰,
F. Soomro¹⁸, D. Souza⁴⁵, B. Souza De Paula², B. Spaan⁹, A. Sparkes⁴⁹, P. Spradlin⁵⁰,

F. Stagni³⁷, S. Stahl¹¹, O. Steinkamp³⁹, S. Stoica²⁸, S. Stone⁵⁸, B. Storaci³⁹, M. Straticiu²⁸, U. Straumann³⁹, V.K. Subbiah³⁷, L. Sun⁵⁶, S. Swientek⁹, V. Syropoulos⁴¹, M. Szczekowski²⁷, P. Szczypka^{38,37}, T. Szumlak²⁶, S. T'Jampens⁴, M. Teklishyn⁷, E. Teodorescu²⁸, F. Teubert³⁷, C. Thomas⁵⁴, E. Thomas³⁷, J. van Tilburg¹¹, V. Tisserand⁴, M. Tobin³⁸, S. Tolk⁴¹, D. Tonelli³⁷, S. Topp-Joergensen⁵⁴, N. Torr⁵⁴, E. Tournefier^{4,52}, S. Tourneur³⁸, M.T. Tran³⁸, M. Tresch³⁹, A. Tsaregorodtsev⁶, P. Tsopelas⁴⁰, N. Tuning⁴⁰, M. Ubeda Garcia³⁷, A. Ukleja²⁷, D. Urner⁵³, U. Uwer¹¹, V. Vagnoni¹⁴, G. Valenti¹⁴, R. Vazquez Gomez³⁵, P. Vazquez Regueiro³⁶, S. Vecchi¹⁶, J.J. Velthuis⁴⁵, M. Veltri^{17,g}, G. Veneziano³⁸, M. Vesterinen³⁷, B. Viaud⁷, D. Vieira², X. Vilasis-Cardona^{35,n}, A. Vollhardt³⁹, D. Volyanskyy¹⁰, D. Voong⁴⁵, A. Vorobyev²⁹, V. Vorobyev³³, C. Voß⁶⁰, H. Voss¹⁰, R. Waldi⁶⁰, R. Wallace¹², S. Wandernoth¹¹, J. Wang⁵⁸, D.R. Ward⁴⁶, N.K. Watson⁴⁴, A.D. Webber⁵³, D. Websdale⁵², M. Whitehead⁴⁷, J. Wicht³⁷, J. Wiechczynski²⁵, D. Wiedner¹¹, L. Wiggers⁴⁰, G. Wilkinson⁵⁴, M.P. Williams^{47,48}, M. Williams⁵⁵, F.F. Wilson⁴⁸, J. Wishahi⁹, M. Witek²⁵, S.A. Wotton⁴⁶, S. Wright⁴⁶, S. Wu³, K. Wyllie³⁷, Y. Xie^{49,37}, Z. Xing⁵⁸, Z. Yang³, R. Young⁴⁹, X. Yuan³, O. Yushchenko³⁴, M. Zangoli¹⁴, M. Zavertyaev^{10,a}, F. Zhang³, L. Zhang⁵⁸, W.C. Zhang¹², Y. Zhang³, A. Zhelezov¹¹, A. Zhokhov³⁰, L. Zhong³, A. Zvyagin³⁷.

¹Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

²Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

³Center for High Energy Physics, Tsinghua University, Beijing, China

⁴LAPP, Université de Savoie, CNRS/IN2P3, Annecy-Le-Vieux, France

⁵Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, Clermont-Ferrand, France

⁶CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France

⁷LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France

⁸LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris, France

⁹Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany

¹⁰Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany

¹¹Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

¹²School of Physics, University College Dublin, Dublin, Ireland

¹³Sezione INFN di Bari, Bari, Italy

¹⁴Sezione INFN di Bologna, Bologna, Italy

¹⁵Sezione INFN di Cagliari, Cagliari, Italy

¹⁶Sezione INFN di Ferrara, Ferrara, Italy

¹⁷Sezione INFN di Firenze, Firenze, Italy

¹⁸Laboratori Nazionali dell'INFN di Frascati, Frascati, Italy

¹⁹Sezione INFN di Genova, Genova, Italy

²⁰Sezione INFN di Milano Bicocca, Milano, Italy

²¹Sezione INFN di Padova, Padova, Italy

²²Sezione INFN di Pisa, Pisa, Italy

²³Sezione INFN di Roma Tor Vergata, Roma, Italy

²⁴Sezione INFN di Roma La Sapienza, Roma, Italy

²⁵Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland

²⁶AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland

²⁷National Center for Nuclear Research (NCBJ), Warsaw, Poland

²⁸Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania

²⁹Petersburg Nuclear Physics Institute (PNPI), Gatchina, Russia

³⁰Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia

³¹Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia

³²Institute for Nuclear Research of the Russian Academy of Sciences (INR RAN), Moscow, Russia

- ³³ *Budker Institute of Nuclear Physics (SB RAS) and Novosibirsk State University, Novosibirsk, Russia*
- ³⁴ *Institute for High Energy Physics (IHEP), Protvino, Russia*
- ³⁵ *Universitat de Barcelona, Barcelona, Spain*
- ³⁶ *Universidad de Santiago de Compostela, Santiago de Compostela, Spain*
- ³⁷ *European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- ³⁸ *Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- ³⁹ *Physik-Institut, Universität Zürich, Zürich, Switzerland*
- ⁴⁰ *Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands*
- ⁴¹ *Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, The Netherlands*
- ⁴² *NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- ⁴³ *Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- ⁴⁴ *University of Birmingham, Birmingham, United Kingdom*
- ⁴⁵ *H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- ⁴⁶ *Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- ⁴⁷ *Department of Physics, University of Warwick, Coventry, United Kingdom*
- ⁴⁸ *STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- ⁴⁹ *School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- ⁵⁰ *School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- ⁵¹ *Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- ⁵² *Imperial College London, London, United Kingdom*
- ⁵³ *School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- ⁵⁴ *Department of Physics, University of Oxford, Oxford, United Kingdom*
- ⁵⁵ *Massachusetts Institute of Technology, Cambridge, MA, United States*
- ⁵⁶ *University of Cincinnati, Cincinnati, OH, United States*
- ⁵⁷ *University of Maryland, College Park, MD, United States*
- ⁵⁸ *Syracuse University, Syracuse, NY, United States*
- ⁵⁹ *Pontificia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to ²*
- ⁶⁰ *Institut für Physik, Universität Rostock, Rostock, Germany, associated to ¹¹*

^a *P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia*

^b *Università di Bari, Bari, Italy*

^c *Università di Bologna, Bologna, Italy*

^d *Università di Cagliari, Cagliari, Italy*

^e *Università di Ferrara, Ferrara, Italy*

^f *Università di Firenze, Firenze, Italy*

^g *Università di Urbino, Urbino, Italy*

^h *Università di Modena e Reggio Emilia, Modena, Italy*

ⁱ *Università di Genova, Genova, Italy*

^j *Università di Milano Bicocca, Milano, Italy*

^k *Università di Roma Tor Vergata, Roma, Italy*

^l *Università di Roma La Sapienza, Roma, Italy*

^m *Università della Basilicata, Potenza, Italy*

ⁿ *LIFAEELS, La Salle, Universitat Ramon Llull, Barcelona, Spain*

^o *Hanoi University of Science, Hanoi, Viet Nam*

^p *Università di Padova, Padova, Italy*

^q *Università di Pisa, Pisa, Italy*

^r *Scuola Normale Superiore, Pisa, Italy*

The B_c^+ meson is composed of two heavy valence quarks, and has a wide range of expected decay modes [1–10]. Prior to LHCb taking data, only a few decay channels, such as $B_c^+ \rightarrow J/\psi \pi^+$ and $B_c^+ \rightarrow J/\psi \mu^+ \nu$ had been observed [11, 12]. For pp collisions at a centre-of-mass energy of 7 TeV, the total B_c^+ production cross-section is predicted to be about $0.4 \mu\text{b}$, one order of magnitude higher than that at the Tevatron [13, 14]. LHCb has thus been able to observe new decay modes, such as $B_c^+ \rightarrow J/\psi \pi^+ \pi^- \pi^+$ [15], $B_c^+ \rightarrow \psi(2S) \pi^+$ [16] and $B_c^+ \rightarrow J/\psi D_s^{(*)+}$ [17], and to measure precisely the mass of the B_c^+ meson [18].

In this paper, we report the first observation of the decay channel $B_c^+ \rightarrow J/\psi K^+$ (inclusion of charge conjugate modes is implied throughout the paper). The J/ψ meson is reconstructed in the dimuon final state. The branching fraction is measured relative to that of the $B_c^+ \rightarrow J/\psi \pi^+$ decay mode, which has identical topology and similar kinematic properties, as shown in Fig. 1. No absolute branching fraction of the B_c^+ meson is known to date. The predicted ratio of branching fractions $\mathcal{B}(B_c^+ \rightarrow J/\psi K^+)/\mathcal{B}(B_c^+ \rightarrow J/\psi \pi^+)$ is dominated by the ratio of the relevant Cabibbo-Kobayashi-Maskawa (CKM) matrix elements $|V_{ud}/V_{us}|^2 \approx 0.05$ [19]. However, after including the decay constants, $f_{K^+(\pi^+)}$, the ratio is enhanced,

$$\frac{\mathcal{B}(B_c^+ \rightarrow J/\psi K^+)}{\mathcal{B}(B_c^+ \rightarrow J/\psi \pi^+)} \approx \left| \frac{V_{us} f_{K^+}}{V_{ud} f_{\pi^+}} \right|^2 = 0.077, \quad (1)$$

where the values of $f_{K^+(\pi^+)}$ are given in Ref. [19]. Taking into account the contributions of the B_c^+ form factor and the kinematics, the theoretical predictions for the ratio of branching fractions lie in the range from 0.054 to 0.088 [2, 3, 5–7, 9, 10]. The large span of these predictions is due to the various models and the uncertainties on the phenomenological parameters. The measurement of $\mathcal{B}(B_c^+ \rightarrow J/\psi K^+)/\mathcal{B}(B_c^+ \rightarrow J/\psi \pi^+)$ therefore provides a test of the theoretical predictions of hadronisation.

The analysis is based on a data sample, corresponding to an integrated luminosity of 1.0 fb^{-1} of pp collisions, collected by the LHCb experiment at a centre-of-mass energy of 7 TeV. The LHCb detector [20] is a single-arm, forward spectrometer covering the pseudorapidity range $2 < \eta < 5$ and is designed for precise measurements in the b and c quark sectors. The detector includes a high precision tracking system consisting of a




Figure 1: Diagram for a $B_c^+ \rightarrow J/\psi \pi^+(K^+)$ decay.

silicon-strip vertex detector surrounding the pp interaction region, a large area silicon-strip detector located upstream of a dipole magnet with a bending power of about 4 Tm, and three stations of silicon-strip detectors and straw drift tubes placed downstream. The combined tracking system has momentum resolution $\Delta p/p$ that varies from 0.4% at 5 GeV/ c to 0.6% at 100 GeV/ c , and impact parameter (IP) resolution of 20 μm for tracks with high transverse momentum (p_{T}). Charged hadrons are identified using two ring-imaging Cherenkov (RICH) detectors and good kaon-pion separation is achieved for tracks with momentum between 5 GeV/ c and 100 GeV/ c [21]. Photon, electron and hadron candidates are identified by a calorimeter system consisting of scintillating-pad and preshower detectors, an electromagnetic calorimeter and a hadronic calorimeter. Muons are identified by a system composed of alternating layers of iron and multiwire proportional chambers. The trigger system [22] consists of a hardware stage, based on information from the calorimeter and muon systems, followed by a two-stage software trigger that applies event reconstruction and reduces the event rate from 1 MHz to around 3 kHz.

In the hardware trigger, events are selected by requiring a single muon or dimuon candidate with high p_{T} . In the software trigger, events are selected by requiring dimuon candidates with invariant mass close to the known J/ψ mass [19] and with decay length significance greater than 3 with respect to the associated primary vertex (PV). For events with several PVs, the one with the smallest χ_{IP}^2 is chosen, where χ_{IP}^2 is defined as the difference in χ^2 of a given PV reconstructed with and without the considered track. The bachelor hadrons (K^+ for $B_c^+ \rightarrow J/\psi K^+$ and π^+ for $B_c^+ \rightarrow J/\psi \pi^+$ decays) are required to be separated from the B_c^+ PV and have $p_{\text{T}} > 0.5$ GeV/ c . The B_c^+ candidates are required to have good vertex quality with vertex fit χ_{vtx}^2 per degree of freedom less than 5, and mass within 500 MeV/ c^2 of the world average value of the B_c^+ mass [19].

A boosted decision tree (BDT) [23] is used for the final event selection. The BDT is trained using a simulated $B_c^+ \rightarrow J/\psi \pi^+$ sample as a proxy for signal and the high-mass sideband ($m_{J/\psi \pi^+} > 6650$ MeV/ c^2) in data for background. The BDT cut value is optimised to maximise the expected $B_c^+ \rightarrow J/\psi K^+$ signal significance. In the simulation, pp collisions are generated using PYTHIA 6.4 [24] with a specific LHCb configuration [25]. The B_c^+ meson production is simulated with the dedicated generator BCVEGPY [26]. Decays of hadronic particles are described by EVTGEN [27], in which final state radiation is generated using PHOTOS [28]. The interaction of the generated particles with the detector and its response are implemented using the GEANT4 toolkit [29] as described in Ref. [30]. The BDT takes the following variables into account: the χ_{IP}^2 of the bachelor hadron and B_c^+ mesons with respect to the PV; the B_c^+ vertex quality; the distance between the B_c^+ decay vertex and the PV; the p_{T} of the B_c^+ candidate; the χ^2 from the B_c^+ decay vertex refit [31], obtained with a constraint on the PV and the reconstructed J/ψ mass; and the cosine of the angle between the momentum of the B_c^+ meson and the direction vector from the PV to the B_c^+ decay vertex. These variables are chosen as they discriminate the signal from the background, and have similar distributions for $B_c^+ \rightarrow J/\psi K^+$ and $B_c^+ \rightarrow J/\psi \pi^+$ decays, ensuring that the systematic uncertainty due to the relative selection efficiency is minimal. After the BDT selection, no event with multiple candidates remains.

The branching fraction ratio is computed as

$$\frac{\mathcal{B}(B_c^+ \rightarrow J/\psi K^+)}{\mathcal{B}(B_c^+ \rightarrow J/\psi \pi^+)} = \frac{N(B_c^+ \rightarrow J/\psi K^+)}{N(B_c^+ \rightarrow J/\psi \pi^+)} \cdot \frac{\epsilon(B_c^+ \rightarrow J/\psi \pi^+)}{\epsilon(B_c^+ \rightarrow J/\psi K^+)}, \quad (2)$$

where N is the signal yield of $B_c^+ \rightarrow J/\psi K^+$ or $B_c^+ \rightarrow J/\psi \pi^+$ decays and ϵ is the total efficiency, which takes into account the geometrical acceptance, detection, reconstruction, selection and trigger effects.

An unbinned maximum likelihood fit is used to determine the yields from the $J/\psi K^+$ mass distribution of the B_c^+ candidates, under the kaon mass hypothesis. The total probability density function for the fit has four components: signals for $B_c^+ \rightarrow J/\psi K^+$ and $B_c^+ \rightarrow J/\psi \pi^+$ decays; the combinatorial background; and the partially reconstructed background.

To discriminate between pion and kaon bachelor tracks, the quantity

$$\text{DLL}_{K\pi} = \ln \mathcal{L}(K) - \ln \mathcal{L}(\pi) \quad (3)$$

is used, where $\mathcal{L}(K)$ and $\mathcal{L}(\pi)$ are the likelihood values provided by the RICH system under the kaon and pion hypotheses, respectively. Since the momentum spectra of the bachelor pions and kaons are correlated with the $\text{DLL}_{K\pi}$, the shapes of the mass distribution used in the fit vary as a function of $\text{DLL}_{K\pi}$. To reduce this dependence and separate the two signals, the $\text{DLL}_{K\pi}$ range is divided into four bins, $\text{DLL}_{K\pi} < -5$, $-5 < \text{DLL}_{K\pi} < 0$, $0 < \text{DLL}_{K\pi} < 5$ and $\text{DLL}_{K\pi} > 5$. The ratio of the total signal yields is defined as $\mathcal{R}_{K^+/\pi^+} = \sum_{i=1}^4 N_{J/\psi K^+}^i / \sum_{i=1}^4 N_{J/\psi \pi^+}^i$, where $N_{J/\psi K^+(\pi^+)}^i$ is the signal yield in each $\text{DLL}_{K\pi}$ bin i . Due to the limited sample size of the $B_c^+ \rightarrow J/\psi K^+$ signal in the bins with $\text{DLL}_{K\pi} < -5$ and $-5 < \text{DLL}_{K\pi} < 0$, their signal yields are fixed, respectively, to be zero and $P \times \sum_{i=1}^4 N_{J/\psi K^+}^i$ where the P is the probability that the kaon from the $B_c^+ \rightarrow J/\psi K^+$ decay has $-5 < \text{DLL}_{K\pi} < 0$, as estimated from simulation.

Figure 2 shows the invariant mass distributions of the B_c^+ candidates, calculated with the kaon mass hypothesis in the four $\text{DLL}_{K\pi}$ bins. In the fit to the B_c^+ mass spectrum, the shape of the $B_c^+ \rightarrow J/\psi K^+$ signal is modelled by a double-sided Crystal Ball (DSCB) function [32] as

$$f(x; M, \sigma, a_l, n_l, a_r, n_r) = \begin{cases} e^{-\frac{a_l^2}{2}} \left(\frac{n_l}{a_l}\right)^{n_l} \left(\frac{n_l}{a_l} - a_l - \frac{x-M}{\sigma}\right)^{-n_l} & \frac{x-M}{\sigma} < -a_l \\ \exp\left[-\frac{1}{2}\left(\frac{x-M}{\sigma}\right)^2\right] & -a_l \leq \frac{x-M}{\sigma} \leq -a_r \\ e^{-\frac{a_r^2}{2}} \left(\frac{n_r}{a_r}\right)^{n_r} \left(\frac{n_r}{a_r} - a_r + \frac{x-M}{\sigma}\right)^{-n_r} & \frac{x-M}{\sigma} > -a_r \end{cases} \quad (4)$$

where the peak position is fixed to that from an independent fit to the $B_c^+ \rightarrow J/\psi \pi^+$ mass distribution, and the tail parameters $a_{l,r}$ and $n_{l,r}$ on both sides are taken from simulation.

As the decay $B_c^+ \rightarrow J/\psi \pi^+$ is reconstructed with the kaon mass replacing the pion mass, the signal is shifted to higher mass values and is modelled by another DSCB function whose




Figure 2: Mass distributions of B_c^+ candidates in four $DLL_{K\pi}$ bins and the superimposed fit results. The solid shaded area (red) represents the $B_c^+ \rightarrow J/\psi K^+$ signal and the hatched area (blue) the $B_c^+ \rightarrow J/\psi \pi^+$ signal. The dot-dashed line (blue) indicates the partially reconstructed background and the dotted (red) the combinatorial background. The solid line (black) represents the sum of the above components and the points with error bars (black) show the data. The labels (a), (b), (c) and (d) correspond to $DLL_{K\pi} < -5$, $-5 < DLL_{K\pi} < 0$, $0 < DLL_{K\pi} < 5$ and $DLL_{K\pi} > 5$ for the bachelor track, respectively.

shape and the relative position to the $B_c^+ \rightarrow J/\psi K^+$ signal are also derived from simulation. Two corrections are applied to the $B_c^+ \rightarrow J/\psi \pi^+$ simulation sample. Firstly, since the resolution of the detector is overestimated, the momenta of charged particles are smeared to make the resolution on the B_c^+ mass in the $B_c^+ \rightarrow J/\psi \pi^+$ simulation sample the same as that of the $J/\psi \pi^+$ mass distribution of the B_c^+ candidates in the data sample. Secondly, the shapes of the $B_c^+ \rightarrow J/\psi \pi^+$ mass distribution in the four $DLL_{K\pi}$ bins depend on the $DLL_{K\pi}$ distribution, which is different in data and simulation. To reduce the effect of this difference, each simulated event is reweighted by a $DLL_{K\pi}$ dependent correction factor, which is derived from a linear fit to the ratio of the $DLL_{K\pi}$ distribution in background-

subtracted data, to that of the simulation sample. The background subtraction [33] is performed with the $J/\psi \pi^+$ mass distribution of the B_c^+ candidates in the data sample with the pion mass hypothesis.

The combinatorial background is modelled as an exponential function with a different freely varying parameter in each $\text{DLL}_{K\pi}$ bin. The contribution of the partially reconstructed background is modelled by an ARGUS function [34]. The contribution of the partially reconstructed background is dominated by events with bachelor pions, which are suppressed in the high-value $\text{DLL}_{K\pi}$ bins, therefore the number of the partially reconstructed events in the $\text{DLL}_{K\pi} > 5$ bin is assumed to be zero. All parameters of the partially reconstructed background are allowed to vary. The observed $B_c^+ \rightarrow J/\psi K^+$ signal yield is 46 ± 12 and the ratio of yields is

$$\mathcal{R}_{K^+/\pi^+} = \frac{N(B_c^+ \rightarrow J/\psi K^+)}{N(B_c^+ \rightarrow J/\psi \pi^+)} = 0.071 \pm 0.020 \text{ (stat)}.$$

The ratio of the total efficiencies computed over the full $\text{DLL}_{K\pi}$ range is

$$\frac{\epsilon(B_c^+ \rightarrow J/\psi K^+)}{\epsilon(B_c^+ \rightarrow J/\psi \pi^+)} = 1.029 \pm 0.007,$$

which is determined from simulation and the uncertainty is due to the finite size of the simulation samples.

The $B_c^+ \rightarrow J/\psi \pi^+$ signal has a long tail that may extend into the high mass region. A systematic uncertainty is assigned due to the choice of fit range, and is determined to be 0.9% by changing the mass window from 6000-6600 MeV/ c^2 to 6200-6700 MeV/ c^2 and comparing the results. To estimate the systematic uncertainty due to the potentially different performance of the BDT on data and simulation, several BDT cut values were tested. A 5.7% spread in the final result is obtained and is propagated to the quoted systematic uncertainty.

To estimate the uncertainty due to the shapes of the $B_c^+ \rightarrow J/\psi K^+$ and $B_c^+ \rightarrow J/\psi \pi^+$ signals, the fit is repeated many times by varying the parameters of the tails of these DSCB functions that were kept constant in the fit within one standard deviation of their values in simulation. A spread of 0.7% is observed. For the $B_c^+ \rightarrow J/\psi \pi^+$ signal the assigned systematic uncertainty is 0.5%.

To estimate the systematic uncertainty due to the choice of signal shape, an alternative $B_c^+ \rightarrow J/\psi \pi^+$ mass shape is used, which is determined from the data sample by subtracting the background in the $J/\psi \pi^+$ mass distribution of the B_c^+ candidates with the pion hypothesis. A 2.7% difference with the ratio obtained with the nominal signal shape is observed.

For the systematic uncertainty due to the choice of the partially reconstructed background shape in each $\text{DLL}_{K\pi}$ bin, the shape is modelled with the ARGUS function convolved with a Gaussian function. The observed 2.3% deviation from the default fit is assigned as the systematic uncertainty.

For the $B_c^+ \rightarrow J/\psi K^+$ yields in the two bins with $\text{DLL}_{K\pi} < 0$, half of the probability estimated from the simulation, namely 1.8%, is taken as systematic uncertainty.

Table 1: Relative systematic uncertainties on the ratio of branching fractions.

Source	Uncertainty (%)
Mass window	0.9
BDT selection	5.7
$B_c^+ \rightarrow J/\psi K^+$ signal model	0.7
$B_c^+ \rightarrow J/\psi \pi^+$ signal model	0.5
Choice of signal shape	2.7
Partially reconstructed background shape	2.3
$B_c^+ \rightarrow J/\psi K^+$ signals in $DLL_{K\pi} < 0$ bins	1.8
$DLL_{K\pi}$ binning choice	1.2
K^+ and π^+ interaction length	2.0
Simulation sample size	0.7
Total	7.5

To estimate the uncertainty due to the choice of the $DLL_{K\pi}$ binning, two other binning choices are tried: $DLL_{K\pi} < -6$, $-6 < DLL_{K\pi} < -1$, $-1 < DLL_{K\pi} < 4$, $DLL_{K\pi} > 4$ and $DLL_{K\pi} < -4$, $-4 < DLL_{K\pi} < 1$, $1 < DLL_{K\pi} < 6$, $DLL_{K\pi} > 6$. The average value of the results with these two binning choices has a 1.2% deviation from the default value, which is taken as the systematic uncertainty.

There is a systematic uncertainty due to the different track reconstruction efficiencies for kaons and pions. Since the simulation does not describe hadronic interactions with detector material perfectly, a 2% uncertainty is assumed, as in Ref. [35].

An uncertainty of 0.7% arises from the statistical uncertainty of the ratio of the total efficiencies, which is due to the finite size of the simulation sample.

The systematic uncertainties are summarised in Table 1. The total systematic uncertainty, obtained as the quadratic sum of the individual uncertainties, is 7.5%.

The asymptotic formula for a likelihood-based test $\sqrt{-2 \ln(\mathcal{L}_B/\mathcal{L}_{S+B})}$ is used to estimate the $B_c^+ \rightarrow J/\psi K^+$ signal significance, where \mathcal{L}_B and \mathcal{L}_{S+B} stand for the likelihood of the background-only hypothesis and the signal and background hypothesis respectively. A deviation from the background-only hypothesis with 5.2 standard deviations is found when only the statistical uncertainty is considered. When taking the systematic uncertainty into account, the total significance of the $B_c^+ \rightarrow J/\psi K^+$ signal is 5.0σ .

In summary, a search for the $B_c^+ \rightarrow J/\psi K^+$ decay is performed using a data sample, corresponding to an integrated luminosity of 1.0 fb^{-1} of pp collisions, collected by the LHCb experiment. The signal yield is 46 ± 12 candidates, and represents the first observation of this decay channel. The branching fraction of $B_c^+ \rightarrow J/\psi K^+$ with respect to that of $B_c^+ \rightarrow J/\psi \pi^+$ is measured as

$$\frac{\mathcal{B}(B_c^+ \rightarrow J/\psi K^+)}{\mathcal{B}(B_c^+ \rightarrow J/\psi \pi^+)} = 0.069 \pm 0.019 \pm 0.005,$$

where the first uncertainty is the statistical and the second is systematic. The measurement

is in agreement with the theoretical predictions [2, 3, 5–7, 9, 10].

Assuming factorization holds, the naïve prediction of the ratio $\mathcal{B}(B_c^+ \rightarrow J/\psi K^+)/\mathcal{B}(B_c^+ \rightarrow J/\psi \pi^+)$ can be compared to other B meson decays with a similar topology

$$\frac{\mathcal{B}(B \rightarrow DK^+)}{\mathcal{B}(B \rightarrow D\pi^+)} = \begin{cases} 0.0646 \pm 0.0043 \pm 0.0025 & \text{for } B_s^0 \rightarrow D_s^- K^+(\pi^+) \\ 0.0774 \pm 0.0012 \pm 0.0019 & \text{for } B^+ \rightarrow \bar{D}^0 K^+(\pi^+) \\ 0.074 \pm 0.009 & \text{for } B^0 \rightarrow D^- K^+(\pi^+) \end{cases} \quad (5)$$

taken from Ref. [19,36,37]. Hence, this measurement of $\mathcal{B}(B_c^+ \rightarrow J/\psi K^+)/\mathcal{B}(B_c^+ \rightarrow J/\psi \pi^+)$ is consistent with naïve factorisation in B decays.

Acknowledgements

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); NSFC (China); CNRS/IN2P3 and Region Auvergne (France); BMBF, DFG, HGF and MPG (Germany); SFI (Ireland); INFN (Italy); FOM and NWO (The Netherlands); SCSR (Poland); MEN/IFA (Romania); MinES, Rosatom, RFBR and NRC ‘‘Kurchatov Institute’’ (Russia); MinECo, XuntaGal and GENCAT (Spain); SNSF and SER (Switzerland); NAS Ukraine (Ukraine); STFC (United Kingdom); NSF (USA). We also acknowledge the support received from the ERC under FP7. The Tier1 computing centres are supported by IN2P3 (France), KIT and BMBF (Germany), INFN (Italy), NWO and SURF (The Netherlands), PIC (Spain), GridPP (United Kingdom). We are thankful for the computing resources put at our disposal by Yandex LLC (Russia), as well as to the communities behind the multiple open source software packages that we depend on.

References

- [1] M. A. Ivanov, J. Korner, and O. Pakhomova, *The nonleptonic decays $B_c^+ \rightarrow D_s^+ \bar{D}^0$ and $B_c^+ \rightarrow D_s^+ D^0$ in a relativistic quark model*, Phys. Lett. **B555** (2003) 189, [arXiv:hep-ph/0212291](#).
- [2] M. A. Ivanov, J. G. Korner, and P. Santorelli, *Exclusive semileptonic and nonleptonic decays of the B_c meson*, Phys. Rev. **D73** (2006) 054024, [arXiv:hep-ph/0602050](#).
- [3] I. Gouz *et al.*, *Prospects for the B_c studies at LHCb*, Phys. Atom. Nucl. **67** (2004) 1559, [arXiv:hep-ph/0211432](#).
- [4] V. Kiselev, A. Kovalsky, and A. Likhoded, *B_c decays and lifetime in QCD sum rules*, Nucl. Phys. **B585** (2000) 353, [arXiv:hep-ph/0002127](#).

- [5] S. Naimuddin *et al.*, *Nonleptonic two-body B_c meson decays*, Phys. Rev. **D86** (2012) 094028.
- [6] C.-H. Chang and Y.-Q. Chen, *The decays of B_c meson*, Phys. Rev. **D49** (1994) 3399.
- [7] D. Ebert, R. Faustov, and V. Galkin, *Weak decays of the B_c meson to charmonium and D mesons in the relativistic quark model*, Phys. Rev. **D68** (2003) 094020, [arXiv:hep-ph/0306306](#).
- [8] D. Ebert, R. Faustov, and V. Galkin, *Weak decays of the B_c meson to B_s and B mesons in the relativistic quark model*, Eur. Phys. J. **C32** (2003) 29, [arXiv:hep-ph/0308149](#).
- [9] A. Abd El-Hady, J. Munoz, and J. Vary, *Semileptonic and nonleptonic B_c decays*, Phys. Rev. **D62** (2000) 014019, [arXiv:hep-ph/9909406](#).
- [10] P. Colangelo and F. De Fazio, *Using heavy quark spin symmetry in semileptonic B_c decays*, Phys. Rev. **D61** (2000) 034012, [arXiv:hep-ph/9909423](#).
- [11] CDF collaboration, F. Abe *et al.*, *Observation of the B_c meson in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV*, Phys. Rev. Lett. **81** (1998) 2432, [arXiv:hep-ex/9805034](#).
- [12] D0 collaboration, V. Abazov *et al.*, *Observation of the B_c meson in the exclusive decay $B_c^+ \rightarrow J/\psi \pi^+$* , Phys. Rev. Lett. **101** (2008) 012001, [arXiv:0802.4258](#).
- [13] C.-H. Chang and X.-G. Wu, *Uncertainties in estimating hadronic production of the meson B_c and comparisons between Tevatron and LHC*, Eur. Phys. J. **C38** (2004) 267, [arXiv:hep-ph/0309121](#).
- [14] Y.-N. Gao *et al.*, *Experimental prospects of the B_c studies of the LHCb experiment*, Chin. Phys. Lett. **27** (2010) 061302.
- [15] LHCb collaboration, R. Aaij *et al.*, *First observation of the decay $B_c^+ \rightarrow J/\psi \pi^+ \pi^- \pi^+$* , Phys. Rev. Lett. **108** (2012) 251802, [arXiv:1204.0079](#).
- [16] LHCb collaboration, R. Aaij *et al.*, *Observation of the decay $B_c^+ \rightarrow \psi(2S) \pi^+$* , Phys. Rev. **D87** (2013) 071103, [arXiv:1303.1737](#).
- [17] LHCb collaboration, R. Aaij *et al.*, *Observation of $B_c^+ \rightarrow J/\psi D_s^+$ and $B_c^+ \rightarrow J/\psi D_s^{*+}$ decays*, [arXiv:1304.4530](#), to appear in Phys. Rev. D.
- [18] LHCb collaboration, R. Aaij *et al.*, *Measurements of B_c^+ production and mass with the $B_c^+ \rightarrow J/\psi \pi^+$ decay*, Phys. Rev. Lett. **109** (2012) 232001, [arXiv:1209.5634](#).
- [19] Particle Data Group, J. Beringer *et al.*, *Review of particle physics (RPP)*, Phys. Rev. **D86** (2012) 010001.
- [20] LHCb collaboration, A. A. Alves Jr. *et al.*, *The LHCb detector at the LHC*, JINST **3** (2008) S08005.

- [21] M. Adinolfi *et al.*, *Performance of the LHCb RICH detector at the LHC*, Eur. Phys. J. **C73** (2013) 2431, [arXiv:1211.6759](#).
- [22] R. Aaij *et al.*, *The LHCb trigger and its performance in 2011*, JINST **8** (2013) P04022, [arXiv:1211.3055](#).
- [23] L. Breiman, J. H. Friedman, R. A. Olshen, and C. J. Stone, *Classification and regression trees*, Wadsworth international group, Belmont, California, USA, 1984.
- [24] T. Sjostrand, S. Mrenna, and P. Z. Skands, *PYTHIA 6.4 physics and manual*, JHEP **05** (2006) 026, [arXiv:hep-ph/0603175](#).
- [25] I. Belyaev *et al.*, *Handling of the generation of primary events in GAUSS, the LHCb simulation framework*, Nuclear Science Symposium Conference Record (NSS/MIC) **IEEE** (2010) 1155.
- [26] C.-H. Chang, J.-X. Wang, and X.-G. Wu, *BCVEGPY2.0: An upgrade version of the generator BCVEGPY with the addition of hadronproduction of the P-wave B_c states*, Comput. Phys. Commun. **174** (2006) 241, [arXiv:hep-ph/0504017](#).
- [27] D. Lange, *The EVTGEN particle decay simulation package*, Nucl. Instrum. Meth. **A462** (2001) 152.
- [28] P. Golonka and Z. Was, *PHOTOS Monte Carlo: a precision tool for QED corrections in Z and W decays*, Eur. Phys. J. **C45** (2006) 97, [arXiv:hep-ph/0506026](#).
- [29] GEANT4 collaboration, J. Allison *et al.*, *Geant4 developments and applications*, IEEE Trans. Nucl. Sci. **53** (2006) 270; GEANT4 collaboration, S. Agostinelli *et al.*, *GEANT4: A simulation toolkit*, Nucl. Instrum. Meth. **A506** (2003) 250.
- [30] M. Clemencic *et al.*, *The LHCb simulation application, GAUSS: design, evolution and experience*, J. Phys.: Conf. Ser. **331** (2011) 032023.
- [31] W. D. Hulsbergen, *Decay chain fitting with a Kalman filter*, Nucl. Instrum. Meth. **A552** (2005) 566, [arXiv:physics/0503191](#).
- [32] T. Skwarnicki, *A study of the radiative cascade transitions between the Upsilon-prime and Upsilon resonances*, PhD thesis, Institute of Nuclear Physics, Krakow, 1986, DESY-F31-86-02.
- [33] M. Pivk and F. R. Le Diberder, *sPlot: a statistical tool to unfold data distributions*, Nucl. Instrum. Meth. **A555** (2005) 356, [arXiv:physics/0402083](#).
- [34] ARGUS collaboration, H. Albrecht *et al.*, *Search for hadronic $b \rightarrow u$ decays*, Phys. Lett. **B241** (1990) 278.

- [35] LHCb collaboration, R. Aaij *et al.*, *Measurements of the branching fractions and CP asymmetries of $B^+ \rightarrow J/\psi \pi^+$ and $B^+ \rightarrow \psi(2S)\pi^+$ decays*, Phys. Rev. **D85** (2012) 091105, arXiv:1203.3592.
- [36] LHCb Collaboration, R. Aaij *et al.*, *Measurements of the branching fractions of the decays $B_s^0 \rightarrow D_s^\mp K^\pm$ and $B_s^0 \rightarrow D_s^- \pi^+$* , JHEP **1206** (2012) 115, arXiv:1204.1237.
- [37] LHCb Collaboration, R. Aaij *et al.*, *Observation of CP violation in B^\pm to DK^\pm decays*, Phys. Lett. **B712** (2012) 203, arXiv:1203.3662.