



Search for the lepton-flavour violating decays $B_s^0 \rightarrow e^\pm \mu^\mp$ and $B^0 \rightarrow e^\pm \mu^\mp$

The LHCb collaboration¹

Abstract

A search for the lepton-flavour violating decays $B_s^0 \rightarrow e^\pm \mu^\mp$ and $B^0 \rightarrow e^\pm \mu^\mp$ is performed with a data sample, corresponding to an integrated luminosity of 1.0 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$, collected by the LHCb experiment. The observed number of $B_s^0 \rightarrow e^\pm \mu^\mp$ and $B^0 \rightarrow e^\pm \mu^\mp$ candidates is consistent with background expectations. Upper limits on the branching fractions of both decays are determined to be $\mathcal{B}(B_s^0 \rightarrow e^\pm \mu^\mp) < 1.1 (1.4) \times 10^{-8}$ and $\mathcal{B}(B^0 \rightarrow e^\pm \mu^\mp) < 2.8 (3.7) \times 10^{-9}$ at 90% (95%) confidence level (C.L.). These limits are a factor of twenty lower than those set by previous experiments. Lower bounds on the Pati-Salam leptoquark masses are also calculated, $M_{\text{LQ}}(B_s^0 \rightarrow e^\pm \mu^\mp) > 101 \text{ TeV}/c^2$ and $M_{\text{LQ}}(B^0 \rightarrow e^\pm \mu^\mp) > 126 \text{ TeV}/c^2$ at 95% C.L., and are a factor of two higher than the previous bounds.

Published on Phys. Rev. Lett. 111, 141801 (2013)

© CERN on behalf of the LHCb collaboration, license CC-BY-3.0.

¹Authors are listed on the following pages.

LHCb collaboration

R. Aaij⁴⁰, B. Adeva³⁶, M. Adinolfi⁴⁵, C. Adrover⁶, A. Affolder⁵¹, Z. Ajaltouni⁵, J. Albrecht⁹, F. Alessio³⁷, M. Alexander⁵⁰, S. Ali⁴⁰, G. Alkhazov²⁹, P. Alvarez Cartelle³⁶, A.A. Alves Jr^{24,37}, S. Amato², S. Amerio²¹, Y. Amhis⁷, L. Anderlini^{17,f}, J. Anderson³⁹, R. Andreassen⁵⁶, J.E. Andrews⁵⁷, R.B. Appleby⁵³, O. Aquines Gutierrez¹⁰, F. Archilli¹⁸, A. Artamonov³⁴, M. Artuso⁵⁸, E. Aslanides⁶, G. Auriemma^{24,m}, M. Baalouch⁵, S. Bachmann¹¹, J.J. Back⁴⁷, C. Baesso⁵⁹, V. Balagura³⁰, W. Baldini¹⁶, R.J. Barlow⁵³, C. Barschel³⁷, S. Barsuk⁷, W. Barter⁴⁶, Th. Bauer⁴⁰, A. Bay³⁸, J. Beddow⁵⁰, F. Bedeschi²², I. Bediaga¹, S. Belogurov³⁰, K. Belous³⁴, I. Belyaev³⁰, E. Ben-Haim⁸, G. Bencivenni¹⁸, S. Benson⁴⁹, J. Benton⁴⁵, A. Berezhnoy³¹, R. Bernet³⁹, M.-O. Bettler⁴⁶, M. van Beuzekom⁴⁰, A. Bien¹¹, S. Bifani⁴⁴, T. Bird⁵³, A. Bizzeti^{17,h}, P.M. Bjørnstad⁵³, T. Blake³⁷, F. Blanc³⁸, J. Blouw¹¹, S. Blusk⁵⁸, V. Bocci²⁴, A. Bondar³³, N. Bondar²⁹, W. Bonivento¹⁵, S. Borghi⁵³, A. Borgia⁵⁸, T.J.V. Bowcock⁵¹, E. Bowen³⁹, C. Bozzi¹⁶, T. Brambach⁹, J. van den Brand⁴¹, J. Bressieux³⁸, D. Brett⁵³, M. Britsch¹⁰, T. Britton⁵⁸, N.H. Brook⁴⁵, H. Brown⁵¹, I. Burducea²⁸, A. Bursche³⁹, G. Busetto^{21,q}, J. Buytaert³⁷, S. Cadeddu¹⁵, O. Callot⁷, M. Calvi^{20,j}, M. Calvo Gomez^{35,n}, A. Camboni³⁵, P. Campana^{18,37}, D. Campora Perez³⁷, A. Carbone^{14,c}, G. Carboni^{23,k}, R. Cardinale^{19,i}, A. Cardini¹⁵, H. Carranza-Mejia⁴⁹, L. Carson⁵², K. Carvalho Akiba², G. Casse⁵¹, L. Castillo Garcia³⁷, M. Cattaneo³⁷, Ch. Cauet⁹, R. Cenci⁵⁷, M. Charles⁵⁴, Ph. Charpentier³⁷, P. Chen^{3,38}, N. Chiapolini³⁹, M. Chrzaszcz²⁵, K. Ciba³⁷, X. Cid Vidal³⁷, G. Ciezarek⁵², P.E.L. Clarke⁴⁹, M. Clemencic³⁷, H.V. Cliff⁴⁶, J. Closier³⁷, C. Coca²⁸, V. Coco⁴⁰, J. Cogan⁶, E. Cogneras⁵, P. Collins³⁷, A. Comerma-Montells³⁵, A. Contu^{15,37}, A. Cook⁴⁵, M. Coombes⁴⁵, S. Coquereau⁸, G. Corti³⁷, B. Couturier³⁷, G.A. Cowan⁴⁹, D.C. Craik⁴⁷, S. Cunliffe⁵², R. Currie⁴⁹, C. D'Ambrosio³⁷, P. David⁸, P.N.Y. David⁴⁰, A. Davis⁵⁶, I. De Bonis⁴, K. De Bruyn⁴⁰, S. De Capua⁵³, M. De Cian¹¹, J.M. De Miranda¹, L. De Paula², W. De Silva⁵⁶, P. De Simone¹⁸, D. Decamp⁴, M. Deckenhoff⁹, L. Del Buono⁸, N. Déleage⁴, D. Derkach⁵⁴, O. Deschamps⁵, F. Dettori⁴¹, A. Di Canto¹¹, H. Dijkstra³⁷, M. Dogaru²⁸, S. Donleavy⁵¹, F. Dordei¹¹, A. Dosil Suárez³⁶, D. Dossett⁴⁷, A. Dovbnya⁴², F. Dupertuis³⁸, P. Durante³⁷, R. Dzhelyadin³⁴, A. Dziurda²⁵, A. Dzyuba²⁹, S. Easo^{48,37}, U. Egede⁵², V. Egorychev³⁰, S. Eidelman³³, D. van Eijk⁴⁰, S. Eisenhardt⁴⁹, U. Eitschberger⁹, R. Ekelhof⁹, L. Eklund^{50,37}, I. El Rifai⁵, Ch. Elsasser³⁹, A. Falabella^{14,e}, C. Färber¹¹, G. Fardell⁴⁹, C. Farinelli⁴⁰, S. Farry⁵¹, V. Fave³⁸, D. Ferguson⁴⁹, V. Fernandez Albor³⁶, F. Ferreira Rodrigues¹, M. Ferro-Luzzi³⁷, S. Filippov³², M. Fiore¹⁶, C. Fitzpatrick³⁷, M. Fontana¹⁰, F. Fontanelli^{19,i}, R. Forty³⁷, O. Francisco², M. Frank³⁷, C. Frei³⁷, M. Frosini^{17,f}, S. Furcas²⁰, E. Furfaro^{23,k}, A. Gallas Torreira³⁶, D. Galli^{14,c}, M. Gandelman², P. Gandini⁵⁸, Y. Gao³, J. Garofoli⁵⁸, P. Garosi⁵³, J. Garra Tico⁴⁶, L. Garrido³⁵, C. Gaspar³⁷, R. Gauld⁵⁴, E. Gersabeck¹¹, M. Gersabeck⁵³, T. Gershon^{47,37}, Ph. Ghez⁴, V. Gibson⁴⁶, L. Giubega²⁸, V.V. Gligorov³⁷, C. Göbel⁵⁹, D. Golubkov³⁰, A. Golutvin^{52,30,37}, A. Gomes², H. Gordon⁵⁴, M. Grabalosa Gándara⁵, R. Graciani Diaz³⁵, L.A. Granado Cardoso³⁷, E. Graugés³⁵, G. Graziani¹⁷, A. Grecu²⁸, E. Greening⁵⁴, S. Gregson⁴⁶, P. Griffith⁴⁴, O. Grünberg⁶⁰, B. Gui⁵⁸, E. Gushchin³², Yu. Guz^{34,37}, T. Gys³⁷, C. Hadjivasiliou⁵⁸, G. Haefeli³⁸, C. Haen³⁷, S.C. Haines⁴⁶, S. Hall⁵², B. Hamilton⁵⁷, T. Hampson⁴⁵, S. Hansmann-Menzemer¹¹, N. Harnew⁵⁴, S.T. Harnew⁴⁵, J. Harrison⁵³, T. Hartmann⁶⁰, J. He³⁷, T. Head³⁷, V. Heijne⁴⁰, K. Hennessy⁵¹, P. Henrard⁵, J.A. Hernandez Morata³⁶, E. van Herwijnen³⁷, A. Hicheur¹, E. Hicks⁵¹, D. Hill⁵⁴, M. Hoballah⁵, C. Hombach⁵³, P. Hopchev⁴, W. Hulsbergen⁴⁰, P. Hunt⁵⁴, T. Huse⁵¹, N. Hussain⁵⁴, D. Hutchcroft⁵¹, D. Hynds⁵⁰, V. Iakovenko⁴³, M. Idzik²⁶, P. Ilten¹², R. Jacobsson³⁷, A. Jaeger¹¹, E. Jans⁴⁰, P. Jaton³⁸, A. Jawahery⁵⁷, F. Jing³, M. John⁵⁴, D. Johnson⁵⁴, C.R. Jones⁴⁶, C. Joram³⁷, B. Jost³⁷, M. Kabbalo⁹, S. Kandybei⁴², W. Kanso⁶, M. Karacson³⁷, T.M. Karbach³⁷, I.R. Kenyon⁴⁴, T. Ketel⁴¹, A. Keune³⁸, B. Khanji²⁰, O. Kochebina⁷, I. Komarov³⁸, R.F. Koopman⁴¹, P. Koppenburg⁴⁰, M. Korolev³¹, A. Kozlinskiy⁴⁰, L. Kravchuk³², K. Kreplin¹¹, M. Kreps⁴⁷, G. Krocker¹¹, P. Krokovny³³, F. Kruse⁹, M. Kucharczyk^{20,25,j}, V. Kudryavtsev³³, T. Kvaratskheliya^{30,37}, V.N. La Thi³⁸, D. Lacarrere³⁷, G. Lafferty⁵³, A. Lai¹⁵, D. Lambert⁴⁹, R.W. Lambert⁴¹, E. Lanciotti³⁷, G. Lanfranchi¹⁸, C. Langenbruch³⁷, T. Latham⁴⁷, C. Lazzeroni⁴⁴, R. Le Gac⁶, J. van Leerdam⁴⁰, J.-P. Lees⁴, R. Lefèvre⁵, A. Leflat³¹, J. Lefrançois⁷, S. Leo²², O. Leroy⁶, T. Lesiak²⁵, B. Leverington¹¹, Y. Li³, L. Li Gioi⁵, M. Liles⁵¹, R. Lindner³⁷, C. Linn¹¹, B. Liu³, G. Liu³⁷, S. Lohn³⁷, I. Longstaff⁵⁰, J.H. Lopes², N. Lopez-March³⁸, H. Lu³, D. Lucchesi^{21,q}, J. Luisier³⁸, H. Luo⁴⁹, F. Machefert⁷, I.V. Machikhiliyan^{4,30}, F. Maciuc²⁸, O. Maev^{29,37}, S. Malde⁵⁴, G. Manca^{15,d}, G. Mancinelli⁶, J. Maratas⁵, U. Marconi¹⁴, P. Marino^{22,s}, R. Märki³⁸, J. Marks¹¹, G. Martellotti²⁴, A. Martens⁸, A. Martín Sánchez⁷, M. Martinelli⁴⁰, D. Martinez Santos⁴¹, D. Martins Tostes², A. Massafferri¹, R. Matev³⁷, Z. Mathe³⁷, C. Matteuzzi²⁰, E. Maurice⁶, A. Mazurov^{16,32,37,e}, B. Mc Skelly⁵¹, J. McCarthy⁴⁴, A. McNab⁵³, R. McNulty¹², B. Meadows^{56,54}, F. Meier⁹, M. Meissner¹¹, M. Merk⁴⁰, D.A. Milanes⁸, M.-N. Minard⁴, J. Molina Rodriguez⁵⁹, S. Monteil⁵, D. Moran⁵³, P. Morawski²⁵, A. Mordà⁶, M.J. Morello^{22,s}, R. Mountain⁵⁸, I. Mous⁴⁰, F. Muheim⁴⁹, K. Müller³⁹, R. Muresan²⁸, B. Muryn²⁶, B. Muster³⁸, P. Naik⁴⁵, T. Nakada³⁸, R. Nandakumar⁴⁸, I. Nasteva¹, M. Needham⁴⁹, S. Neubert³⁷, N. Neufeld³⁷, A.D. Nguyen³⁸, T.D. Nguyen³⁸, C. Nguyen-Mau^{38,o}, M. Nicol⁷, V. Niess⁵, R. Niet⁹, N. Nikitin³¹, T. Nikodem¹¹, A. Nomerotski⁵⁴, A. Novoselov³⁴, A. Oblakowska-Mucha²⁶, V. Obraztsov³⁴, S. Oggero⁴⁰, S. Ogilvy⁵⁰, O. Okhrimenko⁴³, R. Oldeman^{15,d}, M. Orlandea²⁸, J.M. Otalora Goicochea², P. Owen⁵², A. Oyunguren³⁵, B.K. Pal⁵⁸, A. Palano^{13,b}, M. Palutan¹⁸, J. Panman³⁷, A. Papanestis⁴⁸, M. Pappagallo⁵⁰, C. Parkes⁵³, C.J. Parkinson⁵², G. Passaleva¹⁷, G.D. Patel⁵¹, M. Patel⁵², G.N. Patrick⁴⁸, C. Patrignani^{19,i}, C. Pavel-Nicorescu²⁸, A. Pazos Alvarez³⁶, A. Pellegrino⁴⁰, G. Penso^{24,l}, M. Pepe Altarelli³⁷,

S. Perazzini^{14,c}, E. Perez Trigo³⁶, A. Pérez-Calero Yzquierdo³⁵, P. Perret⁵, M. Perrin-Terrin⁶, L. Pescatore⁴⁴, G. Pessina²⁰, K. Petridis⁵², A. Petrolini^{19,i}, A. Phan⁵⁸, E. Picatoste Olloqui³⁵, B. Pietrzyk⁴, T. Pilar⁴⁷, D. Pinci²⁴, S. Playfer⁴⁹, M. Plo Casasus³⁶, F. Polci⁸, G. Polok²⁵, A. Poluektov^{47,33}, E. Polycarpo², A. Popov³⁴, D. Popov¹⁰, B. Popovici²⁸, C. Potterat³⁵, A. Powell⁵⁴, J. Prisciandaro³⁸, A. Pritchard⁵¹, C. Prouve⁷, V. Pugatch⁴³, A. Puig Navarro³⁸, G. Punzi^{22,r}, W. Qian⁴, J.H. Rademacker⁴⁵, B. Rakotomiaramanana³⁸, M.S. Rangel², I. Raniuk⁴², N. Rauschmayr³⁷, G. Raven⁴¹, S. Redford⁵⁴, M.M. Reid⁴⁷, A.C. dos Reis¹, S. Ricciardi⁴⁸, A. Richards⁵², K. Rinnert⁵¹, V. Rives Molina³⁵, D.A. Roa Romero⁵, P. Robbe⁷, D.A. Roberts⁵⁷, E. Rodrigues⁵³, P. Rodriguez Perez³⁶, S. Roiser³⁷, V. Romanovsky³⁴, A. Romero Vidal³⁶, J. Rouvinet³⁸, T. Ruf³⁷, F. Ruffini²², H. Ruiz³⁵, P. Ruiz Valls³⁵, G. Sabatino^{24,k}, J.J. Saborido Silva³⁶, N. Sagidova²⁹, P. Sail⁵⁰, B. Saitta^{15,d}, V. Salustino Guimaraes², C. Salzmann³⁹, B. Sanmartin Sedes³⁶, M. Sannino^{19,i}, R. Santacesaria²⁴, C. Santamarina Rios³⁶, E. Santovetti^{23,k}, M. Sapunov⁶, A. Sarti^{18,l}, C. Satriano^{24,m}, A. Satta²³, M. Savrie^{16,e}, D. Savrina^{30,31}, P. Schaack⁵², M. Schiller⁴¹, H. Schindler³⁷, M. Schlupp⁹, M. Schmelling¹⁰, B. Schmidt³⁷, O. Schneider³⁸, A. Schopper³⁷, M.-H. Schune⁷, R. Schwemmer³⁷, B. Sciascia¹⁸, A. Sciubba²⁴, M. Seco³⁶, A. Semennikov³⁰, K. Senderowska²⁶, I. Sepp⁵², N. Serra³⁹, J. Serrano⁶, P. Seyfert¹¹, M. Shapkin³⁴, I. Shapoval^{16,42}, P. Shatalov³⁰, Y. Shcheglov²⁹, T. Shears^{51,37}, L. Shekhtman³³, O. Shevchenko⁴², V. Shevchenko³⁰, A. Shires⁵², R. Silva Coutinho⁴⁷, M. Sirendi⁴⁶, T. Skwarnicki⁵⁸, N.A. Smith⁵¹, E. Smith^{54,48}, J. Smith⁴⁶, M. Smith⁵³, M.D. Sokoloff⁵⁶, F.J.P. Soler⁵⁰, F. Soomro¹⁸, D. Souza⁴⁵, B. Souza De Paula², B. Spaan⁹, A. Sparkes⁴⁹, P. Spradlin⁵⁰, F. Stagni³⁷, S. Stahl¹¹, O. Steinkamp³⁹, S. Stevenson⁵⁴, S. Stoica²⁸, S. Stone⁵⁸, B. Storaci³⁹, M. Straticic²⁸, U. Straumann³⁹, V.K. Subbiah³⁷, L. Sun⁵⁶, S. Swientek⁹, V. Syropoulos⁴¹, M. Szczekowski²⁷, P. Szczypka^{38,37}, T. Szumlak²⁶, S. T'Jampens⁴, M. Teklishyn⁷, E. Teodorescu²⁸, F. Teubert³⁷, C. Thomas⁵⁴, E. Thomas³⁷, J. van Tilburg¹¹, V. Tisserand⁴, M. Tobin³⁸, S. Tolk⁴¹, D. Tonelli³⁷, S. Topp-Joergensen⁵⁴, N. Torr⁵⁴, E. Tournefier^{4,52}, S. Tourneur³⁸, M.T. Tran³⁸, M. Tresch³⁹, A. Tsaregorodtsev⁶, P. Tsopelas⁴⁰, N. Tuning⁴⁰, M. Ubeda Garcia³⁷, A. Ukleja²⁷, D. Urner⁵³, A. Ustyuzhanin^{52,p}, U. Uwer¹¹, V. Vagnoni¹⁴, G. Valenti¹⁴, A. Vallier⁷, M. Van Dijk⁴⁵, R. Vazquez Gomez¹⁸, P. Vazquez Regueiro³⁶, C. Vázquez Sierra³⁶, S. Vecchi¹⁶, J.J. Velthuis⁴⁵, M. Veltri^{17,g}, G. Veneziano³⁸, M. Vesterinen³⁷, B. Viaud⁷, D. Vieira², X. Vilasis-Cardona^{35,n}, A. Vollhardt³⁹, D. Volyansky¹⁰, D. Voong⁴⁵, A. Vorobyev²⁹, V. Vorobyev³³, C. Vob⁶⁰, H. Voss¹⁰, R. Waldi⁶⁰, C. Wallace⁴⁷, R. Wallace¹², S. Wandernoth¹¹, J. Wang⁵⁸, D.R. Ward⁴⁶, N.K. Watson⁴⁴, A.D. Webber⁵³, D. Websdale⁵², M. Whitehead⁴⁷, J. Wicht³⁷, J. Wiechczynski²⁵, D. Wiedner¹¹, L. Wiggers⁴⁰, G. Wilkinson⁵⁴, M.P. Williams^{47,48}, M. Williams⁵⁵, F.F. Wilson⁴⁸, J. Wimberley⁵⁷, J. Wishahli⁹, M. Witek²⁵, S.A. Wotton⁴⁶, S. Wright⁴⁶, S. Wu³, K. Wyllie³⁷, Y. Xie^{49,37}, Z. Xing⁵⁸, Z. Yang³, R. Young⁴⁹, X. Yuan³, O. Yushchenko³⁴, M. Zangoli¹⁴, M. Zavertyaev^{10,a}, F. Zhang³, L. Zhang⁵⁸, W.C. Zhang¹², Y. Zhang³, A. Zhelezov¹¹, A. Zhokhov³⁰, L. Zhong³, A. Zvyagin³⁷.

¹Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

²Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

³Center for High Energy Physics, Tsinghua University, Beijing, China

⁴LAPP, Université de Savoie, CNRS/IN2P3, Annecy-Le-Vieux, France

⁵Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, Clermont-Ferrand, France

⁶CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France

⁷LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France

⁸LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris, France

⁹Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany

¹⁰Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany

¹¹Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

¹²School of Physics, University College Dublin, Dublin, Ireland

¹³Sezione INFN di Bari, Bari, Italy

¹⁴Sezione INFN di Bologna, Bologna, Italy

¹⁵Sezione INFN di Cagliari, Cagliari, Italy

¹⁶Sezione INFN di Ferrara, Ferrara, Italy

¹⁷Sezione INFN di Firenze, Firenze, Italy

¹⁸Laboratori Nazionali dell'INFN di Frascati, Frascati, Italy

¹⁹Sezione INFN di Genova, Genova, Italy

²⁰Sezione INFN di Milano Bicocca, Milano, Italy

²¹Sezione INFN di Padova, Padova, Italy

²²Sezione INFN di Pisa, Pisa, Italy

²³Sezione INFN di Roma Tor Vergata, Roma, Italy

²⁴Sezione INFN di Roma La Sapienza, Roma, Italy

²⁵Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland

²⁶AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland

²⁷National Center for Nuclear Research (NCBJ), Warsaw, Poland

²⁸Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania

²⁹Petersburg Nuclear Physics Institute (PNPI), Gatchina, Russia

³⁰Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia

- ³¹ *Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia*
- ³² *Institute for Nuclear Research of the Russian Academy of Sciences (INR RAN), Moscow, Russia*
- ³³ *Budker Institute of Nuclear Physics (SB RAS) and Novosibirsk State University, Novosibirsk, Russia*
- ³⁴ *Institute for High Energy Physics (IHEP), Protvino, Russia*
- ³⁵ *Universitat de Barcelona, Barcelona, Spain*
- ³⁶ *Universidad de Santiago de Compostela, Santiago de Compostela, Spain*
- ³⁷ *European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- ³⁸ *Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- ³⁹ *Physik-Institut, Universität Zürich, Zürich, Switzerland*
- ⁴⁰ *Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands*
- ⁴¹ *Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, The Netherlands*
- ⁴² *NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- ⁴³ *Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- ⁴⁴ *University of Birmingham, Birmingham, United Kingdom*
- ⁴⁵ *H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- ⁴⁶ *Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- ⁴⁷ *Department of Physics, University of Warwick, Coventry, United Kingdom*
- ⁴⁸ *STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- ⁴⁹ *School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- ⁵⁰ *School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- ⁵¹ *Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- ⁵² *Imperial College London, London, United Kingdom*
- ⁵³ *School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- ⁵⁴ *Department of Physics, University of Oxford, Oxford, United Kingdom*
- ⁵⁵ *Massachusetts Institute of Technology, Cambridge, MA, United States*
- ⁵⁶ *University of Cincinnati, Cincinnati, OH, United States*
- ⁵⁷ *University of Maryland, College Park, MD, United States*
- ⁵⁸ *Syracuse University, Syracuse, NY, United States*
- ⁵⁹ *Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to ²*
- ⁶⁰ *Institut für Physik, Universität Rostock, Rostock, Germany, associated to ¹¹*
- ^a *P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia*
- ^b *Università di Bari, Bari, Italy*
- ^c *Università di Bologna, Bologna, Italy*
- ^d *Università di Cagliari, Cagliari, Italy*
- ^e *Università di Ferrara, Ferrara, Italy*
- ^f *Università di Firenze, Firenze, Italy*
- ^g *Università di Urbino, Urbino, Italy*
- ^h *Università di Modena e Reggio Emilia, Modena, Italy*
- ⁱ *Università di Genova, Genova, Italy*
- ^j *Università di Milano Bicocca, Milano, Italy*
- ^k *Università di Roma Tor Vergata, Roma, Italy*
- ^l *Università di Roma La Sapienza, Roma, Italy*
- ^m *Università della Basilicata, Potenza, Italy*
- ⁿ *LIFAELS, La Salle, Universitat Ramon Llull, Barcelona, Spain*
- ^o *Hanoi University of Science, Hanoi, Viet Nam*
- ^p *Institute of Physics and Technology, Moscow, Russia*
- ^q *Università di Padova, Padova, Italy*
- ^r *Università di Pisa, Pisa, Italy*
- ^s *Scuola Normale Superiore, Pisa, Italy*

Rare decays that are forbidden in the Standard Model (SM) probe potential contributions from new processes and particles at a scale beyond the reach of direct searches. The decays $B_s^0 \rightarrow e^\pm \mu^\mp$ and $B^0 \rightarrow e^\pm \mu^\mp$ and their charged conjugate processes¹ are forbidden within the SM, in which lepton flavour is conserved. These decays are allowed in some scenarios beyond the SM that include models with heavy singlet Dirac neutrinos [1], supersymmetric models [2] and the Pati-Salam model [3]. The latter predicts a new interaction to mediate transitions between leptons and quarks via exchange of spin-1 gauge bosons, called Pati-Salam leptoquarks (LQ), that carry both colour and lepton quantum numbers.

Current limits from ATLAS [4, 5, 6] and CMS [7, 8, 9] on the masses of first, second or third generation leptoquarks are in the range $[0.4, 0.9] \text{ TeV}/c^2$, depending on the value of the couplings and the decay channel. These leptoquarks arise from a coupling between a quark and lepton of the same generation. The decays $B_s^0 \rightarrow e^\pm \mu^\mp$ and $B^0 \rightarrow e^\pm \mu^\mp$ can be mediated by other leptoquarks which couple leptons and quarks that are not necessarily from the same generation [10, 11], such as when the τ lepton couples to a first or second quark generation.

The previous best upper limits on the branching fraction of these decays come from the CDF collaboration [12], $\mathcal{B}(B_s^0 \rightarrow e^\pm \mu^\mp) < 2.0 (2.6) \times 10^{-7}$ and $\mathcal{B}(B^0 \rightarrow e^\pm \mu^\mp) < 6.4 (7.9) \times 10^{-8}$ at 90% (95%) confidence level (C.L.). These limits correspond to bounds on the masses of the corresponding Pati-Salam leptoquarks of $M_{\text{LQ}}(B_s^0 \rightarrow e^\pm \mu^\mp) > 47.8 (44.9) \text{ TeV}/c^2$ and $M_{\text{LQ}}(B^0 \rightarrow e^\pm \mu^\mp) > 59.3 (56.3) \text{ TeV}/c^2$ at 90 (95)% C.L. [12].

This Letter presents a search for the $B_s^0 \rightarrow e^\pm \mu^\mp$ and $B^0 \rightarrow e^\pm \mu^\mp$ lepton-flavour violating (LFV) decays performed with a data sample, corresponding to an integrated luminosity of 1.0 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$, collected by the LHCb experiment in 2011 at the Large Hadron Collider. To avoid potential bias, events in the signal mass region $[5.1, 5.5] \text{ GeV}/c^2$ were not examined until all analysis choices were finalized.

The LHCb detector is a single-arm forward spectrometer covering the pseudorapidity range $2 < \eta < 5$, and is described in detail in Ref. [13]. Events were simulated for this analysis using the software described in Refs. [14, 15, 16, 17, 18, 19, 20].

The trigger [21] consists of a hardware stage (L0),

¹Inclusion of charge conjugate processes are implied throughout this Letter.

based on information from the calorimeter and muon systems, followed by a software stage (HLT) that applies a full event reconstruction, and is split into two stages called HLT1 and HLT2. Candidate $B_{(s)}^0 \rightarrow e^\pm \mu^\mp$ decays considered in this analysis must satisfy a hardware decision that requires the presence of a muon candidate with transverse momentum $p_T > 1.5 \text{ GeV}/c$.

All tracks considered in the HLT1 are required to have $p_T > 0.5 \text{ GeV}/c$. The muon track of the $B_{(s)}^0 \rightarrow e^\pm \mu^\mp$ candidates is required to have $p_T > 1.0 \text{ GeV}/c$ and impact parameter, $\text{IP} > 0.1 \text{ mm}$. The HLT2 consists of exclusive, cut-based triggers for $B_{(s)}^0$ two-body decays, and inclusive multivariate [21, 22] b -hadron triggers.

The $B^0 \rightarrow K^+ \pi^-$ decay is used as the normalization channel and $B_{(s)}^0 \rightarrow h^+ h'^-$ ($h^{(\prime)} = K, \pi$) decays are used as a control channel, since both have the same event topology as the signal. The $B^0 \rightarrow K^+ \pi^-$ yield is computed from the yield of $B_{(s)}^0 \rightarrow h^+ h'^-$ decays, and the fraction of $B^0 \rightarrow K^+ \pi^-$ in the $B_{(s)}^0 \rightarrow h^+ h'^-$ sample, as described in Ref. [23]. In order to minimize the bias introduced by the trigger requirements, only $B_{(s)}^0 \rightarrow h^+ h'^-$ candidates that are triggered independently of the presence of either of the two signal hadrons at L0 and HLT1 are considered.

The $B_{(s)}^0 \rightarrow e^\pm \mu^\mp$ candidates that pass the trigger selection criteria are further required to have well identified electron and muon [24] candidates. The measured momenta of the electrons are corrected to account for loss of momentum by bremsstrahlung in the detector using the photon energy deposition in the electromagnetic calorimeter [25]. The signal candidates are required to be displaced with respect to any pp collision vertex (PV), and form a secondary vertex (SV) with χ^2 per degree of freedom smaller than 9 and separated from the PV in the downstream direction by a flight distance significance greater than 15.

Only $B_{(s)}^0$ candidates with an impact parameter χ^2 (χ_{IP}^2) less than 25 are considered. The χ_{IP}^2 of a $B_{(s)}^0$ candidate is defined as the difference between the χ^2 of the PV reconstructed with and without the considered candidate. When more than one PV is reconstructed, that giving the smallest χ_{IP}^2 for the $B_{(s)}^0$ candidate is chosen. Only $B_{(s)}^0$ candidates with invariant mass in the range $[4.9, 5.9] \text{ GeV}/c^2$ are kept for further analysis. The selection criteria for the $B_{(s)}^0 \rightarrow h^+ h'^-$ and $B^0 \rightarrow K^+ \pi^-$ candidates are identical to those of the signal, apart from those used for particle identification.

A two-stage multivariate selection based on boosted decision trees (BDT) [26, 27] is applied to the

$B_{(s)}^0 \rightarrow e^\pm \mu^\mp$ candidates following the same strategy as Ref. [28]. The two multivariate discriminants are trained using simulated samples, $B_s^0 \rightarrow e^\pm \mu^\mp$ for signal and $b\bar{b} \rightarrow l^\pm l'^\mp X$ for background (where $l^{(\prime)}$ can either be a μ or an e and X is any other set of particles), which is dominated by simultaneous semileptonic decays of both b and \bar{b} hadrons within the same event.

The requirement on the first multivariate discriminant [28] removes 75 % of the background while retaining 93 % of signal, as determined from simulation using half of the available samples to train and the other half to evaluate the efficiencies. The same selection is applied to the $B^0 \rightarrow K^+ \pi^-$ normalization channel and the efficiencies of this requirement for the signal and normalization channel are equal within 1.2 %, as determined from simulation.

The surviving background mainly comprises random combinations of electrons and muons from semileptonic $b\bar{b} \rightarrow e^\pm \mu^\mp X$ decays. In total 5766 electron-muon pairs pass the trigger, the offline selection and the first multivariate discriminant requirements. The selected candidates are classified in a binned two-dimensional space formed by the electron-muon invariant mass and the output of a second BDT, for which nine variables are employed [28]. The BDT output is independent of the invariant mass for signal inside the search window. The output is transformed such that the signal is approximately uniformly distributed between zero and one, while the background peaks at zero.

The probability for a signal event to have a given BDT value is obtained from data using the $B_{(s)}^0 \rightarrow h^+ h'^-$ sample [29, 30]. Simulated samples of $B_{(s)}^0 \rightarrow e^\pm \mu^\mp$ and $B_{(s)}^0 \rightarrow h^+ h'^-$ decays have been used to check that the distributions of the variables entering in the BDT that do not depend on the bremsstrahlung radiation are in good agreement. Corrections to the BDT shape due to the presence of the radiation emitted by the electron of the $B_{(s)}^0 \rightarrow e^\pm \mu^\mp$ decays have been evaluated using simulation. The number of $B_{(s)}^0 \rightarrow h^+ h'^-$ signal events in each BDT bin is determined by fitting the $h^+ h'^-$ invariant mass distribution. The systematic uncertainty on the signal BDT probability distribution function is taken to be the maximum spread in the fractions of yields going into each bin, obtained by fitting the same $B_{(s)}^0 \rightarrow h^+ h'^-$ dataset with different signal and background fit models. Corrections are applied to the BDT shape in order to take into account the effect of the different trigger requirements used for the signal and the $B_{(s)}^0 \rightarrow h^+ h'^-$

control sample.

The invariant mass line shape of the signal events is described by a Crystal Ball function (CB) [31] with two tails, left and right, defined by two parameters each. The values of the parameters depend on the momentum resolution, the momentum scale and the amount of bremsstrahlung radiation recovered.

The signal shape parameters are obtained from simulation, but need to be reweighted to account for their dependency on the event multiplicity, which affects the amount of bremsstrahlung radiation recovered and differs between data and simulation. We use the number of hits in the scintillating pad detector (N_{SPD}) as a measure of the event multiplicity. The distribution of N_{SPD} for $B_{(s)}^0 \rightarrow e^\pm \mu^\mp$ signal candidates is obtained from a $B^+ \rightarrow J/\psi(\mu^+ \mu^-) K^+$ data sample, which is selected with the same trigger conditions as the signal, ensuring a similar distribution of N_{SPD} . The signal mass shape parameters are determined by reweighting the $B_{(s)}^0 \rightarrow e^\pm \mu^\mp$ simulated events with the N_{SPD} distribution measured in the $B^+ \rightarrow J/\psi(\mu^+ \mu^-) K^+$ sample.

This reweighting technique is used also for a $J/\psi \rightarrow e^+ e^-$ simulated sample and the reweighted parameters are then compared with those obtained with a $J/\psi \rightarrow e^+ e^-$ sample in data. The difference between the mean values of the $J/\psi \rightarrow e^+ e^-$ mass in data and simulation (+0.16%) is applied as a systematic shift to the peak values of the $B^0 \rightarrow e^\pm \mu^\mp$ and $B_s^0 \rightarrow e^\pm \mu^\mp$ invariant mass in simulation. A systematic uncertainty is added to the $B_{(s)}^0 \rightarrow e^\pm \mu^\mp$ mass parameters when the differences in the values of the other mass parameters for the $J/\psi \rightarrow e^+ e^-$ sample in data and SPD-reweighted simulation are larger than their statistical uncertainties.

The signal region, defined by the invariant mass window $[5.1, 5.5] \text{ GeV}/c^2$, retains $(85.0 \pm 0.1_{\text{stat}} \pm 5.0_{\text{syst}})\%$ and $(82.0 \pm 0.1_{\text{stat}} \pm 5.0_{\text{syst}})\%$ of the $B_s^0 \rightarrow e^\pm \mu^\mp$ and $B^0 \rightarrow e^\pm \mu^\mp$ signal decays, respectively. The systematic uncertainties on these fractions are evaluated with pseudo-experiments that fluctuate each parameter of the mass lineshape according to its uncertainty. The width of the corresponding fraction distribution is taken as the systematic uncertainty.

The $B_s^0 \rightarrow e^\pm \mu^\mp$ and $B^0 \rightarrow e^\pm \mu^\mp$ yields are translated into branching fractions according to

$$\begin{aligned} \mathcal{B}(B_{(s)}^0 \rightarrow e^\pm \mu^\mp) &= \frac{\mathcal{B}_{\text{norm}} \epsilon_{\text{norm}} f_d}{N_{\text{norm}} \epsilon_{\text{sig}} f_{d(s)}} \times N_{B_{(s)}^0 \rightarrow e^\pm \mu^\mp} \\ &= \alpha_{B_{(s)}^0} \times N_{B_{(s)}^0 \rightarrow e^\pm \mu^\mp}, \end{aligned} \quad (1)$$

where $N_{\text{norm}} = 10\,120 \pm 920$ is the number of signal events in the normalization channel and is determined from the total yield of the $B_{(s)}^0 \rightarrow h^+h'^-$ channel and the fraction of $B^0 \rightarrow K^+\pi^-$ events in the inclusive sample. The systematic uncertainty is comparable to the statistical one and is dominated by the maximum spread in the yield obtained by fitting the same $B_{(s)}^0 \rightarrow h^+h'^-$ dataset with different fit models [29, 30]. The branching fraction of the normalization channel is $\mathcal{B}_{\text{norm}} = (1.94 \pm 0.06) \times 10^{-5}$ [32] and $N_{B_{(s)}^0 \rightarrow e^\pm\mu^\mp}$ is the number of observed signal events. The factors f_d and f_s indicate the probabilities that a b quark fragments into a B^0 or B_s^0 meson, respectively. We use $f_s/f_d = 0.256 \pm 0.020$ measured in pp collision data at $\sqrt{s} = 7$ TeV [33]. The measured dependence of f_s/f_d on the B meson p_T [33] is found to be negligible for this analysis.

The efficiency $\epsilon_{\text{sig(norm)}}$ for the signal (normalization) channel is the product of the reconstruction efficiency of the final state particles including the geometric detector acceptance, the selection efficiency and the trigger efficiency. The ratios of acceptance, reconstruction and selection efficiencies are computed with simulation. A systematic uncertainty is assigned to these ratios, to take into account the difference between the tracking efficiencies measured in data and predicted in simulation. Reweighting techniques are used to correct distributions in the simulation that do not match those from data, in particular for those variables that depend on N_{SPD} . The trigger efficiency of L0 and HLT1 on signal decays is evaluated using data, while the HLT2 efficiency is evaluated in simulation after validation with control samples. The electron and muon identification efficiencies are evaluated from data using the $B^+ \rightarrow J/\psi(\mu^+\mu^-)K^+$ and $B^+ \rightarrow J/\psi(e^+e^-)K^+$ control samples. The two normalization factors $\alpha_{B_s^0}$ and α_{B^0} are determined to be $\alpha_{B_s^0} = (1.1 \pm 0.2) \times 10^{-9}$ and $\alpha_{B^0} = (2.8 \pm 0.5) \times 10^{-10}$.

The BDT range is divided into eight bins with boundaries at 0.0, 0.25, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9 and 1.0. The number of expected combinatorial background events in each BDT bin and the invariant mass signal region is determined from data by fitting to an exponential function events in the mass sidebands, defined by $[4.9, 5.0]$ GeV/ c^2 and $[5.5, 5.9]$ GeV/ c^2 . The invariant mass distributions of the selected candidates in BDT bins and the binned BDT distributions for the signals and the combinatorial background samples are available in Supplemental Material [34].

In the exponential function both the slope and the normalization are allowed to vary. The systematic un-

certainty on the estimated number of combinatorial background events in the signal regions is determined by fluctuating the number of events observed in the sidebands according to a Poisson distribution, and by varying the exponential slope according to its uncertainty. As a cross-check, two other models, the sum of two exponential functions and a single exponential fitted to the right sideband only, have been used and provide consistent background estimates inside the signal region.

The low-mass sideband and the signal region are potentially polluted by exclusive backgrounds. The background from $B_c^+ \rightarrow J/\psi(\mu^+\mu^-)e^+\nu_e$ and $B_c^+ \rightarrow J/\psi(e^+e^-)\mu^+\nu_\mu$ decays is evaluated assuming the branching fraction value from Ref. [35]. The decays $B^0 \rightarrow \pi^-l^+\nu_l$, $B_{(s)}^0 \rightarrow h^+h'^-$, $B_s^0 \rightarrow K^-l^+\nu_l$, $\Lambda_b^0 \rightarrow pl^-\bar{\nu}_l$ and $B^+ \rightarrow \pi^+l^+l^-$ (where $l^\pm = e^\pm$ or μ^\pm) are potential backgrounds if the hadrons are misidentified as electrons or muons. The $B^0 \rightarrow \pi^-l^+\nu_l$ and $B_{(s)}^0 \rightarrow h^+h'^-$ branching fractions are taken from Ref. [32]. The $B^+ \rightarrow \pi^+l^+l^-$ branching fraction is taken from Ref. [36]. The theoretical estimates of the $\Lambda_b^0 \rightarrow pl^-\bar{\nu}_l$ and $B_s^0 \rightarrow K^-l^+\nu_l$ branching fractions are taken from Refs. [37] and [38], respectively. We use the Λ_b^0 fragmentation fraction $f_{\Lambda_b^0}$ measured by LHCb [39] and account for its p_T dependence.

The mass and BDT distributions of these background modes are evaluated from simulated samples, using the probabilities of misidentifying kaon, pion and proton as muon or electron as functions of momenta and transverse momenta, which are determined from $D^{*+} \rightarrow D^0(\rightarrow K^-\pi^+)\pi^+$ and $\Lambda \rightarrow p\pi^-$ data samples. The yield of the $B_{(s)}^0 \rightarrow h^+h'^- \rightarrow e^+\mu^-$ peaking background in each BDT bin is obtained by multiplying the $B_{(s)}^0 \rightarrow h^+h'^-$ yields obtained by fitting the invariant mass distribution of an inclusive $B_{(s)}^0 \rightarrow h^+h'^-$ sample in BDT bins [29, 30] with the probabilities of misidentifying kaon, pion and proton as muon or electron as functions of momenta and transverse momenta, as determined from control samples. The mass lineshape of the $B_{(s)}^0 \rightarrow h^+h'^- \rightarrow e^+\mu^-$ peaking background is obtained from a simulated sample of doubly-misidentified $B_{(s)}^0 \rightarrow h^+h'^-$ events. Apart from $B_{(s)}^0 \rightarrow h^+h'^-$, all background modes are normalized relative to the $B^+ \rightarrow J/\psi(\mu^+\mu^-)K^+$ decay. We assume $f_u = f_d$ where f_u is the B^+ fragmentation fraction.

The $\Lambda_b^0 \rightarrow pl^-\bar{\nu}_l$ and the $B_c^+ \rightarrow J/\psi(\mu^+\mu^-)e^+\nu_e$ and $B_c^+ \rightarrow J/\psi(e^+e^-)\mu^+\nu_\mu$ modes are the dominant exclusive modes in the range $\text{BDT} > 0.5$, where the combinatorial background is reduced by a factor ~ 500 according to simulation. These decay modes have an

invariant mass distribution that is compatible with an exponential in the region [4.9-5.9] GeV/ c^2 , and hence are taken into account by the exponential fit to the mass sidebands.

In the entire BDT and mass range ([4.9, 5.9] GeV/ c^2), 4.5 ± 0.7 doubly misidentified $B_{(s)}^0 \rightarrow h^+ h'^-$ decays are expected, with $(87.9 \pm 0.1)\%$ lying in the signal mass interval of [5.1, 5.5] GeV/ c^2 .

For each BDT bin we count the number of candidates observed in the signal region, and compare to the expected number of signal and background events.

The systematic uncertainties in the background and signal predictions in each bin are computed by varying the normalization factor, and the mass and BDT shapes within their Gaussian uncertainties.

The results for the $B_s^0 \rightarrow e^\pm \mu^\mp$ and $B^0 \rightarrow e^\pm \mu^\mp$ decays are summarized in Table 1. In the high BDT range, the observed number of candidates is in agreement with the number of expected exclusive backgrounds in the signal region. The compatibility of the observed distribution of events with that expected for a given branching fraction hypothesis is computed with the CL_s method [40].

The expected and observed CL_s values are shown in Fig. 1 for the $B_s^0 \rightarrow e^\pm \mu^\mp$ and $B^0 \rightarrow e^\pm \mu^\mp$ channels, as a function of the assumed branching fraction. The expected and measured limits for $B_s^0 \rightarrow e^\pm \mu^\mp$ and $B^0 \rightarrow e^\pm \mu^\mp$ at 90% and 95% C.L. are shown in Table 2. Note that since the same events are used to set limits for both B_s^0 and B^0 decays, the results are strongly correlated. The inclusion of systematic uncertainties increases the expected $B^0 \rightarrow e^\pm \mu^\mp$ and $B_s^0 \rightarrow e^\pm \mu^\mp$ upper limits by $\sim 20\%$. The systematic uncertainties are dominated by the uncertainty in the interpolation of the background yields inside the signal region. The observed limits are $\sim 1\sigma$ below the expectation due to the lower than expected numbers of observed events in the fourth and last BDT bins.

In the framework of the Pati-Salam model, the relation linking the $B_{(s)}^0 \rightarrow e^\pm \mu^\mp$ branching fractions and the leptoquark mass (M_{LQ}) [10] is

$$\mathcal{B}(B_{(s)}^0 \rightarrow e^\pm \mu^\mp) = \pi \frac{\alpha_S^2(M_{LQ})}{M_{LQ}^4} F_{B_{(s)}^0}^2 m_{B_{(s)}^0}^3 R^2 \frac{\tau_{B_{(s)}^0}}{\hbar}, \quad (2)$$

where

$$R = \frac{m_{B_{(s)}^0}}{m_b} \left(\frac{\alpha_S(M_{LQ})}{\alpha_S(m_t)} \right)^{-\frac{4}{7}} \left(\frac{\alpha_S(m_t)}{\alpha_S(m_b)} \right)^{-\frac{12}{23}}.$$

Table 2: Expected (background only) and observed limits on the $B_{(s)}^0 \rightarrow e^\pm \mu^\mp$ branching fractions.

Mode	Limit	90 % C.L.	95 % C.L.
$B_s^0 \rightarrow e^\pm \mu^\mp$	Expected	1.5×10^{-8}	1.8×10^{-8}
	Observed	1.1×10^{-8}	1.4×10^{-8}
$B^0 \rightarrow e^\pm \mu^\mp$	Expected	3.8×10^{-9}	4.8×10^{-9}
	Observed	2.8×10^{-9}	3.7×10^{-9}

The B^0 and B_s^0 masses, m_{B^0} and $m_{B_s^0}$, and the average lifetimes, τ_{B^0} and $\tau_{B_s^0}$, are taken from Ref. [32]. The factors $F_{B^0} = 0.190 \pm 0.004$ GeV and $F_{B_s^0} = 0.227 \pm 0.004$ GeV are the decay constants of the B^0 and B_s^0 mesons [41], and m_b and m_t are the bottom and top quark masses [32], respectively, computed in the $\overline{\text{MS}}$ scheme [42]. The value of α_s at an arbitrary scale M_{LQ} is determined using the software package RUNDEC [43].

Using the limits on the branching fractions shown in Table 2, we find the following lower bounds for the leptoquark masses if the leptoquark links the τ lepton to the first and second quark generation, $M_{LQ}(B_s^0 \rightarrow e^\pm \mu^\mp) > 107$ (101) TeV/ c^2 and $M_{LQ}(B^0 \rightarrow e^\pm \mu^\mp) > 135$ (126) TeV/ c^2 at 90 (95)% C.L., respectively. When the parameters entering in Eq. 2 are fluctuated within $\pm 1\sigma$, the limits on the leptoquark masses change by $\sim \pm 1$ TeV.

In summary, a search for the lepton-flavour violating decays $B_s^0 \rightarrow e^\pm \mu^\mp$ and $B^0 \rightarrow e^\pm \mu^\mp$ has been performed on a data sample, corresponding to an integrated luminosity of 1.0 fb $^{-1}$, collected in pp collisions at $\sqrt{s} = 7$ TeV. The data are consistent with the background-only hypothesis. Upper limits are set on the branching fractions, $\mathcal{B}(B_s^0 \rightarrow e^\pm \mu^\mp) < 1.1$ (1.4) $\times 10^{-8}$ and $\mathcal{B}(B^0 \rightarrow e^\pm \mu^\mp) < 2.8$ (3.7) $\times 10^{-9}$ at 90 (95)% C.L., that are the most restrictive to date. These limits translate into lower bounds on the leptoquark masses in the Pati-Salam model [10] of $M_{LQ}(B_s^0 \rightarrow e^\pm \mu^\mp) > 107$ (101) TeV/ c^2 and $M_{LQ}(B^0 \rightarrow e^\pm \mu^\mp) > 135$ (126) TeV/ c^2 at 90 (95)% C.L., respectively. These are a factor of two higher than the previous bounds.

Table 1: Expected background (bkg) from the fit to the data sidebands, and expected $B_{(s)}^0 \rightarrow h^+h'^- \rightarrow e^+\mu^-$ events, compared to the number of observed events in the mass signal region, in bins of BDT response.

BDT bin	0.0 – 0.25	0.25 – 0.4	0.4 – 0.5	0.5 – 0.6	0.6 – 0.7	0.7 – 0.8	0.8 – 0.9	0.9 – 1.0
Expected bkg (from fit)	2222 ± 51	$80.9^{+10.1}_{-9.4}$	$20.4^{+5.0}_{-4.5}$	$13.2^{+3.9}_{-3.6}$	$2.1^{+2.9}_{-1.4}$	$3.1^{+1.9}_{-1.4}$	$3.1^{+1.9}_{-1.4}$	$1.7^{+1.4}_{-1.0}$
Expected $B_{(s)}^0 \rightarrow h^+h'^- \rightarrow e^+\mu^-$ bkg	0.67 ± 0.12	0.47 ± 0.09	0.40 ± 0.08	0.37 ± 0.06	0.45 ± 0.08	0.49 ± 0.08	0.57 ± 0.09	0.54 ± 0.12
Observed	2332	90	19	4	3	3	3	1




Figure 1: CL_s as a function of the assumed branching fraction for (left) $B_s^0 \rightarrow e^\pm\mu^\mp$ and (right) $B^0 \rightarrow e^\pm\mu^\mp$ decays. The dashed lines are the medians of the expected CL_s distributions if background only was observed. The yellow (green) area covers, at a given branching fraction, 34%(47.5%) of the expected CL_s distribution on each side of its median. The solid black curves are the observed CL_s . The upper limits at 90 % (95 %) C.L. are indicated by the dotted (solid) vertical lines in blue for the expectation and in red for the observation.

1 Acknowledgements

We thank Diego Guadagnoli for the theory inputs. We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); NSFC (China); CNRS/IN2P3 and Region Auvergne (France); BMBF, DFG, HGF and MPG (Germany); SFI (Ireland); INFN (Italy); FOM and NWO (The Netherlands); SCSR (Poland); MEN/IFA (Romania); MinES, Rosatom, RFBR and NRC ‘‘Kurchatov Institute’’ (Russia); MinECo, XuntaGal and GENCAT (Spain); SNSF and SER (Switzerland); NAS Ukraine (Ukraine); STFC (United Kingdom); NSF (USA). We also acknowledge the support received from the ERC under FP7. The Tier1 computing centres are supported by IN2P3 (France), KIT and BMBF (Ger-

many), INFN (Italy), NWO and SURF (The Netherlands), PIC (Spain), GridPP (United Kingdom). We are thankful for the computing resources put at our disposal by Yandex LLC (Russia), as well as to the communities behind the multiple open source software packages that we depend on.

References

- [1] A. Ilakovic, *Lepton flavor violation in the Standard Model extended by heavy singlet Dirac neutrinos*, Phys. Rev. D **62** (2000) 036010.
- [2] R. A. Diaz *et al.*, *Improving bounds on flavor changing vertices in the two Higgs doublet model from $B^0 - \bar{B}^0$ mixing*, Eur. Phys. J **C41** (2005) 305.
- [3] J. C. Pati and A. Salam, *Lepton number as the fourth color*, Phys. Rev. D **10** (1974) 275.

- [4] ATLAS collaboration, G. Aad *et al.*, *Search for second generation scalar leptoquarks in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector*, Eur. Phys. J. **C72** (2012) 2151, [arXiv:1203.3172](#).
- [5] ATLAS collaboration, G. Aad *et al.*, *Search for first generation scalar leptoquarks in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector*, Phys. Lett. **B709** (2012) 158, [arXiv:1112.4828](#).
- [6] ATLAS collaboration, G. Aad *et al.*, *Search for third generation scalar leptoquarks in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector*, JHEP **06** (2013) 033, [arXiv:1303.0526](#).
- [7] CMS collaboration, S. Chatrchyan *et al.*, *Search for pair production of first- and second-generation scalar leptoquarks in pp collisions at $\sqrt{s} = 7$ TeV*, Phys. Rev. D **86** (2012) 052013, [arXiv:1207.5406](#).
- [8] CMS collaboration, S. Chatrchyan *et al.*, *Search for third-generation leptoquarks and scalar bottom quarks in pp collisions at $\sqrt{s} = 7$ TeV*, JHEP **12** (2012) 055, [arXiv:1210.5627](#).
- [9] CMS collaboration, S. Chatrchyan *et al.*, *Search for pair production of third-generation leptoquarks and top squarks in pp collisions at $\sqrt{s} = 7$ TeV*, Phys. Rev. Lett. **110** (2013) 081801, [arXiv:1210.5629](#).
- [10] G. Valencia and S. Willenbrock, *Quark-lepton unification and rare meson decays*, Phys. Rev. D **50** (1994) 6843.
- [11] M. Blanke *et al.*, *Charged lepton flavour violation and $(g - 2)_\mu$ in the Littlest Higgs Model with T-Parity: a clear distinction from Supersymmetry*, JHEP **05** (2007) 013, [arXiv:0702136](#).
- [12] CDF collaboration, T. Aaltonen *et al.*, *Search for the decays $B_{(s)}^0 \rightarrow e^+\mu^-$ and $B_{(s)}^0 \rightarrow e^+e^-$ in CDF Run II*, Phys. Rev. Lett. **102** (2009) 201801, [arXiv:0901.3803](#).
- [13] LHCb collaboration, A. A. Alves Jr. *et al.*, *The LHCb detector at the LHC*, JINST **3** (2008) S08005.
- [14] T. Sjöstrand, S. Mrenna, and P. Skands, *PYTHIA 6.4 physics and manual*, JHEP **05** (2006) 026, [arXiv:hep-ph/0603175](#).
- [15] D. J. Lange, *The EvtGen particle decay simulation package*, Nucl. Instrum. Meth. **A462** (2001) 152.
- [16] GEANT4 collaboration, J. Allison *et al.*, *GEANT4 developments and applications*, IEEE Trans. Nucl. Sci. **53** (2006) 270.
- [17] GEANT4 collaboration, S. Agostinelli *et al.*, *GEANT4: a simulation toolkit*, Nucl. Instrum. Meth. **A506** (2003) 250.
- [18] P. Golonka and Z. Was, *PHOTOS Monte Carlo: a precision tool for QED corrections in Z and W decays*, Eur. Phys. J. **C45** (2006) 97, [arXiv:hep-ph/0506026](#).
- [19] I. Belyaev *et al.*, *Handling of the generation of primary events in GAUSS, the LHCb simulation framework*, Nuclear Science Symposium Conference Record (NSS/MIC) **IEEE** (2010) 1155.
- [20] M. Clemencic *et al.*, *The LHCb simulation application, GAUSS: design, evolution and experience*, J. of Phys: Conf. Ser. **331** (2011) 032023.
- [21] R. Aaij *et al.*, *The LHCb trigger and its performance in 2011*, JINST **8** (2013) P04022, [arXiv:1211.3055](#).
- [22] V. V. Gligorov and M. Williams, *Efficient, reliable and fast high-level triggering using a bonsai boosted decision tree*, JINST **8** (2013) P02013, [arXiv:1210.6861](#).
- [23] LHCb collaboration, B. Adeva *et al.*, *Roadmap for selected key measurements of LHCb*, [arXiv:0912.4179](#).
- [24] F. Archilli *et al.*, *Performance of the Muon Identification at LHCb*, [arXiv:1306.0249](#), submitted to JINST.
- [25] LHCb collaboration, R. Aaij *et al.*, *Measurement of the $B^0 \rightarrow K^{*0}e^+e^-$ branching fraction at low dilepton mass*, JHEP **05** (2013) 159, [arXiv:1304.3035](#).
- [26] L. Breiman, J. H. Friedman, R. A. Olshen, and C. J. Stone, *Classification and regression trees*, Wadsworth international group, Belmont, California, USA, 1984.
- [27] R. E. Schapire and Y. Freund, *A decision-theoretic generalization of on-line learning and an application to boosting*, Jour. Comp. and Syst. Sc. **55** (1997) 119.

- [28] LHCb collaboration, R. Aaij *et al.*, *First evidence for the decay $B_s^0 \rightarrow \mu^+\mu^-$* , Phys. Rev. Lett. **110** (2012) 021801, [arXiv:1211.2674](#).
- [29] LHCb collaboration, R. Aaij *et al.*, *Search for the rare decays $B_s^0 \rightarrow \mu^+\mu^-$ and $B^0 \rightarrow \mu^+\mu^-$* , Phys. Lett. **B708** (2012) 55, [arXiv:1112.1600](#).
- [30] LHCb collaboration, R. Aaij *et al.*, *Strong constraints on the rare decays $B_s \rightarrow \mu^+\mu^-$ and $B^0 \rightarrow \mu^+\mu^-$* , Phys. Rev. Lett. **108** (2012) 231801, [arXiv:1203.4493](#).
- [31] T. Skwarnicki, *A study of the radiative cascade transitions between the Upsilon-prime and Upsilon resonances*, PhD thesis, Institute of Nuclear Physics, Krakow, 1986, DESY-F31-86-02.
- [32] Particle Data Group, J. Beringer *et al.*, *Review of particle physics*, Phys. Rev. **D86** (2012) 010001.
- [33] LHCb collaboration, R. Aaij *et al.*, *Measurement of the ratio of fragmentation fractions f_s/f_d and dependence on B meson kinematics*, JHEP **04** (2013) 001, [arXiv:1301.5286](#).
- [34] See Supplemental Material at [URL will be inserted by publisher] for the invariant mass distributions of selected candidates in BDT bins and the binned BDT distributions for the signals and the combinatorial background samples.
- [35] CDF collaboration, F. Abe *et al.*, *Observation of the B_c meson in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV*, Phys. Rev. Lett. **81** (1998) 2432, [arXiv:hep-ex/9805034](#).
- [36] LHCb collaboration, R. Aaij *et al.*, *First observation of the decay $B^+ \rightarrow \pi^+\mu^+\mu^-$* , JHEP **12** (2012) 125, [arXiv:1210.2645](#).
- [37] A. Khodjamirian, C. Klein, T. Mannel, and Y.-M. Wang, *Form factors and strong couplings of heavy baryons from QCD light-cone sum rules*, JHEP **09** (2011) 106, [arXiv:1108.2971](#).
- [38] W.-F. Wang and Z.-J. Xiao, *Semileptonic decays $B/B_s \rightarrow (\pi, K)(\ell^+\ell^-, \ell\nu, \nu\bar{\nu})$ in the perturbative QCD approach beyond the leading order*, Phys. Rev. **D86** (2012) 114025, [arXiv:1207.0265](#).
- [39] LHCb collaboration, R. Aaij *et al.*, *Measurement of b-hadron production fractions in 7 TeV pp collisions*, Phys. Rev. **D85** (2012) 032008, [arXiv:1111.2357](#).
- [40] A. L. Read, *Presentation of search results: the CL_s technique*, J. Phys. **G28** (2002) 2693.
- [41] C. Davies, *Standard Model heavy flavor physics on the lattice*, PoS **LATTICE2011** (2011) 019, [arXiv:1203.3862](#).
- [42] W. A. Bardeen, A. Buras, D. Duke, and T. Muta, *Deep inelastic scattering beyond the leading order in asymptotically free gauge theories*, Phys. Rev. **D18** (1978) 3998.
- [43] K. Chetyrkin, J. Kuhn, and M. Steinhauser, *RunDec: a Mathematica package for running and decoupling of the strong coupling and quark masses*, Comput. Phys. Comm. **133** (2000) 43, [arXiv:hep-ph/0004189](#).