



# Precision measurement of $CP$ violation in $B_s^0 \rightarrow J/\psi K^+ K^-$ decays

The LHCb collaboration<sup>†</sup>

## Abstract

The time-dependent  $CP$  asymmetry in  $B_s^0 \rightarrow J/\psi K^+ K^-$  decays is measured using  $pp$  collision data, corresponding to an integrated luminosity of  $3.0 \text{ fb}^{-1}$ , collected with the LHCb detector at centre-of-mass energies of 7 and 8 TeV. In a sample of 96 000  $B_s^0 \rightarrow J/\psi K^+ K^-$  decays, the  $CP$ -violating phase  $\phi_s$  is measured, as well as the decay widths  $\Gamma_L$  and  $\Gamma_H$  of the light and heavy mass eigenstates of the  $B_s^0$ - $\bar{B}_s^0$  system. The values obtained are  $\phi_s = -0.058 \pm 0.049 \pm 0.006$  rad,  $\Gamma_s \equiv (\Gamma_L + \Gamma_H)/2 = 0.6603 \pm 0.0027 \pm 0.0015 \text{ ps}^{-1}$ , and  $\Delta\Gamma_s \equiv \Gamma_L - \Gamma_H = 0.0805 \pm 0.0091 \pm 0.0032 \text{ ps}^{-1}$ , where the first uncertainty is statistical and the second systematic. These are the most precise single measurements of those quantities to date. A combined analysis with  $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$  decays gives  $\phi_s = -0.010 \pm 0.039$  rad. All measurements are in agreement with the Standard Model predictions. For the first time the phase  $\phi_s$  is measured independently for each polarisation state of the  $K^+ K^-$  system and shows no evidence for polarisation dependence.

Submitted to Phys. Rev. Lett.

<sup>†</sup>Authors are listed at the end of this Letter.



The  $CP$ -violating phase  $\phi_s$  arises in the interference between the amplitudes of  $B_s^0$  mesons decaying via  $b \rightarrow c\bar{c}s$  transitions to  $CP$  eigenstates directly and those decaying after oscillation. In the Standard Model (SM), ignoring sub-leading contributions, this phase is predicted to be  $-2\beta_s$ , where  $\beta_s = \arg[-(V_{ts}V_{tb}^*)/(V_{cs}V_{cb}^*)]$  and  $V_{ij}$  are elements of the quark-mixing matrix [1]. Global fits to experimental data give  $-2\beta_s = -0.0363 \pm 0.0013$  rad [2]. This phase could be modified if non-SM particles were to contribute to the  $B_s^0$ - $\bar{B}_s^0$  oscillations [3, 4] and a measurement of  $\phi_s$  significantly different from the SM prediction would provide unambiguous evidence for processes beyond the SM.

The LHCb collaboration has previously reported measurements of  $\phi_s$  using  $B_s^0 \rightarrow J/\psi K^+ K^-$  and  $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$  decays [5, 6]. These measurements were based upon data, corresponding to an integrated luminosity of up to  $1.0 \text{ fb}^{-1}$ , collected in  $pp$  collisions at a centre-of-mass energy of 7 TeV in 2011 at the LHC. The ATLAS, CDF and D0 collaborations have also measured  $\phi_s$  in  $B_s^0 \rightarrow J/\psi K^+ K^-$  decays [7–9]. This Letter extends the LHCb measurements in the  $B_s^0 \rightarrow J/\psi K^+ K^-$  channel by adding data corresponding to  $2.0 \text{ fb}^{-1}$  of integrated luminosity collected at 8 TeV in 2012, and presents the combined results for  $\phi_s$  including the analysis of  $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$  decays from Ref. [10]. For the first time, the  $CP$ -violating phases are measured separately for each polarisation state of the  $K^+ K^-$  system. Knowledge of these parameters is an important step towards the control of loop-induced effects to the decay amplitude, which could potentially be confused with non-SM contributions to  $B_s^0$ - $\bar{B}_s^0$  mixing [11]. The analysis of the  $B_s^0 \rightarrow J/\psi K^+ K^-$  channel reported here is as described in Ref. [6], to which the reader is referred for details, except for the changes described below.

The LHCb detector is a single-arm forward spectrometer covering the pseudorapidity range  $2 < \eta < 5$ , designed for the study of particles containing  $b$  or  $c$  quarks, and is described in Ref. [12]. The trigger [13] consists of a hardware stage, based on information from the calorimeter and muon systems, followed by a software stage, in which all charged particles with transverse momentum greater than 500 (300) MeV/ $c$  are reconstructed for 2011 (2012) data. Further selection requirements are applied offline, as described in Ref. [6], in order to increase the signal purity.

The  $B_s^0 \rightarrow J/\psi K^+ K^-$  decay proceeds predominantly via  $B_s^0 \rightarrow J/\psi \phi$ , in which the  $K^+ K^-$  pair from the  $\phi$  is in a P-wave configuration. The final state is a superposition of  $CP$ -even and  $CP$ -odd states depending upon the relative orbital angular momentum of the  $J/\psi$  and  $\phi$  mesons. The  $J/\psi K^+ K^-$  final state can also be produced with  $K^+ K^-$  pairs in a  $CP$ -odd S-wave configuration [14]. The measurement of  $\phi_s$  requires the  $CP$ -even and  $CP$ -odd components to be disentangled by analysing the distribution of the reconstructed decay angles of the final-state particles. In this analysis the decay angles are defined in the helicity basis,  $\cos \theta_K$ ,  $\cos \theta_\mu$ , and  $\varphi_h$ , as described in Ref. [6].

The invariant mass distributions for  $K^+ K^-$  and  $J/\psi (\rightarrow \mu^+ \mu^-) K^+ K^-$  candidates are shown in Figs. 1(a) and 1(b), respectively. The combinatorial background is modelled with an exponential function and the  $B_s^0$  signal shape is parameterised by a double-sided Hypatia function [15]. The fitted signal yield is  $95\,690 \pm 350$ . In addition to the combinatorial background, studies of the data in sidebands of the  $m(J/\psi K^+ K^-)$  spectrum




Figure 1: (a) Background-subtracted invariant mass distributions of the  $K^+K^-$  system in the selected  $B_s^0 \rightarrow J/\psi K^+K^-$  candidates (black points). The vertical red lines denote the boundaries of the six bins used in the maximum likelihood fit. (b) Distribution of  $m(J/\psi K^+K^-)$  for the data sample (black points) and projection of the maximum likelihood fit (blue line). The  $B_s^0$  signal component is shown by the red dashed line and the combinatorial background by the green long-dashed line. Background from misidentified  $B^0$  and  $\Lambda_b^0$  decays is subtracted, as described in the text.

show contributions from approximately 1700  $B^0 \rightarrow J/\psi K^+\pi^-$  (4800  $\Lambda_b^0 \rightarrow J/\psi pK^-$ ) decays where the pion (proton) is misidentified as a kaon. These background events have complicated correlations between the angular variables and  $m(J/\psi K^+K^-)$ . In order to avoid the need to describe explicitly such correlations in the analysis, the contributions from these backgrounds are statistically subtracted by adding to the data simulated events of these decays with negative weight. Prior to injection, the simulated events are weighted such that the distributions of the relevant variables used in the fit, and their correlations, match those of data.

The principal physics parameters of interest are  $\Gamma_s$ ,  $\Delta\Gamma_s$ ,  $\phi_s$ ,  $|\lambda|$ , the  $B_s^0$  mass difference,  $\Delta m_s$ , and the polarisation amplitudes  $A_k = |A_k|e^{-i\delta_k}$ , where the indices  $k \in \{0, \parallel, \perp, S\}$  refer to the different polarisation states of the  $K^+K^-$  system. The sum  $|A_{\parallel}|^2 + |A_0|^2 + |A_{\perp}|^2$  equals unity and by convention  $\delta_0$  is zero. The parameter  $\lambda$  describes  $CP$  violation in the interference between mixing and decay and is defined by  $\eta_k(q/p)(\bar{A}_k/A_k)$ , where it is assumed to be the same for all polarisation states. The complex parameters  $p = \langle B_s^0 | B_L \rangle$  and  $q = \langle \bar{B}_s^0 | B_L \rangle$  describe the relation between mass and flavour eigenstates and  $\eta_k$  is the  $CP$  eigenvalue of the polarisation state  $k$ . The  $CP$ -violating phase is defined by  $\phi_s \equiv -\arg \lambda$ . In the absence of  $CP$  violation in decay,  $|\lambda| = 1$ .  $CP$  violation in  $B_s^0$ -meson mixing is negligible, following measurements in Ref. [16]. Measurements of the above parameters are obtained from a weighted maximum likelihood fit [17] to the decay-time and helicity angle distributions of the data as described in Ref. [6].

The  $B_s^0$  decay-time distribution is distorted by the trigger selection requirements and

by the track reconstruction algorithms. Corrections are determined from data using the methods described in Ref. [18] and are incorporated in the maximum likelihood fit by a parameterised function, in the case of the trigger, and by per-candidate weights, in the case of the track reconstruction. Both corrections are validated using a sample of  $10^6$  simulated  $B_s^0 \rightarrow J/\psi \phi$  events.

To account for the experimental decay-time resolution, the signal probability density function (PDF) is defined per candidate and is convolved with the sum of two Gaussian functions with a common mean,  $\mu$ , and independent widths  $w_i$ ,  $i \in \{1, 2\}$ . The widths are given by  $w_i \equiv r_i \sigma + s_i \sigma^2$ , where  $r_i$  and  $s_i$  are scale factors for each Gaussian function and  $\sigma$  is the per candidate decay-time uncertainty, estimated by the kinematic fit used to calculate the decay time. The scale factors are determined from the decay-time distribution of a control sample of  $J/\psi K^+ K^-$  candidates that are selected as for signal except for decay-time requirements. The average value of the  $\sigma$  distribution in the sample of prompt candidates is approximately 35 fs and the effective average resolution is 46 fs.

The flavour of the  $B_s^0$  candidate at production is inferred using two independent classes of flavour tagging algorithms, the opposite-side (OS) tagger and the same-side kaon (SSK) tagger, which exploit specific features of the production of  $b\bar{b}$  quark pairs in  $pp$  collisions. The OS tagger algorithm is described in Ref. [6] but is re-calibrated using data sets of flavour-specific decays, yielding a tagging power of  $(1.19 \pm 0.06)\%$  for events with only an OS-tag. The SSK algorithm deduces the signal production flavour by exploiting charge-flavour correlations of the kaons produced during the hadronisation process of the  $\bar{b}$  quark forming the signal  $B_s^0$  meson. The tagging kaon is identified using a selection based on a neural network that gives an effective tagging power of  $(0.84 \pm 0.11)\%$ , corresponding approximately to a 40% improvement in tagging power with respect to that in Refs. [6, 19]. The SSK algorithm is calibrated using a sample of  $B_s^0 \rightarrow D_s^- \pi^+$  decays. For events that have both OS and SSK tagging decisions, corresponding to 26% of the tagged sample, the effective tagging power is  $(1.70 \pm 0.08)\%$ . The combined tagging power of the three independent tagging categories defined above is  $(3.73 \pm 0.15)\%$ .

Due to different  $m(K^+ K^-)$  line shapes of the S- and P-wave contributions, their interferences are suppressed by an effective coupling factor after integrating over a finite  $m(K^+ K^-)$  region. The fit is carried out in six bins of  $m(K^+ K^-)$ , as shown in Fig. 1(a), to allow measurement of the small S-wave amplitude in each bin and to minimise correction factors in the interference terms of the PDF.

The results of the fit are consistent with the measurements reported in Ref. [6] and are reported in Table 1 where the first uncertainty is statistical and the second systematic. The correlation matrix is given in Ref. [20]. In contrast to Ref. [6] the value of  $\Delta m_s$  is unconstrained in this fit, thereby providing an independent measurement of this quantity, which is consistent with the results of Ref. [21]. The projections of the decay time and angular distributions are shown in Fig. 2.

The results reported in Table 1 are obtained with the assumption that  $\phi_s$  and  $|\lambda|$  are independent of the final-state polarisation. This condition can be relaxed to allow the measurement of  $\phi_s^k$  and  $|\lambda^k|$  separately for each polarisation, following the formalism in Ref. [22]. The results of this fit are shown in Table 2 and the statistical correlation matrix

Table 1: Values of the principal physics parameters determined from the polarisation-independent fit.

Parameter	Value
$\Gamma_s$ [ps <sup>-1</sup> ]	$0.6603 \pm 0.0027 \pm 0.0015$
$\Delta\Gamma_s$ [ps <sup>-1</sup> ]	$0.0805 \pm 0.0091 \pm 0.0032$
$ A_\perp ^2$	$0.2504 \pm 0.0049 \pm 0.0036$
$ A_0 ^2$	$0.5241 \pm 0.0034 \pm 0.0067$
$\delta_\parallel$ [rad]	$3.26^{+0.10}_{-0.17} {}^{+0.06}_{-0.07}$
$\delta_\perp$ [rad]	$3.08^{+0.14}_{-0.15} \pm 0.06$
$\phi_s$ [rad]	$-0.058 \pm 0.049 \pm 0.006$
$ \lambda $	$0.964 \pm 0.019 \pm 0.007$
$\Delta m_s$ [ps <sup>-1</sup> ]	$17.711^{+0.055}_{-0.057} \pm 0.011$

Table 2: Values of the polarisation-dependent parameters  $\phi_s^k$  and  $|\lambda^k|$  determined from the polarisation-dependent fit.

Parameter	Value
$ \lambda^0 $	$1.012 \pm 0.058 \pm 0.013$
$ \lambda^\parallel/\lambda^0 $	$1.02 \pm 0.12 \pm 0.05$
$ \lambda^\perp/\lambda^0 $	$0.97 \pm 0.16 \pm 0.01$
$ \lambda^S/\lambda^0 $	$0.86 \pm 0.12 \pm 0.04$
$\phi_s^0$ [rad]	$-0.045 \pm 0.053 \pm 0.007$
$\phi_s^\parallel - \phi_s^0$ [rad]	$-0.018 \pm 0.043 \pm 0.009$
$\phi_s^\perp - \phi_s^0$ [rad]	$-0.014 \pm 0.035 \pm 0.006$
$\phi_s^S - \phi_s^0$ [rad]	$0.015 \pm 0.061 \pm 0.021$

is given in Ref. [20]. There is no evidence for a polarisation-dependent  $CP$  violation arising in  $B_s^0 \rightarrow J/\psi K^+ K^-$  decays.

A summary of systematic uncertainties is reported in Tables 3 and 4. The tagging parameters are constrained in the fit and therefore their associated systematic uncertainties contribute to the statistical uncertainty of each parameter in Table 1. This contribution is 0.004 rad to the statistical uncertainty on  $\phi_s$ ; 0.004 ps<sup>-1</sup> to that of  $\Delta m_s$ ; 0.01 rad to that of  $\delta_\parallel$  and is negligible for all other parameters.

The assumption that the  $m(J/\psi K^+ K^-)$  distribution is independent from the decay time and angles is tested by re-evaluating the signal weights in bins of the decay time and angles and repeating the fit. The difference in fit results is assigned as a systematic uncertainty. The systematic effect from the statistical uncertainty on the signal weights is determined by re-computing them after varying the parameters of the  $m(J/\psi K^+ K^-)$  fit model within their statistical uncertainties, and assigning the difference in fit results as a systematic uncertainty.




Figure 2: Decay-time and helicity-angle distributions for  $B_s^0 \rightarrow J/\psi K^+ K^-$  decays (data points) with the one-dimensional fit projections overlaid. The solid blue line shows the total signal contribution, which is composed of  $CP$ -even (long-dashed red),  $CP$ -odd (short-dashed green) and S-wave (dotted-dashed purple) contributions.

The effect due to the  $b$ -hadron background contributions is evaluated by varying the proportion of simulated background events included in the fit by one standard deviation of their measured fractions. In addition, a further systematic uncertainty is assigned as the difference between the results of the fit to weighted or non-weighted data.

A small fraction of  $B_s^0 \rightarrow J/\psi K^+ K^-$  decays come from the decays of  $B_c^+$  mesons [23]. The effect of ignoring this component in the fit is evaluated using simulated pseudoexperiments where a 0.8% contribution [23, 24] of  $B_s^0$ -from- $B_c^+$  decays is added from a simulated sample of  $B_c^+ \rightarrow B_s^0(\rightarrow J/\psi \phi)\pi^+$  decays. Neglecting the  $B_c^+$  component leads to a bias on  $\Gamma_s$  of  $0.0005 \text{ ps}^{-1}$ , which is added as a systematic uncertainty. Other parameters are unaffected.

The decay angle resolution is found to be of the order of 20 mrad in simulated events. The result of pseudoexperiments shows that ignoring this effect in the fit only leads to small biases in the polarisation amplitudes, which are assigned as systematic uncertainties.

The angular efficiency correction is determined from simulated signal events weighted as in Ref. [6] such that the kinematic distributions of the final state particles match those

Table 3: Statistical and systematic uncertainties for the polarisation-independent result.

Source	$\Gamma_s$ [ps <sup>-1</sup> ]	$\Delta\Gamma_s$ [ps <sup>-1</sup> ]	$ A_\perp ^2$	$ A_0 ^2$	$\delta_\parallel$ [rad]	$\delta_\perp$ [rad]	$\phi_s$ [rad]	$ \lambda $	$\Delta m_s$ [ps <sup>-1</sup> ]
Total stat. uncertainty	0.0027	0.0091	0.0049	0.0034	$^{+0.10}_{-0.17}$	$^{+0.14}_{-0.15}$	0.049	0.019	$^{+0.055}_{-0.057}$
Mass factorisation	–	0.0007	0.0031	0.0064	0.05	0.05	0.002	0.001	0.004
Signal weights (stat.)	0.0001	0.0001	–	0.0001	–	–	–	–	–
<i>b</i> -hadron background	0.0001	0.0004	0.0004	0.0002	0.02	0.02	0.002	0.003	0.001
$B_c^+$ feed-down	0.0005	–	–	–	–	–	–	–	–
Angular resolution bias	–	–	0.0006	0.0001	$^{+0.02}_{-0.03}$	0.01	–	–	–
Ang. efficiency (reweighting)	0.0001	–	0.0011	0.0020	0.01	–	0.001	0.005	0.002
Ang. efficiency (stat.)	0.0001	0.0002	0.0011	0.0004	0.02	0.01	0.004	0.002	0.001
Decay-time resolution	–	–	–	–	–	0.01	0.002	0.001	0.005
Trigger efficiency (stat.)	0.0011	0.0009	–	–	–	–	–	–	–
Track reconstruction (simul.)	0.0007	0.0029	0.0005	0.0006	$^{+0.01}_{-0.02}$	0.002	0.001	0.001	0.006
Track reconstruction (stat.)	0.0005	0.0002	–	–	–	–	–	–	0.001
Length and momentum scales	0.0002	–	–	–	–	–	–	–	0.005
S-P coupling factors	–	–	–	–	0.01	0.01	–	0.001	0.002
Fit bias	–	–	0.0005	–	–	0.01	–	0.001	–
Quadratic sum of syst.	0.0015	0.0032	0.0036	0.0067	$^{+0.06}_{-0.07}$	0.06	0.006	0.007	0.011

Table 4: Statistical and systematic uncertainties for the polarisation-dependent result.

Source	$ \lambda^0 $	$ \lambda^\parallel/\lambda^0 $	$ \lambda^\perp/\lambda^0 $	$ \lambda^S/\lambda^0 $	$\phi_s^0$ [rad]	$\phi_s^\parallel - \phi_s^0$ [rad]	$\phi_s^\perp - \phi_s^0$ [rad]	$\phi_s^S - \phi_s^0$ [rad]
Total stat. uncertainty	0.058	0.12	0.16	0.12	0.053	0.043	0.035	0.061
Mass factorisation	0.010	0.04	0.01	0.03	0.003	0.005	0.003	0.016
<i>b</i> -hadron background	0.002	0.01	–	0.01	0.003	0.001	0.001	0.009
Ang. efficiency (reweighting)	–	–	–	0.02	0.001	0.002	0.001	0.007
Ang. efficiency (stat.)	0.004	0.02	0.01	0.01	0.004	0.007	0.005	0.004
Decay-time resolution	0.006	0.01	–	0.01	0.003	0.002	0.001	0.002
S-P coupling factors	–	–	–	–	–	–	–	0.006
Quadratic sum of syst.	0.013	0.05	0.01	0.04	0.007	0.009	0.006	0.021

in the data. A systematic uncertainty is assigned as the difference between the fit results using angular corrections from weighted or non-weighted simulated events. The limited size of the simulated sample leads to an additional systematic uncertainty.

The systematic uncertainty from the decay time resolution parameters is not included in the statistical uncertainty of each parameter and is now quoted explicitly. It is assigned as the difference of fit parameters obtained from the nominal fit and a fit where the resolution model parameters are calibrated using a sample of simulated prompt- $J/\psi$  events.

The trigger decay-time efficiency model, described in Ref. [6], introduces a systematic



uncertainty that is determined by fixing the value of each model parameter in the fit and subsequently repeating the fit with the parameter values constrained within their statistical uncertainty. The quadratic differences of the uncertainties returned by each fit are then assigned as systematic uncertainties. The systematic effect of the track reconstruction efficiency is evaluated by applying the same techniques on a large simulated sample of  $B_s^0 \rightarrow J/\psi \phi$  decays. The differences between the generation and fitted values of each physics parameter in this sample is assigned as the systematic uncertainty. The limited size of the control sample used to determine the track reconstruction efficiency parameterisation leads to an additional systematic uncertainty.

The uncertainty on the longitudinal coordinate of the LHCb vertex detector is found from survey data and leads to an uncertainty on  $\Gamma_s$  and  $\Delta\Gamma_s$  of 0.020%, with other parameters being unaffected. The momentum scale uncertainty is at most 0.022% [21], which only affects  $\Delta m_s$ .

Different models of the S-wave line-shape and  $m(K^+K^-)$  resolution are used to evaluate the coupling factors in each of the six  $m(K^+K^-)$  bins and the resulting variation of the fit parameters are assigned as systematic uncertainties. Possible biases of the fitting procedure are studied by generating and fitting many simulated pseudoexperiments of equivalent size to the data. The resulting biases are small, and those that are significantly different from zero are assigned as systematic uncertainties.

The systematic correlations between parameters are evaluated by assuming that parameters are fully (anti)correlated when the systematic uncertainty is determined by comparing results obtained from the nominal and a modified fit. Other sources of systematic uncertainty are assumed to have negligible parameter correlations. The combined statistical and systematic correlation matrix is given in Ref. [20].

A measurement of  $\phi_s$  and  $|\lambda|$  by LHCb using  $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$  decays of  $\phi_s^{\pi\pi} = 0.070 \pm 0.068 \pm 0.008$  rad and  $|\lambda^{\pi\pi}| = 0.89 \pm 0.05 \pm 0.01$ , consistent with the measurement reported here, was reported in Ref. [10]. The results from the two analyses are combined by incorporating the  $B_s^0 \rightarrow J/\psi K^+ K^-$  result as a correlated Gaussian constraint in the  $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$  fit, under the assumption that  $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$  and  $B_s^0 \rightarrow J/\psi K^+ K^-$  decays both proceed dominantly via  $b \rightarrow c\bar{c}s$  transitions and the ratio between loop-induced processes and tree diagrams are the same in each mode. The fit accounts for correlations between common parameters and correlations between systematic uncertainties. The combined result is  $\phi_s = -0.010 \pm 0.039$  rad and  $|\lambda| = 0.957 \pm 0.017$ . The correlation between the parameters is about  $-0.02$ .

In conclusion, the  $CP$ -violating phase  $\phi_s$ , and the  $B_s^0$  decay width parameters  $\Gamma_s$  and  $\Delta\Gamma_s$ , are measured using  $B_s^0 \rightarrow J/\psi K^+ K^-$  decays selected from the full LHCb data set from the first LHC operation period. The results are  $\phi_s = -0.058 \pm 0.049 \pm 0.006$  rad,  $|\lambda| = 0.964 \pm 0.019 \pm 0.007$ ,  $\Gamma_s = 0.6603 \pm 0.0027 \pm 0.0015$  ps $^{-1}$  and  $\Delta\Gamma_s = 0.0805 \pm 0.0091 \pm 0.0032$  ps $^{-1}$ . The parameter  $|\lambda|$  is consistent with unity, implying no evidence for  $CP$  violation in  $B_s^0 \rightarrow J/\psi K^+ K^-$  decays. For the first time, the polarisation-dependent  $CP$ -violating parameters are measured and show no significant difference between the four polarisation states. The measurements of  $\phi_s$  and  $|\lambda|$  in  $B_s^0 \rightarrow J/\psi K^+ K^-$  decays are consistent with those measured in  $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$  decays, and the combined results are

$\phi_s = -0.010 \pm 0.039$  rad and  $|\lambda| = 0.957 \pm 0.017$ . The measurement of the  $CP$  violating phase  $\phi_s$  is the most precise to date and is in agreement with the SM prediction [2], in which it is assumed that sub-leading contributions to the decay amplitude are negligible.

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); NSFC (China); CNRS/IN2P3 (France); BMBF, DFG, HGF and MPG (Germany); SFI (Ireland); INFN (Italy); FOM and NWO (The Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MinES and FANO (Russia); MinECo (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); NSF (USA). The Tier1 computing centres are supported by IN2P3 (France), KIT and BMBF (Germany), INFN (Italy), NWO and SURF (The Netherlands), PIC (Spain), GridPP (United Kingdom). We are indebted to the communities behind the multiple open source software packages on which we depend. We are also thankful for the computing resources and the access to software R&D tools provided by Yandex LLC (Russia). Individual groups or members have received support from EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union), Conseil général de Haute-Savoie, Labex ENIGMASS and OCEVU, Région Auvergne (France), RFBR (Russia), XuntaGal and GENCAT (Spain), Royal Society and Royal Commission for the Exhibition of 1851 (United Kingdom).

## References

- [1] M. Kobayashi and T. Maskawa, *CP violation in the renormalizable theory of weak interaction*, Prog. Theor. Phys. **49** (1973) 652; N. Cabibbo, *Unitary symmetry and leptonic decays*, Phys. Rev. Lett. **10** (1963) 531.
- [2] J. Charles *et al.*, *Predictions of selected flavour observables within the Standard Model*, Phys. Rev. **D84** (2011) 033005, [arXiv:1106.4041](https://arxiv.org/abs/1106.4041), with updated results and plots available at <http://ckmfitter.in2p3.fr>.
- [3] A. J. Buras, *Flavour theory: 2009*, PoS **EPS-HEP2009** (2009) 024, [arXiv:0910.1032](https://arxiv.org/abs/0910.1032).
- [4] C.-W. Chiang *et al.*, *New physics in  $B_s^0 \rightarrow J/\psi\phi$ : A general analysis*, JHEP **04** (2010) 031, [arXiv:0910.2929](https://arxiv.org/abs/0910.2929).
- [5] LHCb collaboration, R. Aaij *et al.*, *Measurement of the CP-violating phase  $\phi_s$  in the decay  $B_s^0 \rightarrow J/\psi\phi$* , Phys. Rev. Lett. **108** (2012) 101803, [arXiv:1112.3183](https://arxiv.org/abs/1112.3183).
- [6] LHCb collaboration, R. Aaij *et al.*, *Measurement of CP violation and the  $B_s^0$  meson decay width difference with  $B_s^0 \rightarrow J/\psi K^+ K^-$  and  $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$  decays*, Phys. Rev. **D87** (2013) 112010, [arXiv:1304.2600](https://arxiv.org/abs/1304.2600).

- [7] ATLAS Collaboration, G. Aad *et al.*, *Flavour tagged time dependent angular analysis of the  $B_s \rightarrow J/\psi\phi$  decay and extraction of  $\Delta\Gamma_s$  and the weak phase  $\phi_s$  in ATLAS*, Phys. Rev. **D90** (2014) 052007, [arXiv:1407.1796](#).
- [8] CDF collaboration, T. Aaltonen *et al.*, *Measurement of the bottom-strange meson mixing phase in the full CDF data set*, Phys. Rev. Lett. **109** (2012) 171802, [arXiv:1208.2967](#).
- [9] D0 collaboration, V. M. Abazov *et al.*, *Measurement of the CP-violating phase  $\phi_s^{J/\psi\phi}$  using the flavor-tagged decay  $B_s^0 \rightarrow J/\psi\phi$  in  $8\text{ fb}^{-1}$  of  $p\bar{p}$  collisions*, Phys. Rev. **D85** (2012) 032006, [arXiv:1109.3166](#).
- [10] LHCb collaboration, R. Aaij *et al.*, *Measurement of the CP-violating phase  $\phi_s$  in  $\bar{B}_s^0 \rightarrow J/\psi\pi^+\pi^-$  decays*, Phys. Lett. **B736** (2014) 186, [arXiv:1405.4140](#).
- [11] S. Faller, R. Fleischer, and T. Mannel, *Precision physics with  $B_s^0 \rightarrow J/\psi\phi$  at the LHC: the quest for new physics*, Phys. Rev. **D79** (2009) 014005, [arXiv:0810.4248](#).
- [12] LHCb collaboration, A. A. Alves Jr. *et al.*, *The LHCb detector at the LHC*, JINST **3** (2008) S08005.
- [13] R. Aaij *et al.*, *The LHCb trigger and its performance in 2011*, JINST **8** (2013) P04022, [arXiv:1211.3055](#).
- [14] S. Stone and L. Zhang, *S-waves and the measurement of CP violating phases in  $B_s^0$  decays*, Phys. Rev. **D79** (2009) 074024, [arXiv:0812.2832](#).
- [15] D. Martinez Santos and F. Dupertuis, *Mass distributions marginalized over per-event errors*, Nucl. Instrum. Meth. **A764** (2014) 150, [arXiv:1312.5000](#).
- [16] LHCb collaboration, R. Aaij *et al.*, *Measurement of the flavour-specific CP-violating asymmetry  $a_{\text{sl}}^s$  in  $B_s^0$  decays*, Phys. Lett. **B728** (2014) 607, [arXiv:1308.1048](#).
- [17] Y. Xie, *sFit: A method for background subtraction in maximum likelihood fit*, [arXiv:0905.0724](#).
- [18] LHCb collaboration, R. Aaij *et al.*, *Measurements of the  $B^+$ ,  $B^0$ ,  $B_s^0$  meson and  $\Lambda_b^0$  baryon lifetimes*, JHEP **04** (2014) 114, [arXiv:1402.2554](#).
- [19] LHCb collaboration, *Optimization and calibration of the same-side kaon tagging algorithm using hadronic  $B_s^0$  decays in 2011 data*, LHCb-CONF-2012-033.
- [20] See supplementary material for details.
- [21] LHCb collaboration, R. Aaij *et al.*, *Precision measurement of the  $B_s^0 - \bar{B}_s^0$  oscillation frequency in the decay  $B_s^0 \rightarrow D_s^+\pi^-$* , New J. Phys. **15** (2013) 053021, [arXiv:1304.4741](#).

- [22] X. Liu, W. Wang, and Y. Xie, *Penguin pollution in  $B \rightarrow J/\psi V$  decays and impact on the extraction of the  $B_s - \bar{B}_s$  mixing phase*, Phys. Rev. **D89** (2014) 094010, [arXiv:1309.0313](#).
- [23] LHCb collaboration, R. Aaij *et al.*, *Observation of the decay  $B_c^+ \rightarrow B_s^0 \pi^+$* , Phys. Rev. Lett. **111** (2013) 181801, [arXiv:1308.4544](#).
- [24] V. Kiselev, *Decays of the  $B_c$  meson*, [arXiv:hep-ph/0308214](#).

# 1 Supplementary material for PRL

Table 5: Statistical correlation matrix from the polarisation-independent fit.

	$\Gamma_s$	$\Delta\Gamma_s$	$ A_\perp ^2$	$ A_0 ^2$	$\delta_\parallel$	$\delta_\perp$	$\phi_s$	$ \lambda $	$\Delta m_s$
$\Gamma_s$	+1.00	-0.45	+0.39	-0.31	-0.07	-0.02	+0.01	-0.01	+0.01
$\Delta\Gamma_s$		+1.00	-0.69	+0.65	+0.02	-0.03	-0.08	+0.02	-0.03
$ A_\perp ^2$			+1.00	-0.59	-0.29	-0.10	+0.04	-0.03	+0.00
$ A_0 ^2$				+1.00	-0.02	-0.04	-0.03	+0.02	-0.03
$\delta_\parallel$					+1.00	+0.42	+0.01	+0.05	+0.05
$\delta_\perp$						+1.00	+0.14	-0.17	+0.67
$\phi_s$							+1.00	-0.02	+0.09
$ \lambda $								+1.00	-0.21
$\Delta m_s$									+1.00

Table 6: Total statistical and systematic correlation matrix from the polarisation-independent fit.

	$\Gamma_s$	$\Delta\Gamma_s$	$ A_\perp ^2$	$ A_0 ^2$	$\delta_\parallel$	$\delta_\perp$	$\phi_s$	$ \lambda $	$\Delta m_s$
$\Gamma_s$	+1.00	-0.30	+0.25	-0.09	-0.08	-0.01	+0.01	-0.01	+0.03
$\Delta\Gamma_s$		+1.00	-0.59	+0.36	-0.06	-0.05	-0.08	+0.03	-0.00
$ A_\perp ^2$			+1.00	-0.70	+0.03	+0.08	+0.05	+0.00	+0.02
$ A_0 ^2$				+1.00	-0.38	-0.28	-0.04	-0.01	-0.05
$\delta_\parallel$					+1.00	+0.47	+0.03	+0.00	+0.05
$\delta_\perp$						+1.00	+0.15	-0.18	+0.64
$\phi_s$							+1.00	-0.03	+0.09
$ \lambda $								+1.00	-0.21
$\Delta m_s$									+1.00

Table 7: Statistical correlation matrix for the  $\phi_s^k$  and  $|\lambda^k|$  parameters from the polarisation-dependent fit.

	$ \lambda^0 $	$ \lambda^\parallel/\lambda^0 $	$ \lambda^\perp/\lambda^0 $	$ \lambda^S/\lambda^0 $	$\phi_s^0$	$\phi_s^\parallel - \phi_s^0$	$\phi_s^\perp - \phi_s^0$	$\phi_s^S - \phi_s^0$
$ \lambda^0 $	+1.00	-0.32	-0.59	-0.89	+0.01	-0.08	-0.06	+0.00
$ \lambda^\parallel/\lambda^0 $		+1.00	-0.23	+0.27	+0.00	+0.31	+0.16	+0.10
$ \lambda^\perp/\lambda^0 $			+1.00	+0.53	-0.02	+0.06	-0.29	-0.02
$ \lambda^S/\lambda^0 $				+1.00	-0.01	+0.07	+0.06	+0.22
$\phi_s^0$					+1.00	-0.14	+0.13	+0.14
$\phi_s^\parallel - \phi_s^0$						+1.00	+0.52	+0.11
$\phi_s^\perp - \phi_s^0$							+1.00	+0.08
$\phi_s^S - \phi_s^0$								+1.00

Table 8: Total statistical and systematic correlation matrix for the  $\phi_s^k$  and  $|\lambda^k|$  parameters from the polarisation-dependent fit.

	$ \lambda^0 $	$ \lambda^\parallel/\lambda^0 $	$ \lambda^\perp/\lambda^0 $	$ \lambda^S/\lambda^0 $	$\phi_s^0$	$\phi_s^\parallel - \phi_s^0$	$\phi_s^\perp - \phi_s^0$	$\phi_s^S - \phi_s^0$
$ \lambda^0 $	+1.00	-0.35	-0.56	-0.77	-0.00	-0.09	-0.07	-0.02
$ \lambda^\parallel/\lambda^0 $		+1.00	-0.23	+0.15	+0.02	+0.32	+0.17	+0.14
$ \lambda^\perp/\lambda^0 $			+1.00	+0.51	-0.02	+0.05	-0.29	-0.03
$ \lambda^S/\lambda^0 $				+1.00	-0.03	+0.03	+0.04	+0.18
$\phi_s^0$					+1.00	-0.13	+0.13	+0.14
$\phi_s^\parallel - \phi_s^0$						+1.00	+0.51	+0.12
$\phi_s^\perp - \phi_s^0$							+1.00	+0.09
$\phi_s^S - \phi_s^0$								+1.00

Table 9: S-wave parameter estimates in each  $m(K^+K^-)$  bin from the polarisation-independent fit. Only statistical uncertainties are shown.

Parameter	Value
$F_{S1}$	0.426 $\pm$ 0.054
$F_{S2}$	0.059 $\pm$ 0.017
$F_{S3}$	0.0101 $\pm$ 0.0067
$F_{S4}$	0.0103 $\pm$ 0.0061
$F_{S5}$	0.049 $\pm$ 0.015
$F_{S6}$	0.193 $\pm$ 0.025
$\delta_{S1} - \delta_\perp$	0.84 $\pm$ 0.20
$\delta_{S2} - \delta_\perp$	2.15 $\pm$ 0.28
$\delta_{S3} - \delta_\perp$	0.47 $\pm$ 0.21
$\delta_{S4} - \delta_\perp$	-0.34 $\pm$ 0.17
$\delta_{S5} - \delta_\perp$	-0.59 $\pm$ 0.15
$\delta_{S6} - \delta_\perp$	-0.90 $\pm$ 0.14

## LHCb collaboration

R. Aaij<sup>41</sup>, B. Adeva<sup>37</sup>, M. Adinolfi<sup>46</sup>, A. Affolder<sup>52</sup>, Z. Ajaltouni<sup>5</sup>, S. Akar<sup>6</sup>, J. Albrecht<sup>9</sup>, F. Alessio<sup>38</sup>, M. Alexander<sup>51</sup>, S. Ali<sup>41</sup>, G. Alkhazov<sup>30</sup>, P. Alvarez Cartelle<sup>37</sup>, A.A. Alves Jr<sup>25,38</sup>, S. Amato<sup>2</sup>, S. Amerio<sup>22</sup>, Y. Amhis<sup>7</sup>, L. An<sup>3</sup>, L. Anderlini<sup>17,g</sup>, J. Anderson<sup>40</sup>, R. Andreassen<sup>57</sup>, M. Andreotti<sup>16,f</sup>, J.E. Andrews<sup>58</sup>, R.B. Appleby<sup>54</sup>, O. Aquines Gutierrez<sup>10</sup>, F. Archilli<sup>38</sup>, A. Artamonov<sup>35</sup>, M. Artuso<sup>59</sup>, E. Aslanides<sup>6</sup>, G. Auriemma<sup>25,n</sup>, M. Baalouch<sup>5</sup>, S. Bachmann<sup>11</sup>, J.J. Back<sup>48</sup>, A. Badalov<sup>36</sup>, C. Baesso<sup>60</sup>, W. Baldini<sup>16</sup>, R.J. Barlow<sup>54</sup>, C. Barschel<sup>38</sup>, S. Barsuk<sup>7</sup>, W. Barter<sup>47</sup>, V. Batozskaya<sup>28</sup>, V. Battista<sup>39</sup>, A. Bay<sup>39</sup>, L. Beaucourt<sup>4</sup>, J. Beddow<sup>51</sup>, F. Bedeschi<sup>23</sup>, I. Bediaga<sup>1</sup>, S. Belogurov<sup>31</sup>, K. Belous<sup>35</sup>, I. Belyaev<sup>31</sup>, E. Ben-Haim<sup>8</sup>, G. Bencivenni<sup>18</sup>, S. Benson<sup>38</sup>, J. Benton<sup>46</sup>, A. Berezhnoy<sup>32</sup>, R. Bernet<sup>40</sup>, A. Bertolin<sup>22</sup>, M.-O. Bettler<sup>47</sup>, M. van Beuzekom<sup>41</sup>, A. Bien<sup>11</sup>, S. Bifani<sup>45</sup>, T. Bird<sup>54</sup>, A. Bizzeti<sup>17,i</sup>, P.M. Bjørnstad<sup>54</sup>, T. Blake<sup>48</sup>, F. Blanc<sup>39</sup>, J. Blouw<sup>10</sup>, S. Blusk<sup>59</sup>, V. Bocci<sup>25</sup>, A. Bondar<sup>34</sup>, N. Bondar<sup>30,38</sup>, W. Bonivento<sup>15</sup>, S. Borghi<sup>54</sup>, A. Borgia<sup>59</sup>, M. Borsato<sup>7</sup>, T.J.V. Bowcock<sup>52</sup>, E. Bowen<sup>40</sup>, C. Bozzi<sup>16</sup>, D. Brett<sup>54</sup>, M. Britsch<sup>10</sup>, T. Britton<sup>59</sup>, J. Brodzicka<sup>54</sup>, N.H. Brook<sup>46</sup>, A. Bursche<sup>40</sup>, J. Buytaert<sup>38</sup>, S. Cadeddu<sup>15</sup>, R. Calabrese<sup>16,f</sup>, M. Calvi<sup>20,k</sup>, M. Calvo Gomez<sup>36,p</sup>, P. Campana<sup>18</sup>, D. Campora Perez<sup>38</sup>, A. Carbone<sup>14,d</sup>, G. Carboni<sup>24,l</sup>, R. Cardinale<sup>19,38,j</sup>, A. Cardini<sup>15</sup>, L. Carson<sup>50</sup>, K. Carvalho Akiba<sup>2,38</sup>, RCM Casanova Mohr<sup>36</sup>, G. Casse<sup>52</sup>, L. Cassina<sup>20,k</sup>, L. Castillo Garcia<sup>38</sup>, M. Cattaneo<sup>38</sup>, Ch. Cauet<sup>9</sup>, R. Cenci<sup>23,t</sup>, M. Charles<sup>8</sup>, Ph. Charpentier<sup>38</sup>, M. Chefdeville<sup>4</sup>, S. Chen<sup>54</sup>, S.-F. Cheung<sup>55</sup>, N. Chiapolini<sup>40</sup>, M. Chrzasteczko<sup>40,26</sup>, X. Cid Vidal<sup>38</sup>, G. Ciezarek<sup>41</sup>, P.E.L. Clarke<sup>50</sup>, M. Clemencic<sup>38</sup>, H.V. Cliff<sup>47</sup>, J. Closier<sup>38</sup>, V. Coco<sup>38</sup>, J. Cogan<sup>6</sup>, E. Cogneras<sup>5</sup>, V. Cogoni<sup>15</sup>, L. Cojocariu<sup>29</sup>, G. Collazuol<sup>22</sup>, P. Collins<sup>38</sup>, A. Comerma-Montells<sup>11</sup>, A. Contu<sup>15,38</sup>, A. Cook<sup>46</sup>, M. Coombes<sup>46</sup>, S. Coquereau<sup>8</sup>, G. Corti<sup>38</sup>, M. Corvo<sup>16,f</sup>, I. Counts<sup>56</sup>, B. Couturier<sup>38</sup>, G.A. Cowan<sup>50</sup>, D.C. Craik<sup>48</sup>, A.C. Crocombe<sup>48</sup>, M. Cruz Torres<sup>60</sup>, S. Cunliffe<sup>53</sup>, R. Currie<sup>53</sup>, C. D'Ambrosio<sup>38</sup>, J. Dalseno<sup>46</sup>, P. David<sup>8</sup>, P.N.Y. David<sup>41</sup>, A. Davis<sup>57</sup>, K. De Bruyn<sup>41</sup>, S. De Capua<sup>54</sup>, M. De Cian<sup>11</sup>, J.M. De Miranda<sup>1</sup>, L. De Paula<sup>2</sup>, W. De Silva<sup>57</sup>, P. De Simone<sup>18</sup>, C.-T. Dean<sup>51</sup>, D. Decamp<sup>4</sup>, M. Deckenhoff<sup>9</sup>, L. Del Buono<sup>8</sup>, N. Déléage<sup>4</sup>, D. Derkach<sup>55</sup>, O. Deschamps<sup>5</sup>, F. Dettori<sup>38</sup>, A. Di Canto<sup>38</sup>, A Di Domenico<sup>25</sup>, H. Dijkstra<sup>38</sup>, S. Donleavy<sup>52</sup>, F. Dordei<sup>11</sup>, M. Dorigo<sup>39</sup>, A. Dosil Suárez<sup>37</sup>, D. Dossett<sup>48</sup>, A. Dovbnya<sup>43</sup>, K. Dreimanis<sup>52</sup>, G. Dujany<sup>54</sup>, F. Dupertuis<sup>39</sup>, P. Durante<sup>38</sup>, R. Dzhelyadin<sup>35</sup>, A. Dziurda<sup>26</sup>, A. Dzyuba<sup>30</sup>, S. Easo<sup>49,38</sup>, U. Egede<sup>53</sup>, V. Egorychev<sup>31</sup>, S. Eidelman<sup>34</sup>, S. Eisenhardt<sup>50</sup>, U. Eitschberger<sup>9</sup>, R. Ekelhof<sup>9</sup>, L. Eklund<sup>51</sup>, I. El Rifai<sup>5</sup>, Ch. Elsasser<sup>40</sup>, S. Ely<sup>59</sup>, S. Esen<sup>11</sup>, H.-M. Evans<sup>47</sup>, T. Evans<sup>55</sup>, A. Falabella<sup>14</sup>, C. Färber<sup>11</sup>, C. Farinelli<sup>41</sup>, N. Farley<sup>45</sup>, S. Farry<sup>52</sup>, R. Fay<sup>52</sup>, D. Ferguson<sup>50</sup>, V. Fernandez Albor<sup>37</sup>, F. Ferreira Rodrigues<sup>1</sup>, M. Ferro-Luzzi<sup>38</sup>, S. Filippov<sup>33</sup>, M. Fiore<sup>16,f</sup>, M. Fiorini<sup>16,f</sup>, M. Firlej<sup>27</sup>, C. Fitzpatrick<sup>39</sup>, T. Fiutowski<sup>27</sup>, P. Fol<sup>53</sup>, M. Fontana<sup>10</sup>, F. Fontanelli<sup>19,j</sup>, R. Forty<sup>38</sup>, O. Francisco<sup>2</sup>, M. Frank<sup>38</sup>, C. Frei<sup>38</sup>, M. Frosini<sup>17</sup>, J. Fu<sup>21,38</sup>, E. Furfaro<sup>24,l</sup>, A. Gallas Torreira<sup>37</sup>, D. Galli<sup>14,d</sup>, S. Gallorini<sup>22,38</sup>, S. Gambetta<sup>19,j</sup>, M. Gandelman<sup>2</sup>, P. Gandini<sup>59</sup>, Y. Gao<sup>3</sup>, J. García Pardiñas<sup>37</sup>, J. Garofoli<sup>59</sup>, J. Garra Tico<sup>47</sup>, L. Garrido<sup>36</sup>, D. Gascon<sup>36</sup>, C. Gaspar<sup>38</sup>, U. Gastaldi<sup>16</sup>, R. Gauld<sup>55</sup>, L. Gavardi<sup>9</sup>, G. Gazzoni<sup>5</sup>, A. Geraci<sup>21,v</sup>, E. Gersabeck<sup>11</sup>, M. Gersabeck<sup>54</sup>, T. Gershon<sup>48</sup>, Ph. Ghez<sup>4</sup>, A. Gianelle<sup>22</sup>, S. Gianì<sup>39</sup>, V. Gibson<sup>47</sup>, L. Giubega<sup>29</sup>, V.V. Gligorov<sup>38</sup>, C. Göbel<sup>60</sup>, D. Golubkov<sup>31</sup>, A. Golutvin<sup>53,31,38</sup>, A. Gomes<sup>1,a</sup>, C. Gotti<sup>20,k</sup>, M. Grabalosa Gándara<sup>5</sup>, R. Graciani Diaz<sup>36</sup>, L.A. Granado Cardoso<sup>38</sup>, E. Graugés<sup>36</sup>, E. Graverini<sup>40</sup>, G. Graziani<sup>17</sup>, A. Grecu<sup>29</sup>, E. Greening<sup>55</sup>, S. Gregson<sup>47</sup>, P. Griffith<sup>45</sup>, L. Grillo<sup>11</sup>, O. Grünberg<sup>63</sup>, B. Gui<sup>59</sup>, E. Gushchin<sup>33</sup>, Yu. Guz<sup>35,38</sup>, T. Gys<sup>38</sup>, C. Hadjivasiliou<sup>59</sup>, G. Haefeli<sup>39</sup>, C. Haen<sup>38</sup>,

S.C. Haines<sup>47</sup>, S. Hall<sup>53</sup>, B. Hamilton<sup>58</sup>, T. Hampson<sup>46</sup>, X. Han<sup>11</sup>, S. Hansmann-Menzemer<sup>11</sup>,  
 N. Harnew<sup>55</sup>, S.T. Harnew<sup>46</sup>, J. Harrison<sup>54</sup>, J. He<sup>38</sup>, T. Head<sup>39</sup>, V. Heijne<sup>41</sup>, K. Hennessy<sup>52</sup>,  
 P. Henrard<sup>5</sup>, L. Henry<sup>8</sup>, J.A. Hernando Morata<sup>37</sup>, E. van Herwijnen<sup>38</sup>, M. Heß<sup>63</sup>, A. Hicheur<sup>2</sup>,  
 D. Hill<sup>55</sup>, M. Hoballah<sup>5</sup>, C. Hombach<sup>54</sup>, W. Hulsbergen<sup>41</sup>, N. Hussain<sup>55</sup>, D. Hutchcroft<sup>52</sup>,  
 D. Hynds<sup>51</sup>, M. Idzik<sup>27</sup>, P. Ilten<sup>56</sup>, R. Jacobsson<sup>38</sup>, A. Jaeger<sup>11</sup>, J. Jalocha<sup>55</sup>, E. Jans<sup>41</sup>,  
 P. Jatou<sup>39</sup>, A. Jawahery<sup>58</sup>, F. Jing<sup>3</sup>, M. John<sup>55</sup>, D. Johnson<sup>38</sup>, C.R. Jones<sup>47</sup>, C. Joram<sup>38</sup>,  
 B. Jost<sup>38</sup>, N. Jurik<sup>59</sup>, S. Kandybei<sup>43</sup>, W. Kanso<sup>6</sup>, M. Karacson<sup>38</sup>, T.M. Karbach<sup>38</sup>,  
 S. Karodia<sup>51</sup>, M. Kelsey<sup>59</sup>, I.R. Kenyon<sup>45</sup>, T. Ketel<sup>42</sup>, B. Khanji<sup>20,38,k</sup>, C. Khurewathanakul<sup>39</sup>,  
 S. Klaver<sup>54</sup>, K. Klimaszewski<sup>28</sup>, O. Kochebina<sup>7</sup>, M. Kolpin<sup>11</sup>, I. Komarov<sup>39</sup>, R.F. Koopman<sup>42</sup>,  
 P. Koppenburg<sup>41,38</sup>, M. Korolev<sup>32</sup>, L. Kravchuk<sup>33</sup>, K. Kreplin<sup>11</sup>, M. Kreps<sup>48</sup>, G. Krocker<sup>11</sup>,  
 P. Krokovny<sup>34</sup>, F. Kruse<sup>9</sup>, W. Kucewicz<sup>26,o</sup>, M. Kucharczyk<sup>20,26,k</sup>, V. Kudryavtsev<sup>34</sup>,  
 K. Kurek<sup>28</sup>, T. Kvaratskheliya<sup>31</sup>, V.N. La Thi<sup>39</sup>, D. Lacarrere<sup>38</sup>, G. Lafferty<sup>54</sup>, A. Lai<sup>15</sup>,  
 D. Lambert<sup>50</sup>, R.W. Lambert<sup>42</sup>, G. Lanfranchi<sup>18</sup>, C. Langenbruch<sup>48</sup>, B. Langhans<sup>38</sup>,  
 T. Latham<sup>48</sup>, C. Lazzeroni<sup>45</sup>, R. Le Gac<sup>6</sup>, J. van Leerdam<sup>41</sup>, J.-P. Lees<sup>4</sup>, R. Lefèvre<sup>5</sup>,  
 A. Leflat<sup>32</sup>, J. Lefrançois<sup>7</sup>, O. Leroy<sup>6</sup>, T. Lesiak<sup>26</sup>, B. Leverington<sup>11</sup>, Y. Li<sup>7</sup>,  
 T. Likhomanenko<sup>64</sup>, M. Liles<sup>52</sup>, R. Lindner<sup>38</sup>, C. Linn<sup>38</sup>, F. Lionetto<sup>40</sup>, B. Liu<sup>15</sup>, S. Lohn<sup>38</sup>,  
 I. Longstaff<sup>51</sup>, J.H. Lopes<sup>2</sup>, P. Lowdon<sup>40</sup>, D. Lucchesi<sup>22,r</sup>, H. Luo<sup>50</sup>, A. Lupato<sup>22</sup>, E. Luppi<sup>16,f</sup>,  
 O. Lupton<sup>55</sup>, F. Machefert<sup>7</sup>, I.V. Machikhiliyan<sup>31</sup>, F. Maciuc<sup>29</sup>, O. Maev<sup>30</sup>, S. Malde<sup>55</sup>,  
 A. Malinin<sup>64</sup>, G. Manca<sup>15,e</sup>, G. Mancinelli<sup>6</sup>, A. Mapelli<sup>38</sup>, J. Maratas<sup>5</sup>, J.F. Marchand<sup>4</sup>,  
 U. Marconi<sup>14</sup>, C. Marin Benito<sup>36</sup>, P. Marino<sup>23,t</sup>, R. Märki<sup>39</sup>, J. Marks<sup>11</sup>, G. Martellotti<sup>25</sup>,  
 M. Martinelli<sup>39</sup>, D. Martinez Santos<sup>42</sup>, F. Martinez Vidal<sup>65</sup>, D. Martins Tostes<sup>2</sup>,  
 A. Massafferri<sup>1</sup>, R. Matev<sup>38</sup>, Z. Mathe<sup>38</sup>, C. Matteuzzi<sup>20</sup>, A. Mazurov<sup>45</sup>, M. McCann<sup>53</sup>,  
 J. McCarthy<sup>45</sup>, A. McNab<sup>54</sup>, R. McNulty<sup>12</sup>, B. McSkelly<sup>52</sup>, B. Meadows<sup>57</sup>, F. Meier<sup>9</sup>,  
 M. Meissner<sup>11</sup>, M. Merk<sup>41</sup>, D.A. Milanese<sup>62</sup>, M.-N. Minard<sup>4</sup>, N. Moggi<sup>14</sup>, J. Molina Rodriguez<sup>60</sup>,  
 S. Monteil<sup>5</sup>, M. Morandin<sup>22</sup>, P. Morawski<sup>27</sup>, A. Mordà<sup>6</sup>, M.J. Morello<sup>23,t</sup>, J. Moron<sup>27</sup>,  
 A.-B. Morris<sup>50</sup>, R. Mountain<sup>59</sup>, F. Muheim<sup>50</sup>, K. Müller<sup>40</sup>, M. Mussini<sup>14</sup>, B. Muster<sup>39</sup>,  
 P. Naik<sup>46</sup>, T. Nakada<sup>39</sup>, R. Nandakumar<sup>49</sup>, I. Nasteva<sup>2</sup>, M. Needham<sup>50</sup>, N. Neri<sup>21</sup>,  
 S. Neubert<sup>38</sup>, N. Neufeld<sup>38</sup>, M. Neuner<sup>11</sup>, A.D. Nguyen<sup>39</sup>, T.D. Nguyen<sup>39</sup>, C. Nguyen-Mau<sup>39,q</sup>,  
 M. Nicol<sup>7</sup>, V. Niess<sup>5</sup>, R. Niet<sup>9</sup>, N. Nikitin<sup>32</sup>, T. Nikodem<sup>11</sup>, A. Novoselov<sup>35</sup>, D.P. O’Hanlon<sup>48</sup>,  
 A. Oblakowska-Mucha<sup>27</sup>, V. Obraztsov<sup>35</sup>, S. Ogilvy<sup>51</sup>, O. Okhrimenko<sup>44</sup>, R. Oldeman<sup>15,e</sup>,  
 C.J.G. Onderwater<sup>66</sup>, M. Orlandea<sup>29</sup>, J.M. Otalora Goicochea<sup>2</sup>, A. Otto<sup>38</sup>, P. Owen<sup>53</sup>,  
 A. Oyanguren<sup>65</sup>, B.K. Pal<sup>59</sup>, A. Palano<sup>13,c</sup>, F. Palombo<sup>21,u</sup>, M. Palutan<sup>18</sup>, J. Panman<sup>38</sup>,  
 A. Papanestis<sup>49,38</sup>, M. Pappagallo<sup>51</sup>, L.L. Pappalardo<sup>16,f</sup>, C. Parkes<sup>54</sup>, C.J. Parkinson<sup>9,45</sup>,  
 G. Passaleva<sup>17</sup>, G.D. Patel<sup>52</sup>, M. Patel<sup>53</sup>, C. Patrignani<sup>19,j</sup>, A. Pearce<sup>54</sup>, A. Pellegrino<sup>41</sup>,  
 G. Penso<sup>25,m</sup>, M. Pepe Altarelli<sup>38</sup>, S. Perazzini<sup>14,d</sup>, P. Perret<sup>5</sup>, L. Pescatore<sup>45</sup>, E. Pesen<sup>67</sup>,  
 K. Petridis<sup>53</sup>, A. Petrolini<sup>19,j</sup>, E. Picatoste Olloqui<sup>36</sup>, B. Pietrzyk<sup>4</sup>, T. Pilař<sup>48</sup>, D. Pinci<sup>25</sup>,  
 A. Pistone<sup>19</sup>, S. Playfer<sup>50</sup>, M. Plo Casasus<sup>37</sup>, F. Polci<sup>8</sup>, A. Poluektov<sup>48,34</sup>, I. Polyakov<sup>31</sup>,  
 E. Polcarpo<sup>2</sup>, A. Popov<sup>35</sup>, D. Popov<sup>10</sup>, B. Popovici<sup>29</sup>, C. Potterat<sup>2</sup>, E. Price<sup>46</sup>, J.D. Price<sup>52</sup>,  
 J. Prisciandaro<sup>39</sup>, A. Pritchard<sup>52</sup>, C. Prouve<sup>46</sup>, V. Pugatch<sup>44</sup>, A. Puig Navarro<sup>39</sup>, G. Punzi<sup>23,s</sup>,  
 W. Qian<sup>4</sup>, B. Rachwal<sup>26</sup>, J.H. Rademacker<sup>46</sup>, B. Rakotomiaramanana<sup>39</sup>, M. Rama<sup>23</sup>,  
 M.S. Rangel<sup>2</sup>, I. Raniuk<sup>43</sup>, N. Rauschmayr<sup>38</sup>, G. Raven<sup>42</sup>, F. Redi<sup>53</sup>, S. Reichert<sup>54</sup>,  
 M.M. Reid<sup>48</sup>, A.C. dos Reis<sup>1</sup>, S. Ricciardi<sup>49</sup>, S. Richards<sup>46</sup>, M. Rihl<sup>38</sup>, K. Rinnert<sup>52</sup>,  
 V. Rives Molina<sup>36</sup>, P. Robbe<sup>7</sup>, A.B. Rodrigues<sup>1</sup>, E. Rodrigues<sup>54</sup>, P. Rodriguez Perez<sup>54</sup>,  
 S. Roiser<sup>38</sup>, V. Romanovsky<sup>35</sup>, A. Romero Vidal<sup>37</sup>, M. Rotondo<sup>22</sup>, J. Rouvinet<sup>39</sup>, T. Rul<sup>38</sup>,  
 H. Ruiz<sup>36</sup>, P. Ruiz Valls<sup>65</sup>, J.J. Saborido Silva<sup>37</sup>, N. Sagidova<sup>30</sup>, P. Sail<sup>51</sup>, B. Saitta<sup>15,e</sup>,  
 V. Salustino Guimaraes<sup>2</sup>, C. Sanchez Mayordomo<sup>65</sup>, B. Sanmartin Sedes<sup>37</sup>, R. Santacesaria<sup>25</sup>,



C. Santamarina Rios<sup>37</sup>, E. Santovetti<sup>24,l</sup>, A. Sarti<sup>18,m</sup>, C. Satriano<sup>25,n</sup>, A. Satta<sup>24</sup>,  
D.M. Saunders<sup>46</sup>, D. Savrina<sup>31,32</sup>, M. Schiller<sup>38</sup>, H. Schindler<sup>38</sup>, M. Schlupp<sup>9</sup>, M. Schmelling<sup>10</sup>,  
B. Schmidt<sup>38</sup>, O. Schneider<sup>39</sup>, A. Schopper<sup>38</sup>, M.-H. Schune<sup>7</sup>, R. Schwemmer<sup>38</sup>, B. Sciascia<sup>18</sup>,  
A. Sciubba<sup>25,m</sup>, A. Semennikov<sup>31</sup>, I. Sepp<sup>53</sup>, N. Serra<sup>40</sup>, J. Serrano<sup>6</sup>, L. Sestini<sup>22</sup>, P. Seyfert<sup>11</sup>,  
M. Shapkin<sup>35</sup>, I. Shapoval<sup>16,43,f</sup>, Y. Shcheglov<sup>30</sup>, T. Shears<sup>52</sup>, L. Shekhtman<sup>34</sup>,  
V. Shevchenko<sup>64</sup>, A. Shires<sup>9</sup>, R. Silva Coutinho<sup>48</sup>, G. Simi<sup>22</sup>, M. Sirendi<sup>47</sup>, N. Skidmore<sup>46</sup>,  
I. Skillicorn<sup>51</sup>, T. Skwarnicki<sup>59</sup>, N.A. Smith<sup>52</sup>, E. Smith<sup>55,49</sup>, E. Smith<sup>53</sup>, J. Smith<sup>47</sup>,  
M. Smith<sup>54</sup>, H. Snoek<sup>41</sup>, M.D. Sokoloff<sup>57</sup>, F.J.P. Soler<sup>51</sup>, F. Soomro<sup>39</sup>, D. Souza<sup>46</sup>,  
B. Souza De Paula<sup>2</sup>, B. Spaan<sup>9</sup>, P. Spradlin<sup>51</sup>, S. Sridharan<sup>38</sup>, F. Stagni<sup>38</sup>, M. Stahl<sup>11</sup>,  
S. Stahl<sup>11</sup>, O. Steinkamp<sup>40</sup>, O. Stenyakin<sup>35</sup>, F. Sterpka<sup>59</sup>, S. Stevenson<sup>55</sup>, S. Stoica<sup>29</sup>,  
S. Stone<sup>59</sup>, B. Storaci<sup>40</sup>, S. Stracka<sup>23,t</sup>, M. Straticiuc<sup>29</sup>, U. Straumann<sup>40</sup>, R. Stroili<sup>22</sup>, L. Sun<sup>57</sup>,  
W. Sutcliffe<sup>53</sup>, K. Swientek<sup>27</sup>, S. Swientek<sup>9</sup>, V. Syropoulos<sup>42</sup>, M. Szczekowski<sup>28</sup>,  
P. Szczypka<sup>39,38</sup>, T. Szumlak<sup>27</sup>, S. T'Jampens<sup>4</sup>, M. Teklishyn<sup>7</sup>, G. Tellarini<sup>16,f</sup>, F. Teubert<sup>38</sup>,  
C. Thomas<sup>55</sup>, E. Thomas<sup>38</sup>, J. van Tilburg<sup>41</sup>, V. Tisserand<sup>4</sup>, M. Tobin<sup>39</sup>, J. Todd<sup>57</sup>, S. Tolk<sup>42</sup>,  
L. Tomassetti<sup>16,f</sup>, D. Tonelli<sup>38</sup>, S. Topp-Joergensen<sup>55</sup>, N. Torr<sup>55</sup>, E. Tournefier<sup>4</sup>, S. Tourneur<sup>39</sup>,  
M.T. Tran<sup>39</sup>, M. Tresch<sup>40</sup>, A. Trisovic<sup>38</sup>, A. Tsaregorodtsev<sup>6</sup>, P. Tsopelas<sup>41</sup>, N. Tuning<sup>41</sup>,  
M. Ubeda Garcia<sup>38</sup>, A. Ukleja<sup>28</sup>, A. Ustyuzhanin<sup>64</sup>, U. Uwer<sup>11</sup>, C. Vacca<sup>15</sup>, V. Vagnoni<sup>14</sup>,  
G. Valenti<sup>14</sup>, A. Vallier<sup>7</sup>, R. Vazquez Gomez<sup>18</sup>, P. Vazquez Regueiro<sup>37</sup>, C. Vázquez Sierra<sup>37</sup>,  
S. Vecchi<sup>16</sup>, J.J. Velthuis<sup>46</sup>, M. Veltri<sup>17,h</sup>, G. Veneziano<sup>39</sup>, M. Vesterinen<sup>11</sup>,  
JVVB Viana Barbosa<sup>38</sup>, B. Viaud<sup>7</sup>, D. Vieira<sup>2</sup>, M. Vieites Diaz<sup>37</sup>, X. Vilasis-Cardona<sup>36,p</sup>,  
A. Vollhardt<sup>40</sup>, D. Volyanskyy<sup>10</sup>, D. Voong<sup>46</sup>, A. Vorobyev<sup>30</sup>, V. Vorobyev<sup>34</sup>, C. Voß<sup>63</sup>,  
J.A. de Vries<sup>41</sup>, R. Waldi<sup>63</sup>, C. Wallace<sup>48</sup>, R. Wallace<sup>12</sup>, J. Walsh<sup>23</sup>, S. Wandernoth<sup>11</sup>,  
J. Wang<sup>59</sup>, D.R. Ward<sup>47</sup>, N.K. Watson<sup>45</sup>, D. Websdale<sup>53</sup>, M. Whitehead<sup>48</sup>, D. Wiedner<sup>11</sup>,  
G. Wilkinson<sup>55,38</sup>, M. Wilkinson<sup>59</sup>, M.P. Williams<sup>45</sup>, M. Williams<sup>56</sup>, H.W. Wilschut<sup>66</sup>,  
F.F. Wilson<sup>49</sup>, J. Wimberley<sup>58</sup>, J. Wishahi<sup>9</sup>, W. Wislicki<sup>28</sup>, M. Witek<sup>26</sup>, G. Wormser<sup>7</sup>,  
S.A. Wotton<sup>47</sup>, S. Wright<sup>47</sup>, K. Wyllie<sup>38</sup>, Y. Xie<sup>61</sup>, Z. Xing<sup>59</sup>, Z. Xu<sup>39</sup>, Z. Yang<sup>3</sup>, X. Yuan<sup>3</sup>,  
O. Yushchenko<sup>35</sup>, M. Zangoli<sup>14</sup>, M. Zavertyaev<sup>10,b</sup>, L. Zhang<sup>3</sup>, W.C. Zhang<sup>12</sup>, Y. Zhang<sup>3</sup>,  
A. Zhelezov<sup>11</sup>, A. Zhokhov<sup>31</sup>, L. Zhong<sup>3</sup>.

<sup>1</sup>Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

<sup>2</sup>Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

<sup>3</sup>Center for High Energy Physics, Tsinghua University, Beijing, China

<sup>4</sup>LAPP, Université de Savoie, CNRS/IN2P3, Annecy-Le-Vieux, France

<sup>5</sup>Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, Clermont-Ferrand, France

<sup>6</sup>CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France

<sup>7</sup>LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France

<sup>8</sup>LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris, France

<sup>9</sup>Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany

<sup>10</sup>Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany

<sup>11</sup>Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

<sup>12</sup>School of Physics, University College Dublin, Dublin, Ireland

<sup>13</sup>Sezione INFN di Bari, Bari, Italy

<sup>14</sup>Sezione INFN di Bologna, Bologna, Italy

<sup>15</sup>Sezione INFN di Cagliari, Cagliari, Italy

<sup>16</sup>Sezione INFN di Ferrara, Ferrara, Italy

<sup>17</sup>Sezione INFN di Firenze, Firenze, Italy

<sup>18</sup>Laboratori Nazionali dell'INFN di Frascati, Frascati, Italy

<sup>19</sup>Sezione INFN di Genova, Genova, Italy

- <sup>20</sup> *Sezione INFN di Milano Bicocca, Milano, Italy*
- <sup>21</sup> *Sezione INFN di Milano, Milano, Italy*
- <sup>22</sup> *Sezione INFN di Padova, Padova, Italy*
- <sup>23</sup> *Sezione INFN di Pisa, Pisa, Italy*
- <sup>24</sup> *Sezione INFN di Roma Tor Vergata, Roma, Italy*
- <sup>25</sup> *Sezione INFN di Roma La Sapienza, Roma, Italy*
- <sup>26</sup> *Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*
- <sup>27</sup> *AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland*
- <sup>28</sup> *National Center for Nuclear Research (NCBJ), Warsaw, Poland*
- <sup>29</sup> *Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*
- <sup>30</sup> *Petersburg Nuclear Physics Institute (PNPI), Gatchina, Russia*
- <sup>31</sup> *Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia*
- <sup>32</sup> *Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia*
- <sup>33</sup> *Institute for Nuclear Research of the Russian Academy of Sciences (INR RAN), Moscow, Russia*
- <sup>34</sup> *Budker Institute of Nuclear Physics (SB RAS) and Novosibirsk State University, Novosibirsk, Russia*
- <sup>35</sup> *Institute for High Energy Physics (IHEP), Protvino, Russia*
- <sup>36</sup> *Universitat de Barcelona, Barcelona, Spain*
- <sup>37</sup> *Universidad de Santiago de Compostela, Santiago de Compostela, Spain*
- <sup>38</sup> *European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- <sup>39</sup> *Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- <sup>40</sup> *Physik-Institut, Universität Zürich, Zürich, Switzerland*
- <sup>41</sup> *Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands*
- <sup>42</sup> *Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, The Netherlands*
- <sup>43</sup> *NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- <sup>44</sup> *Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- <sup>45</sup> *University of Birmingham, Birmingham, United Kingdom*
- <sup>46</sup> *H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- <sup>47</sup> *Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- <sup>48</sup> *Department of Physics, University of Warwick, Coventry, United Kingdom*
- <sup>49</sup> *STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- <sup>50</sup> *School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- <sup>51</sup> *School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- <sup>52</sup> *Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- <sup>53</sup> *Imperial College London, London, United Kingdom*
- <sup>54</sup> *School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- <sup>55</sup> *Department of Physics, University of Oxford, Oxford, United Kingdom*
- <sup>56</sup> *Massachusetts Institute of Technology, Cambridge, MA, United States*
- <sup>57</sup> *University of Cincinnati, Cincinnati, OH, United States*
- <sup>58</sup> *University of Maryland, College Park, MD, United States*
- <sup>59</sup> *Syracuse University, Syracuse, NY, United States*
- <sup>60</sup> *Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to <sup>2</sup>*
- <sup>61</sup> *Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China, associated to <sup>3</sup>*
- <sup>62</sup> *Departamento de Física, Universidad Nacional de Colombia, Bogota, Colombia, associated to <sup>8</sup>*
- <sup>63</sup> *Institut für Physik, Universität Rostock, Rostock, Germany, associated to <sup>11</sup>*
- <sup>64</sup> *National Research Centre Kurchatov Institute, Moscow, Russia, associated to <sup>31</sup>*
- <sup>65</sup> *Instituto de Física Corpuscular (IFIC), Universitat de Valencia-CSIC, Valencia, Spain, associated to <sup>36</sup>*
- <sup>66</sup> *Van Swinderen Institute, University of Groningen, Groningen, The Netherlands, associated to <sup>41</sup>*
- <sup>67</sup> *Celal Bayar University, Manisa, Turkey, associated to <sup>38</sup>*

- <sup>a</sup> *Universidade Federal do Triângulo Mineiro (UFMT), Uberaba-MG, Brazil*
- <sup>b</sup> *P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia*
- <sup>c</sup> *Università di Bari, Bari, Italy*
- <sup>d</sup> *Università di Bologna, Bologna, Italy*
- <sup>e</sup> *Università di Cagliari, Cagliari, Italy*
- <sup>f</sup> *Università di Ferrara, Ferrara, Italy*
- <sup>g</sup> *Università di Firenze, Firenze, Italy*
- <sup>h</sup> *Università di Urbino, Urbino, Italy*
- <sup>i</sup> *Università di Modena e Reggio Emilia, Modena, Italy*
- <sup>j</sup> *Università di Genova, Genova, Italy*
- <sup>k</sup> *Università di Milano Bicocca, Milano, Italy*
- <sup>l</sup> *Università di Roma Tor Vergata, Roma, Italy*
- <sup>m</sup> *Università di Roma La Sapienza, Roma, Italy*
- <sup>n</sup> *Università della Basilicata, Potenza, Italy*
- <sup>o</sup> *AGH - University of Science and Technology, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland*
- <sup>p</sup> *LIFAELS, La Salle, Universitat Ramon Llull, Barcelona, Spain*
- <sup>q</sup> *Hanoi University of Science, Hanoi, Viet Nam*
- <sup>r</sup> *Università di Padova, Padova, Italy*
- <sup>s</sup> *Università di Pisa, Pisa, Italy*
- <sup>t</sup> *Scuola Normale Superiore, Pisa, Italy*
- <sup>u</sup> *Università degli Studi di Milano, Milano, Italy*
- <sup>v</sup> *Politecnico di Milano, Milano, Italy*