

Production of strange particles in charged jets in p–Pb and Pb–Pb collisions measured with ALICE at the LHC

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2017 J. Phys.: Conf. Ser. 805 012009

(<http://iopscience.iop.org/1742-6596/805/1/012009>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 131.169.5.251

This content was downloaded on 10/03/2017 at 20:38

Please note that [terms and conditions apply](#).

You may also be interested in:

[Open heavy-flavour measurements in p–Pb and Pb–Pb collisions with ALICE at the LHC](#)
Cristina Terrevoli and ALICE Collaboration

[Recent results on ultra-peripheral Pb–Pb and p–Pb collisions at the ALICE experiment](#)
J D Tapia Takaki

[Mid-rapidity D-meson production in pp, Pb–Pb and p–Pb collisions at the LHC](#)
Jeremy Wilkinson and the ALICE Collaboration

[Measurement of dielectrons in pp, p–Pb and Pb–Pb collisions with ALICE at the LHC](#)
Alberto Caliva and ALICE Collaboration

[D meson-hadron angular correlations in pp and p–Pb collisions with ALICE at the LHC](#)
Fabio Colamaria and the Alice collaboration

[Strangeness production in p–Pb and Pb–Pb collisions with ALICE at LHC](#)
Domenico Colella and ALICE Collaboration

[Charmonium production at mid-rapidity in Pb–Pb and p–Pb collisions with ALICE](#)
Steffen Georg Weber

[Identified charged hadron production in Pb–Pb collisions with event shape engineering](#)
Marco Antonio Tangaro and ALICE collaboration

[Low mass vector meson production in pp and Pb–Pb collisions measured with ALICE at the LHC](#)
Alessandro De Falco and the Alice Collaboration

Production of strange particles in charged jets in p–Pb and Pb–Pb collisions measured with ALICE at the LHC

Alice Zimmermann¹ (for the ALICE Collaboration)

1. Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Im Neuenheimer Feld 226, 69120 Heidelberg, Germany

alice.zimmermann@cern.ch

Abstract. Studies of jet production can provide information about the properties of the hot and dense strongly interacting matter created in ultra-relativistic heavy-ion collisions. Specifically, measurement of strange particles in jets may clarify the role of fragmentation processes in the anomalous baryon to meson ratio at intermediate particle p_T that was observed in lead-lead (Pb–Pb) and, to a lesser extent, in proton-lead (p–Pb) collisions.

In this contribution, measurements are presented of the p_T spectra of $\Lambda(\bar{\Lambda})$ baryons and K_S^0 mesons produced in association with charged jets in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV and p–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV. The analysis is based on data recorded by ALICE at the LHC, exploiting its excellent particle identification capabilities. The baryon/meson ratios of strange particles associated with jets are studied for different event activities in p–Pb collisions and are restricted to central events in Pb–Pb collisions. A comparison is shown to the ratios obtained for inclusive particle production and for particles stemming from the underlying event as well as to PYTHIA proton-proton (pp) simulations.

1. Introduction

The first measurements of the baryon-meson ratio in heavy-ion collisions at the Relativistic Heavy Ion Collider (RHIC) showed an enhanced baryon/meson production at intermediate transverse momentum ($p_T = 3$ GeV/ c) [1, 2] relative to pp collisions. Recent results at LHC energy corroborated this observation [3, 4]. There are several scenarios proposed to explain this, e.g. collective effects and string fragmentation in a hydrodynamically expanding environment [5]. These collective phenomena [6] are a characteristic feature of the Underlying Event (UE) in Pb–Pb collisions, that are independent of the fragmentation process but could have a possible impact on the jet fragments. Interactions between partons, stemming from the fragmentation shower and the Quark-Gluon Plasma, could change the jet pattern, which was already observed as quenching of jets [7]. Some other models consider alternative hadronisation mechanisms, e.g. recombination at low p_T [8], which is a soft process that could favour baryon over meson production. Particles with a momentum larger than $p_T = 4-6$ GeV/ c would, on the other hand, be produced in hard processes via fragmentation, that does not lead to an enhanced production of baryons (compared to production in vacuum). The study of identified particle yields and ratios like $\Lambda(\bar{\Lambda})$ and K_S^0 in inclusive production and comparison to production in jets will allow to trace



V0 UE estimation method	V0 sample
No-jet events	in events without any selected jets (reconstructed jets fulfilling all further requirements)
Outside cones	outside of $2R$ ($R = 0.2, 0.3$)
Random cones	in cones with randomly chosen axis (η, ϕ) and no overlap with selected jets
Median-cluster cones	uses median k_t cluster (similar to k_t alg. for average background estimation)
Perpendicular cones	in cone perpendicular (in ϕ) to jet axis

Table 1. Methods to estimate the V0 particle contribution from the UE

back possible differences in the hadronisation process and its p_T dependence. Since collective features were proposed to also occur in p–Pb collisions, one can compare among the different collision systems (pp, p–Pb and Pb–Pb), to study the onset of this baryon enhancement, that has already been observed for inclusive production in p–Pb collisions [9]. For the analyses presented in this contribution, the p_T spectra of strange particles associated with jets are studied for events with different event multiplicities in p–Pb collisions and for the 10% most central events in Pb–Pb collisions.

2. Experimental setup and analysis strategy

The jet reconstruction is done with the *anti* – k_t jet finder using charged tracks with $p_T > 150$ MeV/ c and requiring a leading particle in each jet to have at minimum $p_T = 5$ GeV/ c . In the following, jets fulfilling all these requirements are referred to as 'selected' jets. The jet energy is corrected on an event-by-event basis for the contribution of the average charged UE [10].

Both p–Pb and Pb–Pb analyses use the decay topology of the strange neutral particles (V0s) for their reconstruction. The decay channels $K_S^0 \rightarrow \pi^+ + \pi^-$ [11] and $\Lambda \rightarrow p + \pi^-$ ($\bar{\Lambda} \rightarrow \bar{p} + \pi^+$) [11] are used. By recombining the charged tracks of opposite curvature, measured in the Inner Tracking System (ITS) and the Time Projection Chamber (TPC), together with a particle mass hypothesis the invariant mass of the V0 candidates is calculated. A set of cuts applied on various observables of the topology of the decay already rejects a sizeable amount of combinatorial background. The particle signal is extracted after a background fit and by applying a bin-counting procedure on the invariant mass distributions in different intervals of p_T^{V0} and $p_{T, \text{jet}}^{\text{ch}}$. The V0 candidate is associated to the jet cone if the distance, which is calculated in azimuthal angle (ϕ) and in pseudo-rapidity (η), to the jet axis is smaller than the jet resolution parameter R :

$$\sqrt{(\phi_{V0} - \phi_{\text{jet}})^2 + (\eta_{V0} - \eta_{\text{jet}})^2} < R \quad (1)$$

The V0 acceptance is chosen to $|\eta^{V0}| < 0.7$, the acceptance for the jets is $|\eta^{\text{jet}}| < |\eta^{V0}| - R = 0.5(0.4)$ for $R = 0.2(0.3)$.

Figure 1 shows the uncorrected K_S^0 and Λ p_T spectra for jets with $R = 0.3$ and for two different jet p_T intervals ($p_{T, \text{jet}}^{\text{ch}} > 10$ GeV/ c , > 20 GeV/ c) in Pb–Pb collisions. The spectra are scaled for better visibility.

Several corrections have to be applied to the raw yields. First, one has to correct for the V0 particle reconstruction efficiency as functions of p_T and η , by using Monte-Carlo simulations. The efficiency is calculated separately for V0s in the inclusive event, the jet cone (JC) and in the UE. Since V0s from the UE contribute quite substantially to the V0 spectra in the JC, especially at low p_T , this has to be accurately subtracted. For this UE V0 correction, we developed several methods (see table 1) to measure the raw V0 p_T spectrum in the UE as shown in Figure 2. The default method are inclusive V0s taken from events, in which no reconstructed jet is selected; this is the so-called "No-jet events" method. To assign a systematic uncertainty to this correction, 4 alternative methods are applied. Compared to the default method (see ratio in Figure 2, bottom), they show a deviation of around 15% for $p_T^{V0} < 4$ GeV/ c . The order of the different correction steps is as follows: After the correction for the V0 reconstruction efficiency is applied,




Figure 1. Uncorrected K_S^0 and Λ p_T spectra for $R = 0.3$ jets in Pb-Pb collisions at 2.76 TeV and in the 10% highest multiplicity class. The jets are reconstructed for two different jet p_T intervals ($p_{T,jet}^{ch} > 10$ GeV/c, > 20 GeV/c) with the *anti* - k_t jet finder using charged tracks with minimum $p_T > 150$ MeV/c.




Figure 2. (top) Uncorrected K_S^0 p_T spectra in UE in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV and in the 10% most central events. The different used methods serve to estimate systematic uncertainty of UE subtraction. (bottom) Each method divided by default (No-jet method), the ratios show a deviation of around 15% for $p_T^{V0} < 4$ GeV/c.

the UE V_0 contribution (using the "no-jets" method) and the feed-down (FD) are subtracted from the V_0 p_T spectrum in the JC. To account for the sizable FD into Λ mostly stemming from $\Xi^{0,-} \rightarrow \Lambda$ and since there is no direct $\Xi^{0,-}$ measurement in jets so far, two extreme scenarios are considered for the V_0 s in the JC, namely the FD contribution from the inclusive particle analysis [3] and the correction derived from PYTHIA pp simulations. The difference is used to assign a systematic uncertainty.

3. Results

Figure 3 shows the Λ/K_S^0 ratio in jets in p-Pb events at 5.02 TeV and in the 10% highest multiplicity class. It is compared to the inclusive Λ/K_S^0 ratio (solid lines) in p-Pb collisions and to the ratio predicted by PYTHIA 8 4C simulations at 5.02 TeV/c (dotted lines, showing spread for the used values of R). The measured ratio in jets shows no visible enhancement at intermediate p_T and is far below the inclusive ratio. Within the systematic uncertainties, the ratio in jets is in agreement with the PYTHIA simulations. Furthermore there is no significant dependence on the $p_{T,jet}^{ch}$ interval or on R visible.

4. Conclusions

The observed Λ/K_S^0 ratio in jets in p-Pb collisions is significantly smaller than the inclusive measurement and compatible with the predictions of a PYTHIA jet simulation. This points to a




Figure 3. Λ/K_S^0 ratio in jets for $p_{T,jet}^{ch} > 10$ GeV/c (left) and $p_{T,jet}^{ch} > 20$ GeV/c (right) measured in p–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV for events in the 0–10% V0A (determined by the VZERO-A detector) multiplicity class [12, 13]. Results of the ratio in jets are compared to the inclusive particle ratio (solid lines) and the expectation from a PYTHIA 8 4C simulation at 5.02 TeV/c (dotted lines, showing spread for the used values of R).

scenario in which the hadronisation of soft particles is disentangled from the hard fragmentation processes. Therefore the baryon enhancement seen in inclusive production (see black markers in Figure 3) at intermediate particle p_T in p–Pb collisions seems to be a feature of the UE and not from fragmentation. Furthermore, the uncorrected spectra for K_S^0 and Λ particles in the JC and in UE in 10% most central events in Pb–Pb collisions have been presented. The Λ/K_S^0 ratio in jets in Pb–Pb collisions will be reported soon.

References

- [1] Agakishiev G *et al.* (STAR Collaboration) 2012 *Phys. Rev. Lett.* **108** 072301 (arXiv:nucl-ex/1107.2955)
- [2] Adams J *et al.* (STAR Collaboration) 2006 (arXiv:nucl-ex/0601042v1)
- [3] Abelev B *et al.* (ALICE Collaboration) 2013 *Phys. Rev. Lett.* **111** 222301 (arXiv:nucl-ex/1307.5530)
- [4] Abelev B *et al.* (ALICE Collaboration) 2014 *Phys. Lett.* **B736** 196207 (arXiv:nucl-ex/1401.1250v4)
- [5] Werner K 2012 *Phys. Rev. Lett.* **109** 102301 (arXiv:nucl-th/1204.1394)
- [6] Aamodt K *et al.* (ALICE Collaboration) 2010 *Phys. Rev. Lett.* **105** (arXiv:nucl-ex/1405.4632)
- [7] Milhano J G 2013 *Nucl. Phys.* **A904-905** 122c–129c (arxiv:hep-ex/1211.2657v1)
- [8] Fries R, Müller B, Nonaka C and Bass S 2003 *Phys. Rev. Lett.* **90** 202303 (arxiv:nucl-th/0301087)
- [9] Abelev B *et al.* (ALICE Collaboration) 2014 *Phys. Lett.* **B728** 25 (arxiv:nucl-ex/1307.6796)
- [10] Cacciari M and Salam G P 2008 *Phys. Lett.* **B659** 119126 (arXiv:hep-ex/0707.1378)
- [11] Olive K A *et al.* (Particle Data Group) 2014 *Chin. Phys.* **C38** 090001
- [12] Aamodt K *et al.* (ALICE Collaboration) 2008 *JINST* **3** S08002 (doi:10.1088/1748-0221/3/08/S08002)
- [13] Abelev B *et al.* (ALICE Collaboration) 2014 *Int. J. Mod. Phys.* **A** 1430044 (arxiv:nucl-ex/1402.4476)