

T-duality, Quotients and Currents for Non-Geometric Closed Strings

Ioannis Bakas^{*1} and Dieter Lüst^{+2,3,4}

¹ *Department of Physics, School of Applied Mathematics and Physical Sciences
National Technical University, 15780 Athens, Greece*

² *Max-Planck-Institut für Physik
Föhringer Ring 6, 80805 München, Germany*

³ *Arnold-Sommerfeld-Center für Theoretische Physik
Department für Physik, Ludwig-Maximilians-Universität München
Theresienstraße 37, 80333 München, Germany*

⁴ *Theory Group, Department of Physics, CERN
CH-1211 Geneva 23, Switzerland*

May 18, 2015

ABSTRACT

We use the canonical description of T-duality as well as the formulation of T-duality in terms of chiral currents to investigate the geometric and non-geometric faces of closed string backgrounds originating from principal torus bundles with constant H-flux. Employing conformal field theory techniques, the non-commutative and non-associative structures among generalized coordinates in the so called Q-flux and R-flux backgrounds emerge by gauging the Abelian symmetries of an enlarged Roček-Verlinde sigma-model and projecting the associated chiral currents of the enlarged theory to the T-dual coset models carrying non-geometric fluxes.

^{*}bakas@mail.ntua.gr

[†]dieter.luest@lmu.de

Contents

1	Introduction	2
2	Canonical T-duality and redefined world-sheet currents	5
2.1	T-duality as canonical transformation	6
2.2	T-duality in terms of chiral currents	9
2.3	Combining the two approaches	12
3	Parabolic flux vacua of closed strings	15
3.1	T-dual coordinates	17
3.2	Currents and generalized coordinates	20
4	Canonical T-duality and non-commutativity	22
4.1	Preliminaries: the case of free fields	22
4.2	Non-commutativity for interacting fields with constant fluxes	24
4.3	Case by case computations	27
4.3.1	Commutators of the H -flux background	27
4.3.2	Commutators of the geometric f -flux model	32
4.3.3	Commutators of the non-geometric Q -flux model	32
4.3.4	Commutators of the non-geometric R -flux model	33
5	Conclusions and discussion	34
A	Monodromies of toroidal fibrations	36
B	Parabolic backgrounds and monodromies of currents	38
B.1	The parabolic backgrounds with constant flux	38
B.2	Monodromies of geometric H -flux model	41
B.3	Monodromies of geometric f -flux model	42
B.4	Monodromies of non-geometric Q -flux model	42
B.5	Monodromies of non-geometric R -flux model	43
C	Some relevant dilogarithmic integrals	44

C.1	Delta-function, step-function and the logarithm	44
C.2	The dilogarithm function	45
C.3	Certain dilogarithmic commutators	46

1 Introduction

Flux vacua of closed string theory have been investigated in great detail in recent years (for topical reviews, see, for instance, [1, 2]). Various superstring backgrounds were brought to light in that context, including examples of geometric as well as non-geometric spaces. Non-geometry is a relatively new concept in string theory whose necessity and importance were unveiled in a series of papers by employing T-duality and its coordinate dependent generalizations [3]-[21], but it should also be noted that non-geometric string constructions appeared long time ago in conformal field theory using free fermions, free bosons and asymmetric orbifolds [22]-[25]. It was found that closed strings on non-geometric flux backgrounds have surprising features in that they exhibit non-commutativity and non-associativity among the closed string coordinates and their momenta [26, 27]. These new algebraic structures were investigated further in a subsequent series of papers [28]-[41], focusing on the derivation of the commutation relations for different string models and their mathematical interpretation. Independent studies by other authors, following a more mathematical line of thought, have also come to the conclusion that non-commutative and non-associative tori arise from geometric flux vacua by a sequence of T-dualities [42, 43].

Non-geometric backgrounds capture properties of string theory that cannot be formulated in terms of conventional compactifications of supergravity. The prime example is provided by the T-dual faces of toroidal flux vacua, known as Q - and R -flux models, but there are also examples of non-geometric spaces that do not have geometric duals [44, 45]. However, it remains a challenge to this day how to define and explore the properties of non-geometric closed string vacua in all generality and find a unifying principle to describe geometry and non-geometry on equal footing. Double geometry and double field theory (DFT) seem to be the most prominent candidates for encompassing string models and their T-dual faces in a democratic way [46]-[68].

There are two different classes of non-geometric spaces. The first class are Q -spaces, which are locally geometric in that they resemble differentiable manifolds locally, but they fail to be globally geometric because their gluing conditions involve T-duality transformations and not just diffeomorphisms. In that case, departure from ordinary geometry is mild and it manifests as non-commutativity of the closed string coordinates. The target space still makes good sense as a fibration though the mon-

odromy of the fibres, as one goes around the base space, is non-trivial. The second class are R -spaces, which fail to be differentiable manifolds globally as well as locally. Then, the concept of target space geometry breaks down completely and the string coordinates exhibit commutation relations of a non-commutative and non-associative algebra.

It is the aim of the present work to provide an alternative conformal field theory (CFT) derivation of the commutation relations among the closed string coordinates of non-geometric flux vacua, focusing attention to the well known class of three-dimensional models with parabolic monodromies, which carry H -, f -, Q - and R -flux and they are interrelated by T-dualities. The commutation relations of the closed string coordinates of these models have already been derived using the operator approach [32] (but see also the computation based on Dirac brackets that was reported in ref. [38]). The approach we take here is entirely based on CFT world-sheet methods and it complements nicely the previous work on the subject that brought to light the non-commutative and non-associative aspects of those flux string models. The present paper also extends and refines the CFT computations of ref. [28], where the non-associativity was derived from the structure of CFT scattering amplitudes among tachyon vertex operators in the R -flux background. The results of ref. [28] led to the introduction of a non-associative tri-product among functions, which was further investigated in [31, 33]. However, as noted in [28], the non-associative structure of CFT scattering amplitudes in the R -flux background is only visible when some of the tachyon momenta are taken off-shell, whereas on-shell the physical scattering amplitudes are associative, as expected in any conformal field theory. This result was analyzed further in the context of double field theory [61], showing, in particular, that the non-associative tri-product acts non-trivially on functions of doubled space only when the strong DFT constraint is violated.

In this paper, we compute directly the underlying commutation relations of some suitably defined *generalized* string coordinates of the H -, f -, Q - and R -flux backgrounds, without making reference to off-shell nor on-shell amplitudes. For this purpose, we start from the ungauged non-linear sigma-model of Roček and Verlinde [69], which provides the "parent" theory in our framework. It contains chirally conserved $U(1)$ world-sheet currents \mathcal{J}^I and $\tilde{\mathcal{J}}^I$, which transform in a simple linear way under T-duality transformations. The associated T-dual "child" theories are obtained by gauging the associated axial/vector like $U(1)$ symmetries. The $U(1)$ currents \mathcal{J}^I and $\tilde{\mathcal{J}}^I$ are not anymore chirally conserved when they are expressed in terms of the target space coordinates X^I of the associated "child" theories, but they still transform linearly under T-duality. Actually, as will be seen later in detail, the non-commutative and/or non-associative structures that arise in the presence of non-geometric fluxes refer to generalized string coordinates \mathcal{X}^I , which are defined by the world-sheet integrals of

the Roček-Verlinde currents \mathcal{J}^I and $\bar{\mathcal{J}}^I$ as $\mathcal{X}^I = \int \mathcal{J}^I dz$ and $\bar{\mathcal{X}}^I = \int \bar{\mathcal{J}}^I d\bar{z}$ with prescribed integration paths. The generalized coordinates also transform nicely under T-duality, mapping the commutative geometry of geometric flux backgrounds to the non-commutative geometry of the non-geometric flux backgrounds via T-duality, as it has already been advocated in ref. [27].

It should be noted at this point that the currents \mathcal{J}^I and $\bar{\mathcal{J}}^I$ used in this paper are refined compared to the choice of currents made in ref. [28]. It should also be emphasized that the non-commutative and/or non-associative algebras that we are discussing here do not arise among the sigma-model coordinates X^I of the child theories with non-geometric fluxes, but they rather emerge by projecting the chiral currents of the parent theory to the child gauged sigma-models and expressing the generalized coordinates \mathcal{X}^I as world-sheet integrals of those non-chiral currents. We further note that the algebra of currents in the parent Roček-Verlinde theory is completely commutative and associative, unlike the non-trivial commutation relations that can emerge in the quotient. All these put on firm basis the existing results for toroidal flux models and they also provide the means to explore generalizations to other non-geometric backgrounds of current interest, including the Q- and R-frames of Wess-Zumino-Witten models on group manifolds (see, for instance, [15, 67, 68]).

The material of this paper is organized as follows. In section 2, we recall the description of T-duality as canonical transformation [70]-[74] on the coordinates X^I and momenta P_I of the world-sheet sigma-model, hereby introducing the notion of dual coordinates \tilde{X}^I and dual momenta \tilde{P}^I , which, in general, are expressed non-locally in terms of the original phase space variables. We also consider the chiral currents of the parent Roček-Verlinde model and combine them with the canonical approach to T-duality, hereby expressing those currents in terms of the (dual) coordinates of the gauged child theories. The action of T-duality as automorphism of the currents in the underlying conformal field theory turns out to be useful for string backgrounds without isometries, giving rise to non-geometries. In section 3, we focus on toroidal flux models originating from T^3 with constant H -flux and explain how T-duality is used at work to obtain the geometric and non-geometric faces of closed string spaces. We also define the generalized coordinates \mathcal{X}^I of the parabolic flux backgrounds via the projected currents of the parent Roček-Verlinde theory. In section 4, we confirm the emergence of non-commutative and non-associative structures in the commutation relations of the generalized closed strings coordinates and momenta of the Q-flux and R-flux models, respectively, by conformal field theory methods. Finally, in section 5, we present our conclusions and discuss some open problems for future work. There are also three appendices summarizing the basic definitions and properties of the monodromies arising in T^2 fibrations over S^1 (Appendix A), the salient features of the parabolic H - f - Q - and R -flux models, which are detached from the main text

(Appendix B), as well as some dilogarithmic integrals, based on the definitions and properties of the dilogarithm function found in ref. [75], which are repeatedly used in the text for the computation of the commutation relations (Appendix C).

Certain aspects of the canonical approach to string non-commutativity were also discussed recently in ref. [76], following a complementary route. There, the monodromies of toroidal fibrations are not built into the closed string boundary conditions a priori, but they rather come out a posteriori. In this sense, there is partial overlap of their results with our section 4, but our presentation is more natural for the purposes of string theory. Our work also emphasizes for the first time the use of generalized string coordinates.

2 Canonical T-duality and redefined world-sheet currents

Our starting point is the world-sheet action of a general non-linear sigma-model with metric G and anti-symmetric tensor field B couplings that provide the background for string propagation,

$$S = \frac{1}{2\pi\alpha'} \int dzd\bar{z} (G_{IJ}(X) + B_{IJ}(X)) \partial X^I \bar{\partial} X^J . \quad (2.1)$$

The inverse string tension is $\alpha' \sim l_s^2$ and it can be normalized to 1. The target space coordinates X^I are two-dimensional fields labeled by $I = 1, 2, \dots, d$. The action (2.1) is often used to describe the entire space in which the string propagates, but it can also be part of it when it is tensored together with other blocks to maintain conformal invariance upon quantization. There is also a dilaton field $\Phi(X)$ that enters in the renormalization group analysis of the sigma-model, but this is suppressed here.

In this section, we review the T-duality rules of the sigma-model in the presence of an isometry in target space [70] and use them to discuss the general features of the commutation relations among the coordinates in the T-dual faces of string backgrounds. Explicit calculations will be performed in subsequent sections for specific models. We focus on the interpretation of T-duality as canonical transformation in the phase space of the two-dimensional sigma-model (2.1) [71, 72, 73], as well as on the axial-vector quotient description of T-dual backgrounds that follows by gauging chiral symmetries [69]. These two approaches are complementary to each, leading to the important notion of dual coordinates. The dual coordinates are expressed non-locally in terms of the original fields, accounting for non-commutativity/non-associativity in the applications that will be considered later.

Note that the duality rules are applicable to all sigma-models, including, for ex-

ample, principal chiral models, irrespective of conformal invariance. However, if the original model is conformal, the dual model will also be conformal for appropriate choice of the dilaton field. In the latter case, T-duality acts as a solution generating symmetry at the fixed points of the renormalization group equations and it is promoted to an exact symmetry of string theory relating two conformal field theories with different target space geometries. Further details and applications of target space duality in string theory can be found in the report [74], and references therein, which also discuss the continuous T-duality rules and their discrete variants for general backgrounds admitting n commuting isometries in terms of $O(n, n)$ group elements.

2.1 T-duality as canonical transformation

Suppose that the background does not depend on the coordinate X^1 , i.e., $\partial/\partial X^1$ is an isometry of the world-sheet action in adapted coordinates of target space. The components of the corresponding Killing vector field $\xi = k^I \partial/\partial X^I$ are $k^I = (1, 0, \dots, 0)$ and $k^2 = G_{IJ} k^I k^J$ is the length-squared of ξ . Setting $z = (\tau + \sigma)/2$ and $\bar{z} = (\tau - \sigma)/2$, so that $\partial = \partial/\partial\tau + \partial/\partial\sigma$ and $\bar{\partial} = \partial/\partial\tau - \partial/\partial\sigma$, the Lagrangian density takes the following form

$$\mathcal{L} = \frac{V}{2} \left((\dot{X}^1)^2 - (X^{1'})^2 \right) + \dot{X}^1 (J_1 + \bar{J}_1) - X^{1'} (J_1 - \bar{J}_1) + U, \quad (2.2)$$

where

$$V = G_{11} = k^2, \quad U = \frac{1}{2} (G_{ij} + B_{ij}) \partial X^i \bar{\partial} X^j \quad (2.3)$$

with i and j taking all other values of I and J but 1 and

$$J_1(z, \bar{z}) = \frac{1}{2} (G_{1i} - B_{1i}) \partial X^i, \quad \bar{J}_1(z, \bar{z}) = \frac{1}{2} (G_{1i} + B_{1i}) \bar{\partial} X^i. \quad (2.4)$$

Next, we apply Legendre transformation to the pair of variables (X^1, \dot{X}^1) . The momentum conjugate to the coordinate X^1 is

$$P_1 = V \dot{X}^1 + J_1 + \bar{J}_1 \quad (2.5)$$

and, hence, the Hamiltonian density (or better to say the *Routhian*, since the transformation is performed only with respect to the cyclic coordinate X^1 that does not appear explicitly in the Lagrangian) is

$$\mathcal{H} = \frac{1}{2V} (P_1)^2 + \frac{V}{2} (X^{1'})^2 + \frac{1}{2V} (J_1 + \bar{J}_1)^2 - \frac{1}{V} P_1 (J_1 + \bar{J}_1) + X^{1'} (J_1 - \bar{J}_1) - U. \quad (2.6)$$

Let us now perform a canonical transformation $(X^1, P_1) \rightarrow (\tilde{X}_1, \tilde{P}^1)$ generated by the

function

$$\mathcal{F} = \frac{1}{2} \oint_{S^1} (X^1 \tilde{X}'_1 - X^{1'} \tilde{X}_1) d\sigma \quad (2.7)$$

so that

$$P_1 = \frac{\delta \mathcal{F}}{\delta X^1} = \tilde{X}'_1, \quad \tilde{P}^1 = -\frac{\delta \mathcal{F}}{\delta \tilde{X}_1} = X^{1'}. \quad (2.8)$$

We call \tilde{X}_1 and \tilde{P}^1 the dual coordinate and momentum to X^1 and P_1 , respectively. Then, in terms of the dual variables, the Hamiltonian takes the form

$$\tilde{\mathcal{H}} = \frac{V}{2} (\tilde{P}^1)^2 + \frac{1}{2V} (\tilde{X}'_1)^2 + \frac{1}{2V} (J_1 + \bar{J}_1)^2 - \frac{1}{V} \tilde{X}'_1 (J_1 + \bar{J}_1) + \tilde{P}^1 (J_1 - \bar{J}_1) - U. \quad (2.9)$$

Varying $\tilde{\mathcal{H}}$ with respect to \tilde{P}^1 , we find that the dual velocity and momentum are related to each other as follows,

$$\dot{\tilde{X}}_1 = V \tilde{P}^1 + J_1 - \bar{J}_1. \quad (2.10)$$

Finally, we perform the inverse Legendre transformation with respect to $(\tilde{X}_1, \tilde{P}^1)$ and arrive at the dual Lagrangian

$$\tilde{\mathcal{L}} = \frac{1}{2V} \left((\dot{\tilde{X}}_1)^2 - (\tilde{X}'_1)^2 \right) - \frac{1}{V} \dot{\tilde{X}}_1 (J_1 - \bar{J}_1) + \frac{1}{V} \tilde{X}'_1 (J_1 + \bar{J}_1) + \tilde{U}, \quad (2.11)$$

where

$$\tilde{U} = U - \frac{2}{V} J_1 \bar{J}_1. \quad (2.12)$$

The form of the dual Lagrangian $\tilde{\mathcal{L}}$ is identical to the original Lagrangian \mathcal{L} provided that the background fields of the dual non-linear sigma-model with target space coordinates (\tilde{X}_1, X^i) are taken to be

$$\begin{aligned} \tilde{G}_{11} &= \frac{1}{G_{11}}, & \tilde{G}_{1i} &= \frac{B_{1i}}{G_{11}}, & \tilde{G}_{ij} &= G_{ij} - \frac{G_{1i} G_{1j} - B_{1i} B_{1j}}{G_{11}}, \\ \tilde{B}_{1i} &= \frac{G_{1i}}{G_{11}}, & \tilde{B}_{ij} &= B_{ij} - \frac{G_{1i} B_{1j} - B_{1i} G_{1j}}{G_{11}}. \end{aligned} \quad (2.13)$$

These are precisely Buscher rules for T-duality with respect to a Killing vector field ζ in adapted coordinates [70], which are formulated as canonical transformation in the phase space of the two-dimensional non-linear sigma-model [71, 72, 73]. A more compact way to describe the T-duality transformation along X^1 is provided by combining the metric and anti-symmetric tensor fields as $E_{IJ} = G_{IJ} + B_{IJ}$ and letting

$$\tilde{E}_{11} = \frac{1}{E_{11}}, \quad \tilde{E}_{1i} = \frac{E_{1i}}{E_{11}}, \quad \tilde{E}_{i1} = -\frac{E_{i1}}{E_{11}}, \quad \tilde{E}_{ij} = E_{ij} - \frac{E_{i1} E_{1j}}{E_{11}}. \quad (2.14)$$

Conformal invariance of the world-sheet sigma-model also requires that the corresponding dilaton field transforms as $\tilde{\Phi} = \Phi - \log V$.

The effect of T-duality can be neatly described by a non-local redefinition of the target space coordinate associated to the Killing isometry. For this purpose, we write down the transformation of the derivatives of the Killing coordinate

$$\partial X^1 = \frac{1}{E_{11}}(\partial \tilde{X}_1 - E_{i1}\partial X^i), \quad \bar{\partial} X^1 = -\frac{1}{E_{11}}(\bar{\partial} \tilde{X}_1 + E_{1i}\bar{\partial} X^i) \quad (2.15)$$

by combining the canonical transformation (2.8) with the expressions for the momenta (2.5) and (2.10). Substituting the expressions (2.14), we arrive at the integral formula

$$X^1(z, \bar{z}) = \int^z (\tilde{E}_{11}\partial \tilde{X}_1 + \tilde{E}_{i1}\partial X^i) dz' - \int^{\bar{z}} (\tilde{E}_{11}\bar{\partial} \tilde{X}_1 + \tilde{E}_{1i}\bar{\partial} X^i) d\bar{z}' \quad (2.16)$$

expressing X^1 in terms of the dual background. There is an analogous formula expressing \tilde{X}_1 in terms of the original background, as

$$\partial \tilde{X}_1 = E_{11}\partial X^1 + E_{i1}\partial X^i, \quad \bar{\partial} \tilde{X}_1 = -(E_{11}\bar{\partial} X^1 + E_{1i}\bar{\partial} X^i), \quad (2.17)$$

which integrates to

$$\tilde{X}_1(z, \bar{z}) = \int^z (E_{11}\partial X^1 + E_{i1}\partial X^i) dz' - \int^{\bar{z}} (E_{11}\bar{\partial} X^1 + E_{1i}\bar{\partial} X^i) d\bar{z}'. \quad (2.18)$$

These expressions show that T-duality acts, in general, in a rather complicated *non-local* way on the target space coordinates.

For constant backgrounds, i.e., for a free two-dimensional conformal field theory with $E_{IJ} = \delta_{IJ}$, the action of T-duality looks very simple. For compactifications on a circle, the left- and right-moving coordinates are chiral,

$$X_L^1(z) = \int^z \partial X^1(z', \bar{z}') dz', \quad X_R^1(\bar{z}) = \int^{\bar{z}} \bar{\partial} X^1(z', \bar{z}') d\bar{z}', \quad (2.19)$$

and they transform as follows, by specializing equation (2.18),

$$\tilde{X}_{1L} = X_L^1, \quad \tilde{X}_{1R} = -X_R^1. \quad (2.20)$$

Hence, one immediately obtains the well-known T-duality relation among the coordinates X^1 and \tilde{X}_1 of the two dual models,

$$X^1 = X_L^1 + X_R^1 \longrightarrow \tilde{X}_1 = X_L^1 - X_R^1. \quad (2.21)$$

In all other cases, one can still introduce variables X_L^1 and X_R^1 by adding and subtracting the integral expressions (2.16) and (2.18), thus decomposing the coordinates X^1 and \tilde{X}_1 as

$$X^1 = X_L^1 + X_R^1, \quad \tilde{X}_1 = X_L^1 - X_R^1. \quad (2.22)$$

The effect of T-duality is still summarized by equation (2.21), as for the case of free fields. Note, however, that X_L^1 and X_R^1 are not chiral and anti-chiral world-sheet functions, in general, because $\partial X^1(z, \bar{z})$ and $\bar{\partial} X^1(z, \bar{z})$ are not separately conserved on general backgrounds. Although the general case of non-constant background fields $E_{IJ}(X^i)$ looks much more involved, it can also be understood in terms of appropriately defined chiral currents, as will be seen in the next subsection using a self-dual sigma-model.

If we were considering a canonical transformation with generating function $-\mathcal{F}$, flipping the sign of \mathcal{F} in defining equation (2.7), \tilde{X}_1 would be replaced by $-\tilde{X}_1$ and Buscher rules would be a variant of equations (2.13), picking a minus sign in the transformation law of \tilde{G}_{1i} and \tilde{B}_{1i} . Then, \tilde{E}_{1i} and \tilde{E}_{i1} (but not E_{1i} and E_{i1} , since X^1 remains inert) should flip signs in the corresponding expressions (2.16) and (2.18) relating X^1 with \tilde{X}_1 and T-duality would manifest as $\tilde{X}_{1L} = -X_L^1$ and $\tilde{X}_{1R} = X_R^1$ instead of $\tilde{X}_{1L} = X_L^1$ and $\tilde{X}_{1R} = -X_R^1$ as a matter of convention.

2.2 T-duality in terms of chiral currents

According to Roček and Verlinde, any pair of dual sigma-models can be represented as quotients of it, showing that T-duality transformations act as automorphism on the algebra of conformal fields [69]. The construction is reminiscent of modern day double field theory, though here it is taken on the world-sheet, since the master action includes the Killing coordinate and its dual counterpart on equal footing. We outline the main steps and results that will be used later in the calculations.

First, let us consider a variant of the two-dimensional non-linear sigma-model (2.1), which is naturally defined by the action

$$S' = \frac{1}{2\pi\alpha'} \int dzd\bar{z} (g_{ij}(X) + b_{ij}(X)) \partial X^i \bar{\partial} X^j. \quad (2.23)$$

Here, the target space is $(d-1)$ -dimensional with coordinates X^i instead of $X^I = (X^1, X^i)$ that appear in the original action S . The metric g_{ij} and the anti-symmetric tensor field b_{ij} are not identical to G_{ij} and B_{ij} of the original action, by restricting them on the orbits of the Killing vector field $\xi = \partial/\partial X^1$, but they will be related to them later.

Next, we enlarge the target space by introducing two additional coordinates X_L^1

and X_R^1 , which are a priori unrelated to the Killing coordinate X^1 of S , and define a new $(d + 1)$ -dimensional "parent" sigma-model

$$S_{\text{LR}} = S' + \frac{1}{2\pi\alpha'} \int dzd\bar{z} \left[\partial X_L^1 \bar{\partial} X_L^1 + \partial X_R^1 \bar{\partial} X_R^1 + 2B(X) \partial X_R^1 \bar{\partial} X_L^1 + 2G_i^L(X) \partial X^i \bar{\partial} X_L^1 + 2G_i^R(X) \bar{\partial} X^i \partial X_R^1 \right] \quad (2.24)$$

whose couplings $B(X)$, $G_i^L(X)$ and $G_i^R(X)$ depend only on the $d - 1$ coordinates X^i . With some additional ingredients that will be introduced shortly, X_L^1 and X_R^1 help to reconstruct the coordinate X^1 and its dual \tilde{X}_1 , making T-duality manifest. The enlarged action (2.24) admits an $U(1)_L \times U(1)_R$ symmetry under $\delta X_L^1 = \alpha_L(z)$ and $\delta X_R^1 = \alpha_R(\bar{z})$ with corresponding currents

$$\begin{aligned} \mathcal{J}^1(z) &= \partial X_L^1 + B(X) \partial X_R^1 + G_i^L(X) \partial X^i, \\ \bar{\mathcal{J}}^1(\bar{z}) &= \bar{\partial} X_R^1 + B(X) \bar{\partial} X_L^1 + G_i^R(X) \bar{\partial} X^i \end{aligned} \quad (2.25)$$

that are chirally conserved on the world-sheet, i.e.,

$$\bar{\partial} \mathcal{J}^1(z) = 0 = \partial \bar{\mathcal{J}}^1(\bar{z}). \quad (2.26)$$

Gauging the symmetry $U(1)_L \times U(1)_R$ completes the construction of the master action. It is achieved with the aid of a gauge field with components (A, \bar{A}) that couple to X_L^1 , X_R^1 and X^i as follows,

$$S_{\text{gauged}}^{(\pm)} = S_{\text{LR}} + \frac{1}{2\pi\alpha'} \int dzd\bar{z} \left[A \bar{\mathcal{J}}^1 \pm \bar{A} \mathcal{J}^1 + \frac{1}{2} (1 \pm B(X)) A \bar{A} \right]. \quad (2.27)$$

The different signs correspond to the two possible ways to gauge the symmetry, employing either axial or vector gauging, $\delta X_L^1 = \alpha$ and $\delta X_R^1 = \mp \alpha$. The resulting gauged action (2.27) is self-dual in that it encompasses the original non-linear sigma-model S and its T-dual \tilde{S} on equal footing.

Indeed, integrating out the gauge fields, we obtain the original action S for one gauging and the dual action \tilde{S} for the other gauging. The resulting d -dimensional

dual "child" actions are¹

$$S = S' + \frac{1}{2\pi\alpha'} \int dzd\bar{z} \left[\frac{1+B}{1-B} \partial X^1 \bar{\partial} X^1 + \frac{2}{1-B} G_i^R \partial X^1 \bar{\partial} X^i + \frac{2}{1-B} G_i^L \bar{\partial} X^1 \partial X^i + \frac{2}{1-B} G_i^L G_j^R \partial X^i \bar{\partial} X^j \right] \quad (2.28)$$

and

$$\tilde{S} = S' + \frac{1}{2\pi\alpha'} \int dzd\bar{z} \left[\frac{1-B}{1+B} \partial \tilde{X}_1 \bar{\partial} \tilde{X}_1 - \frac{2}{1+B} G_i^R \partial \tilde{X}_1 \bar{\partial} X^i + \frac{2}{1+B} G_i^L \bar{\partial} \tilde{X}_1 \partial X^i - \frac{2}{1+B} G_i^L G_j^R \partial X^i \bar{\partial} X^j \right]. \quad (2.29)$$

The identification is exact provided that the Killing coordinates of S and \tilde{S} are chosen to be

$$X^1 = X_L^1 + X_R^1, \quad \tilde{X}_1 = X_L^1 - X_R^1 \quad (2.30)$$

and they are required to have the same periodicity. Thus, X_L^1 and X_R^1 reconstruct the Killing coordinate and its dual, as in the canonical formulation of T-duality. The couplings appearing in the action S_{LR} should also be adjusted as

$$G_{11} = \frac{1+B}{1-B}, \quad G_{1i} \mp B_{1i} = \frac{2}{1-B} G_i^{R,L}(X) \quad (2.31)$$

and

$$G_{ij} = g_{ij} + \frac{1-B}{2} (G_{1i} G_{1j} - B_{1i} B_{1j}), \quad B_{ij} = -b_{ij} - \frac{1-B}{2} (G_{1i} B_{1j} - B_{1i} G_{1j}), \quad (2.32)$$

hereby expressing $B(X)$, $G_i^{R,L}(X)$, $g_{ij}(X)$ and $b_{ij}(X)$ in terms of the background fields $G_{IJ}(X)$ and $B_{IJ}(X)$ of the d -dimensional target space associated to the action S .

With these identifications in mind, we obtain the T-duality rules for the components of the metric G_{IJ} and the anti-symmetric tensor field B_{IJ} by comparing the two actions S and \tilde{S} ,

$$\begin{aligned} \tilde{G}_{11} &= \frac{1}{G_{11}}, \quad \tilde{G}_{1i} = \frac{B_{1i}}{G_{11}}, \quad \tilde{G}_{ij} = G_{ij} - \frac{G_{1i} G_{1j} - B_{1i} B_{1j}}{G_{11}}, \\ \tilde{B}_{1i} &= \frac{G_{1i}}{G_{11}}, \quad \tilde{B}_{ij} = B_{ij} - \frac{G_{1i} B_{1j} - B_{1i} G_{1j}}{G_{11}}, \end{aligned} \quad (2.33)$$

which are exactly the transformation rules (2.13). Looking at the expressions (2.28),

¹We assume that the world-sheet is topologically trivial to avoid unnecessary complications that arise by the presence of a total derivative term $\partial X^1 \bar{\partial} \tilde{X}_1 - \partial \tilde{X}_1 \bar{\partial} X^1$ in the Lagrangian, which is omitted. Of course, higher genera should be employed to establish T-duality as an exact symmetry of theory to all orders in string perturbation theory, but they are beyond the scope of the present discussion.

(2.29) and (2.30), it follows that the T-duality rules (2.33) can be neatly summarized as

$$B \rightarrow -B, \quad G_i^{L,R} \rightarrow \pm G_i^{L,R}, \quad X_L^1 \rightarrow X_L^1, \quad X_R^1 \rightarrow -X_R^1. \quad (2.34)$$

An immediate consequence is that the chiral currents (2.25) transform under duality as

$$\mathcal{J}^1(z) \rightarrow \mathcal{J}^1(z), \quad \tilde{\mathcal{J}}^1(\bar{z}) \rightarrow -\tilde{\mathcal{J}}^1(\bar{z}). \quad (2.35)$$

Thus, T-duality leaves the left-moving current $\mathcal{J}_1(z)$ invariant and acts by flipping the sign of the right-moving current $\tilde{\mathcal{J}}_1(\bar{z})$, as in the case of free fields. If the enlarged sigma-model (2.24) is conformally invariant in the quantum regime, the pair of dual sigma-models S and \tilde{S} will also be conformal; in that case, the resulting models are equivalent as quantum field theories. The duality among sigma-models, as defined above, is a valid classical operation for non-conformal backgrounds as well, but in those cases the two models are not equivalent as quantum field theories.

The previous discussion generalizes to backgrounds with n commuting isometries, in which case the enlarged action S_{LR} involves $d + n$ target space coordinates, encompassing the Killing coordinates and their dual counterparts in a unified way. S_{LR} exhibits an $U(1)_L^n \times U(1)_R^n$ chiral symmetry with associated currents $\mathcal{J}^a(z)$ and $\tilde{\mathcal{J}}^a(\bar{z})$ for each $a = 1, 2, \dots, n$. Gauging all symmetries axially or vectorially results into two sigma-models that are related by a general $O(n, n)$ T-duality transformation. Further details can be found in the report [74] and references therein.

It is tempting to use the enlarged action S_{LR} for the world-sheet description of double field theory, which also encompasses the coordinates and their dual counterparts on equal footing in target space. Work in this direction is in progress.

2.3 Combining the two approaches

Let us now explain the different meaning of the Roček-Verlinde currents in the enlarged parent theory S_{LR} and in the reduced child theories S and \tilde{S} obtained by quotient. The key point is that X_L^1 and X_R^1 are independent fields of the enlarged theory, but they are interrelated by duality in the reduced theories. The form of the currents in S and \tilde{S} will be obtained by eliminating \tilde{X} in favor of X , and vice versa, through the canonical formulation of T-duality, hereby combining the two approaches.

As explained before, the parent currents (2.25) are chiral by the classical equations of motion of the enlarged two-dimensional non-linear sigma-model (2.24) and they

take the following form

$$\begin{aligned}\mathcal{J}^1(z) &= \partial X_L^1 + \frac{G_{11}-1}{G_{11}+1} \partial X_R^1 + \frac{1}{G_{11}+1} E_{1i} \partial X^i, \\ \bar{\mathcal{J}}^1(\bar{z}) &= \bar{\partial} X_R^1 + \frac{G_{11}-1}{G_{11}+1} \bar{\partial} X_L^1 + \frac{1}{G_{11}+1} E_{i1} \bar{\partial} X^i.\end{aligned}\quad (2.36)$$

They can be rewritten in terms of the target space coordinate $X^1 = X_L^1 + X_R^1$ and $\tilde{X}_1 = X_L^1 - X_R^1$, assumed to be independent in S_{LR} , as follows,

$$\begin{aligned}\mathcal{J}^1(z) &= \frac{1}{G_{11}+1} \left(G_{11} \partial X^1 + \partial \tilde{X}_1 + E_{1i} \partial X^i \right), \\ \bar{\mathcal{J}}^1(\bar{z}) &= \frac{1}{G_{11}+1} \left(G_{11} \bar{\partial} X^1 - \bar{\partial} \tilde{X}_1 + E_{i1} \bar{\partial} X^i \right).\end{aligned}\quad (2.37)$$

Then, it is immediately obvious that $\mathcal{J}^1(z)$ remains invariant and $\bar{\mathcal{J}}^1(\bar{z})$ flips sign under T-duality, since $\tilde{G}_{11} = 1/G_{11}$, $\tilde{E}_{1i} = E_{1i}/G_{11}$ and $\tilde{E}_{i1} = -E_{i1}/G_{11}$.

The coordinates X_L^1 and X_R^1 are not chiral, in general, but one can always define chiral coordinates $\mathcal{X}_L^1(z)$ and $\mathcal{X}_R^1(\bar{z})$ in the enlarged theory S_{LR} , letting

$$\mathcal{J}^1(z) = \partial \mathcal{X}_L^1(z), \quad \bar{\mathcal{J}}^1(\bar{z}) = \bar{\partial} \mathcal{X}_R^1(\bar{z}). \quad (2.38)$$

Then, T-duality is reformulated as automorphism

$$\tilde{\mathcal{X}}_{1L}(z) = \mathcal{X}_L^1(z), \quad \tilde{\mathcal{X}}_{1R}(\bar{z}) = -\mathcal{X}_R^1(\bar{z}). \quad (2.39)$$

Likewise, one can introduce the following combinations of chiral components

$$\mathcal{X}^1 = \mathcal{X}_L^1 + \mathcal{X}_R^1, \quad \tilde{\mathcal{X}}_1 = \mathcal{X}_L^1 - \mathcal{X}_R^1 \quad (2.40)$$

and restate T-duality in direct analogy with free fields, as

$$\mathcal{X}^1 = \mathcal{X}_L^1 + \mathcal{X}_R^1 \longrightarrow \tilde{\mathcal{X}}_1 = \mathcal{X}_L^1 - \mathcal{X}_R^1. \quad (2.41)$$

These new fields are non-locally expressed in terms of the target space coordinates $X^I = (X^1, X^i)$ of the original sigma-model (2.1). The free fields \mathcal{X}^1 and $\tilde{\mathcal{X}}_1$ exist only in the master theory S_{LR} for general backgrounds. Of course, for ordinary toroidal backgrounds we have $\mathcal{X}^1 = X^1$ and $\tilde{\mathcal{X}}_1 = \tilde{X}_1$, in which case these fields are well defined in the coset theories as well.

Next, using the non-local relations (2.15) and (2.17) among X^1 and \tilde{X}_1 , we can eliminate one coordinate in favor of the other and express the parent Roček-Verlinde

currents in the original sigma-model frame S as

$$\mathcal{J}^1 = \frac{2}{G_{11} + 1} \left(G_{11} \partial X^1 + G_{1i} \partial X^i \right), \quad \bar{\mathcal{J}}^1 = \frac{2}{G_{11} + 1} \left(G_{11} \bar{\partial} X^1 + G_{1i} \bar{\partial} X^i \right) \quad (2.42)$$

and in the dual sigma-model frame \tilde{S} as

$$\mathcal{J}^1 = \frac{2}{\tilde{G}_{11} + 1} \left(\tilde{G}_{11} \partial \tilde{X}^1 + \tilde{G}_{1i} \partial \tilde{X}^i \right), \quad \bar{\mathcal{J}}^1 = -\frac{2}{\tilde{G}_{11} + 1} \left(\tilde{G}_{11} \bar{\partial} \tilde{X}^1 + \tilde{G}_{1i} \bar{\partial} \tilde{X}^i \right). \quad (2.43)$$

It is immediately obvious that these "child" currents are not chiral fields of the original sigma-model nor its dual, in general. Thus, as expected by taking the quotient, the currents \mathcal{J}^1 and $\bar{\mathcal{J}}^1$ are not bona fide conformal fields of the coset theory S nor its dual \tilde{S} . The physical operators of the coset conformal field theory S or \tilde{S} should necessarily commute with the BRST charge of the gauged sigma-model (2.27) and they should be defined up to BRST commutators. T-duality acts by automorphisms only in the self-dual theory S_{LR} , whereas after the gauging the currents (2.42) of one reduced theory are mapped to the currents (2.43) of the other.

Let us illustrate this point by considering the well known example of the $SU(2)$ WZW model and its gauged variant $SU(2)/U(1)$. In this case, the $SU(2)$ group elements are parametrized canonically as

$$g = e^{\frac{i}{2} X_L \sigma_3} e^{\frac{i}{2} \theta \sigma_1} e^{\frac{i}{2} X_R \sigma_3} \quad (2.44)$$

and the action of the $SU(2)$ WZW model for the coordinates θ , X_L and X_R is the enlarged action S_{LR} , which is subsequently gauged. The corresponding Roček-Verlinde currents for the $U(1)_L \times U(1)_R$ symmetry that is gauged are (dropping the index 1 for convenience)

$$\mathcal{J}(z) = \partial X_L + \cos(2\theta) \partial X_R, \quad \bar{\mathcal{J}}(\bar{z}) = \bar{\partial} X_R + \cos(2\theta) \bar{\partial} X_L \quad (2.45)$$

and they correspond to the Cartan generators of the $SU(2)_L \times SU(2)_R$ current algebra of the $SU(2)$ WZW model. Here, we have $B = \cos(2\theta)$, accounting for the anti-symmetric tensor field of the $SU(2)$ model in the X_L and X_R directions, whereas $G_i^{L,R} = 0$. Setting $X = X_L + X_R$, $\tilde{X} = X_L - X_R$ and noting that $G_{11} = \cot^2 \theta = 1/\tilde{G}_{11}$, we obtain by gauging the two dual faces of the $SU(2)/U(1)$ WZW model described by the actions S and \tilde{S} with respective target space metrics

$$ds^2 = d\theta^2 + \cot^2 \theta dX^2 \quad (2.46)$$

and

$$d\tilde{s}^2 = d\theta^2 + \tan^2 \theta d\tilde{X}^2. \quad (2.47)$$

There is also a corresponding dilaton, required by conformal invariance, but no anti-symmetric tensor field.

The currents (2.45) are chiral, i.e., $\bar{\partial}\mathcal{J}(z) = 0 = \partial\bar{\mathcal{J}}(\bar{z})$ by the equations of motion of the $SU(2)$ WZW model, and they can be rewritten as

$$\mathcal{J}(z) = \cos^2\theta \partial X + \sin^2\theta \partial \tilde{X}, \quad \bar{\mathcal{J}}(\bar{z}) = \cos^2\theta \bar{\partial} X - \sin^2\theta \bar{\partial} \tilde{X}, \quad (2.48)$$

treating X and \tilde{X} as independent fields of the enlarged theory S_{LR} . After gauging, X and \tilde{X} are not independent, but they are dual to each other related by $\partial\tilde{X} = \cot^2\theta \partial X$ and $\bar{\partial}\tilde{X} = -\cot^2\theta \bar{\partial} X$. Then, in the reduced theory S the two currents take the form

$$\mathcal{J}(z, \bar{z}) = 2\cos^2\theta \partial X, \quad \bar{\mathcal{J}}(z, \bar{z}) = 2\cos^2\theta \bar{\partial} X, \quad (2.49)$$

whereas in the reduced theory \tilde{S} they are expressed as

$$\mathcal{J}(z, \bar{z}) = 2\sin^2\theta \partial \tilde{X}, \quad \bar{\mathcal{J}}(z, \bar{z}) = -2\sin^2\theta \bar{\partial} \tilde{X}. \quad (2.50)$$

It can be easily seen, using the classical equations of motion following from S or \tilde{S} , respectively, that these are no longer chiral after gauging, hence, their dependence on both z and \bar{z} indicated above. Also, it becomes obvious that the dual $\bar{\mathcal{J}}$ of S is the \mathcal{J} of \tilde{S} and the dual $\bar{\mathcal{J}}$ of \tilde{S} is the $-\mathcal{J}$ of S , unlike the left and right components of the currents (2.48) of the ungauged $SU(2)$ WZW model, which are self-dual and anti-self-dual, respectively.

Similar considerations apply to the toroidal flux vacua of closed string theory that will be considered next in detail.

3 Parabolic flux vacua of closed strings

Let us first consider the case of a 3-torus, T^3 , with $G_{IJ} = \delta_{IJ}$ (taking all radii equal to 1 to simplify the presentation), constant dilaton and background H -flux given by the field strength of the anti-symmetric tensor field,

$$H_{IJK} = \partial_I B_{JK} + \partial_K B_{IJ} + \partial_J B_{KI}. \quad (3.1)$$

Then, the original sigma-model action (2.1) takes the form

$$S = \frac{1}{2\pi\alpha'} \int dzd\bar{z} (\delta_{IJ} + B_{IJ}(X)) \partial X^I \bar{\partial} X^J \quad (3.2)$$

and the classical equations of motion are

$$\partial\bar{\partial}X^I = \frac{1}{2}H^I{}_{JK}\partial X^J\bar{\partial}X^K. \quad (3.3)$$

More generally, we can consider flux backgrounds which are T^3 fibrations over a base manifold M , but, here, M is taken to be a point to avoid unnecessary technicalities.

When the H -flux is not zero, the theory (3.2) is not conformally invariant at the quantum level. Indeed, to lowest order in α' , the beta functions of the metric, anti-symmetric tensor field and dilaton are, respectively,

$$\begin{aligned} \beta(G_{IJ}) &= -\frac{1}{4}H_I{}^{KL}H_{JKL}, & \beta(B_{IJ}) &= -\frac{1}{2}\partial_K H^K{}_{IJ}, \\ \beta(\Phi) &= -\frac{23}{6} - \frac{1}{24}H_{IJK}H^{IJK}. \end{aligned} \quad (3.4)$$

The theory becomes conformal (though subcritical) only when the H -flux is constant and very weak, so that $\mathcal{O}(H^2)$ terms are negligible. In the following, we choose to work in this dilute flux approximation to implement T-dualities and denote the corresponding conformal field theory where all operations will be made by CFT_H .

The defining background, which is called H -flux vacuum of closed string theory, exhibits Abelian isometries, but their number depends on the choice of the anti-symmetric tensor field. B_{IJ} is taken to be linear in the toroidal coordinates, so that H_{IJK} is constant. A particularly convenient choice is

$$B_{12} = H X^3, \quad B_{23} = 0 = B_{31} \quad (3.5)$$

so that CFT_H exhibits two commuting isometries labeled by $a = 1, 2$, whereas $I = 1, 2, 3$. Another convenient choice is

$$B_{12} = \frac{H}{2} X^3, \quad B_{23} = 0, \quad B_{31} = \frac{H}{2} X^2, \quad (3.6)$$

which accounts for only one isometry associated to the coordinate X^1 . Finally, there is yet another (more symmetric) choice

$$B_{12} = \frac{H}{3} X^3, \quad B_{23} = \frac{H}{3} X^1, \quad B_{31} = \frac{H}{3} X^2 \quad (3.7)$$

in which case none of the isometries is manifest and Buscher's rules, which involve the field B rather than H , are not directly applicable. Yet, the various choices of the 2-form field B are related by gauge transformations $B \rightarrow B + d\Lambda$ for appropriate choices

of Λ and the T-dual faces of the model should be gauge independent.

In the following, we stick to the choice (3.5) for B , which makes manifest the maximum number of isometries of the model. The H -flux background and its T-dual faces belong in the class of toroidal fibrations and they exhibit non-trivial monodromies. These mathematical aspects of the model, as well as the description of the so called f -flux, Q -flux and R -flux backgrounds that arise by successive T-duality transformations are discussed in detail in Appendices A and B, respectively, together with the monodromies of their currents that encode non-trivial closed string boundary conditions.

3.1 T-dual coordinates

Consider the solution to the classical equations of motion of the H -flux model in terms of free fields, which to linear order in H reads

$$X^I(z, \bar{z}) = X_{0,L}^I(z) + X_{0,R}^I(\bar{z}) + \frac{1}{2} H^I{}_{JK} X_{0,L}^J(z) X_{0,R}^K(\bar{z}) \quad (3.8)$$

for all $I = 1, 2, 3$. This expression is independent of the gauge choice for the 2-form field B , which only affects the form of the dual coordinates. Our task is to find similar expressions for the dual coordinates in terms of free fields for the choice (3.5).

Recall that the dual coordinate to X^1 is defined by the following general relations

$$\partial \tilde{X}_1 = E_{11} \partial X^1 + E_{1i} \partial X^i, \quad \bar{\partial} \tilde{X}_1 = -(E_{11} \bar{\partial} X^1 + E_{1i} \bar{\partial} X^i). \quad (3.9)$$

For the model at hand we have $E_{11} = 1$ and $E_{1i} = -E_{i1} = B_{1i}$ and so

$$\partial \tilde{X}_1 = \partial X^1 - H X^3 \partial X^2, \quad \bar{\partial} \tilde{X}_1 = -\bar{\partial} X^1 - H X^3 \bar{\partial} X^2. \quad (3.10)$$

Substituting the free field realization (3.8) of the toroidal coordinates, we obtain

$$\partial \tilde{X}_1 = \partial X_{0,L}^1 - H X_{0,L}^3 \partial X_{0,L}^2 - \frac{H}{2} \partial (X_{0,L}^2 X_{0,R}^3 + X_{0,L}^3 X_{0,R}^2), \quad (3.11)$$

$$\bar{\partial} \tilde{X}_1 = -\bar{\partial} X_{0,R}^1 - H X_{0,R}^3 \bar{\partial} X_{0,R}^2 - \frac{H}{2} \bar{\partial} (X_{0,L}^2 X_{0,R}^3 + X_{0,L}^3 X_{0,R}^2). \quad (3.12)$$

Integration is straightforward and it yields the following result for the dual coordinate

$$\begin{aligned} \tilde{X}_1(z, \bar{z}) &= X_{0,L}^1(z) - X_{0,R}^1(\bar{z}) - \frac{H}{2} \left(X_{0,L}^2(z) X_{0,R}^3(\bar{z}) + X_{0,L}^3(z) X_{0,R}^2(\bar{z}) \right) - \\ &H \int^z X_{0,L}^3(w) \partial X_{0,L}^2(w) dw - H \int^{\bar{z}} X_{0,R}^3(\bar{w}) \bar{\partial} X_{0,R}^2(\bar{w}) d\bar{w}, \end{aligned} \quad (3.13)$$

whereas the corresponding expression for the original coordinate X^1 , following from eq. (3.8), is

$$X^1(z, \bar{z}) = X_{0,L}^1(z) + X_{0,R}^1(\bar{z}) + \frac{H}{2} \left(X_{0,L}^2(z) X_{0,R}^3(\bar{z}) - X_{0,L}^3(z) X_{0,R}^2(\bar{z}) \right). \quad (3.14)$$

Next, we define the fields

$$X_L^1(z, \bar{z}) = \frac{1}{2}(X^1 + \tilde{X}_1), \quad X_R^1(z, \bar{z}) = \frac{1}{2}(X^1 - \tilde{X}_1) \quad (3.15)$$

in terms of which T-duality reads

$$X_L^1(z, \bar{z}) \rightarrow X_L^1(z, \bar{z}), \quad X_R^1(z, \bar{z}) \rightarrow -X_R^1(z, \bar{z}). \quad (3.16)$$

Their free field realization follows from above, leading to the expressions

$$\begin{aligned} X_L^1(z, \bar{z}) &= X_{0,L}^1(z) - \frac{H}{2} X_{0,L}^3(z) X_{0,R}^2(\bar{z}) - \frac{H}{2} \int^z X_{0,L}^3(w) \partial X_{0,L}^2(w) dw - \\ &\quad \frac{H}{2} \int^{\bar{z}} X_{0,R}^3(\bar{w}) \bar{\partial} X_{0,R}^2(\bar{w}) d\bar{w}, \end{aligned} \quad (3.17)$$

$$\begin{aligned} X_R^1(z, \bar{z}) &= X_{0,R}^1(\bar{z}) + \frac{H}{2} X_{0,L}^2(z) X_{0,R}^3(\bar{z}) + \frac{H}{2} \int^z X_{0,L}^3(w) \partial X_{0,L}^2(w) dw + \\ &\quad \frac{H}{2} \int^{\bar{z}} X_{0,R}^3(\bar{w}) \bar{\partial} X_{0,R}^2(\bar{w}) d\bar{w}. \end{aligned} \quad (3.18)$$

Thus, the effect of T-duality, going from the H -flux to the f -flux background, can be neatly expressed in terms of free fields as follows,

$$X_{0,L}^1(z) \rightarrow X_{0,L}^1(z), \quad (3.19)$$

$$\begin{aligned} X_{0,R}^1(\bar{z}) &\rightarrow -X_{0,R}^1(\bar{z}) - H X_{0,L}^2(z) X_{0,R}^3(\bar{z}) \\ &\quad - H \int^z X_{0,L}^3(w) \partial X_{0,L}^2(w) dw - H \int^{\bar{z}} X_{0,R}^3(\bar{w}) \bar{\partial} X_{0,R}^2(\bar{w}) d\bar{w}, \end{aligned} \quad (3.20)$$

leaving $X_{0,L}^i(z)$ and $X_{0,R}^i(\bar{z})$ inert for the remaining toroidal coordinates $i = 2, 3$.

Let us now prepare to pass to the Q -flux model by performing T-duality along the direction X^2 of the f -flux model. On the f -flux background there is an off-diagonal component of the metric $G_{12} = fX^3$, identifying $H \equiv f$, whereas all diagonal components are 1 as before. Furthermore, the 2-form field vanishes. The dual coordinate to X^2 is defined by specializing the defining relations (3.9) to the f -flux model whose coordinates are \tilde{X}_1, X^2, X^3 . We have

$$\partial \tilde{X}_2 = \partial X^2 + H X^3 \partial \tilde{X}_1, \quad \bar{\partial} \tilde{X}_2 = -\bar{\partial} X^2 - H X^3 \bar{\partial} \tilde{X}_1. \quad (3.21)$$

Employing equation (3.10) we can eliminate \tilde{X}_1 and express everything in terms of the coordinates X^I of the original H -flux model. To linear order in $H \equiv f \equiv Q$ the result is

$$\partial\tilde{X}_2 = \partial X^2 + HX^3\partial X^1, \quad \bar{\partial}\tilde{X}_2 = -\bar{\partial}X^2 + HX^3\bar{\partial}X^1. \quad (3.22)$$

Note that these are precisely the defining relations of \tilde{X}_2 if we were to perform a T-duality of the H -flux model along the X^2 direction alone. Thus, the dual coordinates can be invariantly defined using the H -flux model and it does not matter if another duality has been performed first.

According to equation (3.8), the original coordinate X^2 admits the following free field realization,

$$X^2(z, \bar{z}) = X_{0,L}^2(z) + X_{0,R}^2(\bar{z}) - \frac{H}{2} \left(X_{0,L}^1(z)X_{0,R}^3(\bar{z}) - X_{0,L}^3(z)X_{0,R}^1(\bar{z}) \right). \quad (3.23)$$

Then, comparing (3.10) to (3.22) and (3.14) to (3.23), we see that the corresponding expressions are simply related by exchanging the coordinate indices 1 and 2 and flipping the sign of H at the same time. Implementing these operations to equations (3.17) and (3.18), we immediately obtain the following result for the left- and right-components of X^2 and \tilde{X}_2 , respectively,

$$\begin{aligned} X_L^2(z, \bar{z}) &= X_{0,L}^2(z) + \frac{H}{2} X_{0,L}^3(z)X_{0,R}^1(\bar{z}) + \frac{H}{2} \int^z X_{0,L}^3(w)\partial X_{0,L}^1(w)dw + \\ &\quad \frac{H}{2} \int^{\bar{z}} X_{0,R}^3(\bar{w})\bar{\partial} X_{0,R}^1(\bar{w})d\bar{w}, \end{aligned} \quad (3.24)$$

$$\begin{aligned} X_R^2(z, \bar{z}) &= X_{0,R}^2(\bar{z}) - \frac{H}{2} X_{0,L}^1(z)X_{0,R}^3(\bar{z}) - \frac{H}{2} \int^z X_{0,L}^3(w)\partial X_{0,L}^1(w)dw - \\ &\quad \frac{H}{2} \int^{\bar{z}} X_{0,R}^3(\bar{w})\bar{\partial} X_{0,R}^1(\bar{w})d\bar{w}. \end{aligned} \quad (3.25)$$

Finally, combining the above, we see that the effect of T-duality, going from the H -flux to the Q -flux background, can be neatly expressed in terms of free fields as

$$X_{0,L}^1(z) \rightarrow X_{0,L}^1(z) - HX_{0,L}^3(z)X_{0,R}^2(\bar{z}) - H \int^{\bar{z}} X_{0,R}^3(\bar{w})\bar{\partial} X_{0,R}^2(\bar{w})d\bar{w}, \quad (3.26)$$

$$X_{0,R}^1(\bar{z}) \rightarrow -X_{0,R}^1(\bar{z}) - HX_{0,L}^2(z)X_{0,R}^3(\bar{z}) - H \int^z X_{0,L}^3(w)\partial X_{0,L}^2(w)dw \quad (3.27)$$

and

$$X_{0,L}^2(z) \rightarrow X_{0,L}^2(z) + HX_{0,L}^3(z)X_{0,R}^1(\bar{z}) + H \int^{\bar{z}} X_{0,R}^3(\bar{w})\bar{\partial} X_{0,R}^1(\bar{w})d\bar{w}, \quad (3.28)$$

$$X_{0,R}^2(\bar{z}) \rightarrow -X_{0,R}^2(\bar{z}) + HX_{0,L}^1(z)X_{0,R}^3(\bar{z}) + H \int^z X_{0,L}^3(w)\partial X_{0,L}^1(w)dw, \quad (3.29)$$

whereas $X_{0,L}^3(z)$ and $X_{0,R}^3(\bar{z})$ remain inert.

3.2 Currents and generalized coordinates

The Roček-Verlinde approach of gauging the isometries does not require conformality and, as such, it is also applicable to tori with large H -flux. Here, we are limiting ourselves to CFT_H . Then, the corresponding currents (2.42) take the simple form

$$\mathcal{J}^a = \partial X^a, \quad \bar{\mathcal{J}}^a = \bar{\partial} X^a \quad (3.30)$$

in terms of the toroidal coordinates $a = 1, 2$. Using the solution (3.8) of the classical equations of motion of the H -flux model with very weak constant flux, the Roček-Verlinde parent currents take following form in terms of free fields,

$$\begin{aligned} \mathcal{J}^I(z, \bar{z}) &= \partial X^I(z, \bar{z}) = \partial X_{0,L}^I(z) + \frac{1}{2} H^I{}_{JK} \partial X_{0,L}^J(z) X_{0,R}^K(\bar{z}) \\ \bar{\mathcal{J}}^I(z, \bar{z}) &= \bar{\partial} X^I(z, \bar{z}) = \bar{\partial} X_{0,R}^I(\bar{z}) + \frac{1}{2} H^I{}_{JK} X_{0,L}^J(z) \bar{\partial} X_{0,R}^K(\bar{z}). \end{aligned} \quad (3.31)$$

Here, the index I is meant to be $a = 1, 2$, but for later use we also extend the definition to encompass $\mathcal{J}^3(z, \bar{z}) := \partial X^3(z, \bar{z})$ and $\bar{\mathcal{J}}^3(z, \bar{z}) := \bar{\partial} X^3(z, \bar{z})$, although the latter have no analogue in the enlarged theory S_{LR} .

The currents \mathcal{J}^a and $\bar{\mathcal{J}}^a$ are chiral fields of the enlarged parent theory S_{LR} , i.e., $\bar{\partial} \mathcal{J}^a = 0 = \partial \bar{\mathcal{J}}^a$, and, thus, they can be readily used to describe the dual faces of CFT_H by the automorphisms $\mathcal{J}^a \rightarrow \mathcal{J}^a$ and $\bar{\mathcal{J}}^a \rightarrow -\bar{\mathcal{J}}^a$. However, these currents are not chiral in the context of the reduced child theory CFT_H , since there are obstructions of order H , which arise by the field equations (3.3). They rather satisfy the non-chiral conservation laws

$$\bar{\partial} \mathcal{J}^I - \partial \bar{\mathcal{J}}^I = 0, \quad (3.32)$$

which are immediate and obvious for all $I = 1, 2, 3$. The monodromy properties of those currents are described in Appendix B for all T-dual faces of the constant flux vacua.

As emphasized before, this difference is attributed to the simple fact that the Roček-Verlinde currents do not commute with the BRST charge of the gauged sigma-model associated to the enlarged theory S_{LR} . Our results about non-commutativity and/or non-associativity of the (appropriately defined) coordinates of the dual faces of CFT_H that will be discussed later rely on this subtle point. In fact, the cohomological interpretation of the resulting algebraic structures in terms of 2- and 3-cocycles can be solely described in terms of the BRST (i.e., Lie algebra) coboundary operator and it is inherited in the reduced theory by gauging. Thus, non-commutativity and/or non-

associativity arise only in the reduced theory and not in the enlarged theory, which is manifestly commutative and associative. This interpretation, which will be pursued later in the discussions, ties up nicely with the idea that the enlarged theory S_{LR} can be used further to provide the world-sheet description of double field theory.

The coordinates X^I do not transform in a simple linear way under T-duality. However, we can define *generalized* coordinates $\mathcal{X}^I(z, \bar{z}) = \mathcal{X}_L^I(z, \bar{z}) + \mathcal{X}_R^I(z, \bar{z})$ that transform nicely under T-duality, as dictated by eq. (2.39). The generalized coordinates are integrals of the Roček-Verlinde currents \mathcal{J}^I and $\bar{\mathcal{J}}^I$, given by eq.(3.31), of the following form,

$$\mathcal{X}_L^I(z, \bar{z}) = \int_a^z \mathcal{J}^I(z', \bar{z}') dz', \quad \mathcal{X}_R^I(z, \bar{z}) = \int_{\bar{a}}^{\bar{z}} \bar{\mathcal{J}}^I(z', \bar{z}') d\bar{z}'. \quad (3.33)$$

Here a and \bar{a} are some complex integration constants. The expressions above involve integration over the world-sheet coordinate σ in the (τ, σ) -plane in radial quantization, which is adopted hereafter, and, therefore, they should be defined as circular integrals in the complex z -plane, keeping $|z|$ and $|\bar{z}|$ fixed (and, thus, it is also implicitly assumed that $|z| = |a|$ and $|\bar{z}| = |\bar{a}|$). The corresponding fields \mathcal{X}^I should be regarded as the proper generalized string coordinates of the σ -model to first order in perturbation theory. The commutation relations of the coordinates \mathcal{X}^I and \mathcal{X}^J , and not of X^I and X^J , and their dual counterparts, will be in focus later.

The different meaning of $(\mathcal{J}^I, \bar{\mathcal{J}}^I)$ in CFT_H and S_{LR} is also reflected in their operator product expansions. In CFT_H , we use the free field 2-point functions

$$\langle \partial X_{0,L}^I(z) \partial X_{0,L}^{I'}(w) \rangle = \frac{\delta^{I,I'}}{2(z-w)^2}, \quad \langle \bar{\partial} X_{0,R}^I(\bar{z}) \bar{\partial} X_{0,R}^{I'}(\bar{w}) \rangle = \frac{\delta^{I,I'}}{2(\bar{z}-\bar{w})^2} \quad (3.34)$$

to obtain the following expansions, to linear order in H ,

$$\mathcal{J}^I(z, \bar{z}) \mathcal{J}^{I'}(w, \bar{w}) = \frac{\delta^{I,I'}}{2(z-w)^2} + \frac{1}{4} H^{II'}{}_K \frac{\bar{z}-\bar{w}}{(z-w)^2} \bar{\mathcal{J}}^K(w, \bar{w}) + \dots, \quad (3.35)$$

$$\bar{\mathcal{J}}^I(z, \bar{z}) \bar{\mathcal{J}}^{I'}(w, \bar{w}) = \frac{\delta^{I,I'}}{2(\bar{z}-\bar{w})^2} + \frac{1}{4} H^{II'}{}_K \frac{z-w}{(\bar{z}-\bar{w})^2} \mathcal{J}^K(w, \bar{w}) + \dots, \quad (3.36)$$

$$\mathcal{J}^I(z, \bar{z}) \bar{\mathcal{J}}^{I'}(w, \bar{w}) = \frac{1}{4} H^{II'}{}_K \left(\frac{\mathcal{J}^K(w, \bar{w})}{\bar{z}-\bar{w}} - \frac{\bar{\mathcal{J}}^K(w, \bar{w})}{z-w} \right) + \dots, \quad (3.37)$$

where dots denote non-singular terms. There is no holomorphic factorization of the current algebra because of the mixed terms involving $X_{0,L}^I(z)$ and $X_{0,R}^I(\bar{z})$ in the free field realization (3.31) via the fluxes. In the enlarged theory S_{LR} , on the other hand, holomorphic factorization is guaranteed by construction, leading to the operation

product expansion of a $U(1)_L \times U(1)_R$ current algebra for each generator,

$$\mathcal{J}^I(z)\mathcal{J}^{I'}(w) = \frac{\delta^{I,I'}}{2(z-w)^2}, \quad \bar{\mathcal{J}}^I(\bar{z})\bar{\mathcal{J}}^{I'}(\bar{w}) = \frac{\delta^{I,I'}}{2(\bar{z}-\bar{w})^2}, \quad \mathcal{J}^I(z)\bar{\mathcal{J}}^{I'}(\bar{w}) = 0, \quad (3.38)$$

up to non-singular terms that are irrelevant.

Strictly speaking, the index I should be restricted to the Killing directions 1 and 2, but their operator product expansion gives rise to \mathcal{J}^3 and $\bar{\mathcal{J}}^3$ in the context of CFT_H . These additional currents are not associated to an isometry and they are the ones that obstruct holomorphic factorization. Note, however, that it is not consistent with the free field realization (3.31) to project them out by forming an ideal of the algebra. Their appearance in the same multiplet with the ordinary Roček-Verlinde currents should be held responsible for the emergence of the non-commutative and non-associative structure when T-duality is formally performed in the third non-Killing direction to yield the non-geometric R -flux vacuum of closed string theory.

4 Canonical T-duality and non-commutativity

This section is devoted to the derivation of the non-commutative and non-associative relations among the generalized coordinates appearing in the different T-dual faces of the constant flux background. As such, it contains our main results, whereas all other material in the paper can be regarded as prerequisite for their derivation.

4.1 Preliminaries: the case of free fields

First, we briefly discuss the commutation relations among free fields, using the formalism of conformal field theory, as warm up, while preparing the ground for the derivation of the commutation relations among the generalized coordinates of the flux backgrounds by similar techniques. Recall that the free fields $X_0^I(z, \bar{z}) = X_{0,L}^I(z) + X_{0,R}^I(\bar{z})$ possesses the following mode expansion

$$\begin{aligned} X_{0,L}^I(z) &= x_{0,L}^I - ip_L^I \log z + i \sum_{n \neq 0} \frac{1}{n} \alpha_n^I z^{-n}, \\ X_{0,R}^I(\bar{z}) &= x_{0,R}^I - ip_R^I \log \bar{z} + i \sum_{n \neq 0} \frac{1}{n} \bar{\alpha}_n^I \bar{z}^{-n}, \end{aligned} \quad (4.39)$$

introducing complex coordinates as $z = e^{\tau - i\sigma}$ and $\bar{z} = e^{\tau + i\sigma}$. All commutators are derived from the free field propagators of the left- and right-moving fields, which are

taken to be

$$\begin{aligned}
\langle X_{0,L}^I(z) X_{0,L}^J(w) \rangle &= \frac{1}{2} \delta^{IJ} \log(z-w), \\
\langle X_{0,R}^I(\bar{z}) X_{0,R}^J(\bar{w}) \rangle &= \frac{1}{2} \delta^{IJ} \log(\bar{z}-\bar{w}), \\
\langle X_{0,L}^I(z) X_{0,R}^J(\bar{w}) \rangle &= 0
\end{aligned} \tag{4.40}$$

with the appropriate normalization of the string tension.

Let us compute the equal time commutator of the left-moving coordinate $X_{0,L}^I(z)$ with its derivative $\partial_w X_{0,L}^J(w)$. Time ordering becomes radial ordering in the complex z -plane, and, thus, the equal time commutator $[X_{0,L}^I(\tau, \sigma), \partial_w X_{0,L}^J(\tau, \sigma')]$ is defined as follows, taking into account the δ -prescription,

$$\begin{aligned}
[X_{0,L}^I(\tau, \sigma), \partial_w X_{0,L}^J(\tau, \sigma')] &= [X_{0,L}^I(z), \partial_w X_{0,L}^J(w)]_{|z|=|w|} = \\
&= \lim_{\delta \rightarrow 0_+} \left((X_{0,L}^I(z) \partial_w X_{0,L}^J(w))_{|z|=|w|+\delta} - (\partial_w X_{0,L}^J(w) X_{0,L}^I(z))_{|z|=|w|-\delta} \right). \tag{4.41}
\end{aligned}$$

Using the two-point function (4.40), we obtain by contraction that

$$[X_{0,L}^I(z), \partial_w X_{0,L}^J(w)]_{|z|=|w|} = -\frac{1}{2} \delta^{IJ} \lim_{\delta \rightarrow 0_+} \left(\frac{1}{z-w} \Big|_{|z|=|w|+\delta} - \frac{1}{z-w} \Big|_{|z|=|w|-\delta} \right) \tag{4.42}$$

and so, choosing $z = e^{\pm\delta-i\sigma}$, $w = e^{-i\sigma'}$, i.e., $\tau = \pm\delta$, $\tau' = 0$ and $\varphi = \sigma - \sigma'$, the above commutator works out to be

$$\begin{aligned}
[X_{0,L}^I(z), \partial_w X_{0,L}^J(w)]_{|z|=|w|} &= -\frac{1}{2} \delta^{IJ} \lim_{\delta \rightarrow 0_+} \left(\frac{e^{-\delta+i\sigma}}{1-e^{-\delta+i\varphi}} + \frac{e^{i\sigma'}}{1-e^{-\delta-i\varphi}} \right) \\
&= -\frac{1}{2} \delta^{IJ} \left(e^{i\sigma} \sum_{m=0}^{\infty} e^{im\varphi} + e^{i\sigma'} \sum_{m=0}^{\infty} e^{-im\varphi} \right) \\
&= -\pi \delta^{IJ} e^{i\sigma'} \delta(\varphi) = i\pi \delta^{IJ} \delta(z-w). \tag{4.43}
\end{aligned}$$

Next, we compute the equal time commutator of $X_{0,L}^I(z)$ with $X_{0,L}^J(w)$, representing $X_{0,L}^J(w)$ by the integral

$$X_{0,L}^J(w) = \int^w d\tilde{w} \partial_{\tilde{w}} X_{0,L}^J(\tilde{w}) \tag{4.44}$$

in view of the generalizations that will be considered later. The integral goes over some path in the complex w -plane which is not a closed contour. The computation of the equal time commutator dictates choosing a path of constant time, thus integrating

along the circular direction in the complex plane by rewriting

$$X_{0,L}^I(w) = -i \int^{\sigma'} d\tilde{\sigma}' e^{-i\tilde{\sigma}'} \partial_{\tilde{w}} X_{0,L}^I(\tilde{w}). \quad (4.45)$$

Then, the commutator $[X_{0,L}^I(\tau, \sigma), X_{0,L}^J(\tau, \sigma')]$ is computed in steps as follows,

$$\begin{aligned} [X_{0,L}^I(\tau, \sigma), X_{0,L}^J(\tau, \sigma')] &= [X_{0,L}^I(z), X_{0,L}^J(w)]_{|z|=|w|} \\ &= [X_{0,L}^I(z), \int^w d\tilde{w} \partial_{\tilde{w}} X_{0,L}^J(\tilde{w})]_{|z|=|w|} \\ &= -i \int^{\sigma'} d\tilde{\sigma}' e^{-i\tilde{\sigma}'} [X_{0,L}^I(z), \partial_{\tilde{w}} X_{0,L}^J(\tilde{w})]_{|z|=|\tilde{w}|} \\ &= i\pi \delta^{IJ} \int^\varphi d\tilde{\varphi} \delta(\tilde{\varphi}) = \frac{i\pi}{2} \delta^{IJ} \epsilon(\varphi), \end{aligned} \quad (4.46)$$

using also the identity (C.2) found in Appendix C. The same result is obtained directly from the logarithmic expression (4.40) for the propagator, without making use of the integral formula above.

Using the same techniques, one obtains similar expressions for the commutators for the right-moving free fields

$$[X_{0,R}^I(\tau, \sigma), X_{0,R}^J(\tau, \sigma')] = [X_{0,R}^I(\bar{z}), X_{0,R}^J(\bar{w})]_{|\bar{z}|=|\bar{w}|} = -\frac{i\pi}{2} \delta^{IJ} \epsilon(\varphi). \quad (4.47)$$

Finally, combining the expressions (4.46) and (4.47), the equal time commutators among $X_0^I(z, \bar{z}) = X_{0,L}^I(z) + X_{0,R}^I(\bar{z})$ and the dual coordinates $\tilde{X}_0^J(z, \bar{z}) = X_{0,L}^J(z) - X_{0,R}^J(\bar{z})$ work out to be

$$\begin{aligned} [X_0^I(\tau, \sigma), X_0^J(\tau, \sigma')] &= [X_{0,L}^I(z), X_{0,L}^J(w)]_{|z|=|w|} + [X_{0,R}^I(\bar{z}), X_{0,R}^J(\bar{w})]_{|\bar{z}|=|\bar{w}|} = 0, \\ [\tilde{X}_0^I(\tau, \sigma), \tilde{X}_0^J(\tau, \sigma')] &= [X_{0,L}^I(z), X_{0,L}^J(w)]_{|z|=|w|} + [X_{0,R}^I(\bar{z}), X_{0,R}^J(\bar{w})]_{|\bar{z}|=|\bar{w}|} = 0, \\ [X_0^I(\tau, \sigma), \tilde{X}_0^J(\tau, \sigma')] &= [X_{0,L}^I(z), X_{0,L}^J(w)]_{|z|=|w|} - [X_{0,R}^I(\bar{z}), X_{0,R}^J(\bar{w})]_{|\bar{z}|=|\bar{w}|} \\ &= i\pi \delta^{IJ} \epsilon(\varphi), \end{aligned} \quad (4.48)$$

as expected. Since $\epsilon(0) = 0$, we have $[X_0^I(\tau, \sigma), \tilde{X}_0^J(\tau, \sigma')] = 0$ for $\sigma = \sigma'$.

4.2 Non-commutativity for interacting fields with constant fluxes

We are now in position to extract the general structure of the commutation relations among the closed string coordinates for toroidal backgrounds with constant flux by

conformal field theory techniques based on the formalism introduced in the previous sections. Analogous calculations were performed earlier, first in ref. [27] and then in [32], using the world-sheet operator formalism of parabolic models (but see also [38] for the computation of Dirac brackets among the coordinates), and they were subsequently extended to backgrounds with elliptic monodromies in ref. [29]. As will be seen here, the CFT treatment is more efficient, making the origin of non-commutativity more transparent compared to the previous approaches. The main advantage of the CFT formalism is that T-duality acts in a simple way, flipping the sign of the right-moving generalized coordinates.

The σ -model coordinates $X^I(z, \bar{z})$ are not anymore free fields, but they interact with each other in the presence of fluxes. Since the left- and right-moving coordinates $X_L^I(z, \bar{z})$ and $X_R^I(z, \bar{z})$ are not directly related to currents that transform linearly under duality transformations, we were led to consider the *generalized* coordinates \mathcal{X}^I , as discussed in the previous section, which are the integrals of the Roček-Verlinde currents $\mathcal{J}_L^I(z, \bar{z})$ and $\tilde{\mathcal{J}}_R^I(z, \bar{z})$. We have

$$\mathcal{X}^I(z, \bar{z}) = \mathcal{X}_L^I(z, \bar{z}) + \mathcal{X}_R^I(z, \bar{z}), \quad (4.49)$$

letting

$$\mathcal{X}_L^I(z, \bar{z}) = \int_a^z \mathcal{J}_L^I(z', \bar{z}') dz', \quad \mathcal{X}_R^I(z, \bar{z}) = \int_{\bar{a}}^{\bar{z}} \tilde{\mathcal{J}}_R^I(z', \bar{z}') d\bar{z}'. \quad (4.50)$$

As explained earlier, T-duality in the I -direction acts on the currents $\mathcal{J}_L^I(z, \bar{z})$ and $\tilde{\mathcal{J}}_R^I(z, \bar{z})$ as well as on the corresponding generalized coordinates $\mathcal{X}_L^I(z, \bar{z})$ and $\mathcal{X}_R^I(z, \bar{z})$ in a simple linear way,

$$\begin{aligned} \mathcal{J}_L^I(z, \bar{z}) &\longrightarrow \mathcal{J}_L^I(z, \bar{z}), & \tilde{\mathcal{J}}_R^I(z, \bar{z}) &\longrightarrow -\tilde{\mathcal{J}}_R^I(z, \bar{z}), \\ \mathcal{X}_L^I(z, \bar{z}) &\longrightarrow \mathcal{X}_L^I(z, \bar{z}), & \mathcal{X}_R^I(z, \bar{z}) &\longrightarrow -\mathcal{X}_R^I(z, \bar{z}). \end{aligned} \quad (4.51)$$

This does not mean that the T-dual faces of the interacting theory are self-dual, but it is a statement that T-duality maps the generalized coordinates of one flux background to the generalized coordinates of another and vice-versa.

It follows from the definition of the coordinates $\mathcal{X}_L^I(z, \bar{z})$ as integrals of the non-chiral currents $\mathcal{J}_L^I(z, \bar{z})$ that such integrals are not anymore path-independent, but, in fact, they depend on the chosen path in the complex plane; the same is also true for the corresponding right-moving objects. Hence, it is necessary to specify the path of integration in order to assign a well defined meaning to the generalized coordinates \mathcal{X}_L^I and \mathcal{X}_R^I . Here, we apply the radial quantization for the computation of the commutators by choosing circular paths around the origin of the complex plane, keeping $|z|$ and $|\bar{z}|$ fixed in the integrals, as in the case of free fields. Then, with this prescription

in mind, which we adopt in the following, the generalized coordinates are defined as follows,

$$\mathcal{X}_L^I(z, \bar{z}) = -i \int^\sigma d\tilde{\sigma} e^{-i\tilde{\sigma}} \mathcal{J}_L^I(z, \bar{z}), \quad \mathcal{X}_R^I(z, \bar{z}) = i \int^\sigma d\tilde{\sigma} e^{i\tilde{\sigma}} \bar{\mathcal{J}}_R^I(z, \bar{z}), \quad (4.52)$$

letting $z = e^{\tau-i\sigma}$ and $\bar{z} = e^{\tau+i\sigma}$. It follows immediately that the non-holomorphic terms of $\mathcal{J}_L^I(z, \bar{z})$, which depend on \bar{z} , also contribute to the integral over z ; similar remarks apply to the holomorphic part of the currents $\bar{\mathcal{J}}_R^I(z, \bar{z})$.

Let us now consider the equal time commutators among the generalized coordinates $\mathcal{X}^I(z, \bar{z})$ and $\mathcal{X}^J(w, \bar{w})$,

$$\begin{aligned} [\mathcal{X}^I(\tau, \sigma), \mathcal{X}^J(\tau, \sigma')] &= [\mathcal{X}^I(z, \bar{z}), \mathcal{X}^J(w, \bar{w})]_{|z|=|w|} = \\ &= \lim_{\delta \rightarrow 0^+} \left((\mathcal{X}^I(z, \bar{z}) \mathcal{X}^J(w, \bar{w}))_{|z|=|w|+\delta} - (\mathcal{X}^J(w, \bar{w}) \mathcal{X}^I(z, \bar{z}))_{|z|=|w|-\delta} \right). \end{aligned} \quad (4.53)$$

It is convenient to split these commutators in four different pieces, namely

$$\begin{aligned} \Theta_{LL}^{IJ}(\tau, \sigma, \sigma') &:= [\mathcal{X}_L^I(z, \bar{z}), \mathcal{X}_L^J(w, \bar{w})]_{|z|=|w|}, \\ \Theta_{RR}^{IJ}(\tau, \sigma, \sigma') &:= [\mathcal{X}_R^I(z, \bar{z}), \mathcal{X}_R^J(w, \bar{w})]_{|z|=|w|}, \\ \Theta_{LR}^{IJ}(\tau, \sigma, \sigma') &:= [\mathcal{X}_L^I(z, \bar{z}), \mathcal{X}_R^J(w, \bar{w})]_{|z|=|w|}, \\ \Theta_{RL}^{IJ}(\tau, \sigma, \sigma') &:= [\mathcal{X}_R^I(z, \bar{z}), \mathcal{X}_L^J(w, \bar{w})]_{|z|=|w|}, \end{aligned} \quad (4.54)$$

letting $z = e^{\tau-i\sigma}$ and $w = e^{\tau'-i\sigma'}$. Note that apart from the left- and right-moving commutators Θ_{LL}^{IJ} and Θ_{RR}^{IJ} , which are in general non-vanishing, the cross terms Θ_{LR}^{IJ} and Θ_{RL}^{IJ} can also be non-zero, because the operator products of $\mathcal{J}_L^I(z, \bar{z})$ with $\mathcal{J}_R^J(z, \bar{z})$ (and, likewise, of $\bar{\mathcal{J}}_R^I(z, \bar{z})$ with $\mathcal{J}_L^J(z, \bar{z})$) contain non-vanishing poles that contribute to their expressions. Adding the left- and right-moving pieces together, we obtain the following general structure for the commutation relations between $\mathcal{X}^I = \mathcal{X}_L^I + \mathcal{X}_R^I$ and $\mathcal{X}^J = \mathcal{X}_L^J + \mathcal{X}_R^J$,

$$\Theta^{IJ} = [\mathcal{X}^I, \mathcal{X}^J] = \Theta_{LL}^{IJ} + \Theta_{RR}^{IJ} + \Theta_{LR}^{IJ} + \Theta_{RL}^{IJ}. \quad (4.55)$$

As emphasized before, $\mathcal{X}_{L,R}^I(z, \bar{z})$ are the appropriate generalized string coordinates that transform nicely under T-duality. Then, starting from a given flux background and dualizing in the I -direction, the four commutators pieces transform, following (4.51), as

$$\Theta_{LL}^{IJ} \rightarrow \Theta_{LL}^{IJ}, \quad \Theta_{RR}^{IJ} \rightarrow -\Theta_{RR}^{IJ}, \quad \Theta_{LR}^{IJ} \rightarrow \Theta_{LR}^{IJ}, \quad \Theta_{RL}^{IJ} \rightarrow -\Theta_{RL}^{ij} \quad (4.56)$$

and, as a result, the total commutators in the dual background take the simple form

$$\Theta_{\text{dual}}^{IJ} = [\tilde{\mathcal{X}}^I, \mathcal{X}^J] = \Theta_{LL}^{IJ} - \Theta_{RR}^{IJ} + \Theta_{LR}^{IJ} - \Theta_{RL}^{IJ}. \quad (4.57)$$

Thus, starting from the H -flux background and applying two consecutive T-duality transformations in the 1- and 2-directions, we obtain in steps:

(i) Geometric H -flux background:

$$\Theta_H^{12} = \Theta_{LL}^{12} + \Theta_{RR}^{12} + \Theta_{LR}^{12} + \Theta_{RL}^{12}. \quad (4.58)$$

(ii) Geometric f -flux background: after a T-duality in the 1-direction, we obtain

$$\Theta_f^{12} = \Theta_{LL}^{12} - \Theta_{RR}^{12} + \Theta_{LR}^{12} - \Theta_{RL}^{12}. \quad (4.59)$$

(iii) Non-geometric Q -flux background: another T-duality in the 2-direction yields

$$\Theta_Q^{12} = \Theta_{LL}^{12} + \Theta_{RR}^{12} - \Theta_{LR}^{12} - \Theta_{RL}^{12}. \quad (4.60)$$

As will be seen next, some of these commutators are zero and others are non-vanishing, leading to non-commutativity in the corresponding flux vacua. The same method leads to non-associativity by making another step in the 3-direction to reach the non-geometric R -flux background.

4.3 Case by case computations

We will compute the individual pieces of the commutators among the generalized coordinates for all T-dual faces of the parabolic flux model, case by case, using the technical material contained in Appendix C.

4.3.1 Commutators of the H -flux background

Let us now apply this formalism to extract the commutators of the H -flux background. In the following, we compute the off-diagonal commutators between the generalized coordinates in the 1- and 2-directions of the H -flux background to linear order in the fluxes. The diagonal commutators between the generalized coordinates and their dual

counterparts in the same directions are identical to those found for the free fields in the previous subsection.

The left- and right-moving currents of the H -flux background in the 1- and 2- directions take the following form, according to eq. (3.31),

$$\begin{aligned}
\mathcal{J}_L^1(z, \bar{z}) &= \partial X_{0,L}^1(z) + \frac{1}{2}H(X_{0,R}^3(\bar{z}) \partial X_{0,L}^2(z) - X_{0,R}^2(\bar{z}) \partial X_{0,L}^3(z)) , \\
\mathcal{J}_L^2(z, \bar{z}) &= \partial X_{0,L}^2(z) - \frac{1}{2}H(X_{0,R}^3(\bar{z}) \partial X_{0,L}^1(z) - X_{0,R}^1(\bar{z}) \partial X_{0,L}^3(z)) , \\
\bar{\mathcal{J}}_R^1(z, \bar{z}) &= \bar{\partial} X_{0,R}^1(\bar{z}) + \frac{1}{2}H(X_{0,L}^2(z) \bar{\partial} X_{0,R}^3(\bar{z}) - X_{0,L}^3(z) \bar{\partial} X_{0,R}^2(\bar{z})) , \\
\bar{\mathcal{J}}_R^2(z, \bar{z}) &= \bar{\partial} X_{0,R}^2(\bar{z}) - \frac{1}{2}H(X_{0,L}^1(z) \bar{\partial} X_{0,R}^3(\bar{z}) - X_{0,L}^3(z) \bar{\partial} X_{0,R}^1(\bar{z})) . \quad (4.61)
\end{aligned}$$

From these currents we obtain the following *generalized* coordinates in the 1- and 2- directions of the three-torus,

$$\begin{aligned}
\mathcal{X}_L^1(z, \bar{z}) &= X_{0,L}^1(z) + \frac{1}{2}H \int^z dz' (X_{0,R}^3(\bar{z}') \partial' X_{0,L}^2(z') - X_{0,R}^2(\bar{z}') \partial' X_{0,L}^3(z')) , \\
\mathcal{X}_R^1(z, \bar{z}) &= X_{0,R}^1(\bar{z}) + \frac{1}{2}H \int^{\bar{z}} d\bar{z}' (X_{0,L}^2(z') \bar{\partial}' X_{0,R}^3(\bar{z}') - X_{0,L}^3(z') \bar{\partial}' X_{0,R}^2(\bar{z}')) , \\
\mathcal{X}_L^2(z, \bar{z}) &= X_{0,L}^2(z) - \frac{1}{2}H \int^z dz' (X_{0,R}^3(\bar{z}') \partial' X_{0,L}^1(z') - X_{0,R}^1(\bar{z}') \partial' X_{0,L}^3(z')) , \\
\mathcal{X}_R^2(z, \bar{z}) &= X_{0,R}^2(\bar{z}) - \frac{1}{2}H \int^{\bar{z}} d\bar{z}' (X_{0,L}^1(z') \bar{\partial}' X_{0,R}^3(\bar{z}') - X_{0,L}^3(z') \bar{\partial}' X_{0,R}^1(\bar{z}')) . \quad (4.62)
\end{aligned}$$

We start with the equal time commutator piece Θ_{LL}^{12} to linear order in H , which is

$$\begin{aligned}
\Theta_{LL}^{12}(z, w, \bar{z}, \bar{w}) &= [\mathcal{X}_L^1(z, \bar{z}), \mathcal{X}_L^2(w, \bar{w})] \\
&= -\frac{1}{2}H \left[X_{0,L}^1(z), \int^w \partial' X_{0,L}^1(z') X_{0,R}^3(\bar{z}') dz' \right] \\
&\quad - \frac{1}{2}H \left[X_{0,L}^2(w), \int^z \partial' X_{0,L}^2(z') X_{0,R}^3(\bar{z}') dz' \right] . \quad (4.63)
\end{aligned}$$

Its computation proceeds by taking single contractions of the operators that appear in the two terms above - there are no higher contractions - which effectively means that the commutators correspond to classical Poisson (respectively Dirac brackets) prior to

quantization. The following contractions contribute to Θ_{LL}^{12} ,

$$\begin{aligned} \Theta_{LL}^{12}(z, w, \bar{z}, \bar{w}) &= - \left[\frac{1}{2} H \int^w \left(\langle X_{0,L}^1(z) \partial' X_{0,L}^1(z') \rangle X_{0,R}^3(\bar{z}') \right) dz' \right. \\ &\quad \left. + \frac{1}{2} H \int^z \left(\langle X_{0,L}^2(w) \partial' X_{0,L}^2(z') \rangle X_{0,R}^3(\bar{z}') \right) dz' \right]_{|z|=|w|+\delta} \\ &\quad - \text{the other half of the commutator} \end{aligned} \quad (4.64)$$

It can be seen that the 1-point function of the operator $X_{0,R}^3(\bar{z}')$ remains in the commutator after the contraction. This looks rather unusual and one may naively expect that the net result for the commutator is zero. Note, however, that we are computing these expressions in a sector of the string Hilbert space where the left- and right-moving momenta p_L^3 and p_R^3 are not zero. For vanishing oscillator numbers, the string level matching constraint implies that

$$p_L^3 = \pm p_R^3. \quad (4.65)$$

Here, we concentrate on states with non-trivial winding number \tilde{p}^3 , having

$$\tilde{p}^3 = p_L^3 - p_R^3 \quad \text{and} \quad p_L^3 = -p_R^3. \quad (4.66)$$

This, in turn, implies that the momentum operators $\hat{p}_{L,R}^3$ possess non-vanishing matrix elements in the winding states,

$$\langle \tilde{p}^3 | \hat{p}_L^3 | \tilde{p}^3 \rangle = \frac{\tilde{p}^3}{2}, \quad \langle \tilde{p}^3 | \hat{p}_R^3 | \tilde{p}^3 \rangle = -\frac{\tilde{p}^3}{2}. \quad (4.67)$$

The existence of winding states is closely related to the non-trivial monodromy properties of the flux backgrounds, which are outlined in Appendices A and B. Specifically, the monodromy induces non-trivial transformations of the coordinates X^1 , X^2 and of the dual coordinates \tilde{X}_1 , \tilde{X}_2 when going around the third circular direction by making the shift

$$X^3 \rightarrow X^3 + 2\pi \quad (4.68)$$

and selects new closed string boundary conditions. In sectors with non-vanishing winding number \tilde{p}^3 , the shift $\sigma \rightarrow \sigma + 2\pi$ acts on X^3 in the standard way,

$$X^3(e^{-2i\pi}z, e^{2i\pi}\bar{z}) = X^3(z, \bar{z}) - 2\pi\tilde{p}^3, \quad (4.69)$$

whereas, for non-vanishing fluxes, the currents $\mathcal{J}_{L,R}^I$, with $I = 1, 2$, undergo non-

trivial transformation given in terms a flux dependent monodromy matrix \mathcal{M}_{IJ} , as

$$\mathcal{J}_{L,R}^I(e^{-2i\pi z}, e^{2i\pi \bar{z}}) = \mathcal{M}_{IJ} \mathcal{J}_{L,R}^J(z, \bar{z}). \quad (4.70)$$

The upshot of this discussion is that the commutators can be non-zero in string sectors with non-vanishing winding number. The physical origin of this behavior should be attributed to the modified closed string boundary conditions, which reflect the simple fact that non-local winding strings can capture the non-trivial global properties of the flux background, thus giving rise to non-commutativity among the generalized coordinates. We shall examine how this is precisely realized in all T-dual faces of the parabolic flux model. With these explanations in mind, let us now continue with the evaluation of the commutator pieces in the H -flux background and then use the results to extract the total commutators in all other T-dual faces of the model.

Using the mode expansion of free fields, the 1-point function of $X_{0,R}^3(\bar{z}')$, which is given by (4.64), turns out to be

$$\langle \tilde{p}^3 | X_{0,R}^3(\bar{z}') | \tilde{p}^3 \rangle = i \frac{\tilde{p}^3}{2} \log \bar{z}' \quad (4.71)$$

in sectors with non-trivial winding number. Furthermore, using the 2-point functions

$$\langle X_{0,L}^1(z) \partial' X_{0,L}^1(z') \rangle = -\frac{1}{2(z-z')}, \quad \langle X_{0,L}^2(w) \partial' X_{0,L}^2(z') \rangle = -\frac{1}{2(w-z')}, \quad (4.72)$$

we arrive at the following intermediate result for Θ_{LL}^{12} ,

$$\begin{aligned} \Theta_{LL}^{12}(z, w, \bar{z}, \bar{w}) &= \frac{i}{8} \tilde{p}^3 H \left[\int^w \left(\frac{\log \bar{z}'}{z-z'} \right) dz' + \int^z \left(\frac{\log \bar{z}'}{w-z'} \right) dz' \right]_{|z|=|w|+\delta} \\ &- \text{the other half of the commutator.} \end{aligned} \quad (4.73)$$

These integrals are evaluated in Appendix C (see, in particular, eq. (C.16)). Hence, we obtain

$$\Theta_{LL}^{12}(z, w, \bar{z}, \bar{w}) = -\frac{i\pi}{4} H \tilde{p}^3 \left(\frac{\varphi^2}{2\pi} - \frac{1}{\pi} \mathcal{L}i_2(e^{i\varphi}) \right). \quad (4.74)$$

Setting $\varphi = 0$, it yields

$$\Theta_{LL}^{12}(\tau, \sigma) = i H \tilde{p}^3 \frac{\pi^2}{12}. \quad (4.75)$$

Likewise, we obtain the following result for Θ_{RR} ,

$$\Theta_{RR}^{12}(z, w, \bar{z}, \bar{w}) = -\frac{i\pi}{4} H \tilde{p}^3 \left(\frac{\varphi^2}{2\pi} - \frac{1}{\pi} \mathcal{L}i_2(e^{i\varphi}) \right), \quad (4.76)$$

which for $\varphi = 0$ yields

$$\Theta_{RR}^{12}(\tau, \sigma) = iH \tilde{p}^3 \frac{\pi^2}{12}. \quad (4.77)$$

Next, we compute the commutator piece Θ_{LR} , which is given by the expression

$$\begin{aligned} \Theta_{LR}^{12}(z, \bar{z}, w, \bar{w}) &= [\mathcal{X}_L^1(z, \bar{z}), \mathcal{X}_R^2(w, \bar{w})] \\ &= -\frac{1}{2} H \left[X_{0,L}^1(z), \int^{\bar{w}} X_{0,L}^1(z') \bar{\partial}' X_{0,R}^3(\bar{z}') d\bar{z}' \right] \\ &\quad + \frac{1}{2} H \left[X_{0,R}^2(\bar{w}), \int^z \partial' X_{0,L}^3(z') X_{0,R}^2(\bar{z}') dz' \right]. \end{aligned} \quad (4.78)$$

Taking the single contractions, we find

$$\begin{aligned} \Theta_{LR}^{12}(z, w, \bar{z}, \bar{w}) &= \left[-\frac{1}{2} H \int^{\bar{w}} \langle X_{0,L}^1(z) X_{0,L}^1(z') \rangle \bar{\partial}' X_{0,R}^3(\bar{z}') d\bar{z}' \right. \\ &\quad \left. + \frac{1}{2} H \int^z \partial' X_{0,L}^3(z') \langle X_{0,R}^2(\bar{w}) X_{0,R}^2(\bar{z}') \rangle dz' \right]_{|z|=|w|+\delta} \\ &\quad - \text{the other half of the commutator} \\ &= \left[-\frac{i}{8} H \tilde{p}^3 \left(\int^{\bar{w}} dz' \frac{\log(z-z')}{\bar{z}'} + \int^z dz' \frac{\log(\bar{w}-\bar{z}')}{z'} \right) \right]_{|z|=|w|+\delta} \\ &\quad - \text{the other half of the commutator.} \end{aligned} \quad (4.79)$$

These integrals are evaluated in Appendix C (see, in particular, eq. (C.18)) and so we have

$$\Theta_{LR}^{12}(z, w, \bar{z}, \bar{w}) = \frac{i\pi}{4} H \tilde{p}^3 \left(\frac{\varphi^2}{2\pi} - \frac{1}{\pi} \mathcal{L}i_2(e^{i\varphi}) \right). \quad (4.80)$$

Setting $\varphi = 0$, it yields

$$\Theta_{LR}^{12}(\tau, \sigma) = -iH \tilde{p}^3 \frac{\pi^2}{12}. \quad (4.81)$$

Likewise, we obtain the following result for Θ_{RL} ,

$$\Theta_{RL}^{12}(z, w, \bar{z}, \bar{w}) = \frac{i\pi}{4} H \tilde{p}^3 \left(\frac{\varphi^2}{2\pi} - \frac{1}{\pi} \mathcal{L}i_2(e^{i\varphi}) \right), \quad (4.82)$$

which yields for $\varphi = 0$

$$\Theta_{RL}^{12}(\tau, \sigma) = -iH\tilde{p}^3 \frac{\pi^2}{12}. \quad (4.83)$$

Finally, adding up all four individual pieces, it comes out that the total commutator of the generalized coordinates in the 1- and 2-directions vanishes in the H -flux background; in fact, it vanishes even before taking the limit $\varphi = 0$. Thus, we end up with

$$\Theta_H^{12}(\tau, \sigma) = \Theta_{LL}^{12} + \Theta_{RR}^{12} + \Theta_{LR}^{12} + \Theta_{RL}^{12} = 0 \quad (4.84)$$

and commutativity is manifest.

4.3.2 Commutators of the geometric f -flux model

The geometric f -flux background is obtained from the H -flux face by a T-duality transformation in the 1-direction, flipping

$$\mathcal{X}_L^1(z, \bar{z}) \longrightarrow \mathcal{X}_L^1(z, \bar{z}), \quad \mathcal{X}_R^1(z, \bar{z}) \longrightarrow -\mathcal{X}_R^1(z, \bar{z}). \quad (4.85)$$

In the computation of the corresponding commutators, we are still using the free fields $X_{0,L}^1(z)$ and $X_{0,R}^1(\bar{z})$ of the H -flux background, and, therefore, we do not have to repeat here the entire CFT computation but we just have to implement the flip of signs of the individual commutator pieces in the appropriate way, as given by eq. (4.59). Thus, we obtain, setting $H \equiv f$,

$$\Theta_f^{12}(\tau, \sigma) = \Theta_{LL}^{12} - \Theta_{RR}^{12} + \Theta_{LR}^{12} - \Theta_{RL}^{12} = 0 \quad (4.86)$$

in the winding sector of string theory with non-vanishing \tilde{p}^3 . Commutativity is maintained.

4.3.3 Commutators of the non-geometric Q -flux model

The non-geometric Q -flux background follows by performing another T-duality in the 2-direction,

$$\mathcal{X}_L^2(z, \bar{z}) \longrightarrow \mathcal{X}_L^2(z, \bar{z}), \quad \mathcal{X}_R^2(z, \bar{z}) \longrightarrow -\mathcal{X}_R^2(z, \bar{z}). \quad (4.87)$$

Now, for the first time, the total commutator is non-vanishing, since

$$\begin{aligned}\Theta_Q^{12}(z, w, \bar{z}, \bar{w}) &= \Theta_{LL}^{12} + \Theta_{RR}^{12} - \Theta_{LR}^{12} - \Theta_{RL}^{12} \\ &= -i\pi Q \tilde{p}^3 \left(\frac{\varphi^2}{2\pi} - \frac{1}{\pi} \mathcal{L}i_2(e^{i\varphi}) \right)\end{aligned}\quad (4.88)$$

with $H \equiv f \equiv Q$. Setting $\varphi = 0$, it yields

$$\Theta_Q^{12}(\tau, \sigma) = \frac{i\pi^2}{3} Q \tilde{p}^3. \quad (4.89)$$

Thus, the generalized coordinates \mathcal{X}^1 and \mathcal{X}^2 of the Q -flux background are non-commutative. The parameter of non-commutativity is provided by the winding number \tilde{p}^3 in the third direction of the torus.

4.3.4 Commutators of the non-geometric R -flux model

The final T-duality with respect to the 3-direction is no longer described by the Buscher rules (2.14) and (2.15), since the background depends explicitly on X^3 . An alternative prescription is provided in the language of conformal field theory by the automorphism of the operator algebra, acting as

$$\mathcal{X}_L^3 \rightarrow \mathcal{X}_L^3, \quad \mathcal{X}_R^3 \rightarrow -\mathcal{X}_R^3, \quad \tilde{p}^3 \rightarrow p_3. \quad (4.90)$$

Thus, starting from the H -flux background and dualizing it in all three directions, we arrive to the non-geometric R -flux background, which is completely left-right asymmetric in all three directions. It is not anymore a geometric manifold, even in a local coordinate neighborhood. The commutator of the generalized coordinates in the 1- and 2-directions becomes now

$$\Theta_R^{12}(\tau, \sigma) = [\mathcal{X}^1(\tau, \sigma), \mathcal{X}^2(\tau, \sigma)] = \frac{i\pi^2}{3} R p_3 \quad (4.91)$$

setting $H \equiv f \equiv Q \equiv R$.

In this case, the parameter of non-commutativity is provided by the momentum number p_3 , which, in turn leads to non-associativity as violation of Jacobi identity among the generalized coordinates.

This completes the derivation of the commutators in all T-dual faces of the parabolic flux model, confirming by CFT techniques the emergence of non-commutative and/or non-associative structures in non-geometric backgrounds.

5 Conclusions and discussion

In this paper we confirmed the non-commutative and non-associative algebraic structure of the parabolic flux compactifications with constant geometric and non-geometric fluxes, using the approach of canonical T-duality in conformal field theory. Combining this approach with the gauging procedure of Roček and Verlinde, we found that the non-commutative and non-associative relations arise in a subtle way. Namely, these structures do not refer to the coordinates X^I and the coordinates \tilde{X}^I of the different T-dual faces of the model, but they rather refer to generalized coordinates \mathcal{X}^I and their dual counterparts $\tilde{\mathcal{X}}^I$ associated to the conserved currents of the underlying Roček-Verlinde "parent" theory. The resulting commutation relations are summarized in the table below for all T-dual faces of the parabolic flux models.

T-dual frames	Commutators	Three-brackets
H -flux	$[\tilde{\mathcal{X}}^I, \tilde{\mathcal{X}}^J] \sim H\epsilon^{IJK}\tilde{p}^K$	$[\tilde{\mathcal{X}}^1, \tilde{\mathcal{X}}^2, \tilde{\mathcal{X}}^3] \sim H$
f -flux	$[\mathcal{X}^I, \tilde{\mathcal{X}}^J] \sim f\epsilon^{IJK}\tilde{p}^K$	$[\mathcal{X}^1, \tilde{\mathcal{X}}^2, \tilde{\mathcal{X}}^3] \sim f$
Q -flux	$[\mathcal{X}^I, \mathcal{X}^J] \sim Q\epsilon^{IJK}\tilde{p}^K$	$[\mathcal{X}^1, \mathcal{X}^2, \tilde{\mathcal{X}}^3] \sim Q$
R -flux	$[\mathcal{X}^I, \mathcal{X}^J] \sim R\epsilon^{IJK}p^K$	$[\mathcal{X}^1, \mathcal{X}^2, \mathcal{X}^3] \sim R$

The third column of this table contains the non-trivial three-brackets among the generalized coordinates, which reflect the non-associative tri-products for each of the four flux backgrounds. This non-associative algebraic structure was further analyzed in refs. [31, 33, 34, 35, 36, 37] in terms of 3-cocycles, star-products, tri-products and related physics topics such as membrane sigma-models, magnetic monopole backgrounds, and their quantization.

Since the generalized coordinates \mathcal{X}^I and their T-dual counterparts $\tilde{\mathcal{X}}^I$ do not correspond to bona-fide conformal fields of the underlying CFT, there is no conflict with the associativity of CFT amplitudes satisfied by on-shell physical fields. Our computations do not require staying off-shell or going on-shell, and, hence, the non-associativity of the phase space variables $(\mathcal{X}^I, \tilde{\mathcal{X}}^I, p^I, \tilde{p}^I)$ is not forbidden on general grounds. In our view, non-geometric flux backgrounds should fit in a broad "geometrical" picture in which the physical on-shell closed string fields, whose operator algebra is fully associative, nevertheless live on a non-commutative and possibly non-associative phase space. Then, as advocated in refs. [26, 27, 35], this space possesses a minimal volume, which is set by the R -flux deformation parameter together with the non-vanishing string length upon quantization. This picture is also consistent with the role of non-associative commutation relations and tri-products in double field theory [61]. Physical fields that satisfy the strong constraint in double field theory do not

display any non-associative product structure. Still, they can live in a non-associative doubled phase space with non-associative doubled coordinates, since there is no need for the doubled coordinates to be physical fields satisfying the strong constraint.

It is certainly interesting to explore in more detail the relations between the doubled CFT approach of Roček and Verlinde on the string world-sheet and the doubled target space description of double field theory. In fact, it is tempting to speculate that the "parent" world-sheet theory of Roček and Verlinde is just the proper world-sheet analogue of target space double field theory (for a different world-sheet description of double field theory see the recent work [77, 78]). Then, in this context, the chirally conserved world-sheet currents \mathcal{J}^I and $\bar{\mathcal{J}}^I$ giving rise to the generalized coordinates \mathcal{X}^I and $\bar{\mathcal{X}}^I$ seem to correspond to the doubled coordinates of double field theory. Projecting these currents to the "child" non-linear sigma-models by gauging seems to be very similar in vain to the different ways of applying the strong constraint in double field theory.

Finally, it would be interesting to investigate other non-geometric backgrounds by similar methods, focusing, in particular, to the novel class of non-geometric spaces that do not admit geometric duals, such as the asymmetric orbifold conformal field theories [29, 44] and the closely related double elliptic spaces that arose recently by dimensional reduction of double field theory [45] as well as to solutions of double field theory for non-geometric group manifolds [15, 67, 68]. We hope to return to these issues elsewhere.

Acknowledgements

This work was partially supported by the ERC Advanced Grant "Strings and Gravity" (Grant.No. 32004) and by the DFG cluster of excellence "Origin and Structure of the Universe". This research is also implemented (I.B.) under the "ARISTEIA" action of the operational programme education and lifelong learning and is co-funded by the European Social Fund (ESF) and National Resources of Greece. I.B. is grateful to the hospitality extended to him at the Arnold Sommerfeld Center for Theoretical Physics and the Max Planck Institute for Physics in Munich during the course of this work, whereas D.L. likes to thank the CERN theory unit for hospitality. We thank David Andriot, Ralph Blumenhagen, Ioannis Florakis, Falk Hassler, Olaf Hohm, Magdalena Larfors, Peter Patalong, Felix Rennecke and Stefan Theisen for fruitful discussions.

A Monodromies of toroidal fibrations

The monodromies specify the gluing conditions of the fibre when going around the base space of a given fibration. Here, in view of the applications that are discussed in this paper, we summarize the monodromy transformations of a three-dimensional fibration with a one-dimensional circle S^1 with coordinate X^3 serving as the base and a two-dimensional torus T^2 with coordinates $X^i = (X^1, X^2)$ as fibre.

Let G_{IJ} and B_{IJ} be the components of the metric and anti-symmetric fields on T^2 , which depend on the base point and they are conveniently combined as $E_{IJ} = G_{IJ} + B_{IJ}$. Then, in this case, the monodromy transformations are given in terms of $O(2, 2; \mathbb{R})$ transformations, which act on the background fields of the T^2 in the following way,

$$E(X^3 + 2\pi) = \mathcal{M}_{O(2,2)} E(X^3) = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} CE(X^3) + D \\ AE(X^3) + B \end{pmatrix}^{-1}, \quad (\text{A.1})$$

by going around the base space. Here, $\mathcal{M}_{O(2,2)}$ is an element of the group $O(2, 2; \mathbb{R})$ represented as

$$\mathcal{M}_{O(2,2)} = \begin{pmatrix} A & B \\ C & D \end{pmatrix}, \quad (\text{A.2})$$

where A, B, C and D are 2-dimensional matrices that satisfy the defining relations

$$A^t C + C^t A = 0, \quad B^t D + D^t B = 0, \quad A^t D + C^t B = \mathbb{1}. \quad (\text{A.3})$$

The right-hand side of equation (A.1) provides the T-dual background $\tilde{E}(X^3)$ of $E(X^3)$ with respect to the two commuting isometries generated by the vector fields tangent to the fibre T^2 . Thus, when the matrix element $\mathcal{M} \in O(2, 2; \mathbb{R})$ is not trivial and the gluing condition of the fibration involves a genuine T-duality transformation, the space fails to be globally geometric. Likewise, if the transition function among different patches involves a T-duality transformation that cannot be undone by a diffeomorphism, the space will fail to be locally geometric and the notion of T-folds has to be used as substitute to ordinary differentiable manifolds. The monodromies of different fibrations are essential for selecting the correct closed string boundary conditions in the geometric and non-geometric spaces that are discussed in this paper.

It is convenient to parametrize the background fields of the fibre by two complex scalars, known as the complex structure τ and the complexified Kähler class ρ of T^2 , setting

$$\tau = \frac{G_{12}}{G_{11}} + i \frac{V}{G_{11}}, \quad \rho = -B_{12} + i V, \quad (\text{A.4})$$

where $V = \sqrt{\det G}$ denotes the volume element of the two-torus. Then, the $O(2, 2)$

group splits naturally as follows,

$$O(2, 2; \mathbb{R}) \simeq SL(2, \mathbb{R})_\tau \times SL(2, \mathbb{R})_\rho. \quad (\text{A.5})$$

The group factor $SL(2, \mathbb{R})_\tau$ corresponds to reparametrization of the torus, acting as modular transformations of its complex structure with generators $T_\tau : \tau \rightarrow \tau + 1$ and $S_\tau : \tau \rightarrow -1/\tau$. It reads

$$\tau \rightarrow \frac{a\tau + b}{c\tau + d}, \quad \text{with } ad - bc = 1. \quad (\text{A.6})$$

The other factor $SL(2, \mathbb{R})_\rho$ contains the shift of the B -field by a constant, $T_\rho : \rho \rightarrow \rho + 1$, as well as the T-duality transformation $S_\rho : \rho \rightarrow -1/\rho$ that reverses the volume of the torus $V \rightarrow 1/V$ when $B_{12} = 0$. Its full action is given by

$$\rho \rightarrow \frac{a'\rho + b'}{c'\rho + d'}, \quad \text{with } a'd' - b'c' = 1. \quad (\text{A.7})$$

The embedding of $SL(2, \mathbb{R})_\tau \times SL(2, \mathbb{R})_\rho$ in $O(2, 2; \mathbb{R})$ is provided by the following identification with the matrices A, B, C and D in equation (A.2),

$$A = a' \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \quad B = b' \begin{pmatrix} -b & a \\ -d & c \end{pmatrix}, \quad C = c' \begin{pmatrix} -c & -d \\ a & b \end{pmatrix}, \quad D = d' \begin{pmatrix} d & -c \\ -b & a \end{pmatrix}. \quad (\text{A.8})$$

More explicitly, the matrix elements $SO(2, 2; \mathbb{R}) \simeq SL(2, \mathbb{R})_\tau \times SL(2, \mathbb{R})_\rho$ factorize as

$$\mathcal{M}_{SO(2,2)} = \begin{pmatrix} a & b & 0 & 0 \\ c & d & 0 & 0 \\ 0 & 0 & d & -c \\ 0 & 0 & -b & a \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} a' & b' & 0 & 0 \\ c' & d' & 0 & 0 \\ 0 & 0 & d' & -c' \\ 0 & 0 & -b' & a' \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, \quad (\text{A.9})$$

explaining the identification (A.8).

We also note for completeness that there are two discrete transformations that can be appended to the previous description: the \mathbb{Z}_2 transformation $(\tau, \rho) \rightarrow (\rho, \tau)$ that exchanges complex and Kähler structures and another \mathbb{Z}_2 transformation $(\tau, \rho) \rightarrow (-\bar{\tau}, -\bar{\rho})$. Both can be intertwined with world-sheet parity.

The group elements of each $SL(2, \mathbb{R})$ factor, which are represented by the 2×2 matrices

$$M_\tau = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \quad M_\rho = \begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix}, \quad (\text{A.10})$$

are classified according to their eigenvalues into elliptic, parabolic or hyperbolic ele-

ments. Since the eigenvalues of M (it can be either M_τ or M_ρ) satisfy the characteristic polynomial equation

$$\lambda^2 - \text{tr}(M)\lambda + 1 = 0, \quad (\text{A.11})$$

we have the following three distinct cases:

- If $|\text{tr}(M)| < 2$, the two eigenvalues will be complex conjugate and M is called elliptic element.
- If $|\text{tr}(M)| = 2$, the two eigenvalues will be real and equal and M is called parabolic element.
- If $|\text{tr}(M)| > 2$, the two eigenvalues will be real and unequal and M is called hyperbolic element.

Elliptic group elements are characterized by the property $M^n = \mathbb{1}$ for some finite integer n , like

$$M = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad (\text{A.12})$$

which has $M^4 = \mathbb{1}$, whereas parabolic elements, like

$$M = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad (\text{A.13})$$

do not share this property.

In this paper we are only concerned with parabolic group elements M , and, for that reason, the corresponding closed string backgrounds associated to T^2 fibrations over S^1 with constant flux are called parabolic.

B Parabolic backgrounds and monodromies of currents

In this appendix we give a brief account of the toroidal vacua of closed string theory with constant fluxes and determine the monodromy properties of their currents in geometric as well as in non-geometric faces.

B.1 The parabolic backgrounds with constant flux

Recall that the constant flux backgrounds in three dimensions come in four different versions, which are related to each other by successive T-duality transformations. Concretely, we consider the compactification on a three-dimensional torus T^3 , which

can be viewed as an $\mathcal{F} = T^n$ fibration over the $(3 - n)$ -dimensional base space $\mathcal{B} = T^{3-n}$ for $n = 0, 1, 2, 3$. In the following, we consider all four different cases. The starting point, provided by $n = 0$, is a flat three-torus T^3 parametrized by $X^{I=1,2,3}$ with periodic identifications $X^I \sim X^I + 2\pi r^I$, where r^I are the three radii of T^3 , which are all set equal to 1 for convenience. There is also a 3-form H -flux $H_3 = H dX^1 \wedge dX^2 \wedge dX^3$ with constant H . The metric and the corresponding B -field are taken to be

$$G_{IJ} = \delta_{IJ}, \quad B_{12} = HX^3, \quad B_{13} = 0 = B_{23}. \quad (\text{B.1})$$

The H -flux has to obey a topological quantization condition, which is given, in general, by (see, for instance, [1])

$$\frac{1}{4\pi^2 \alpha'} \int_{T^3} H = k, \quad k \in \mathbb{Z}. \quad (\text{B.2})$$

Since the volume of the unit three-torus is $(2\pi)^3$, it follows immediately that the quantization condition for H takes the following form, in units $\alpha' = 1$,

$$H = \frac{k}{2\pi}, \quad k \in \mathbb{Z}. \quad (\text{B.3})$$

For the discussion of T-duality transformations as well as for the CFT computation of the commutators, we consider the simplest case $k = 1$.

The quantization condition (B.3) appears to be incompatible with the dilute flux approximation used in the definition of CFT_H , but this is illusive because we have normalized the radii of T^3 as well as α' to 1 for notational convenience. H has units $[L]^{-1}$ and it can be made very small for appropriate choice of radii as can be seen by reinstating the parameters in natural units in which $\alpha' = l_s^2$.

Performing a T-duality transformation on the one-dimensional circle fibre $\mathcal{F} = T_{X^1}^1$ with coordinate X^1 , one obtains the Heisenberg nilmanifold, which is a twisted torus without B -field corresponding to the case $n = 1$. This background is topologically distinct from T^3 , since it has a different homology group. Its Levi-Civita connection is related to the structure constants f_{JK}^I of the Heisenberg group, which are the geometric fluxes. We also note in this respect that the flat 3-torus corresponds to Bianchi-I geometry, whereas the twisted torus to Bianchi-II. The background fields of the twisted torus have the following form, setting $H \equiv f$,

$$G = \begin{pmatrix} 1 & fX^3 & 0 \\ fX^3 & 1 + (fX^3)^2 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad B = 0. \quad (\text{B.4})$$

Performing another T-duality in the X^2 -direction of the two-dimensional torus fibre $\mathcal{F} = T_{X_1, X_2}^2$, one obtains the so-called Q -space that corresponds to the case $n = 2$. This background is again a T^2 -fibration, but the corresponding metric and B -field are defined only locally and not globally. Setting $H \equiv f \equiv Q$ for notational purposes, G and B take the form

$$G = g(X^3) \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \frac{1}{g(X^3)} \end{pmatrix}, \quad B = g(X^3) \begin{pmatrix} 0 & -QX^3 & 0 \\ QX^3 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad g(X^3) = \frac{1}{1 - (QX^3)^2}. \quad (\text{B.5})$$

The Q -space is not a Riemannian manifold because the fibre \mathcal{F} is glued with a T-duality transformation when transporting it once around the base \mathcal{B} and not with a standard diffeomorphism.

The three backgrounds we have discussed so far take particularly simple form to linear order in the fluxes, summarized in the table below

Backgrounds	Target space fields
Torus + H -flux	$G = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & HX^3 & 0 \\ -HX^3 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$
Twisted torus	$G = \begin{pmatrix} 1 & fX^3 & 0 \\ fX^3 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} + \mathcal{O}(f^2), \quad B = 0$
Non-geom.	$G = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} + \mathcal{O}(Q^2), \quad B = \begin{pmatrix} 0 & -QX^3 & 0 \\ QX^3 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \mathcal{O}(Q^2)$

Thus, to linear order, one sees that the Q -flux background looks locally identical to the H -flux background, up to a sign flip in the flux. However, these two background are not the same globally, since the Q -flux background is glued together utilizing a T -duality transformation.

Finally, one can also consider applying a T-duality transformation to the entire three-dimensional torus, which is seen as a fibration with fibre $\mathcal{F} = T_{X_1, X_2, X_3}^3$ over a base point corresponding to the case $n = 3$. This last T-duality looks somewhat problematic, since the X_3 -direction is no longer a Killing isometry of the background and the standard Buscher rules cannot be applied as they stand. Nevertheless, this final step can be performed in the context of the underlying CFT by making a sign flip in the corresponding coordinate, hereby mimicking the action of T-duality on the currents, ending up with a "space" that is left-right asymmetric in all three directions. This defines the so called R -space, setting $H \equiv f \equiv Q \equiv R$ for notational purposes,

which is non-geometric locally as well as globally.

B.2 Monodromies of geometric H -flux model

The monodromies of the original H -flux background are given by shifts of the B -field, which act on τ and ρ in the following way under $X^3 \rightarrow X^3 + 2\pi$,

$$\rho(X^3 + 2\pi) = \rho(X^3) - \pi H, \quad \tau(X^3 + 2\pi) = \tau(X^3). \quad (\text{B.6})$$

This gauge transformation of the B -field corresponds to the $O(2,2)$ monodromy transformation in the 1, 2-directions given by the group elements

$$\mathcal{M}_{O(2,2)} = \begin{pmatrix} 1 & 0 & 0 & \pi H \\ 0 & 1 & -\pi H & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad (\text{B.7})$$

which indeed act on the Kähler and complex moduli ρ and τ as above.

The shift of the base coordinate $X^3 \rightarrow X^3 + 2\pi$ induces the following closed string boundary condition on the left- and right-moving closed string coordinates $X_L^3(z)$ and $X_R^3(\bar{z})$,

$$X_L^3(e^{2i\pi}z) = X_L^3(z) - \pi \tilde{p}^3, \quad X_R^3(e^{-2i\pi}\bar{z}) = X_R^3(\bar{z}) - \pi \tilde{p}^3, \quad (\text{B.8})$$

where $\tilde{p}^3 = p_L^3 - p_R^3$ is the winding number of the closed string in the third direction. Then, using the winding shifts of $X_{L,R}^3$, it follows from eq.(4.61) that the associated closed string boundary conditions of the CFT currents \mathcal{J}^I and $\bar{\mathcal{J}}^I$ to linear order in the flux are

$$\begin{aligned} \mathcal{J}^1(e^{-2i\pi}z, e^{2i\pi}\bar{z}) &= \mathcal{J}^1(z, \bar{z}) - \frac{\pi}{2} H \tilde{p}^3 \mathcal{J}^2(z, \bar{z}), \\ \bar{\mathcal{J}}^1(e^{-2i\pi}z, e^{2i\pi}\bar{z}) &= \bar{\mathcal{J}}^1(z, \bar{z}) + \frac{\pi}{2} H \tilde{p}^3 \bar{\mathcal{J}}^2(z, \bar{z}), \\ \mathcal{J}^2(e^{-2i\pi}z, e^{2i\pi}\bar{z}) &= \mathcal{J}^2(z, \bar{z}) + \frac{\pi}{2} H \tilde{p}^3 \mathcal{J}^1(z, \bar{z}), \\ \bar{\mathcal{J}}^2(e^{-2i\pi}z, e^{2i\pi}\bar{z}) &= \bar{\mathcal{J}}^2(z, \bar{z}) - \frac{\pi}{2} H \tilde{p}^3 \bar{\mathcal{J}}^1(z, \bar{z}), \\ \mathcal{J}^3(e^{-2i\pi}z, e^{2i\pi}\bar{z}) &= \mathcal{J}^3(z, \bar{z}), \\ \bar{\mathcal{J}}^3(e^{-2i\pi}z, e^{2i\pi}\bar{z}) &= \bar{\mathcal{J}}^3(z, \bar{z}). \end{aligned} \quad (\text{B.9})$$

Thus, the monodromies of the H -flux background act in a left-right asymmetric way on the currents, transforming the currents into the dual currents and vice versa.

B.3 Monodromies of geometric f -flux model

Perform a T-duality transformation in the 1-direction we obtain the parabolic f -flux background with geometric flux. The corresponding monodromies are given as shifts of the complex structure τ under $X^3 \rightarrow X^3 + 2\pi$,

$$\rho(X^3 + 2\pi) = \rho(X^3), \quad \tau(X^3 + 2\pi) = \tau(X^3) - \pi f, \quad (\text{B.10})$$

associated to the $O(2,2)$ monodromy transformation in the 1, 2-directions with group elements

$$\mathcal{M}_{O(2,2)} = \begin{pmatrix} 1 & -\pi f & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & \pi f & 1 \end{pmatrix}. \quad (\text{B.11})$$

Since T-duality acts in this case as $\bar{\mathcal{J}}^1 \rightarrow -\bar{\mathcal{J}}^1$, the corresponding closed string boundary conditions of the currents \mathcal{J}^I take the following form,

$$\begin{aligned} \mathcal{J}^1(e^{-2i\pi}z, e^{2i\pi}\bar{z}) &= \mathcal{J}^1(z, \bar{z}) - \frac{\pi}{2}f\check{p}^3 \mathcal{J}^2(z, \bar{z}), \\ \bar{\mathcal{J}}^1(e^{-2i\pi}z, e^{2i\pi}\bar{z}) &= \bar{\mathcal{J}}^1(z, \bar{z}) - \frac{\pi}{2}f\check{p}^3 \bar{\mathcal{J}}^2(z, \bar{z}), \\ \mathcal{J}^2(e^{-2i\pi}z, e^{2i\pi}\bar{z}) &= \mathcal{J}^2(z, \bar{z}) + \frac{\pi}{2}f\check{p}^3 \mathcal{J}^1(z, \bar{z}), \\ \bar{\mathcal{J}}^2(e^{-2i\pi}z, e^{2i\pi}\bar{z}) &= \bar{\mathcal{J}}^2(z, \bar{z}) + \frac{\pi}{2}f\check{p}^3 \bar{\mathcal{J}}^1(z, \bar{z}), \\ \mathcal{J}^3(e^{-2i\pi}z, e^{2i\pi}\bar{z}) &= \mathcal{J}^3(z, \bar{z}), \\ \bar{\mathcal{J}}^3(e^{-2i\pi}z, e^{2i\pi}\bar{z}) &= \bar{\mathcal{J}}^3(z, \bar{z}). \end{aligned} \quad (\text{B.12})$$

Thus, for the f -flux background, the monodromies act in left-right symmetric way on the currents, meaning that the currents and the dual currents do not mix with each other.

B.4 Monodromies of non-geometric Q -flux model

Next, performing another T-duality in the 2-direction brings us to the non-geometric background with Q -flux. The monodromy of the torus is now determined by the following transformation of ρ and τ under $X^3 \rightarrow X^3 + 2\pi$,

$$\rho(X^3 + 2\pi) = \frac{\rho}{1 - \pi Q \rho(X^3)}, \quad \tau(X^3 + 2\pi) = \tau(X^3), \quad (\text{B.13})$$

which is an $O(2,2)$ transformation in the 1, 2-directions with group elements

$$\mathcal{M}_{O(2,2)} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & \pi Q & 1 & 0 \\ -\pi Q & 0 & 0 & 1 \end{pmatrix}. \quad (\text{B.14})$$

Here, T-duality is performed by flipping the signs $\bar{\mathcal{J}}^1 \rightarrow -\bar{\mathcal{J}}^1$ and $\bar{\mathcal{J}}^2 \rightarrow -\bar{\mathcal{J}}^2$, and, thus, the closed string boundary conditions of the currents are taking the form

$$\begin{aligned} \mathcal{J}^1(e^{-2i\pi}z, e^{2i\pi}\bar{z}) &= \mathcal{J}^1(z, \bar{z}) - \frac{\pi}{2}Q\tilde{p}^3\mathcal{J}^2(z, \bar{z}), \\ \bar{\mathcal{J}}^1(e^{-2i\pi}z, e^{2i\pi}\bar{z}) &= \bar{\mathcal{J}}^1(z, \bar{z}) + \frac{\pi}{2}Q\tilde{p}^3\bar{\mathcal{J}}^2(z, \bar{z}), \\ \mathcal{J}^2(e^{-2i\pi}z, e^{2i\pi}\bar{z}) &= \mathcal{J}^2(z, \bar{z}) + \frac{\pi}{2}Q\tilde{p}^3\mathcal{J}^1(z, \bar{z}), \\ \bar{\mathcal{J}}^2(e^{-2i\pi}z, e^{2i\pi}\bar{z}) &= \bar{\mathcal{J}}^2(z, \bar{z}) - \frac{\pi}{2}Q\tilde{p}^3\bar{\mathcal{J}}^1(z, \bar{z}), \\ \mathcal{J}^3(e^{-2i\pi}z, e^{2i\pi}\bar{z}) &= \mathcal{J}^3(z, \bar{z}), \\ \bar{\mathcal{J}}^3(e^{-2i\pi}z, e^{2i\pi}\bar{z}) &= \bar{\mathcal{J}}^3(z, \bar{z}). \end{aligned} \quad (\text{B.15})$$

The monodromies are again left-right asymmetric, as for the H -flux model. In fact, they agree with the monodromies of H -flux background to linear order in the fluxes, up to a minus sign in the flux, as expected from the previous general discussion.

B.5 Monodromies of non-geometric R -flux model

Finally, coming to the R -flux model, the closed string boundary conditions for the currents \mathcal{J}^I are determined by the momentum number p_3 and they take the following form,

$$\begin{aligned} \mathcal{J}^1(e^{-2i\pi}z, e^{2i\pi}\bar{z}) &= \mathcal{J}^1(z, \bar{z}) - \frac{\pi}{2}Rp_3\mathcal{J}^2(z, \bar{z}), \\ \bar{\mathcal{J}}^1(e^{-2i\pi}z, e^{2i\pi}\bar{z}) &= \bar{\mathcal{J}}^1(z, \bar{z}) + \frac{\pi}{2}Rp_3\bar{\mathcal{J}}^2(z, \bar{z}), \\ \mathcal{J}^2(e^{-2i\pi}z, e^{2i\pi}\bar{z}) &= \mathcal{J}^2(z, \bar{z}) + \frac{\pi}{2}Rp_3\mathcal{J}^1(z, \bar{z}), \\ \bar{\mathcal{J}}^2(e^{-2i\pi}z, e^{2i\pi}\bar{z}) &= \bar{\mathcal{J}}^2(z, \bar{z}) - \frac{\pi}{2}Rp_3\bar{\mathcal{J}}^1(z, \bar{z}), \\ \mathcal{J}^3(e^{-2i\pi}z, e^{2i\pi}\bar{z}) &= \mathcal{J}^3(z, \bar{z}), \\ \bar{\mathcal{J}}^3(e^{-2i\pi}z, e^{2i\pi}\bar{z}) &= \bar{\mathcal{J}}^3(z, \bar{z}). \end{aligned} \quad (\text{B.16})$$

This concludes the description of how the monodromies act on the currents in all T-dual faces of the toroidal flux model. Judging from the action on the ρ and τ moduli, the corresponding $SL(2, \mathbb{R})$ group elements are parabolic (i.e., of infinite order) using the nomenclature of Appendix A.

Summarizing, the chain of T-duality transformations of the toroidal background with constant H -flux, encoded in the diagram

$$H_{IJK} \xrightarrow{T_I} f_{JK}^I \xrightarrow{T_J} Q_K^{IJ} \xrightarrow{T_K} R^{IJK}; \quad I, J, K = 1, \dots, 3, \quad (\text{B.17})$$

provide the parabolic flux models of closed strings that are in focus in the present work.

C Some relevant dilogarithmic integrals

In this appendix we collect some useful expressions involving the dilogarithm function based on series expansions and integral formulae (see, for instance, [75]). We also evaluate certain commutators used in the main text based on the dilogarithm function.

C.1 Delta-function, step-function and the logarithm

The delta-function on a circle of unit radius is defined through the infinite series

$$\delta(\varphi) = \sum_{m \in \mathbb{Z}} e^{im\varphi}. \quad (\text{C.1})$$

Likewise, the step-function, which is the integral of the delta-function,

$$\epsilon(\varphi) = \frac{1}{2} \frac{d}{d\varphi} \epsilon(\varphi), \quad (\text{C.2})$$

is represented by the infinite series

$$\epsilon(\varphi) = -\frac{i}{\pi} \sum_{m \neq 0} \frac{1}{m} e^{im\varphi} + \frac{\varphi}{\pi} \quad (\text{C.3})$$

so that $\epsilon(\varphi) = 1$ for $\varphi > 0$ (i.e., for $\sigma > \sigma'$), $\epsilon(\varphi) = -1$ for $\varphi < 0$ (i.e., for $\sigma < \sigma'$) and $\epsilon(\varphi) = 0$ for $\varphi = 0$ (i.e., for $\sigma = \sigma'$). Finally, the logarithm is represented by the infinite series

$$\log(1 - e^{i\varphi}) = - \sum_{m=1}^{\infty} \frac{1}{m} e^{im\varphi}. \quad (\text{C.4})$$

Hence, one obtains the following relation between the logarithm, the step-function and the delta-function,

$$\begin{aligned}
\epsilon(\varphi) &= \frac{i}{\pi} \log |1 - e^{i\varphi}|^2 + \frac{\varphi}{\pi}, \\
\delta(\varphi) &= \frac{i}{2\pi} \frac{d}{d\varphi} \log |1 - e^{i\varphi}|^2 + \frac{1}{2\pi} \\
&= -\frac{1}{2\pi} \left(\frac{e^{i\varphi}}{1 - e^{i\varphi}} + \frac{e^{-i\varphi}}{1 - e^{-i\varphi}} - 1 \right). \tag{C.5}
\end{aligned}$$

It useful to introduce a regulator $e^{-\delta}$ with $\delta > 0$ when writing infinite series, as in the example

$$\lim_{\delta \rightarrow 0^+} \frac{1}{1 - e^{-\delta + i\varphi}} = \sum_{m=0}^{\infty} e^{im\varphi}. \tag{C.6}$$

The δ -prescription is implicitly assumed hereafter.

C.2 The dilogarithm function

The dilogarithm function follows by integrating the logarithm through the defining relation

$$\log(1 - e^{i\varphi}) = i \frac{d}{d\varphi} Li_2(e^{i\varphi}), \tag{C.7}$$

leading to the expression

$$\begin{aligned}
Li_2(e^{i\varphi}) &= -i \int^{\varphi} d\tilde{\varphi} \log(1 - e^{i\tilde{\varphi}}) \\
&= \sum_{m=1}^{\infty} \frac{1}{m^2} e^{im\varphi}. \tag{C.8}
\end{aligned}$$

When these relations are written in terms of the complex variable $z = e^{i\varphi}$ they assume the form

$$\begin{aligned}
Li_2(z) &= -i \int_0^z d\tilde{z} \frac{\log(1 - \tilde{z})}{\tilde{z}} \\
&= \sum_{m=1}^{\infty} \frac{1}{m^2} z^m. \tag{C.9}
\end{aligned}$$

Finally, we define the closely related function $\mathcal{L}i_2(e^{i\varphi})$ as follows

$$\begin{aligned}\mathcal{L}i_2(e^{i\varphi}) &= Li_2(e^{i\varphi}) + Li_2(e^{-i\varphi}) = -i \int^\varphi d\tilde{\varphi} \log\left(\frac{1 - e^{i\tilde{\varphi}}}{1 - e^{-i\tilde{\varphi}}}\right) \\ &= \sum_{m \neq 0}^{\infty} \frac{1}{m^2} e^{im\varphi}.\end{aligned}\tag{C.10}$$

C.3 Certain dilogarithmic commutators

We are now in position to evaluate certain commutators that appear in the main text and help to establish the non-commutative structure of the non-geometric flux backgrounds. First consider the following (prototype) commutator

(i) $[X(z), \int^w d\tilde{w} \frac{X(\tilde{w})}{\tilde{w}}]:$

This example is used as prototype for the computation of all other commutators in the following. Using the δ -prescription, it assumes the form

$$\begin{aligned}\left[X(z), \int^w d\tilde{w} \frac{X(\tilde{w})}{\tilde{w}}\right]_{|z|=|w|} &= \\ \frac{1}{2} \lim_{\delta \rightarrow 0^+} \left(\int^w d\tilde{w} \frac{\log(z - \tilde{w})}{\tilde{w}} \Big|_{|z|=|w|+\delta} - \int^w d\tilde{w} \frac{\log(z - \tilde{w})}{\tilde{w}} \Big|_{|z|=|w|-\delta} \right).\end{aligned}\tag{C.11}$$

Choosing $z = e^{\pm\delta - i\sigma}$, $w = e^{-i\sigma'}$ and $\tilde{w} = e^{-i\tilde{\sigma}'}$ and setting $\varphi = \sigma - \sigma'$ and $\tilde{\varphi} = \sigma - \tilde{\sigma}'$, we note that integration takes place over a circular path of unit modulus. Since $d\tilde{w} = i\tilde{w}d\tilde{\varphi}$, the expression above becomes

$$\begin{aligned}\left[X(z), \int^w d\tilde{w} \frac{X(\tilde{w})}{\tilde{w}}\right]_{|z|=|w|} &= \frac{\pi}{2} \int^\varphi d\tilde{\varphi} \left(1 + \frac{\tilde{\varphi}}{\pi} - \frac{i}{\pi} \sum_{m \neq 0} \frac{1}{m} e^{im\tilde{\varphi}} \right) \\ &= \frac{\pi}{2} \left(\varphi + \frac{\varphi^2}{2\pi} - \frac{1}{\pi} \sum_{m \neq 0} \frac{1}{m^2} e^{im\varphi} \right) \\ &= \frac{\pi}{2} \left(\varphi + \frac{\varphi^2}{2\pi} - \frac{1}{\pi} Li_2(e^{i\varphi}) - \frac{1}{\pi} Li_2(e^{-i\varphi}) \right) \\ &= \frac{\pi}{2} \left(\varphi + \frac{\varphi^2}{2\pi} - \frac{1}{\pi} \mathcal{L}i_2(e^{i\varphi}) \right).\end{aligned}\tag{C.12}$$

Thus, for $\varphi = 0$, we obtain the result

$$\left[X(z), \int^w d\tilde{w} \frac{X(\tilde{w})}{\tilde{w}}\right]_{|z|=|w|} = -\frac{\pi^2}{6}.\tag{C.13}$$

(ii) $[X(z), \int^w d\tilde{w} \partial_{\tilde{w}} X(\tilde{w}) \log \tilde{w}]$:

Using the δ -prescription, as before, this particular commutator is given by

$$\begin{aligned} & \left[X(z), \int^w d\tilde{w} \partial_{\tilde{w}} X(\tilde{w}) \log \tilde{w} \right]_{|z|=|w|} = \\ & = -\frac{1}{2} \lim_{\delta \rightarrow 0_+} \left(\int^w d\tilde{w} \frac{1}{z - \tilde{w}} \log \tilde{w} \Big|_{|z|=|w|+\delta} - \int^w d\tilde{w} \frac{1}{z - \tilde{w}} \log \tilde{w} \Big|_{|z|=|w|-\delta} \right). \end{aligned} \quad (\text{C.14})$$

Integration by parts bring it into the form of the previous example, noting that the boundary contributions arising from partial integration cancel from the two terms of the commutator. Thus, we obtain the result

$$\begin{aligned} & \left[X(z), \int^w d\tilde{w} \partial_{\tilde{w}} X(\tilde{w}) \log \tilde{w} \right]_{|z|=|w|} = \\ & = -\frac{1}{2} \lim_{\delta \rightarrow 0_+} \left(\int^w d\tilde{w} \frac{\log(z - \tilde{w})}{\tilde{w}} \Big|_{|z|=|w|+\delta} - \int^w d\tilde{w} \frac{\log(z - \tilde{w})}{\tilde{w}} \Big|_{|z|=|w|-\delta} \right) \\ & = -\frac{\pi}{2} \left(\varphi + \frac{\varphi^2}{2\pi} - \frac{1}{\pi} \mathcal{L}i_2(e^{i\varphi}) \right). \end{aligned} \quad (\text{C.15})$$

(iii) $[X(z), \int^w d\tilde{w} \partial_{\tilde{w}} X(\tilde{w}) \log \bar{\tilde{w}}]$:

This example as well as the next one involve non-homomorphic expressions. This particular commutator is relevant for computing Θ_{LL} in the main text, whereas its complex conjugate expression is relevant one for computing Θ_{RR} . Since we are always integrating over circular paths, we use $\log \bar{\tilde{w}} = -\log \tilde{w}$ to cast this commutator into the form

$$\begin{aligned} & \left[X(z), \int^w d\tilde{w} \partial_{\tilde{w}} X(\tilde{w}) \log \bar{\tilde{w}} \right]_{|z|=|w|} = \\ & = \frac{1}{2} \lim_{\delta \rightarrow 0_+} \left(\int^w d\tilde{w} \frac{1}{z - \tilde{w}} \log \tilde{w} \Big|_{|z|=|w|+\delta} - \int^w d\tilde{w} \frac{1}{z - \tilde{w}} \log \tilde{w} \Big|_{|z|=|w|-\delta} \right) \\ & = \frac{\pi}{2} \left(\varphi + \frac{\varphi^2}{2\pi} - \frac{1}{\pi} \mathcal{L}i_2(e^{i\varphi}) \right). \end{aligned} \quad (\text{C.16})$$

(iv) $[X(z), \int^{\bar{w}} d\bar{\tilde{w}} \frac{X(\tilde{w})}{\tilde{w}}]$:

This last example is relevant for computing Θ_{LR} and Θ_{RL} in the main text. As before,

we have

$$\begin{aligned} & \left[X(z), \int^{\bar{w}} d\bar{w} \frac{X(\tilde{w})}{\bar{w}} \right]_{|z|=|w|} = \\ & \frac{1}{2} \lim_{\delta \rightarrow 0_+} \left(\int^{\bar{w}} d\bar{w} \frac{\log(z - \tilde{w})}{\bar{w}} \Big|_{|z|=|w|+\delta} - \int^{\bar{w}} d\bar{w} \frac{\log(z - \tilde{w})}{\bar{w}} \Big|_{|z|=|w|-\delta} \right) \end{aligned} \quad (\text{C.17})$$

Since $d\bar{w} = -i\bar{w}d\tilde{\varphi}$, it follows that

$$\begin{aligned} \left[X(z), \int^{\bar{w}} d\bar{w} \frac{X(\tilde{w})}{\bar{w}} \right]_{|z|=|w|} &= -\frac{\pi}{2} \int d\tilde{\varphi} \left(1 + \frac{\tilde{\varphi}}{\pi} - \frac{i}{\pi} \sum_{m \neq 0} \frac{1}{m} e^{im\tilde{\varphi}} \right) \\ &= -\frac{\pi}{2} \left(\varphi + \frac{\varphi^2}{2\pi} - \frac{1}{\pi} \sum_{m \neq 0} \frac{1}{m^2} e^{im\varphi} \right) \\ &= -\frac{\pi}{2} \left(\varphi + \frac{\varphi^2}{2\pi} - \frac{1}{\pi} \mathcal{L}i_2(e^{i\varphi}) \right). \end{aligned} \quad (\text{C.18})$$

References

- [1] M. Grana, "Flux Compactifications in String Theory: A Comprehensive Review", Phys. Rept. **423** (2006) 91 [arXiv:hep-th/0509003].
- [2] R. Blumenhagen, B. Körs, D. Lüst and S. Stieberger, "Four-dimensional String Compactifications with D-Branes, Orientifolds and Fluxes", Phys. Rept. **445** (2007) 1 [hep-th/0610327].
- [3] S. Hellerman, J. McGreevy and B. Williams, "Geometric Constructions of Nongeometric String Theories", JHEP **0401** (2004) 024 [arXiv:hep-th/0208174].
- [4] A. Dabholkar and C. Hull, "Duality Twists, Orbifolds, and Fluxes", JHEP **0309** (2003) 054 [arXiv:hep-th/0210209].
- [5] S. Kachru, M.B. Schulz, P.K. Tripathy and S.P. Trivedi, "New Supersymmetric String Compactifications", JHEP **0303** (2003) 061 [arXiv:hep-th/0211182].
- [6] A. Flournoy, B. Wecht and B. Williams, "Constructing Non-geometric Vacua in String Theory", Nucl. Phys. **B706** (2005) 127 [arXiv:hep-th/0404217];
- [7] C. M. Hull, "A Geometry for Non-geometric String Backgrounds", JHEP **0510** (2005) 065 [arXiv:hep-th/0406102].
- [8] G. Dall'Agata and S. Ferrara, "Gauged Supergravity Algebras from Twisted Tori Compactifications with Fluxes", Nucl. Phys. **B717** (2005) 223 [arXiv:hep-th/0502066].
- [9] C. M. Hull and R.A. Reid-Edwards, "Flux Compactifications of String Theory on Twisted Tori", Fortsch. Phys. **57**, 862 (2009) [arXiv:hep-th/0503114];
"Non-geometric Backgrounds, Doubled Geometry and Generalised T-duality", JHEP **0909** (2009) 014 [arXiv:0902.4032 [hep-th]].
- [10] A. Flournoy and B. Williams, "Non-geometry, Duality Twists and the Worldsheet", JHEP **0601** (2006) 166 [arXiv:hep-th/0511126].
- [11] J. Shelton, W. Taylor and B. Wecht, "Non-geometric Flux Compactifications", JHEP **0510** (2005) 085 [arXiv:hep-th/0508133];
"Generalized Flux Vacua", JHEP **0702** (2007) 095 [arXiv:hep-th/0607015].
- [12] A. Dabholkar and C. Hull, "Generalised T-duality and Non-geometric Backgrounds", JHEP **0605** (2006) 009 [arXiv:hep-th/0512005].
- [13] W. Schulgin and J. Troost, "Backreacted T-folds and Non-geometric Regions in Configuration Space," JHEP **0812** (2008) 098 [arXiv:0808.1345 [hep-th]].

- [14] J. McOrist, D.R. Morrison and S. Sethi, "Geometries, Non-Geometries, and Fluxes", *Adv. Theor. Math. Phys.* **14** (2010) 1515 [arXiv:1004.5447 [hep-th]].
- [15] M.B. Schulz, "T-folds, Doubled Geometry, and the SU(2) WZW Model", *JHEP* **1206** (2012) 158 [arXiv:1106.6291 [hep-th]].
- [16] D. Andriot, M. Larfors, D. Lüst, and P. Patalong, "A Ten-dimensional Action for Non-geometric Fluxes", *JHEP* **1109** (2011) 134 [arXiv:1106.4015 [hep-th]].
- [17] D. Andriot, O. Hohm, M. Larfors, D. Lüst, and P. Patalong, "A Geometric Action for Non-geometric Fluxes", *Phys. Rev. Lett.* **108** (2012) 261602 [arXiv:1202.3060 [hep-th]];
 "Non-Geometric Fluxes in Supergravity and Double Field Theory", *Fortschr. Phys.* **60** (2012) 1150 [arXiv:1204.1979 [hep-th]].
- [18] R. Blumenhagen, A. Deser, E. Plauschinn and F. Rennecke, "Non-geometric Strings, Symplectic Gravity and Differential Geometry of Lie Algebroids", *JHEP* **1302** (2013) 122 [arXiv:1211.0030 [hep-th]].
- [19] R. Blumenhagen, A. Deser, E. Plauschinn, F. Rennecke and C. Schmid, "The Intriguing Structure of Non-geometric Frames in String Theory", *Fortsch. Phys.* **61** (2013) 893 [arXiv:1304.2784 [hep-th]].
- [20] R. Blumenhagen, X. Gao, D. Herschmann and P. Shukla, "Dimensional Oxidation of Non-geometric Fluxes in Type II Orientifolds", *JHEP* **1310** (2013) 201 [arXiv:1306.2761 [hep-th]].
- [21] D. Andriot and A. Betz, " β -supergravity: A Ten-dimensional Theory with Non-geometric Fluxes and its Geometric Framework", *JHEP* **1312** (2013) 083 [arXiv:1306.4381 [hep-th]];
 "NS-branes, Source Corrected Bianchi Identities and More on Backgrounds with Non-geometric Fluxes", *JHEP* **1407** (2014) 059 [arXiv:1402.5972 [hep-th]];
 "Supersymmetry with Non-geometric Fluxes or a β -twist in Generalized Geometry and Dirac Operator", *JHEP* **1504** (2015) 006 [arXiv:1411.6640 [hep-th]].
- [22] H. Kawai, D. C. Lewellen and S. H. H. Tye, "Construction of Fermionic String Models in Four-Dimensions", *Nucl. Phys. B* **288** (1987) 1.
- [23] W. Lerche, D. Lüst and A. N. Schellekens, "Chiral Four-Dimensional Heterotic Strings from Self-dual Lattices", *Nucl. Phys. B* **287** (1987) 477.
- [24] I. Antoniadis, C. P. Bachas and C. Kounnas, "Four-Dimensional Superstrings", *Nucl. Phys. B* **289** (1987) 87.

- [25] K. S. Narain, M. H. Sarmadi and C. Vafa, "Asymmetric Orbifolds", Nucl. Phys. B **288** (1987) 551.
- [26] R. Blumenhagen and E. Plauschinn, "Non-associative Gravity in String Theory?", J. Phys. **A44** (2011) 015401 [arXiv:1010.1263 [hep-th]].
- [27] D. Lüst, "T-duality and Closed String Non-commutative (Doubled) Geometry", JHEP **1012** (2010) 084 [arXiv:1010.1361 [hep-th]].
- [28] R. Blumenhagen, A. Deser, D. Lüst, E. Plauschinn and F. Rennecke, "Non-geometric Fluxes, Asymmetric Strings and Nonassociative Geometry", J. Phys. **A44** (2011) 385401 [arXiv:1106.0316 [hep-th]].
- [29] C. Condeescu, I. Florakis and D. Lüst, "Asymmetric Orbifolds, Non-Geometric Fluxes and Non-Commutativity in Closed String Theory", JHEP **1204** (2012) 121 [arXiv:1202.6366 [hep-th]].
- [30] D. Lüst, "Twisted Poisson Structures and Non-commutative/Non-associative Closed String Geometry", PoS CORFU **2011** (2011) 086 [arXiv:1205.0100 [hep-th]].
- [31] D. Mylonas, P. Schupp and R. J. Szabo, "Membrane Sigma-Models and Quantization of Non-Geometric Flux Backgrounds", JHEP **1209** (2012) 012 [arXiv:1207.0926 [hep-th]].
- [32] D. Andriot, M. Larfors, D. Lüst and P. Patalong, "(Non-)commutative Closed String on T-dual Toroidal Backgrounds", JHEP **1306** (2013) 021 [arXiv:1211.6437 [hep-th]].
- [33] I. Bakas and D. Lüst, "3-Cocycles, Non-Associative Star-Products and the Magnetic Paradigm of R -Flux String Vacua", JHEP **1401** (2014) 171 [arXiv:1309.3172 [hep-th]].
- [34] L. Castellani, "Higher Form Gauge Fields and Their Non-associative Symmetry Algebras", JHEP **1409** (2014) 055 [arXiv:1310.7185 [hep-th]].
- [35] D. Mylonas, P. Schupp and R. J. Szabo, "Non-Geometric Fluxes, Quasi-Hopf Twist Deformations and Non-associative Quantum Mechanics", J. Math. Phys. **55** (2014) 122301 [arXiv:1312.1621 [hep-th]].
- [36] D. Mylonas, P. Schupp and R. J. Szabo, "Non-associative Geometry and Twist Deformations in Non-geometric String Theory", PoS ICMP **2013** (2013) 007 [arXiv:1402.7306 [hep-th]].
- [37] D. Mylonas and R. J. Szabo, "Non-associative Field Theory on Non-Geometric Spaces", Fortsch. Phys. **62** (2014) 727 [arXiv:1404.7304 [hep-th]].

- [38] C. D. A. Blair, "Non-commutativity and Non-associativity of the Doubled String in Non-geometric Backgrounds", preprint arXiv:1405.2283 [hep-th].
- [39] G. E. Barnes, A. Schenkel and R. J. Szabo, "Non-associative Geometry in Quasi-Hopf Representation Categories I: Bimodules and Their Internal Homomorphisms", J. Geom. Phys. **89** (2014) 111 [arXiv:1409.6331 [math.QA]].
- [40] A. Deser, "Star Products on Graded Manifolds and α' -corrections to Courant Algebroids from String Theory", preprint arXiv:1412.5966 [hep-th].
- [41] P. Aschieri and R. J. Szabo, "Tri-products, Non-associative Star Products and Geometry of R -flux String Compactifications", preprint arXiv:1504.03915 [hep-th].
- [42] V. Mathai and J. Rosenberg, "T-duality for Torus Bundles via Non-commutative Topology", Commun. Math. Phys. **253** (2005) 705 [arXiv:hep-th/0401168];
"T-duality for Torus Bundles with H -fluxes via Non-commutative Topology, II: The High-dimensional Case and the T-duality Group", Adv. Theor. Math. Phys. **10** (2006) 123 [arXiv:hep-th/0508084].
- [43] P. Bouwknegt, K. Hannabuss and V. Mathai, "Non-associative Tori and Applications to T-duality", Commun. Math. Phys. **264** (2006) 41 [arXiv:hep-th/0412092].
- [44] C. Condeescu, I. Florakis, C. Kounnas and D. Lüst, "Gauged Supergravities and Non-geometric Q/R -fluxes from Asymmetric Orbifold CFT's", JHEP **1310** (2013) 057 [arXiv:1307.0999 [hep-th]].
- [45] F. Hassler and D. Lüst, "Consistent Compactification of Double Field Theory on Non-geometric Flux Backgrounds", JHEP **1405** (2014) 085 [arXiv:1401.5068 [hep-th]].
- [46] A. A. Tseytlin, "Duality Symmetric Formulation of String World-Sheet Dynamics", Phys. Lett. B **242** (1990) 163.
- [47] W. Siegel, "Superspace Duality in Low-energy Superstrings", Phys. Rev. D **48** (1993) 2826 [hep-th/9305073].
- [48] C. M. Hull, "Doubled Geometry and T-Folds", JHEP **0707** (2007) 080 [arXiv:hep-th/0605149].
- [49] C. Hull and B. Zwiebach, "Double Field Theory", JHEP **0909** (2009) 099 [arXiv:0904.4664 [hep-th]];
"The Gauge Algebra of Double Field Theory and Courant Brackets", JHEP **0909** (2009) 090 [arXiv:0908.1792 [hep-th]].

- [50] O. Hohm, C. Hull and B. Zwiebach, "Background Independent Action for Double Field Theory", JHEP **1007** (2010) 016 [arXiv:1003.5027 [hep-th]];
 "Generalized Metric Formulation of Double Field Theory", JHEP **1008** (2010) 008 [arXiv:1006.4823 [hep-th]].
- [51] D. S. Berman, H. Godazgar, M. J. Perry and P. West, "Duality Invariant Actions and Generalised Geometry", JHEP **1202** (2012) 108 [arXiv:1111.0459 [hep-th]].
- [52] G. Aldazabal, W. Baron, D. Marques and C. Nunez, "The Effective Action of Double Field Theory", JHEP **1111** (2011) 052 (Erratum JHEP **1111** (2011) 109) [arXiv:1109.0290 [hep-th]].
- [53] O. Hohm and B. Zwiebach, "On the Riemann Tensor in Double Field Theory", JHEP **1205** (2012) 126 [arXiv:1112.5296 [hep-th]];
 "Large Gauge Transformations in Double Field Theory", JHEP **1302** (2013) 075 [arXiv:1207.4198 [hep-th]];
 "Towards an Invariant Geometry of Double Field Theory", J. Math. Phys. **54** (2013) 032303 [arXiv:1212.1736 [hep-th]];
 "Green-Schwarz Mechanism and α' -deformed Courant Brackets", JHEP **1501** (2015) 012 [arXiv:1407.0708 [hep-th]];
 "Double Field Theory at Order α' ", JHEP **1411** (2014) 075 [arXiv:1407.3803 [hep-th]].
- [54] D. S. Berman, C. D. A. Blair, E. Malek and M. J. Perry, "The $O(D, D)$ Geometry of String Theory", Int. J. Mod. Phys. A **29** (2014) 15, 1450080 [arXiv:1303.6727 [hep-th]].
- [55] D. Geissbuhler, D. Marques, C. Nunez and V. Penas, "Exploring Double Field Theory", JHEP **1306** (2013) 101 [arXiv:1304.1472 [hep-th]].
- [56] G. Aldazabal, D. Marques and C. Nunez, "Double Field Theory: A Pedagogical Review", Class. Quant. Grav. **30** (2013) 163001 [arXiv:1305.1907 [hep-th]].
- [57] D. S. Berman and K. Lee, "Supersymmetry for Gauged Double Field Theory and Generalised Scherk-Schwarz Reductions", Nucl. Phys. B **881** (2014) 369 [arXiv:1305.2747 [hep-th]].
- [58] D. S. Berman and D. C. Thompson, "Duality Symmetric String and M-Theory", Phys. Rept. **566** (2014) 1 [arXiv:1306.2643 [hep-th]].
- [59] O. Hohm, W. Siegel and B. Zwiebach, "Doubled α' -geometry", JHEP **1402** (2014) 065 [arXiv:1306.2970 [hep-th]].

- [60] O. Hohm, D. Lüst and B. Zwiebach, "The Spacetime of Double Field Theory: Review, Remarks, and Outlook", *Fortsch. Phys.* **61** (2013) 926 [arXiv:1309.2977 [hep-th]].
- [61] R. Blumenhagen, M. Fuchs, F. Hassler, D. Lüst and R. Sun, "Non-associative Deformations of Geometry in Double Field Theory", *JHEP* **1404** (2014) 141 [arXiv:1312.0719 [hep-th]].
- [62] D. S. Berman, M. Cederwall and M. J. Perry, "Global Aspects of Double Geometry", *JHEP* **1409** (2014) 066 [arXiv:1401.1311 [hep-th]].
- [63] A. Betz, R. Blumenhagen, D. Lüst and F. Rennecke, "A Note on the CFT Origin of the Strong Constraint of DFT", *JHEP* **1405** (2014) 044 [arXiv:1402.1686 [hep-th]].
- [64] M. Cederwall, "The Geometry Behind Double Geometry", *JHEP* **1409** (2014) 070 [arXiv:1402.2513 [hep-th]];
"T-duality and Non-geometric Solutions from Double Geometry", *Fortsch. Phys.* **62** (2014) 942 [arXiv:1409.4463 [hep-th]].
- [65] G. Papadopoulos, "Seeking the Balance: Patching Double and Exceptional Field Theories", *JHEP* **1410** (2014) 089 [arXiv:1402.2586 [hep-th]];
"C-spaces, Generalized Geometry and Double Field Theory", preprint arXiv:1412.1146 [hep-th].
- [66] C. M. Hull, "Finite Gauge Transformations and Geometry in Double Field Theory", *JHEP* **1504** (2015) 109 [arXiv:1406.7794 [hep-th]].
- [67] R. Blumenhagen, F. Hassler and D. Lüst, "Double Field Theory on Group Manifolds", *JHEP* **1502** (2015) 001 [arXiv:1410.6374 [hep-th]].
- [68] R. Blumenhagen, P. d. Bosque, F. Hassler and D. Lüst, "Generalized Metric Formulation of Double Field Theory on Group Manifolds", preprint arXiv:1502.02428 [hep-th].
- [69] M. Rocek and E. P. Verlinde, "Duality, Quotients, and Currents", *Nucl. Phys.* **B373** (1992) 630 [arXiv:hep-th/9110053].
- [70] T. Buscher, "A Symmetry of the String Background Field Equations", *Phys. Lett.* **B194** (1987) 59;
"Path Integral Derivation of Quantum Duality in Non-linear Sigma-Models", *Phys. Lett.* **B201** (1988) 466.
- [71] A. Giveon, E. Rabinovici and G. Veneziano, "Duality in String Background Space", *Nucl. Phys.* **B322** (1989) 167.

- [72] E. Alvarez, L. Alvarez-Gaume and Y. Lozano, "A Canonical Approach to Duality Transformations", Phys. Lett. **B336**, 183 (1994) [arXiv:hep-th/9406206].
- [73] I. Bakas and K. Sfetsos, "T-duality and World-Sheet Supersymmetry", Phys. Lett. **B349** (1995) 448 [arXiv:hep-th/9502065].
- [74] A. Giveon, M. Porrati and E. Rabinovici, "Target Space Duality in String Theory", Phys. Rept. **244** (1994) 77 [arXiv:hep-th/9401139].
- [75] D. Zagier, "The Dilogarithm Function" in *Frontiers in Number Theory, Physics and Geometry II*, P. Cartier, B. Julia, P. Moussa, P. Vanhove (eds.), p. 3-65, Springer-Verlag, Berlin, 2006.
- [76] Lj. Davidović, B. Nicolić and B. Sazdović, "Canonical Approach to the Closed String Noncommutativity", Eur. Phys. J. C **74** (2014) 1, 2734 [arXiv:1307.6158 [hep-th]].
- [77] S. Groot Nibbelink and P. Patalong, "A Lorentz Invariant Doubled World-sheet Theory", Phys. Rev. D **87** (2013) 4, 041902 [arXiv:1207.6110 [hep-th]].
- [78] S. Groot Nibbelink, F. Kurz and P. Patalong, "Renormalization of a Lorentz Invariant Doubled World-sheet Theory", JHEP **1410** (2014) 114 [arXiv:1308.4418 [hep-th]].