

CERN L3 013

AB

c1



CERN LIBRARIES, GENEVA

CERN LIBRARIES, GENEVA



CM-P00065282

Test of QED in $e^+e^- \rightarrow \gamma\gamma$ at LEP

The L3 Collaboration

ABSTRACT

We have measured the cross-section of the reaction $e^+e^- \rightarrow \gamma\gamma$ at center of mass energies around the Z^0 mass. The results are in good agreement with QED predictions. For the QED cutoff parameters the limit values of $\Lambda_+ > 103$ GeV and $\Lambda_- > 118$ GeV are found. For the decays $Z^0 \rightarrow \gamma\gamma$, $Z^0 \rightarrow \pi^0\gamma$, $Z^0 \rightarrow \eta\gamma$ and $Z^0 \rightarrow \gamma\gamma\gamma$ we find upper limits of 2.9×10^{-4} , 2.9×10^{-4} , 4.1×10^{-4} and 1.2×10^{-4} , respectively. All limits are at 95% C.L.

1. Introduction

At LEP-energies, the reaction $e^+e^- \rightarrow \gamma\gamma$ provides a clean test of QED. In contrast to lepton pair production it is in lowest order not affected by weak interaction effects and hadronic vacuum polarisation. In the absence of rare or theoretically forbidden decays, such as $Z^0 \rightarrow \gamma\gamma$, this reaction can only proceed via the exchange of a virtual electron. On the other hand, deviations from QED in the Z^0 region, could yield information on non-Standard Model properties of this boson.

The first order differential cross-section for $e^+e^- \rightarrow \gamma\gamma$ is predicted by the Born-term of QED. Deviations from QED are generally parametrized by introducing cutoff parameters Λ_{\pm} , i.e. by generalizing the lowest order QED differential cross section to [1,2,3]:

$$\frac{d\sigma}{d\Omega} = \frac{\alpha^2}{s} \cdot \frac{1 + \cos^2\theta}{1 - \cos^2\theta} \cdot \left[1 \pm \frac{s^2}{2\Lambda_{\pm}^4} (1 - \cos^2\theta) \right]. \quad (1)$$

At LEP-energies higher order terms are important however and radiative corrections [4] are necessary before comparing the above expression with data. We report here on a comparison between our data on $e^+e^- \rightarrow \gamma\gamma$ obtained in the Z^0 -region and the QED prediction for this reaction calculated up to order α^3 .

The same reaction $e^+e^- \rightarrow \gamma\gamma$ can also be used to set limits for the mass of an excited virtual electron (e^*) [3] and various forbidden or rare Z^0 decay modes.

2. The L3 Detector

The L3 detector covers 99% of 4π . The detector includes a central vertex chamber, a precise electromagnetic calorimeter composed of BGO crystals, a uranium and brass hadron calorimeter with proportional wire chamber readout, a high accuracy muon chamber system, and a ring of scintillation counters. These detectors are installed in a 12 m diameter magnet which provides a uniform field of 0.5 T along the beam direction. The luminosity is determined by measuring small angle Bhabha events in two forward calorimeters consisting of BGO crystals. A detailed description of each detector subsystem, and its performance, is given in ref. [5].

The fine segmentation of the electromagnetic calorimeter allows the measurement of photon and electron showers with an angular resolution of 2 mrad. For the present analysis we only use data collected in the polar angle θ region:

$$\frac{\pi}{4} < \theta < \frac{3\pi}{4}.$$

3. Event Selection

Events collected from the 1990 LEP-running period (March to June) corresponding to 2.24 pb^{-1} are used for the present analysis.

The selection of the $e^+e^- \rightarrow \gamma\gamma$ candidates is primarily based on the number of shower peaks in the BGO-calorimeter and on their measured energies. We require:

- i) at least two but no more than twelve shower peaks.
- ii) two of them should have an energy between 35 GeV and 55 GeV.

The first cut eliminates a major fraction of the hadronic events. The second one eliminates cosmic ray interactions, hadronic and $\tau\tau$ events.

iii) next, we eliminate all events in which both major shower peaks have a matching track in the vertex chamber within 50 mrad in transverse projection.

The result is a sample of 35 candidates with no matching vertex chamber tracks and a sample of 4 candidates with one matching vertex chamber track. Further examination of the latter sample indicates that only one of them is a recognizable γ -conversion in the beam pipe with both decay tracks clearly separated. It should be noted that from the material in front of the TEC we expect 0.7 γ -conversions in our sample. The remaining 3 events are classified as $e^+e^- \rightarrow e^+e^-$ events with one track detected and are eliminated from the sample. The contamination in the remaining sample where $e^+e^- \rightarrow e^+e^-$ with both tracks undetected is less than 0.005 events.

Fig. 1 shows a plot of $(E_1 + E_2)/\sqrt{s}$ versus $(\theta_1 + \theta_2 - \pi)$ for the centers of the shower peaks of the remaining 36 events. The figure shows the presence of 4 events with $(E_1 + E_2) < 0.9\sqrt{s}$, which combine a relatively large acollinearity with a low total energy. Initial state radiation produces events of this type. It affects the angular distribution in equation 1, therefore a cut is imposed requiring $(E_1 + E_2) > 0.9\sqrt{s}$. Applying the same cut to a Monte Carlo sample (of about 1000 $e^+e^- \rightarrow \gamma\gamma(\gamma)$ events) indicates that it has an acceptance of 89% in the polar angle region under consideration; the number of events removed by the cut agrees with this acceptance.

After the above cut we are left with a sample of 32 events.

4. Analysis $e^+e^- \rightarrow \gamma\gamma$

Table 1 presents the cross sections observed for $e^+e^- \rightarrow \gamma\gamma$ for three different center of mass energy regions and for the polar region under consideration. Within this angular range the efficiency of the electromagnetic calorimeter is $(98 \pm 1)\%$. Also given are the corresponding integrated luminosities and the number of events.

TABLE 1

Region	Energy (GeV)	Luminosity (pb ⁻¹)	Events	Visible Cross-section (pb)
below Z	89.50	0.567	8	16.2 ± 5.4
on Z	91.28	1.202	18	17.2 ± 3.8
above Z	93.17	0.471	6	14.7 ± 5.6
all	91.22	2.240	32	16.4 ± 2.7

Integration of the lowest order QED differential cross section between $\pi/4$ and $3\pi/4$ yields a prediction of 16.4 pb at 91.2 GeV. Applying the required radiative corrections, up to order α^3 , using a program written by Berends and Kleiss [4] changes the QED-prediction to 16.8 pb, in agreement with the results presented in table 1.

Fig.2 shows a comparison between the radiatively corrected QED angular dependence and our data points. The $\cos \theta$ -dependence of the data is consistent with QED. To test the agreement with QED we used the binning-free Smirnov-Cramér-von Mises test [6]. Parametrizing a possible QED-deviation as indicated in the introduction, and varying the cutoff Λ -parameters, one obtains the following limits at the 95% confidence level:

$$\begin{aligned}\Lambda_+ &> 103 \text{ GeV} \\ \Lambda_- &> 118 \text{ GeV.}\end{aligned}$$

Table 2 shows a comparison between these limits and previously obtained e^+e^- -results [7].

TABLE 2

Experiment	Λ_+ (GeV)	Λ_- (GeV)
L3 (this experiment)	103	118
OPAL	82	89
CELLO	59	44
JADE	61	57
MARK J	72	65
PLUTO	46	-
TASSO	61	56
HRS	59	59
MAC	66	67
AMY	65	-
TOPAZ	94	59

It is possible to parametrize a modification of the electron propagator in the reaction $e^+e^- \rightarrow \gamma\gamma$ in terms of the exchange of a virtual excited electron e^* [3].

The two parameters that enter into consideration then are: λ , the ratio of the e^* to e coupling, and M_{e^*} , the mass of the e^* . Assuming $\lambda = 1$ we find at 95% C.L.:

$$M_{e^*} > 83 \text{ GeV}.$$

This result is in agreement with limits recently obtained [8,9,10].

5. Rare Z^0 Decays

Any deviation from the QED-prediction for the $e^+e^- \rightarrow \gamma\gamma$ cross-section could also be due to rare Z^0 -decays such as the theoretically forbidden $Z^0 \rightarrow \gamma\gamma$ [11] or the decays $Z^0 \rightarrow \pi^0\gamma$, $Z^0 \rightarrow \eta\gamma$, in which the π^0 or η would be indistinguishable from a γ . A difference between such Z^0 -decays and QED-events is the angular distribution. The QED-reaction is strongly peaked forward whereas the Z^0 -decays show essentially a $(1 + \cos^2\theta)$ -dependence. Upper limits for the Z^0 -decays can thus be obtained by determining the amount of extra contribution of the type $(1 + \cos^2\theta)$ our $\gamma\gamma$ -sample is able to accommodate in the θ -region under consideration. Taking into account the geometrical acceptances and the η -decay modes one arrives at detection efficiencies of 62%, 62% and 43% for $Z^0 \rightarrow \gamma\gamma$, $Z^0 \rightarrow \pi^0\gamma$ and $Z^0 \rightarrow \eta\gamma$ respectively. Using these numbers one derives [6] at 95% confidence level the following upper limits:

$$\begin{aligned} \text{B.R. } (Z^0 \rightarrow \gamma\gamma) &< 2.9 \times 10^{-4} \\ \text{B.R. } (Z^0 \rightarrow \pi^0\gamma) &< 2.9 \times 10^{-4} \\ \text{B.R. } (Z^0 \rightarrow \eta\gamma) &< 4.1 \times 10^{-4}. \end{aligned}$$

As an extension of the $e^+e^- \rightarrow \gamma\gamma$ study, a search was made for the higher order QED process $e^+e^- \rightarrow \gamma\gamma\gamma$. The selection criteria used were the following:

- i) at least three shower peaks, together having an energy greater than $0.5\sqrt{s}$,
- ii) an acoplanarity between these clusters of less than 9 degrees,
- iii) no matching tracks in the vertex chamber.

No candidate events are found. The acceptance of the above cuts, for the process $Z^0 \rightarrow \gamma\gamma\gamma$, roughly corresponds to the 3γ acceptance of the BGO-barrel i.e. 35%. Using Poisson-statistics we derive:

$$\text{B.R. } (Z^0 \rightarrow \gamma\gamma\gamma) < 1.2 \times 10^{-4}$$

at the 95% confidence level.

The Standard Model predicts a $\text{B.R.}(Z^0 \rightarrow \gamma\gamma\gamma)$ of 8×10^{-10} [12]. However in some composite models branching ratio's as high as 2×10^{-4} have been predicted [13].

Acknowledgments

We wish to thank CERN for its hospitality and help. We want particularly to express our gratitude to the LEP division: it is their excellent achievements which made this experiment possible. We acknowledge the support of all the funding agencies which contributed to this experiment.

The L3 Collaboration:

B.Adeva,¹⁵ O.Adriani,¹³ M.Aguilar-Benitez,²¹ H.Akbari,⁵ J.Alcaraz,²¹ A.Aloisio,²³ G.Alverson,⁹
 M.G.Alviggi,²³ Q.An,¹⁶ H.Anderhub,³⁶ A.L.Anderson,¹² V.P.Andreev,¹⁴ T.Angelov,¹² L.Antonov,³¹
 D.Antreasyan,⁷ P.Arce,²¹ A.Arefiev,²² T.Azemoon,³ T.Aziz,⁸ P.V.K.S.Baba,¹⁶ P.Bagnaia,²⁷
 J.A.Bakken,²⁶ L.Baksay,²⁹ R.C.Ball,³ S.Banerjee,^{8,16} J.Bao,⁵ L.Barone,²⁷ A.Bay,¹⁷ U.Becker,¹²
 J.Behrens,³⁶ S.Beingessner,⁴ Gy.L.Bencze,^{10,29} J.Berdugo,²¹ P.Berges,¹² B.Bertucci,²⁷ B.L.Betev,³¹
 A.Biland,³⁶ R.Bizzarri,²⁷ J.J.Blaising,⁴ P.Blömeke,¹ B.Blumenfeld,⁵ G.J.Bobbink,² M.Bocciolini,¹³
 W.Böhlen,³⁴ A.Böhm,¹ T.Böhringer,¹⁹ B.Borgia,²⁷ D.Bourilkov,³¹ M.Bourquin,¹⁷ D.Boutigny,⁴
 J.G.Branson,²⁸ I.C.Brock,²⁵ F.Bruyan,¹⁵ C.Buisson,²⁰ A.Bujak,³³ J.D.Burger,¹² J.P.Burq,²⁰
 J.Busenitz,³² X.D.Cai,¹⁶ C.Camps,¹ M.Capelli,³ F.Carbonara,²³ F.Carminati,¹³ A.M.Cartacci,¹³
 M.Cerrada,²¹ F.Cesaroni,²⁷ Y.H.Chang,¹² U.K.Chaturvedi,¹⁶ M.Chemarin,²⁰ A.Chen,³⁸ C.Chen,⁶
 G.M.Chen,⁶ H.F.Chen,¹⁸ H.S.Chen,⁶ M.Chen,¹² M.L.Chen,³ G.Chiefari,²³ C.Y.Chien,⁵ C.Civinini,¹³
 I.Clare,¹² R.Clare,¹² G.Coignet,⁴ N.Colino,¹⁵ V.Commichau,¹ G.Conforto,¹³ A.Contin,¹⁵ F.Crijns,²
 X.Y.Cui,¹⁶ T.S.Dai,¹² R.D'Alessandro,¹³ R.de Asmundis,²³ A.Degré,⁴ K.Deiters,^{15,35} E.Dénes,¹⁰
 P.Denes,²⁶ F.DeNotaristefani,²⁷ M.Dhina,³⁶ D.DiBitonto,³² M.Diemoz,²⁷ F.Diez-Hedo,¹⁵
 H.R.Dimitrov,³¹ C.Dionisi,²⁷ F.Dittus,²⁴ R.Dolin,¹² E.Drago,²³ T.Driever,² D.Duchesneau,¹⁷
 P.Duinker,^{2,15} I.Duran,^{15,21} H.El Mamouni,²⁰ A.Engler,²⁵ F.J.Eppling,¹² F.C.Erné,² P.Extermann,¹⁷
 R.Fabbretti,³⁶ G.Faber,¹² S.Falciano,^{15,27} Q.Fan,^{16,6} S.J.Fan,³⁰ M.Fabre,³⁶ J.Fay,²⁰ J.Fehlmann,³⁶
 H.Fenker,⁹ T.Ferguson,²⁵ G.Fernandez,²¹ F.Ferroni,^{27,15} H.Fesefeldt,¹ J.Field,¹⁷ G.Finocchiaro,²⁷
 P.H.Fisher,⁵ G.Forconi,¹⁷ T.Foreman,² K.Freudenreich,³⁶ W.Friebel,³⁵ M.Fukushima,¹² M.Gailloud,¹⁹
 Yu.Galaktionov,²² E.Gallo,¹³ S.N.Ganguli,³ P.Garcia-Abia,²¹ S.S.Gau,³⁸ S.Gentile,²⁷ M.Glaubman,⁹
 S.Goldfarb,³ Z.F.Gong,^{16,18} E.Gonzalez,²¹ A.Gordeev,²² P.Göttlicher,¹ D.Goujon,¹⁷ C.Goy,⁴
 G.Gratta,²⁴ C.Grinnell,¹² M.Gruenewald,²⁴ M.Guanziroli,¹⁶ A.Gurtu,⁸ H.R.Gustafson,³ L.J.Gutay,³³
 H.Haan,¹ S.Hancke,¹ K.Hangarter,¹ M.Harris,¹⁵ A.Hasan,¹⁶ C.F.He,³⁰ A.Heavey,²⁶ T.Hebbeker,¹
 M.Hebert,²⁸ G.Herten,¹² U.Herten,¹ A.Hervé,¹⁵ K.Hilgers,¹ H.Hofer,³⁶ H.Hoorani,¹⁶ L.S.Hsu,³⁸ G.Hu,¹⁶
 G.Q.Hu,³⁰ B.Ille,²⁰ M.M.Ilyas,¹⁶ V.Innocente,^{23,15} E.Isiksal,³⁶ E.Jagei,¹⁶ B.N.Jin,⁶ L.W.Jones,³
 P.Kaaret,²⁶ R.A.Khan,¹⁶ Yu.Kamyshkov,²² Y.Karyotakis,^{4,15} M.Kaur,¹⁶ S.Khokhar,¹⁶ V.Khoze,¹⁴
 D.Kirkby,²⁴ W.Kitte,² A.Klimentov,²² A.C.König,² O.Kornadt,¹ V.Koutsenko,²² R.W.Kraemer,²⁵
 T.Kramer,¹² V.R.Krastev,³¹ W.Krenz,¹ J.Krizmanic,⁵ A.Kuhn,³⁴ K.S.Kumar,¹¹ V.Kumar,¹⁶
 A.Kunin,²² A.van Laak,¹ V.Lalieu,¹⁷ G.Landi,¹³ K.Lanius,^{15,35} W.Lange,³⁵ D.Lanskei,¹ S.Lanzano,²³
 P.Lebrun,²⁰ P.Lecomte,³⁶ P.Lecoq,¹⁵ P.Le Coultre,³⁶ I.Leedom,⁹ J.M.Le Goff,¹⁵ A.Leike,³⁵ L.Leistam,¹⁵
 R.Leiste,³⁵ M.Lenti,¹³ J.Lettry,³⁶ P.M.Levchenko,¹⁴ X.Leytens,² C.Li,¹⁸ H.T.Li,⁶ J.F.Li,¹⁶ L.Li,³⁶
 P.J.Li,³⁰ Q.Li,¹⁶ X.G.Li,⁶ J.Y.Liao,³⁰ Z.Y.Lin,¹⁸ F.L.Linde,²⁵ D.Linnhofer,¹⁵ R.Liu,¹⁶ Y.Liu,¹⁶
 W.Lohmann,³⁵ S.Lökös,²⁹ E.Longo,²⁷ Y.S.Lu,⁶ J.M.Lubbers,² K.Lübelsmeyer,¹ C.Luci,¹⁵ D.Luckey,^{7,12}
 L.Ludovici,²⁷ X.Lue,³⁶ L.Luminari,²⁷ W.G.Ma,¹⁸ M.MacDermott,³⁶ R.Magahiz,²⁹ M.Maire,⁴
 P.K.Malhotra,⁸ R.Malik,¹⁶ A.Malinin,²² C.Maña,^{15,21} D.N.Mao,³ Y.F.Mao,⁸ M.Maolinbay,³⁶
 P.Marchesini,¹⁶ A.Marchionni,¹³ J.P.Martin,²⁰ L.Martinez,²¹ F.Marzano,²⁷ G.G.G.Massaro,²
 T.Matsuda,¹² K.Mazumdar,⁸ P.McBride,¹¹ T.McMahon,³³ D.McNally,³⁶ Th.Meinholtz,¹ M.Merk,²
 L.Merola,²³ M.Meschini,¹³ W.J.Metzger,² Y.Mi,¹⁶ M.Micke,¹ U.Micke,¹ G.B.Mills,³ Y.Mir,¹⁶
 G.Mirabelli,²⁷ J.Mnich,¹ M.Möller,¹ L.Montanet,¹⁵ B.Monteleoni,¹³ G.Morand,¹⁷ R.Morand,⁴
 S.Morganti,²⁷ V.Morgunov,²² R.Mount,²⁴ E.Nagy,^{10,15} M.Napolitano,²³ H.Newman,²⁴ M.A.Niaz,¹⁶
 L.Niessen,¹ W.D.Nowak,³⁵ H.Nowak,³⁵ S.Nowak,³⁵ D.Pandoulas,¹ G.Passaleva,¹³ G.Paternoster,²³
 S.Patricelli,²³ Y.J.Pei,¹ D.Perret-Gallix,⁴ J.Perrier,¹⁷ A.Pevsner,⁵ M.Pieri,¹³ P.A.Piroué,²⁶
 V.Plyaskin,²² M.Pohl,³⁶ V.Pojidaev,²² N.Produit,¹⁷ J.M.Qian,^{12,16} K.N.Qureshi,¹⁶ R.Raghavan,⁸
 G.Rahal-Callot,³⁶ P.Razis,³⁶ K.Read,²⁶ D.Ren,³⁶ Z.Ren,¹⁶ S.Reucroft,⁹ T.Riemann,³⁵ O.Rind,³
 C.Rippich,²⁵ H.A.Rizvi,¹⁶ B.P.Roe,³ M.Röhner,¹ S.Röhner,¹ Th.Rombach,¹ L.Romero,²¹ J.Rose,¹
 S.Rosier-Lees,⁴ R.Rosmalen,² Ph.Rosselet,¹⁹ J.A.Rubio,^{15,21} W.Ruckstuhl,¹⁷ H.Rykaczewski,³⁶
 M.Sachwitz,³⁵ J.Salicio,²¹ J.M.Salicio,²¹ G.Sartorelli,^{7,16} G.Sauvage,⁴ A.Savin,²² V.Schegelsky,¹⁴
 D.Schmitz,¹ P.Schmitz,¹ M.Schneegans,⁴ M.Schöntag,¹ H.Schopper,³⁷ D.J.Schotanus,² H.J.Schreiber,³⁵
 R.Schulte,¹ S.Schulte,¹ K.Schultze,¹ J.Schütte,¹¹ J.Schwenke,¹ G.Schwering,¹ C.Sciacca,²³ R.Sehgal,¹⁶
 P.G.Seiler,³⁶ J.C.Sens,² I.Sheer,²⁸ V.Shevchenko,²² S.Shevchenko,²² X.R.Shi,²⁵ K.Shrakov,²²
 V.Shoutko,²² E.Shumilov,²² N.Smirnov,¹⁴ A.Sopczak,^{24,28} C.Souyri,⁴ C.Spartiotis,⁵ T.Spickermann,¹
 B.Spiess,³⁴ P.Spillantini,¹³ R.Starosta,¹ M.Steuer,^{7,12} D.P.Stickland,²⁶ B.Stöhr,³⁶ H.Stone,¹⁷
 K.Strauch,¹¹ B.C.Stringfellow,³³ K.Sudhakar,^{8,1} G.Sultanov,¹⁵ R.L.Sumner,²⁶ H.Suter,³⁶
 R.B.Sutton,²⁵ J.D.Swain,¹⁶ A.A.Syed,¹⁶ X.W.Tang,⁶ E.Tarkovsky,²² L.Taylor,⁹ J.M.Thenard,⁴
 E.Thomas,¹⁶ C.Timmermans,² Samuel C.C.Ting,¹² S.M.Ting,¹² Y.P.Tong,³⁸ F.Tonisch,³⁵
 M.Tonutti,¹ S.C.Tonwar,⁸ J.Tóth,¹⁰ G.Trowitzsch,³⁵ K.L.Tung,⁶ J.Ulbricht,³⁴ L.Urbàñ,¹⁰ U.Uwer,¹
 E.Valente,²⁷ R.T.Van de Walle,² H.van der Graaf,² I.Vetlitsky,²² G.Viertei,³⁶ P.Vikas,¹⁶ U.Vikas,¹⁶

M.Vivargent^{4,12} H.Vogel²⁵ H.Vogt³⁵ M.Vollmar¹ G.Von Dardel¹⁵ I.Vorobiev²² A.A.Vorobyov¹⁴
An.A.Vorobyov¹⁴ L.Vuilleumier¹⁹ M.Wadhwa¹⁶ W.Walk¹⁵ W.Wallraff¹ C.R.Wang¹⁸ G.H.Wang²⁵
J.H.Wang⁶ Q.F.Wang¹¹ X.L.Wang¹⁸ Y.F.Wang¹³ Z.Wang¹⁶ Z.M.Wang^{16,18} J.Weber³⁶ R.Weill¹⁹
T.J.Wenaus¹² J.Wenninger¹⁷ M.White¹² R.Wilhelm² C.Willmott²¹ F.Wittgenstein¹⁵ D.Wright²⁶
R.J.Wu⁶ S.L.Wu¹⁶ S.X.Wu¹⁶ Y.G.Wu⁶ B.Wyslouch^{12,15} Z.Z.Xu¹⁸ Z.L.Xue³⁰ D.S.Yan³⁰
B.Z.Yang¹⁸ C.G.Yang⁶ G.Yang¹⁶ K.S.Yang⁶ Q.Y.Yang⁶ Z.Q.Yang³⁰ Q.Ye¹⁶ C.H.Ye¹⁶ S.C.Yeh³⁸
Z.W.Yin³⁰ J.M.You¹⁶ C.Zaccardelli²⁴ L.Zehnder³⁶ M.Zeng¹⁶ Y.Zeng¹ D.Zhang²⁸ D.H.Zhang²
S.Y.Zhang⁶ Z.P.Zhang¹⁸ J.F.Zhou¹ R.Y.Zhu²⁴ A.Zichichi^{15,16} J.Zoll¹⁵

- 1 I. Physikalisches Institut, RWTH, Aachen, Federal Republic of Germany§
III. Physikalisches Institut, RWTH, Aachen, Federal Republic of Germany§
2 National Institute for High Energy Physics, NIKHEF, Amsterdam; NIKHEF-H and University of Nijmegen, Nijmegen, The Netherlands
3 University of Michigan, Ann Arbor, United States of America
4 Laboratoire de Physique des Particules, LAPP, Annecy, France
5 Johns Hopkins University, Baltimore, United States of America
6 Institute of High Energy Physics, IHEP, Beijing, China
7 INFN-Sezione di Bologna, Italy
8 Tata Institute of Fundamental Research, Bombay, India
9 Northeastern University, Boston, United States of America
10 Central Research Institute for Physics of the Hungarian Academy of Sciences, Budapest, Hungary
11 Harvard University, Cambridge, United States of America
12 Massachusetts Institute of Technology, Cambridge, United States of America
13 INFN Sezione di Firenze and University of Firenze, Italy
14 Leningrad Nuclear Physics Institute, Gatchina, Soviet Union
15 European Laboratory for Particle Physics, CERN, Geneva, Switzerland
16 World Laboratory, FBLJA Project, Geneva, Switzerland
17 University of Geneva, Geneva, Switzerland
18 Chinese University of Science and Technology, USTC, Hefei, China
19 University of Lausanne, Lausanne, Switzerland
20 Institut de Physique Nucléaire de Lyon, IN2P3-CNRS/Université Claude Bernard, Villeurbanne, France
21 Centro de Investigaciones Energeticas, Medioambientales y Tecnologicas, CIEMAT, Madrid, Spain
22 Institute of Theoretical and Experimental Physics, ITEP, Moscow, Soviet Union
23 INFN-Sezione di Napoli and University of Naples, Italy
24 California Institute of Technology, Pasadena, United States of America
25 Carnegie Mellon University, Pittsburgh, United States of America
26 Princeton University, Princeton, United States of America
27 INFN-Sezione di Roma and University of Roma, "La Sapienza", Italy
28 University of California, San Diego, United States of America
29 Union College, Schenectady, United States of America
30 Shanghai Institute of Ceramics, SIC, Shanghai, China
31 Central Laboratory of Automation and Instrumentation, CLANP, Sofia, Bulgaria
32 University of Alabama, Tuscaloosa, Alabama, United States of America
33 Purdue University, West Lafayette, Indiana, United States of America
34 Paul Scherrer Institut, PSI, Würenlingen, Switzerland
35 High Energy Physics Institute, Zeuthen-Berlin, German Democratic Republic
36 Eidgenössische Technische Hochschule, ETH Zürich Switzerland
37 University of Hamburg, Federal Republic of Germany
38 High Energy Physics Group, Taiwan, China

§ Supported by the German Bundesministerium für Forschung und Technologie

REFERENCES

- [1] S. Drell, Ann. Phys. (N.Y.) **4** (1958) 75.
- [2] F.E. Low, Phys. Rev. Lett. **14** (1965) 238.
- [3] A. Litke, Harvard Univ., Ph.D. Thesis (1970) unpublished.
- [4] F.A. Berends and R. Kleiss, Nucl. Phys. **B186** (1981) 22.
- [5] L3 Collaboration, B. Adeva *et al.*, Nucl. Instr. and Meth. **A289** (1990) 35.
- [6] W.T. Eadie *et al.*, Statistical Methods in Experimental Physics, p.268. North Holland (1971).
- [7] OPAL Collaboration, Phys. Lett. **B241** (1990) 133;
CELLO Collaboration, H.J. Behrend *et al.*, Phys. Lett. **B123** (1983) 127;
JADE Collaboration, W. Bartel *et al.*, Z. Phys. **C19** (1983) 197;
MARK J Collaboration, B. Adeva *et al.*, Phys. Lett. **B152** (1985) 439;
H.S. Chen, La Rivista del Nuovo Cimento, **1** (1988) 11. PLUTO Collaboration, Ch. Berger *et al.*, Phys. Lett. **B94** (1980) 87;
TASSO Collaboration, M. Althoff *et al.*, Z. Phys. **C26** (1984) 337;
HRS Collaboration, M. Derrick *et al.*, Phys. Lett. **B166** (1986) 468; Phys. Rev. **D34** (1986) 3286;
MAC Collaboration, E. Fernandez *et al.*, Phys. Rev. **D35** (1987) 1;
AMY Collaboration, S.K. Kim *et al.*, Phys. Lett. **B223** (1989) 476;
TOPAZ Collaboration, I. Adachi *et al.*, Phys. Lett. **B200** (1988) 391;
- [8] Aleph Collaboration, D. Decamp *et al.*, Phys. Lett. **B236** (1990) 501.
- [9] Opal Collaboration, M.Z. Akrawy *et al.*, CERN-EP/90-49.
- [10] L3 Collaboration, B. Adeva *et al.*, "Mass Limit for Excited Electrons and Muons from Z^0 -Decay", To be published in Physics Letters.
- [11] C.N. Yang, Phys. Rev. **77** (1950) 242.
- [12] M. Laursen *et al.*, Phys. Rev. **D23** (1981) 2795.
- [13] Compositeness Working Group (F.M. Renard *et al.*), CERN report 89-08, Volume 2, p.185.

FIGURE CAPTIONS

Fig.1. The $\gamma\gamma$ energy sum versus $|\theta_1 + \theta_2 - \pi|$ with the solid line showing the energy cut $(E_1 + E_2) > 0.9\sqrt{s}$.

Fig.2. The measured number of events versus $(|\cos \theta_1| + |\cos \theta_2|)/2$. The solid curves show the QED prediction and the expectations with the cutoff parameters $\Lambda_+ = 103$ GeV and $\Lambda_- = 118$ GeV (95% confidence level limit).

Figure 1




Figure 2

