

Nuclear Matrix Elements for the $\beta\beta$ Decay of ^{76}Ge

B.A. Brown¹, D.L. Fang², and M. Horoi³

¹ Department of Physics and Astronomy and National Superconducting Cyclotron Laboratory Michigan State University, East Lansing, Michigan 48824-1321, USA

² College of Physics, Jilin University, Changchun 130012, China

³ Department of Physics, Central Michigan University, Mount Pleasant, Michigan 48859, USA

Abstract

The nuclear matrix elements for two-neutrino double-beta ($2\nu\beta\beta$) and zero-neutrino double-beta ($0\nu\beta\beta$) decay of ^{76}Ge are evaluated in terms of the configuration interaction (CI), quasiparticle random phase approximation (QRPA) and interacting boson model (IBM) methods. We show that the decomposition of the matrix elements in terms of intermediate states in ^{74}Ge is dominated by ground state of this nucleus. We consider corrections to the CI results that arise from configurations admixtures involving orbitals outside of the CI configuration space by using results from QRPA, many-body-perturbation theory, and the connections to related observables. The CI two-neutrino matrix element is reduced due to the inclusion of spin-orbit partners, and to many-body correlations connected with Gamow-Teller beta decay. The CI zero-neutrino matrix element for the heavy neutrino is enhanced due to particle-particle correlations that are connected with the odd-even oscillations in the nuclear masses. The CI zero-neutrino matrix element for the light neutrino contains both types of correlations that approximately cancel each other.

Many properties of the active neutrinos are measured, but it is not yet established whether they are Dirac or Majorana type particles and their absolute masses are not known. Left-right symmetric extensions to the standard model provide an explanation for the non-zero masses of the left-handed light neutrinos and also predict the existence of right-handed heavy neutrinos [1]. Neutrinoless double-beta ($0\nu\beta\beta$) decay of nuclei provides unique information and constraints on these neutrino properties [2], [3], [4], [5], [6]. The $\beta\beta$ decay process and the associated nuclear matrix elements (NME) have been investigated by using several approaches including the quasiparticle random phase approximation (QRPA), [7], [8] the configuration interaction (CI) model, the interacting boson model (IBM), the generator coordinate method [9], and the projected Hartree-Fock Bogoliubov model [10].

Assuming contributions from the light left-handed (ν) neutrino-exchange mechanism and the heavy right-handed (N) neutrino-exchange mechanism, the decay rate of a neutrinoless double-beta decay process can be written as [4], [11]

$$\left[T_{1/2}^{0\nu}\right]^{-1} = G^{0\nu} \left(|M^{0\nu}|^2 |\eta_\nu|^2 + |M^{0N}|^2 |\eta_N|^2 \right), \quad (1)$$

where $G^{0\nu}$ is the phase space factor [12], [13], M are the nuclear matrix elements (NME), and η are combinations of the neutrino masses [11], [4].

Since the experimental decay rate is proportional to the square of the calculated nuclear matrix elements, it is important to calculate these matrix elements with good accuracy to be able to determine the absolute scale of neutrino masses. However, the theoretical methods used give results that differ from one another by factors of up to 2-3. It is important to understand the nuclear structure aspects of these matrix elements and why the models give differing results.




Fig. 1: The $\beta\beta$ decay of ^{76}Ge . The top part shows the conversion of two neutrons (blue on the left) into two protons (red on the right) in $2\nu\beta\beta$ decay. The bottom part shows the $0\nu\beta\beta$ decay where a virtual neutrino is exchanged.

In this talk we discuss the NME for the $\beta\beta$ decay of ^{76}Ge obtained with the CI, QRPA and IBM-2 methods. We will show that all of these methods have deficiencies. Some of the deficiencies can be addressed with many-body perturbation theory (MBPT) approaches, and connections to other observables.

The nuclear matrix elements can be presented as a sum of Gamow-Teller (M_{GT}), Fermi (M_F), and Tensor (M_T) matrix elements (see, for example, Refs. [14], [15]),

$$M = M_{GT} - \left(\frac{g_V}{g_A} \right)^2 M_F + M_T, \quad (2)$$

where g_V and g_A are the vector and axial constants, correspondingly. We use $g_V = 1$ and $g_A = 1.27$. The M_α are matrix elements of scalar two-body potentials. The Gamow-Teller has the form $V_{GT}(r, A, \mu) \sigma_1 \cdot \sigma_2 \tau_1^- \tau_2^-$ and the Fermi has the form $V_F(r, A, \mu) \tau_1^- \tau_2^-$, where τ^- are the isospin lowering operators. The neutrino potentials depend on the relative distance between the two decaying nucleons, r , the mass number A , and the closure energy μ [16]. The radial forms are given explicitly in [14]. For the heavy-neutrino exchange, the potential does not depend on μ . For the light neutrino matrix element the closure approximation is good to within 10% [17].

The operators for M_{GT} are given to a good approximation by $f(r) \sigma_1 \cdot \sigma_2 \tau_1^- \tau_2^-$, where $f(r)^{2\nu} = 1$ (in closure), $f(r)^{0\nu} = a/r$ and $f(r)^{0N} = b \delta(r)$ where the constants a and b depend on A , μ and the SRC. The results discussed below follow from the expansions of the many-body matrix elements for these three operators in terms of the particle-hole (ph) in ^{76}As or particle-particle (pp) intermediate states in ^{74}Ge [18]. The expansion over pp states ($J_m = J_{pp}$) in ^{74}Ge is shown in Fig. 1. The left-hand column




Fig. 2: Nuclei involved in the calculations for the double-beta decay of ^{76}Ge .

shows the light-neutrino results (ν) for the sum of the GT, F and T contributions. The middle column shows the light-neutrino results for GT contribution only. The right-hand column shows the heavy-neutrino results (N) for the sum of the GT, F and T contribution. The bottom row shows the running sums for 0^+ intermediate states. The middle row shows the running sums for 0^+ and 2^+ intermediate states. The top row shows the running sums for all intermediate states. The red dots are the exact results for the sum over all intermediate states. This shows that the $J_{pp}=0^+$ contribution is completely dominated by a path through the ground state of ^{74}Ge . There is some cancellation from higher J_{pp} values mainly coming from 2^+ that is spread over intermediate states up to about 6 MeV.

The expansion of the NME over intermediate J values obtained from the QRPA calculations is shown in Fig. 2. This figure shows the different types of correlations for the three operators. The 2ν decay is completely determined by the 1^+ states of the ph channel and its expansion over pp is complicated. The heavy neutrino is dominated by the 0^+ states of the pp channel and its expansion over ph states is complicated. The light neutrino NME is some where between these and but looks simplest in the pp channel.

The 2ν tensor NME is zero and the Fermi NME is zero since isospin is conserved. For 0ν and $0N$ the Fermi and tensor parts are both relatively small, and we define a correction factor for these given by $R_{GT} = M/M_{GT}$, where M contains all three terms of Eq. 2. The CI calculations give $R_{GT}^{0\nu} = 1.10(3)$. Larger values of 1.23 for QRPA [19] and 1.33 for IBM-2 [20] were obtained with the older calculations. But more recently, it was found that the 2ν Fermi matrix element was not zero because isospin was being treated incorrectly in QRPA [21] and IBM-2 [22]. After this was corrected the new $M_F^{2\nu}$ values are now zero in all methods. The new results for $R_{GT}^{0\nu}$ are 1.10 [21] and 1.19 [8] for QRPA, and 1.04 [22] for IBM-2. Taking these results into account we adopt a correction factor from the tensor plus Fermi contributions of $R_{GT}^{0\nu} = 1.12(7)$. The ratio for the heavy neutrino is 1.20 for CI, 1.26 for QRPA [8] and 1.00 for IBM-2 [22]. The adopted correction factor is $R_{GT}^{0N} = 1.13(13)$.

In the following we first focus on results for M_{GT} . At the end, the total matrix element M will be obtained from M_{GT} via a product of correction factors R given by $M = [M_{GT}(\text{CI})][R_V][R_S][R_{GT}]$. R_{GT} is defined above. We start with the use of short-range correlations (SRC) [15] based on the CD-Bonn potential [23]. At the end we will give a value and error for the correction to this, R_S , based on a range of assumptions about the SRC. R_V represents the correction coming from a “vertical” expansion of the CI model space that includes the effect of orbitals below and above those in $jj44$. R_V is the main focus of attention of this talk. The orbitals, model spaces and expansion methods are shown in Fig. 3.

The model space for CI and IBM-2 is $jj44$ that consists of the four valence orbitals $0f_{5/2}$, $1p_{3/2}$,




Fig. 3: Expansion of the NME over intermediate states ^{74}Ge .

$1p_{1/2}$ and $0g_{9/2}$ for protons and neutrons. The model space for QRPA are the 21 orbitals with oscillator quanta $N \leq 5$ where $N = 2n + \ell$ for protons and neutrons. The QRPA results are also given when the evaluation of the NME are restricted to $jj44$ and to fp ($jj44$ plus $0f_{7/2}$ and $0g_{7/2}$). In addition to our own CI calculations with the JUN45 [24] and $jj4bpn$ [25] Hamiltonians, we will show results from the $gcn28:50$ Hamiltonian [26] for 2ν [27], 0ν [28] and $0N$ [29].

The method and parameters used for the QRPA calculations [30] are similar to those used in [21]. For the particle-particle channel in order to restore the isospin symmetry, we follow the formalism introduced in [31], [21], by separately fitting the $T = 0$ and $T = 1$ parts of the interaction. For the $T = 1$ part, $g_{pp}^{T=1} = 0.985$ is taken to give $M_F^{2\nu} = 0$. For the $T = 0$ particle-particle channel, two parameter sets were used: (a) $g_{pp}^{T=0} = 0.673$ reproduces the experimental value for $M_{GT}^{2\nu}$, and (b) $g_{pp}^{T=0} = 0.643$ gives a value for $M_{GT}^{2\nu}$ that is a factor of $(1/0.75)^2$ larger than experiment, anticipating that there may be MBPT corrections beyond QRPA that could reduce the strength to low-lying states.

Results for the $2\nu\beta\beta$ NME are shown in Fig. 4. This NME is completely determined by $J_{ph}^\pi = 1^+$ intermediate states in ^{76}As . In CI the summation over intermediate including the energy denominator (Eq. 2 in [27]) is obtained with the strength-function method [32]. The IBM-2 result is not shown because it uses an approximation for the NME based on the closure result for the operator $\sigma_1 \cdot \sigma_2 \tau_1^- \tau_2^-$ together with average closure energies from other methods (Eq. 16 in Ref. [22]). Experiment is reduced by a factor of about $R_V^{2\nu} = 0.45$ compared to CI. $R_V^{2\nu} = M^{2\nu}/M^{2\nu}(\text{CI})$ denotes the correction beyond the $jj44$ model space, due to a “vertical” expansion that includes correlations from orbitals below and above the $jj44$ model space. The QRPA results for $jj44$ and fp show that part of this reduction




Fig. 4: The NME obtained in the QRPA calculations expressed in terms of their contributions from the J_{ph} states in ^{76}As on the left-hand side and the J_{pp} states in ^{74}Ge on the right-hand side. The color code is blue for the results obtained in $jj44$, black for the results obtained in fpg and red for the results obtained in the 21-orbit model spaces.

is due to the missing spin-orbit partners in the $jj44$ model space. The particle-hole correlations are dominated by a strong repulsive interaction in the 1^+ channel. Relative to the non-interacting single-particle distribution, Gamow-Teller strength is reduced in low-lying states and shifted into the giant Gamow-Teller resonance. As shown by the QRPA results for $jj44$ and fpg , both spin-orbit partners are important for the reduction. A similar behavior was observed for CI in the case of ^{136}Xe [33].

Beyond QRPA, it is known that two-particle two-hole (2p-2h) admixtures into the model space wavefunctions are important for Gamow-Teller beta decay. The experimental Gamow-Teller strength is observed to be reduced by a factor of $R'_V = 0.5 - 0.6$ relative to the CI calculations in the sd [35] and pf [36] model spaces. Also the strength extracted from charge-exchange reactions for the total Gamow-Teller strength up to about 25 MeV in excitation energy is reduced by this factor relative to QRPA [37] and the $3(N - Z)$ Ikeda sum rule [38]. Arima et al. [39] and Towner [40] have explained this reduction using MBPT in terms of 2p-2h admixtures into the model-space wavefunctions. Earlier calculations claimed that the reduction in GT strength was due to Δ excitations [41] in the nucleus. However, calculations with a realistic $N\Delta\pi$ interaction vertex have shown that the influence of Δ (and other mesonic-exchange currents) is small [39], [40]. These results are compared to the empirical sd results in Fig. 13 of [35]. In order to conserve the Ikeda sum rule, the reduction in low-lying B(GT) strength is associated with a spreading of strength to high excitation energy [42] that gets removed from the 2ν NME due to the energy denominator in the summation over intermediate states. In summary, relative to CI in the $jj44$ model space, reductions due to a spin-orbit complete model space, together with 2p-2h admixtures are required for the $2\nu\beta\beta$ NME. The observed factor of $R_V = 0.45$ is consistent with expectations.

The results for 0ν (heavy neutrino) are shown in Fig. 5. In addition to our own QRPA results, we show the QRPA result from [8]. The J_{pp} intermediate states are dominated by the 0^+ ground state




Fig. 5: The $jj44$ model space used for the IBM and CI calculations and the vertical expansion terms considered in QRPA and MBPT for the factor R_V .

of ^{74}Ge (see Ref. [18] for details on the analysis). In QRPA the NME increases by a factor of $R_V^{0N} = 1.9$ as the number of orbitals included in the sums increases from $jj44$ to full (21 orbitals). This is due to the strong pairing (particle-particle) part of the Hamiltonians and the resulting increase in the number of coherent pairs contributing to the $0N$ NME. The pairing also gives rise to the odd-even staggering of the nuclear binding energies quantified by the pairing energies D [43], [44]. For the germanium isotopes the experimental pairing energies are a factor of 1.45 larger than that obtained with the first-order expectation value of the CD-Bonn Hamiltonian. Based on the average of this result and the increase observed in QRPA, we will use $R_V^{0N} = 1.65(25)$.

The results for $0\nu\beta\beta$ (light neutrino) are shown in Fig. 6. The largest term in the 0ν NME is from the $J_{pp}^\pi = 0^+$ ground state of ^{74}Ge [18]. In QRPA the NME is nearly constant as the number of orbitals included in the sums increase. Qualitatively this is due to a competition between the reduction from the particle-hole channel observed for 2ν and the enhancement due to the particle-particle channel observed for $0N$. The connection of the 0ν matrix elements with pairing has been previously discussed [45]. The new point of our analysis is that the increase expected from pairing coming from MBPT beyond the $jj44$ model space is cancelled by the reduction from the ph -type correlations.

Contributions from states with $J_{pp} > 0$ cancel part of the NME from $J_{pp} = 0^+$. Within $jj44$ the reduction is dominated by the $J_{pp} = 2^+$ states [18]. For the 0ν NME within $jj44$, one finds $R_{pp}^{0\nu} = \{M_{GT}^{0\nu}/[M_{GT}^{0\nu}(J_{pp} = 0^+)]\} = 0.53$ for CI [18], 0.90 for IBM-2 [20] and 0.72 for QRPA. The reason for these differences may be due to the truncation within $jj44$ made by IBM-2 and QRPA. For the $0N$ NME this ratio is $R_{pp}^{0N} = 0.89$ in CI [18]; the cancellation from higher J_{pp} is much less, and the result is dominated by the $J_{pp} = 0^+$ contribution and its connection to pairing is discussed above. In the $jj44$ model space the agreement between the $0N$ NME (Fig. 5) for CI, QRPA and IBM-2 is much better than that for 0ν (Fig. 6) since the cancellation from higher J_{pp} terms is small.

Holt and Engel [46] considered the effect of 2p-2h admixtures beyond the $jj44$ model space by treating the effective transition operator in MBPT. They found a 20% increase of the 0ν NME for ^{76}Ge . Part of these MBPT contributions go beyond QRPA. At present this is the best estimate for the




Fig. 6: Nuclear matrix elements for $2\nu\beta\beta$ decay of ^{76}Ge . The top point in green is the experimental value [34]. The QRPA results are shown for $g_{pp}^{T=0} = 0.673$ (red dots) and $g_{pp}^{T=0} = 0.643$ (red crosses). The CI results are shown for the JUN45 (dot), jj44bpn (cross) and gcn28:50 (triangle) Hamiltonians.




Fig. 7: The $0N$ NME for heavy neutrino decay of ^{76}Ge . See caption to Fig. 4. The QRPA point with the triangle is from Ref. [8].




Fig. 8: The 0ν NME for the light-neutrino decay of ^{76}Ge . See captions to Figs. 2 and 3.

correction beyond CI in the $jj44$ model space. We will use $R_V^{0\nu} = 1.2(2)$ with a generously large value of 20% for its uncertainty.

The results shown above are based on the CD-Bonn SRC. This is the weakest of several SRC that have been used [15]. The strongest is the AV18 SRC, and the UCOM [47] SRC is about half way between. For our final result we use the average of CD-Bonn and AV18 with an error that encompasses both. The result is that the $0N$ NME are multiplied by $R_S^{0N} = 0.80(20)$ and the 0ν NME are multiplied by $R_S^{0\nu} = 0.97(3)$, where R_S is the SRC correction relative to the CD-Bonn starting point.

Finally, we combine all of the factors discussed above in the form $M = [M_{GT}(CI)][R_V][R_S][R_{GT}]$. Based on the experimental value [34] for 2ν the NME is,

$$M^{2\nu} = 0.140(5) = [0.31(3)][0.45][1][1]. \quad (3)$$

The second term is the empirical correction for R_V due to mixing beyond the $jj44$ model space. The error in the CI NME reflects the spread obtained with the three different Hamiltonians used (Fig. 4).

For $0N$,

$$M^{0N} = [155(10)][1.65(25)][0.80(20)][1.13(13)] = 232(80), \quad (4)$$

where the CI value is from Fig. 5. The error for $0N$ is dominated by the SRC correction. Finally For 0ν ,

$$M^{0\nu} = [3.0(3)][1.2(2)][0.97(3)][1.12(7)] = 3.9(8), \quad (5)$$

where the CI value is from Fig. 6. The error for 0ν is dominated by an estimated uncertainty of 20% in the correction beyond $jj44$. Comparison to previous values must take into account the isospin correction for QRPA and IBM discussed above, and the choice of SRC (in our R_S factor). The range is from 2.8 for CI [28] to 4.7 for IBM-2 [22] and 5.3 for QRPA [8]. Our result is in between these, but it is not an average since we have made comments on the deficiencies of all of these models. Using Eq. 1 with the experimental limit of the half-life ($T_{1/2}^{0\nu} > 3 \times 10^{25}$ yr [48]), and the phase space factor from [12], we obtain $|\eta_V| m_e c^2 < 0.3$ eV.

Sometimes the 2ν correction factor (0.45 in this case) is expressed in terms of an effective g_A value ($g'_A = 0.85$ in this case). Since the factor $(g_A)^4$ appears inside the phase-space factor of Eq. 1, one might think that the decay rate for 0ν and $0N$ could be reduced by a factor of $(g'_A/1.27)^4 = 0.20$ [22], [49]. However, this g'_A is only for a specific operator associated with a specific observable ($2\nu\beta\beta$ decay) relative to a specific model (CI in $jj44$ in this case). The operators involved in 0ν and $0N$ beta decay are different (short ranged), and corrections beyond CI cannot be expressed in terms of an overall change in g_A . It is better to express the renormalizations in terms of factors such as R_V that are operator and model space dependent.

The model-space truncation contributions to R_{pp} should be understood. The error for the R_{GT} correction could be reduced if reasons for the variations within the models is understood. The error for the R_V correction could be reduced if the MBPT results such as those in [46] should be expanded to include the renormalization of the separate effects in the ph and pp channels in order to compare to the results found previously relative to the $jj44$ model space. This includes the reduction in Gamow-Teller beta decay strength [39], [40], and the enhancements of the pairing strength seen in the D values. The basic division between CI and its MBPT corrections from all other orbitals can be checked by no-core and ab-initio CI in lighter nuclei where they are tractable. Other methods such as in-medium SRG [50] and coupled cluster [51] can be used in place of MBPT, and at this level the division between short-range renormalization, R_S , and long-range renormalization, R_V , might be merged. The CI results for the $A = 76$ region can be further checked against spectroscopic observables (occupations number are in good agreement with CI [28]) including two-nucleon transfer. Future calculations should be presented in terms of changes relative to various contributions we have discussed, and evaluations for other cases of interest [52] should be made.

We acknowledge correspondence from Jose Barea, Scott Bogner, Mark Caprio, Javier Menendez, Alfredo Poves and Fedor Simkovic. Support from the NUCLEI SciDAC Collaboration under U.S. Department of Energy Grant No. DE-SC0008529 is acknowledged. MH and BAB also acknowledge U.S. NSF Grant No. PHY-1404442.

References

- [1] J. C. Pati and A. Salam, Phys. Rev. D **10**, 275 (1974); R. N. Mohapatra and J. C. Pati, Phys. Rev. D **11**, 2558 (1975); W.-Y. Keung and G. Senjanovic, Phys. Rev. Lett. **50**, 1427 (1983).
- [2] M. Doi et al., Prog. Theor. Phys. **66**, 1739 (1981); **69**, 602 (1983).
- [3] W. C. Haxton and G. J. Stephenson, Prog. Part. Nucl. Phys. **12**, 409 (1984).
- [4] J. D. Vergados, H. Ejiri, and F. Simkovic, Rep. Prog. Phys. **75**, 106301 (2012).
- [5] F.T. Avignone, S.R. Elliott, and J. Engel, Rev. Mod. Phys. **80**, 481 (2008).

- [6] T. Tomoda, Rep. Prog. Phys. **54**, 53 (1991).
- [7] A. Faessler, M. Gonzalez, S. Kovalenko, and Fedor Simkovic, Phys. Rev. D **90**, 096010 (2014).
- [8] J. Hyvarinen and J. Suhonen, Phys. Rev. **91**, 024613 (2015).
- [9] T.R. Rodriguez and G. Martinez-Pinedo, Phys. Rev. Lett. **105**, 252503 (2010).
- [10] P.K. Rath, R. Chandra, K. Chaturvedi, P.K. Raina, and J.G. Hirsch, Phys. Rev. C **82**, 064310 (2010).
- [11] M. Horoi, Phys. Rev. C **87**, 014320 (2013).
- [12] J. Kotila and F. Iachello, Phys. Rev. C **85**, 034316 (2012).
- [13] S. Stoica and M. Mirea, Phys. Rev. C **88**, 037303 (2013).
- [14] R. A. Senkov and M. Horoi, Phys. Rev. C **88**, 064312 (2013).
- [15] M. Horoi and S. Stoica, Phys. Rev. C **81**, 024321 (2010).
- [16] R. A. Senkov, M. Horoi and B. A. Brown, Phys. Rev. C **89**, 054304 (2014).
- [17] R. A. Senkov and M. Horoi, Phys. Rev. C **90**, 051301(R) (2014).
- [18] B. A. Brown, M. Horoi and R. A. Sen'kov, Phys. Rev. Lett. **113**, 262501 (2014).
- [19] M. S. Yousef, V. Rodin, A. Faessler and F. Simkovic, Phys. Rev. C **79**, 014314 (2009).
- [20] J. Barea and F. Iachello, Phys. Rev. C **79**, 044301 (2009).
- [21] F. Simkovic, V. Rodin, A. Faessler and P. Vogel, Phys. Rev. C **87**, 045501 (2013).
- [22] J. Barea, J. Kotila and F. Iachello, Phys. Rev. C **91**, 034304 (2015).
- [23] R. Machleidt, Phys. Rev. C **63**, 024001 (2001).
- [24] M. Honma, T. Otsuka, T. Mizusaki, and M. Hjorth-Jensen, Phys. Rev. C **80**, 064323 (2009).
- [25] B. Cheal et al., Phys. Rev. Lett. **104**, 252502 (2010).
- [26] A. Gniady, E. Caurier, and F. Nowacki (unpublished).
- [27] E. Caurier, F. Nowacki and A. Poves, Phys. Lett. B **711**, 62 (2012).
- [28] J. Menendez, A. Poves, E. Caurier and F. Nowacki, Phys. Rev. C **80**, 048501 (2009), and A. Poves, private communication.
- [29] M. Blennow, E. Fernandez-Martinez, J. Lopez-Pavon, and J. Menendez, JHEP07, 096 (2010); and J. Menendez, private communication.
- [30] The single-particle energies are taken from a Woods-Saxon potential with Coulomb corrections. All of the residual interactions for QRPA are obtained from solutions of Bruckner G-matrix based on the CD-Bonn one-boson exchange nucleon-nucleon potential. We solve the BCS equations with the CD-Bonn pairing interactions adjusted to give the experimental five-point mass pairing gap. The renormalization factors are $g_{pair}^p = 0.858$ and $g_{pair}^n = 0.978$ for ^{76}Ge , and $g_{pair}^p = 0.894$ and $g_{pair}^n = 1.008$ for ^{76}Se . For the renormalization of the QRPA residual interactions, we use $g_{ph} = 1.0$ for the particle-hole channel.
- [31] V. Rodin and A. Faessler, Phys. Rev. C **84**, 014322 (2011).
- [32] M. Horoi, AIP Conf. Proc. **1304**, 106 (2010).
- [33] M. Horoi and B. A. Brown, Phys. Rev. Lett. **110**, 222502 (2013).
- [34] A. S. Barabash, Phys. Rev. C **81**, 035501 (2010).
- [35] B. A. Brown and B. H. Wildenthal, Annu. Rev. Nucl. Part. Sci. **38**, 29 (1988).
- [36] G. Martinez-Pinedo, A. Poves, E. Caurier, and A. P. Zuker, Phys. Rev. C **53**, R2602 (1996).
- [37] E. Litvinova, B. A. Brown, D. L. Fang, T. Marketin and R. G. T. Zegers, Phys. Lett B **730**, 307 (2014).
- [38] B. D. Anderson, T. C. Chittarakarn, A. R. Baldwin, C. Lebo, R. Madey, J. W. Watson, B. A. Brown and C. C. Foster, Phys. Rev. C **31**, 1161 (1985), and references therein.
- [39] A. Arima, K. Shimizu, W. Bentz, and H. Hyuga, Adv. Nucl. Phys. **18**, 1 (1987).
- [40] I. S. Towner, Phys. Rep. **155**, 263 (1987).
- [41] M. Rho, Nucl. Phys. A **231**, 493 (1974).
- [42] G. F. Bertsch and I. Hamamoto, Phys. Rev. C **26**, 1323 (1982); G. F. Bertsch and H. Esbensen,

- Rep. Prog. Phys. **50**, 607 (1987).
- [43] B. A. Brown, Phys. Rev. Lett. **111**, 162502 (2013).
 - [44] B. A. Brown; Journal of Physics: Conference Series **445**, 012010 (2013).
 - [45] E. Caurier, J. Menendez, F. Nowacki, and A. Poves, Phys. Rev. Lett. **100**, 052503 (2008).
 - [46] J. D. Holt and J. Engel, Phys. Rev. C **87**, 064315 (2013).
 - [47] R. Roth, H. Hergert, P. Papakonstantinou, T. Neff and H. Feldmeier, Phys. Rev. C **72**, 034002 (2005).
 - [48] M. Agostini, et al., Phys. Rev. Lett. **111**, 122503 (2013).
 - [49] J. Menendez, D. Gazit and A. Schwenk, Phys. Rev. Lett. **107**, 062501 (2011).
 - [50] S. K. Bogner, H. Hergert, J. D. Holt, A. Schwenk, S. Binder, A. Calci, J. Langhammer, and R. Roth, Phys. Rev. Lett. **113**, 142501 (2014).
 - [51] G. R. Jansen, J. Engel, G. Hagen, P. Navratil, and A. Signoracci, Phys. Rev. Lett. **113**, 142502 (2014).
 - [52] http://science.energy.gov/~media/np/nsac/pdf/docs/2014/NLDBD_Report_2014_Final.pdf