




The Compact Muon Solenoid Experiment  
**Conference Report**

Mailing address: CMS CERN, CH-1211 GENEVA 23, Switzerland



06 September 2016 (v2, 14 September 2016)

# D0 Meson Production in Heavy Ion Collisions in CMS experiment

Jing Wang for the CMS Collaboration

## Abstract

The measurement of heavy flavour production is a powerful tool to study the properties of the high-density QCD medium created in heavy-ion collisions as heavy quarks are sensitive to the transport properties of the medium and may interact with the QCD matter differently from light quarks. In particular, the comparison between the nuclear modification factors of light and heavy flavoured particles provides insights into the expected flavour dependence of in-medium parton energy loss. With the CMS detector, the D0 meson production is studied in pp and PbPb collisions at 2.76 and 5.02 TeV. In this talk, the nuclear modification factor of D0 meson are presented and compared to the charged hadron nuclear modification factor and theoretical calculations.

Presented at *SQM2016 Strangeness in Quark Matter 2016*

# D<sup>0</sup> Meson Production in Heavy Ion Collisions in CMS experiment

**Jing Wang for the CMS Collaboration**

Massachusetts Institute of Technology, 77 Massachusetts Ave, Cambridge, MA, US

E-mail: [jing.wang@cern.ch](mailto:jing.wang@cern.ch)

**Abstract.** The measurement of heavy flavour production is a powerful tool to study the properties of the high-density QCD medium created in heavy-ion collisions as heavy quarks are sensitive to the transport properties of the medium and may interact with the QCD matter differently from light quarks. In particular, the comparison between the nuclear modification factors of light and heavy flavoured particles provides insights into the expected flavour dependence of in-medium parton energy loss. With the CMS detector, the D<sup>0</sup> meson production is studied in pp and PbPb collisions at 2.76 and 5.02 TeV. In this talk, the nuclear modification factor of D<sup>0</sup> meson are presented and compared to the charged hadron nuclear modification factor and theoretical calculations.

## 1. Introduction


Heavy quarks are considered to be produced mainly via initial hard scatterings in heavy ion collisions. Therefore, they propagate through the medium and carry information about the system at early stage. That makes heavy flavor hadrons good probes of the properties of QCD matter and important tools to study the mechanism of interaction with the matter. The interactions with the medium, the energy loss of a parton, happens via elastic and inelastic collisions, and via gluon bremsstrahlung. At small angles, the radiative energy loss is thought to be suppressed, the manifestation of the ‘dead cone effect’ [1]. If only this effect would be dominant, the nuclear modification factor of heavy flavor hadrons should be larger (less suppression) than that of light hadrons. This effect is also expected to have a strong  $p_T$  dependence, with the highest contribution at low transverse momentum of the heavy meson.

The measurements of the D<sup>0</sup> meson production were conducted by the Compact Muon Solenoid (CMS) [2] detector in PbPb collisions at 2.76 TeV [3] and 5.02 TeV [4]. The talk presented the analysis technique for extracting the D<sup>0</sup> signal, and the D<sup>0</sup> nuclear modification factor.

## 2. D<sup>0</sup> in PbPb at 2.76 TeV

Prompt D<sup>0</sup> production was studied with the minimum bias PbPb data collected with CMS detector during LHC Run I in 2011. Specifically, the decay channel  $D^0 \rightarrow K\pi$  is analyzed in the rapidity range  $|y| < 1$  and  $p_T$  range from 2.5-40 GeV/c. Figure 1 shows the prompt D<sup>0</sup>  $R_{AA}$  in PbPb collisions at 2.76 TeV as a function of  $p_T$  for centrality 0-20% and the comparison with the result of ALICE [5]. The pp reference used for this result was built using both FONLL [6] and data extrapolation: in the range  $p_T < 16$  GeV/c range, the ALICE collaboration measurement

of prompt  $D^0$  cross-section at 7 TeV [7] was rescaled to 2.76 TeV using FONLL input, following the same procedure used in Ref. [8].



**Figure 1.** Prompt  $D^0$   $R_{AA}$  as a function of  $p_T$  for centrality 0-20% in PbPb collisions at 2.76 TeV by CMS collaboration. The results from ALICE collaboration are also shown for comparison [5]

The same analysis method, described in Section 3, was used for both 2.76 TeV and 5 TeV results.


### 3. $D^0$ in pp and PbPb at 5.02 TeV

Both pp and PbPb data at 5.02 TeV are collected by CMS collaboration during LHC Run II in 2015. It is the first  $D^0$  measurement reaching  $p_T = 100$  GeV/c. The 5 TeV measurement is performed in the same rapidity interval as the 2.76 TeV one, in the same decay channel, in two centrality classes, 0-10% and 0-100%.

Events selected online by a minimum bias trigger (about 2 billion events for pp and 150 million events for PbPb) are used for the results at low  $p_T$  ( $< 20$  GeV/c), while events selected by a dedicated  $D^0$  High-Level-Trigger (HLT) are used for the result at  $p_T > 20$  GeV/c. The  $D^0$  HLT path starts from events tagged by energy deposited in the calorimeters. A full track reconstruction is performed for these events, and pairs of tracks with  $p_T > 2$  GeV/c (pp) or 8 GeV/c (PbPb) are analyzed. An event is selected if the pair passed a loose selection on their common vertex displacement. This  $D^0$  trigger worked well and allows reaching very high  $p_T$   $D^0$ .

Offline,  $D^0$  candidates are reconstructed by pairing two oppositely charged tracks, which are afterwards required to pass a kinematic filter. Topological selections based on pointing angle, 3D decay length normalized by its error and  $D^0$  vertex probability are applied to reduce combinatorial background. The pointing angle is defined as the angle between  $D^0$  momentum and the vector from primary vertex and secondary vertex in transverse plane. Figure 2 shows the invariant mass distribution for selected D candidates, for  $5 < p_T < 6$  GeV/c, for pp and PbPb analysis.

The signal shape is modeled by two Gaussians with same mean but different widths. The combinatorial background, generated by randomly combining pairs of tracks not originating



**Figure 2.** Invariant mass distributions of  $D^0$  candidates and their charge conjugates for selected  $p_T$  intervals of pp (left) and PbPb (right) in centrality 0-100%

from a  $D^0$  meson decay, is modeled by a third order polynomial. With the choice of not using particle identification, one track is assigned the mass of a  $K$ , and the other the mass of a  $\pi$ . An additional Gaussian function is used to describe the invariant mass shape of  $D^0$  candidates with incorrect mass assignment from the exchange of  $K$  and  $\pi$  designation.

The raw yields are then corrected for detector acceptance and reconstruction efficiency effects, calculated using PYTHIA, and PYTHIA events embedded in HYDJET PbPb events. In addition, the fraction of  $D^0$  coming from  $c$ -quark fragmentation, are determined by fitting data distribution of the distance of closest approach (DCA) of  $D^0$ , with a prompt and nonprompt shape from Monte Carlo.


Prompt  $D^0$   $R_{AA}$  as a function of  $p_T$  in PbPb collisions at 5.02 TeV for centrality classes 0-100% (left) and 0-10% (right) are shown in Figure 3. The measurements are compared with predictions from some theoretical calculations [9, 10, 11, 12, 13, 14, 15, 16, 17], and most of them describe data well. Comparison with charge particles  $R_{AA}$  in PbPb collisions at 5.02 TeV in the same centrality classes [18] is also shown in Figure 3.

#### 4. Summary

The measurements of prompt  $D^0$  nuclear modification factors in 5.02 TeV PbPb collisions from CMS collaboration are presented.  $D^0$  production presents strong suppression. No significant difference of  $R_{AA}$  is observed between  $D^0$  and charged hadrons, in the kinematic region studied. These results put strong constraints on theoretical calculations in a wide kinematic range. With LHC Run II data, there are various exciting studies concerning heavy flavors. For  $c$  quark related studies, there is an effort to reach prompt  $D^0$   $R_{AA}$  at very low  $p_T$ , and  $D^0$   $v_n$  measurements are also conducted. Also, the results of measurements of B mesons nuclear modification factor are expected to come soon.

#### References

- [1] Armesto N, Salgado C A and Wiedemann U A 2004 *Phys. Rev. D* **69**(11) 114003 URL <http://link.aps.org/doi/10.1103/PhysRevD.69.114003>
- [2] Chatrchyan S *et al.* (CMS Collaboration) 2008 *JINST* **3** S08004
- [3] 2015 Nuclear Modification Factor of prompt  $D^0$  in PbPb Collisions at  $\sqrt{s_{NN}} = 2.76$  TeV Tech. Rep. CMS-PAS-HIN-15-005 CERN Geneva URL <http://cds.cern.ch/record/2055466>



**Figure 3.** Prompt  $D^0$   $R_{AA}$  as function of  $p_T$  in PbPb collisions at 5.02 TeV for centrality classes 0-100% (left) and 0-10% (right)

- [4] 2016  $D^0$  meson nuclear modification factor in PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV Tech. Rep. CMS-PAS-HIN-16-001 CERN Geneva URL <http://cds.cern.ch/record/2157844>
- [5] Abalev B e a (ALICE Collaboration) 2012 *JHEP* **09** 112 (*Preprint arXiv:nucl-ex/1203.2160*)
- [6] Cacciari M, Greco M and Nason P 1998 *Journal of High Energy Physics* **1998** 007 URL <http://stacks.iop.org/1126-6708/1998/i=05/a=007>
- [7] Abelev B *et al.* (ALICE Collaboration) 2012 *JHEP* **1201** 128 (*Preprint arXiv:hep-ex/1111.1553*)
- [8] Averbeck R, Bastid N, del Valle Z C, Crochet P, Dainese A and Zhang X 2011 (*Preprint arXiv:hep-ph/1107.3243*)
- [9] Xu J, Liao J and Gyulassy M 2016 *JHEP* **02** 169 (*Preprint arXiv:hep-ph/1508.00552*)
- [10] Xu J, Buzzatti A and Gyulassy M 2014 *JHEP* **08** 063 (*Preprint arXiv:hep-ph/1402.2956*)
- [11] Jiechen X, Jinfeng L and Gyulassy M 2015 *Chinese Physics Letters* **32** 092501 URL <http://stacks.iop.org/0256-307X/32/i=9/a=092501>
- [12] Djordjevic M and Djordjevic M 2015 *Phys. Rev. C* **92**(2) 024918 URL <http://link.aps.org/doi/10.1103/PhysRevC.92.024918>
- [13] Cao S, Luo T, Qin G Y and Wang X N (*Preprint arXiv:nucl-th/1605.06447v1*)
- [14] Song T, Berrehrah H, Cabrera D, Torres-Rincon J M, Tolos L, Cassing W and Bratkovskaya E 2015 *Phys. Rev. C* **92**(1) 014910 URL <http://link.aps.org/doi/10.1103/PhysRevC.92.014910>
- [15] Song T, Berrehrah H, Cabrera D, Cassing W and Bratkovskaya E 2016 *Phys. Rev. C* **93**(3) 034906 URL <http://link.aps.org/doi/10.1103/PhysRevC.93.034906>
- [16] Kang Z B, Lashof-Regas R, Ovanesyan G, Saad P and Vitev I 2015 *Phys. Rev. Lett.* **114**(9) 092002 URL <http://link.aps.org/doi/10.1103/PhysRevLett.114.092002>
- [17] Chien Y T, Emerman A, Kang Z B, Ovanesyan G and Vitev I 2016 *Phys. Rev. D* **93**(7) 074030 URL <http://link.aps.org/doi/10.1103/PhysRevD.93.074030>
- [18] 2016 Measurement of the charged particle nuclear modification factor in PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV Tech. Rep. CMS-PAS-HIN-15-015 CERN Geneva URL <https://cds.cern.ch/record/2155301>