

Single-Transverse-Spin-Asymmetry studies with a fixed-target experiment using the LHC beams (AFTER@LHC)

J.P. Lansberg^{1*}, M. Anselmino², R. Arnaldi², S.J. Brodsky³, V. Chambert¹, C. Da Silva⁴, J.P. Didelez¹, M.G. Echevarria⁵, E.G. Ferreira⁶, F. Fleuret⁷, Y. Gao⁸, B. Genolini¹, C. Hadjidakis¹, I. Hřivnáčová¹, D. Kikola⁹, A. Klein⁴, A. Kurepin¹⁰, A. Kusina¹¹, C. Lorcé¹², F. Lyonnet¹³, L. Massacrier¹, A. Nass¹⁴, C. Pisano¹⁵, P. Robbe¹⁶, I. Schienbein¹¹, M. Schlegel¹⁷, E. Scomparin², J. Seixas¹⁸, H.S. Shao¹⁹, A. Signori²⁰, E. Steffens²¹, N. Topilskaya¹⁰, B. Trzeciak²², U.I. Uggerhøj²³, A. Uras²⁴, R. Ulrich²⁵, and Z. Yang⁸.

¹ IPNO, Univ. Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, Orsay, France

² Dip. di Fisica and INFN Sez. Torino, Via P. Giuria 1, Torino, Italy

³ SLAC National Accelerator Laboratory, Stanford University, Menlo Park, USA

⁴ LANL, P-25, Los Alamos National Laboratory, Los Alamos, NM 87545, USA

⁵ ECM, Universitat de Barcelona, Barcelona, Spain

⁶ Dept. de Física de Partícules, USC, Santiago de Compostella, Spain

⁷ LLR, École Polytechnique, CNRS/IN2P3, Palaiseau, France

⁸ CHEP, Department of Engineering Physics, Tsinghua University, Beijing, China

⁹ Faculty of Physics, Warsaw University of Technology, Warsaw, Poland

¹⁰ Institute for Nuclear Research, Russian Academy of Sciences, Moscow, Russia

¹¹ LPSC, Université Grenoble-Alpes, CNRS/IN2P3, 38026 Grenoble, France

¹² CPhT, Ecole Polytechnique, CNRS, Université Paris-Saclay, Palaiseau, France

¹³ Southern Methodist University, Dallas, TX 75275, USA

¹⁴ Institut für Kernphysik, Forschungszentrum Jülich, Jülich, Germany

¹⁵ Dipartimento di Fisica, Università degli Studi di Pavia, Pavia, Italy

¹⁶ LAL, Univ. Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, Orsay, France

¹⁷ Institute for Theoretical Physics, Tübingen U., Tübingen, Germany

¹⁸ LIP and IST, Lisbon, Portugal

¹⁹ Theoretical Physics Department, CERN, CH-1211 Geneva 23, Switzerland

²⁰ Nikhef and Dept. of Physics and Astronomy, VU Amsterdam, Amsterdam, The Netherlands

²¹ Physics Institute, Friedrich-Alexander University Erlangen-Nürnberg, Erlangen, Germany

²² Institute for Subatomic Physics, Utrecht University, Utrecht, The Netherlands

²³ Department of Physics and Astronomy, University of Aarhus, Denmark

²⁴ IPNL, Université Claude Bernard Lyon-I, CNRS/IN2P3, Villeurbanne, France

²⁵ Institut für Kernphysik, Karlsruhe Institute of Technology (KIT), Karlsruhe, Germany

We discuss the potential of AFTER@LHC to measure single-transverse-spin asymmetries in open-charm and bottomonium production. With a HERMES-like hydrogen polarised target, such measurements over a year can reach precisions close to the per cent level. This is particularly remarkable since these analyses can probably not be carried out anywhere else.

*XXIV International Workshop on Deep-Inelastic Scattering and Related Subjects
11-15 April, 2016, DESY Hamburg, Germany*

*Speaker. Email: Jean-Philippe.Lansberg@in2p3.fr

1. Introduction

Continuous efforts have been made in the last decades to advance our knowledge of the internal structure of the nucleon and, in particular, that of the constituent dynamics, the quarks and the gluons. However, it remains largely unknown with a limited understanding of the proton and neutron spin structure, namely how they bind into a spin- $\frac{1}{2}$ object. There are two types of quark and gluon contributions to the nucleon spin: their spin and their Orbital Angular Momentum (OAM). For a $+\frac{1}{2}$ helicity nucleon, one has $\frac{1}{2} = \frac{1}{2}\Delta\Sigma + \Delta G + \mathcal{L}_g + \mathcal{L}_q$ where $\frac{1}{2}\Delta\Sigma$ refers to the combined spin contribution of quarks and antiquarks, ΔG the gluon spin, and $\mathcal{L}_{q,g}$ the quark and gluon OAM contributions (see *e.g.* [1, 2]).

This equation applies whatever the energy scale we look at the nucleons and this strongly motivates the scale-evolution study of the individual contributions.

The spin crisis of the 80's has evolved into a puzzle, *i.e.* to determine how large the quark and gluon contributions to the nucleon spin are, to disentangle them and eventually to explain them from first principles in QCD. Recent experimental analyses pointed at $\Delta\Sigma$ as low as 0.25 [3] and that ΔG could¹ reach 0.2. There is still ample room –rather, need– for \mathcal{L}_q and \mathcal{L}_g , which have not yet been measured. This highlights how important studies of the transverse motion of quarks and gluons inside the proton are. Indirect² information on the orbital motion of the partons bound inside hadrons can be accessed via Single (Transverse) Spin Asymmetries (S(T)SA) in different hard-scattering processes, in particular with a transversely polarised hadron (see [6, 7] for recent reviews) to probe the Sivers effect [8, 9]³. This effect is naturally connected [10, 11] to the transverse motion of the partons inside the polarised nucleons (see also [12]).

As of today, the Sivers effect can be approached via two dual formalisms [13]. One extends the collinear parton model of Bjorken with quark-gluon or gluon-gluon correlation functions [14, 15, 16]. It is called the Collinear Twist-3 (CT3) formalism. Another, referred to as the Transverse-Momentum Dependent (TMD) factorisation (see *e.g.* [17, 18, 19]), uses a complete tri-dimensional mapping of the parton momentum. It can also deal with all the possible spin-spin and spin-orbit correlations between the hadron and the partons. Both approaches encode the re-scatterings⁴ of the quarks and gluons with the hadron remnants [20, 21, 22] which are believed to generate the STSAs, also referred to A_N . Both formalisms have their preferred range of applicability. As for the CT3 observables, the AFTER@LHC physics case adds up heavy-flavour and bottomonium production, whose STSAs are essentially unknown, to the usual list of A_N studies which includes single hadron [23], Drell Yan (DY) pair [24, 25, 26] or isolated photon [27] production. For TMD observables, one usually looks at processes immune to final-state radiations which allows one to control the transverse momentum of the initial partons by measuring momentum imbalances. In the case of AFTER@LHC, the possible measurements go well beyond DY produc-

¹The latter has only been probed for $x > 0.05$ [4, 5]

²To measure the parton OAM, observables which are sensitive to the parton position and momentum are in principle required. A well known example of such objects are Generalised Parton Distributions (GPDs) accessible in exclusive processes.

³It is connected to left-right asymmetries in the parton distributions with respect to the plane formed by the proton momentum and spin directions

⁴They are accounted by gauge links in the definition of the TMD PDFs and explicitly considered in the hard-scattering coefficient for CT3.

tion and include pseudo-scalar quarkonium, quarkonium-pair and other associated production of colourless particles.

It is noteworthy that the TMD factorisation approach also allows one to investigate the structure of hadrons in a tridimensional momentum space (see [19] and references therein) in a rigorous and systematic way and it is not restricted to the study of STSAs. It provides theoretical tools to directly study not only the transverse-momentum distributions of the partons but also their polarisation in both polarised and unpolarised nucleons. Such studies are in particular related to azimuthal asymmetries in the final state. At AFTER@LHC [28], one can study them, for instance, via pseudo-scalar quarkonium production [29, 30], associated quarkonium production [31, 32] and, of course, DY pair production [33, 34, 35]. We guide the reader to [36, 37] for more details.

With its high luminosity, a highly polarised target and an access towards the large momentum fraction x^\uparrow in the target, AFTER@LHC is probably the best set-up [28, 38] to carry out an inclusive set of A_N and azimuthal asymmetry measurements both to improve existing analyses and to perform studies which are simply impossible elsewhere. Let us recall that nearly nothing is known from the experimental side about the gluon Sivers effect (see e.g. [39] for a recent review). The polarisation of not only hydrogen but also deuterium and helium targets allows for an even more ambitious spin program bearing on the neutron and spin 1 bound states [40].

2. Experimental implementation

Two approaches for a fixed-target experiment at the LHC with a polarised target can be considered: one with a bent-crystal extracted beam on a conventional polarised target, the other with an internal polarised gas target. Both have the virtue of being parasitic for they do not affect the LHC performances. In the following, we will focus on the latter option⁵ which can directly be combined with existing detectors such as LHCb or ALICE. In particular, we will rely on the expected performances of a target system like the one of the HERMES experiment at DESY-HERA [42], as proposed in [43].

Beam	Target	A	Areal density (θ) (cm^{-2})	\mathcal{L} ($\mu\text{b}^{-1}\cdot\text{s}^{-1}$)	$\int \mathcal{L}$ ($\text{fb}^{-1}\cdot\text{y}^{-1}$)
p	H	1	2.5×10^{14}	900	9
p	D	2	3.2×10^{14}	1200	12

Table 1: Expected luminosities with an internal gas-target inspired by the HERMES experiment.

In such a case, the luminosity is given by the product of the particle current I and the areal density of the polarised storage cell target, θ . The density results from many parameters, the flux of the polarised source injected into the target cell and the cell geometry are the most relevant ones [44]. For a LHC compatible set-up [44] and taking the HERMES-target-source flux, one

⁵Such an internal gas-target option is in fact currently used by the LHCb collaboration but as a luminosity monitor [41] (SMOG) initially designed to study the transverse size of the beam. Despite its limited pressure (about 10^{-7} mbar), it acts as an ideal demonstrator of such a solution over extended periods of time, without any interferences on the other LHC experiments. LHCb performed pilot runs with p and Pb beams on a Ne gas target in 2012 and 2013. These pilot runs were followed by longer runs in 2015: p on Ne (12 hours), He (8 hours) and Ar (3 days) as well as Pb on Ar (1 week) and p on Ar (a few hours). As for now, the current gas pressure is limited by the pumping system.

can expect a density of $2.5 \times 10^{14} \text{ cm}^{-2}$ for a 1 m long cell for H at 300 K and, for D, $3.2 \times 10^{14} \text{ cm}^{-2}$. This is well below the densities which might affect the proton beam life time. The resulting luminosities with $I = 3.14 \times 10^{18} p^+ \text{ s}^{-1}$ for the p^+ beam case are shown in Table 1. The limit of such a solution is essentially set by the number of minimum bias collisions by fill, that is the number of proton “consumed” by the target as compared to the total number stored in the LHC.

In terms of the polarisation, by reusing a target like that of HERMES, the effective polarisation of H or D [43] can be as high as 0.8. ^3He can also be used. In the following, we will take an effective polarisation of $P = 0.6$ as a working hypothesis which, from the number above, is definitely a conservative assumption.

3. Selected figures-of-merit for STSAs

In this section, we quickly discuss figures-of-merit for open charm and bottomonium production. Those for Drell-Yan and J/ψ can be found in [38]. They both show that A_N can be measured close to the per cent level in regions where the expected effects can be as large as ten or even twenty per cent.

3.1 STSAs in open (anti)-charm production

Some years ago, it was argued [45] that the study of STSAs in open heavy-flavour production in the RHIC energy domain gives a direct access to the tri-gluon correlations functions appearing in the CT3 approach. Such correlations can be related to the gluon Sivvers functions under some assumptions. If STSAs for charm quarks and anti-quarks can separately be measured, this will also be a unique probe of C -parity odd twist-3 tri-gluon correlators [46, 47].




Figure 1: Projected P_T dependence of A_N for D^0 at $y_{c.m.s.} = 0$ ($y_{c.m.s.} = 2.25$) on the left (middle).

Fig. 1 (left & middle) shows the projected statistical precision⁶ for D^0 meson STSAs at AFTER@LHC. It definitely lies at the per cent level. The expected magnitude is from [49, 50]. As for now, such measurements are not planned elsewhere, certainly not in the large x^\uparrow region where the STSAs are expected to be the largest.

⁶In view of very low level of background in various D^0 LHCb analyses, we neglected it here. We have also assumed LHCb-like acceptance and performances, see [48].

3.2 STSAs in bottomonium production

Bottomonia can also provide much information on the Siverson effects in the gluon sector, especially via STSAs analyses of the $\Upsilon(1S)$, $\Upsilon(2S)$, $\Upsilon(3S)$ which contain different amounts of P -wave feed-downs. We believe that STSAs of $\Upsilon(2S)$ and $\Upsilon(3S)$ would not be accessible anywhere else than at AFTER@LHC. Since only one particle is observed, these STSAs are preferably treated in the CT3 approach that was, for instance, applied to the η_c case [51] which could also be measured at AFTER@LHC. As for now, quantitative predictions for these STSAs are lacking even though the qualitative conclusion of [52] should apply and the observation of a non-zero STSA should already provide us with new information both on the quarkonium-production mechanisms [53, 54, 55] and on the gluon Siverson effect [39]. For the figure-of-merit displayed on Fig. 2, we have thus set the central value of the points to $A_N = 0$. It clearly shows that, even for the least produced $\Upsilon(3S)$, the statistical precision is better than 5% taking into account a full background simulation as discussed in [56].




Figure 2: Projected x_F dependence of the statistical uncertainty on A_N for $\Upsilon(1S)$, $\Upsilon(2S)$, $\Upsilon(3S)$ corresponding to 3 bins in rapidity ([2:3],[3:4],[4:5]).

4. Conclusion

The combination of the TeV LHC proton beam, of a polarised internal gas-target system inspired by that of the HERMES experiment and a detector like LHCb (or ALICE) opens the way to a number of outstanding STSA measurements at the per cent level, most of which cannot be carried out at other facilities.

As concluding remarks, we would like to emphasise two important facts: (i) at AFTER@LHC, these STSAs would naturally be measured at large x^\uparrow where the (quark and gluon) Siverson effect is expected to be the largest and (ii) such measurements have to be measured in pp collisions as mandatory complementary pieces of information to similar studies in lepton-induced reactions to perform quantitative tests of the generalised universality of the TMD-related observables, deeply connected to QCD.

Acknowledgements

This research was supported in part by the French P2IO Excellence Laboratory, the French

CNRS via the grants FCPPL-Quarkonium4AFTER & Défi Inphyniti–Théorie LHC France and by the Department of Energy, contract DE–AC02–76SF00515.

References

- [1] E. Leader and C. Lorce, “The angular momentum controversy: What’s it all about and does it matter?,” *Phys. Rept.* **541** (2014) 163–248, [arXiv:1309.4235 \[hep-ph\]](#).
- [2] M. Wakamatsu, “Is gauge-invariant complete decomposition of the nucleon spin possible?,” *Int. J. Mod. Phys.* **A29** (2014) 1430012, [arXiv:1402.4193 \[hep-ph\]](#).
- [3] D. de Florian, R. Sassot, M. Stratmann, and W. Vogelsang, “Global Analysis of Helicity Parton Densities and Their Uncertainties,” *Phys. Rev. Lett.* **101** (2008) 072001, [arXiv:0804.0422 \[hep-ph\]](#).
- [4] STAR Collaboration, L. Adamczyk *et al.*, “Precision Measurement of the Longitudinal Double-spin Asymmetry for Inclusive Jet Production in Polarized Proton Collisions at $\sqrt{s} = 200$ GeV,” *Phys. Rev. Lett.* **115** (2015) 092002, [arXiv:1405.5134 \[hep-ex\]](#).
- [5] D. de Florian, R. Sassot, M. Stratmann, and W. Vogelsang, “Evidence for polarization of gluons in the proton,” *Phys. Rev. Lett.* **113** (2014) 012001, [arXiv:1404.4293 \[hep-ph\]](#).
- [6] U. D’Alesio and F. Murgia, “Azimuthal and Single Spin Asymmetries in Hard Scattering Processes,” *Prog. Part. Nucl. Phys.* **61** (2008) 394–454, [arXiv:0712.4328 \[hep-ph\]](#).
- [7] V. Barone, F. Bradamante, and A. Martin, “Transverse-spin and transverse-momentum effects in high-energy processes,” *Prog. Part. Nucl. Phys.* **65** (2010) 267–333, [arXiv:1011.0909 \[hep-ph\]](#).
- [8] D. W. Sivers, “Single Spin Production Asymmetries from the Hard Scattering of Point-Like Constituents,” *Phys. Rev.* **D41** (1990) 83.
- [9] D. W. Sivers, “Hard scattering scaling laws for single spin production asymmetries,” *Phys. Rev.* **D43** (1991) 261–263.
- [10] M. Burkardt, “Impact parameter dependent parton distributions and transverse single spin asymmetries,” *Phys. Rev.* **D66** (2002) 114005, [arXiv:hep-ph/0209179 \[hep-ph\]](#).
- [11] M. Burkardt, “Chromodynamic lensing and transverse single spin asymmetries,” *Nucl. Phys.* **A735** (2004) 185–199, [arXiv:hep-ph/0302144 \[hep-ph\]](#).
- [12] K.-F. Liu and C. Lorce, “The Parton Orbital Angular Momentum: Status and Prospects,” *Eur. Phys. J.* **A52** (2016) 160, [arXiv:1508.00911 \[hep-ph\]](#).
- [13] Y. Koike, W. Vogelsang, and F. Yuan, “On the Relation Between Mechanisms for Single-Transverse-Spin Asymmetries,” *Phys. Lett.* **B659** (2008) 878–884, [arXiv:0711.0636 \[hep-ph\]](#).
- [14] A. V. Efremov and O. V. Teryaev, “On Spin Effects in Quantum Chromodynamics,” *Sov. J. Nucl. Phys.* **36** (1982) 140. [*Yad. Fiz.*36,242(1982)].
- [15] A. V. Efremov and O. V. Teryaev, “QCD Asymmetry and Polarized Hadron Structure Functions,” *Phys. Lett.* **B150** (1985) 383.
- [16] J.-w. Qiu and G. F. Sterman, “Single transverse spin asymmetries,” *Phys. Rev. Lett.* **67** (1991) 2264–2267.
- [17] J. Collins, *Foundations of perturbative QCD*. Cambridge University Press, 2013.
- [18] M. G. Echevarria, A. Idilbi, and I. Scimemi, “Factorization Theorem For Drell-Yan At Low q_T And Transverse Momentum Distributions On-The-Light-Cone,” *JHEP* **07** (2012) 002, [arXiv:1111.4996 \[hep-ph\]](#).
- [19] R. Angeles-Martinez *et al.*, “Transverse Momentum Dependent (TMD) parton distribution functions: status and prospects,” *Acta Phys. Polon.* **B46** (2015) 2501–2534, [arXiv:1507.05267 \[hep-ph\]](#).
- [20] S. J. Brodsky, D. S. Hwang, and I. Schmidt, “Initial state interactions and single spin asymmetries in Drell-Yan processes,” *Nucl. Phys.* **B642** (2002) 344–356, [arXiv:hep-ph/0206259 \[hep-ph\]](#).

- [21] J. C. Collins, “Leading twist single transverse-spin asymmetries: Drell-Yan and deep inelastic scattering,” *Phys. Lett.* **B536** (2002) 43–48, [arXiv:hep-ph/0204004](#) [hep-ph].
- [22] S. J. Brodsky, D. S. Hwang, and I. Schmidt, “Final state interactions and single spin asymmetries in semiinclusive deep inelastic scattering,” *Phys. Lett.* **B530** (2002) 99–107, [arXiv:hep-ph/0201296](#) [hep-ph].
- [23] J.-w. Qiu and G. F. Sterman, “Single transverse spin asymmetries in hadronic pion production,” *Phys. Rev.* **D59** (1999) 014004, [arXiv:hep-ph/9806356](#) [hep-ph].
- [24] N. Hammon, O. Teryaev, and A. Schafer, “Single spin asymmetry for the Drell-Yan process,” *Phys. Lett.* **B390** (1997) 409–412, [arXiv:hep-ph/9611359](#) [hep-ph].
- [25] D. Boer, P. J. Mulders, and O. V. Teryaev, “Single spin asymmetries from a gluonic background in the Drell-Yan process,” *Phys. Rev.* **D57** (1998) 3057–3064, [arXiv:hep-ph/9710223](#) [hep-ph].
- [26] D. Boer and P. J. Mulders, “Color gauge invariance in the Drell-Yan process,” *Nucl. Phys.* **B569** (2000) 505–526, [arXiv:hep-ph/9906223](#) [hep-ph].
- [27] J.-w. Qiu and G. F. Sterman, “Single transverse spin asymmetries in direct photon production,” *Nucl. Phys.* **B378** (1992) 52–78.
- [28] S. J. Brodsky, F. Fleuret, C. Hadjidakis, and J. P. Lansberg, “Physics Opportunities of a Fixed-Target Experiment using the LHC Beams,” *Phys. Rept.* **522** (2013) 239–255, [arXiv:1202.6585](#) [hep-ph].
- [29] D. Boer and C. Pisano, “Polarized gluon studies with charmonium and bottomonium at LHCb and AFTER,” *Phys. Rev.* **D86** (2012) 094007, [arXiv:1208.3642](#) [hep-ph].
- [30] A. Signori, “Gluon TMDs in quarkonium production,” *Few Body Syst.* **57** (2016) 651–655, [arXiv:1602.03405](#) [hep-ph].
- [31] W. J. den Dunnen, J. P. Lansberg, C. Pisano, and M. Schlegel, “Accessing the Transverse Dynamics and Polarization of Gluons inside the Proton at the LHC,” *Phys. Rev. Lett.* **112** (2014) 212001, [arXiv:1401.7611](#) [hep-ph].
- [32] J.-P. Lansberg and H.-S. Shao, “Double-quarkonium production at a fixed-target experiment at the LHC (AFTER@LHC),” *Nucl. Phys.* **B900** (2015) 273–294, [arXiv:1504.06531](#) [hep-ph].
- [33] T. Liu and B.-Q. Ma, “Azimuthal asymmetries in lepton-pair production at a fixed-target experiment using the LHC beams (AFTER),” *Eur. Phys. J.* **C72** (2012) 2037, [arXiv:1203.5579](#) [hep-ph].
- [34] M. Anselmino, U. D’Alesio, and S. Melis, “Transverse single-spin asymmetries in proton-proton collisions at the AFTER@LHC experiment in a TMD factorisation scheme,” *Adv. High Energy Phys.* **2015** (2015) 475040, [arXiv:1504.03791](#) [hep-ph].
- [35] K. Kanazawa, Y. Koike, A. Metz, and D. Pitonyak, “Transverse single-spin asymmetries in proton-proton collisions at the AFTER@LHC experiment,” *Adv. High Energy Phys.* **2015** (2015) 257934, [arXiv:1502.04021](#) [hep-ph].
- [36] J. P. Lansberg *et al.*, “Spin physics and TMD studies at A Fixed-Target Experiment at the LHC (AFTER@LHC),” *EPJ Web Conf.* **85** (2015) 02038, [arXiv:1410.1962](#) [hep-ex].
- [37] L. Massacrier *et al.*, “Studies of Transverse-Momentum-Dependent distributions with A Fixed-Target Experiment using the LHC beams (AFTER@LHC),” *Int. J. Mod. Phys. Conf. Ser.* **40** (2016) 1660107, [arXiv:1502.00984](#) [nucl-ex].
- [38] J.-P. Lansberg *et al.*, “Physics case for a polarised target for AFTER@LHC,” *PoS PSTP2015* (2016) 042, [arXiv:1602.06857](#) [nucl-ex].
- [39] D. Boer, C. Lorce, C. Pisano, and J. Zhou, “The gluon Sivers distribution: status and future prospects,” *Adv. High Energy Phys.* **2015** (2015) 371396, [arXiv:1504.04332](#) [hep-ph].
- [40] D. Boer, S. Cotogno, T. van Daal, P. J. Mulders, A. Signori, and Y.-J. Zhou, “Gluon and Wilson loop TMDs for hadrons of spin ≤ 1 ,” [arXiv:1607.01654](#) [hep-ph].
- [41] M. Ferro-Luzzi, “Proposal for an absolute luminosity determination in colliding beam experiments using vertex detection of beam-gas interactions,” *Nucl. Instrum. Meth.* **A553** (2005) 388–399.
- [42] HERMES Collaboration, A. Airapetian *et al.*, “The HERMES polarized hydrogen and deuterium gas

- target in the HERA electron storage ring,” *Nucl. Instrum. Meth.* **A540** (2005) 68–101, [arXiv:physics/0408137 \[physics\]](#).
- [43] C. Barschel, P. Lenisa, A. Nass, and E. Steffens, “A Gas Target Internal to the LHC for the Study of pp Single-Spin Asymmetries and Heavy Ion Collisions,” *Adv. High Energy Phys.* **2015** (2015) 463141.
- [44] E. Steffens, “Estimation of the performance of a HERMES-type gas target internal to the LHC,” *PoS PSTP2015* (2015) 019.
- [45] Z.-B. Kang, J.-W. Qiu, W. Vogelsang, and F. Yuan, “Accessing tri-gluon correlations in the nucleon via the single spin asymmetry in open charm production,” *Phys. Rev.* **D78** (2008) 114013, [arXiv:0810.3333 \[hep-ph\]](#).
- [46] X.-D. Ji, “Gluon correlations in the transversely polarized nucleon,” *Phys. Lett.* **B289** (1992) 137–142.
- [47] H. Beppu, Y. Koike, K. Tanaka, and S. Yoshida, “Contribution of Twist-3 Multi-Gluon Correlation Functions to Single Spin Asymmetry in Semi-Inclusive Deep Inelastic Scattering,” *Phys. Rev.* **D82** (2010) 054005, [arXiv:1007.2034 \[hep-ph\]](#).
- [48] D. Kikola, “Prospects for Open Heavy Flavor Measurements in Heavy Ion and p + A Collisions in a Fixed-Target Experiment at the LHC,” *Adv. High Energy Phys.* **2015** (2015) 783134.
- [49] M. Anselmino, M. Boglione, U. D’Alesio, E. Leader, and F. Murgia, “Accessing Sivers gluon distribution via transverse single spin asymmetries in p(transv. polarized) p \rightarrow D X processes at RHIC,” *Phys. Rev.* **D70** (2004) 074025, [arXiv:hep-ph/0407100 \[hep-ph\]](#).
- [50] U. D’Alesio, F. Murgia, and C. Pisano, “Towards a first estimate of the gluon Sivers function from A_N data in pp collisions at RHIC,” *JHEP* **09** (2015) 119, [arXiv:1506.03078 \[hep-ph\]](#).
- [51] A. Schafer and J. Zhou, “Transverse single spin asymmetry in hadronic $\eta_{c,b}$ production,” *Phys. Rev.* **D88** (2013) 014008, [arXiv:1302.4600 \[hep-ph\]](#).
- [52] F. Yuan, “Heavy Quarkonium Production in Single Transverse Polarized High Energy Scattering,” *Phys. Rev.* **D78** (2008) 014024, [arXiv:0801.4357 \[hep-ph\]](#).
- [53] A. Andronic *et al.*, “Heavy-flavour and quarkonium production in the LHC era: from proton-proton to heavy-ion collisions,” *Eur. Phys. J.* **C76** (2016) 107, [arXiv:1506.03981 \[nucl-ex\]](#).
- [54] N. Brambilla *et al.*, “Heavy quarkonium: progress, puzzles, and opportunities,” *Eur. Phys. J.* **C71** (2011) 1534, [arXiv:1010.5827 \[hep-ph\]](#).
- [55] J. P. Lansberg, “ J/ψ , ψ' and Υ production at hadron colliders: A Review,” *Int. J. Mod. Phys.* **A21** (2006) 3857–3916, [arXiv:hep-ph/0602091 \[hep-ph\]](#).
- [56] L. Massacrier, B. Trzeciak, F. Fleuret, C. Hadjidakis, D. Kikola, J. P. Lansberg, and H.-S. Shao, “Feasibility studies for quarkonium production at a fixed-target experiment using the LHC proton and lead beams (AFTER@LHC),” *Adv. High Energy Phys.* **2015** (2015) 986348, [arXiv:1504.05145 \[hep-ex\]](#).