



Observation of the decays $\Lambda_b^0 \rightarrow \chi_{c1} p K^-$ and $\Lambda_b^0 \rightarrow \chi_{c2} p K^-$

The LHCb collaboration[†]

Abstract

The first observation of the decays $\Lambda_b^0 \rightarrow \chi_{c1} p K^-$ and $\Lambda_b^0 \rightarrow \chi_{c2} p K^-$ is reported using a data sample corresponding to an integrated luminosity of 3.0 fb^{-1} , collected by the LHCb experiment in pp collisions at centre-of-mass energies of 7 and 8 TeV. The following ratios of branching fractions are measured

$$\frac{\mathcal{B}(\Lambda_b^0 \rightarrow \chi_{c1} p K^-)}{\mathcal{B}(\Lambda_b^0 \rightarrow J/\psi p K^-)} = 0.242 \pm 0.014 \pm 0.013 \pm 0.009,$$

$$\frac{\mathcal{B}(\Lambda_b^0 \rightarrow \chi_{c2} p K^-)}{\mathcal{B}(\Lambda_b^0 \rightarrow J/\psi p K^-)} = 0.248 \pm 0.020 \pm 0.014 \pm 0.009,$$

$$\frac{\mathcal{B}(\Lambda_b^0 \rightarrow \chi_{c2} p K^-)}{\mathcal{B}(\Lambda_b^0 \rightarrow \chi_{c1} p K^-)} = 1.02 \pm 0.10 \pm 0.02 \pm 0.05,$$

where the first uncertainty is statistical, the second systematic and the third due to the uncertainty on the branching fractions of the $\chi_{c1} \rightarrow J/\psi \gamma$ and $\chi_{c2} \rightarrow J/\psi \gamma$ decays. Using both decay modes, the mass of the Λ_b^0 baryon is also measured to be

$$m_{\Lambda_b^0} = 5619.44 \pm 0.28 \pm 0.26 \text{ MeV}/c^2,$$

where the first and second uncertainties are statistical and systematic, respectively.

Published as Phys. Rev. Lett. 119, 062001 (2017)

© CERN on behalf of the LHCb collaboration, licence CC-BY-4.0.

[†]Authors are listed at the end of this paper.

Since the birth of the quark model, it has been speculated that hadrons could be formed from multi-quark states beyond the well-studied quark-antiquark (meson) and three-quark (baryon) combinations [1–3]. Using a six-dimensional amplitude analysis of the $\Lambda_b^0 \rightarrow J/\psi p K^-$ decay mode, the LHCb collaboration observed the $P_c(4380)^+$ and $P_c(4450)^+$ states [4,5], which are consistent with $uudc\bar{c}$ hidden-charm pentaquarks decaying to $J/\psi p$. Many phenomenological models describing the dynamics of these states have been proposed, including meson-baryon molecules [6–8], compact pentaquarks [9–11] and kinematic rescattering effects [12–16]. In particular, the authors of Ref. [12] noted the closeness of the $P_c(4450)^+$ mass to the $\chi_{c1}p$ threshold and proposed that, if the $P_c(4450)^+$ state is a rescattering effect, then it would not appear as an enhancement near the $\chi_{c1}p$ threshold in the $\Lambda_b^0 \rightarrow \chi_{c1}p K^-$ decay mode, an approach recently challenged in Ref. [17]. This Letter presents an initial stage in the investigation of this hypothesis by making the first observation of $\Lambda_b^0 \rightarrow \chi_{c1}p K^-$ and $\Lambda_b^0 \rightarrow \chi_{c2}p K^-$ decays and measurements of their branching fractions relative to the $\Lambda_b^0 \rightarrow J/\psi p K^-$ decay. Throughout this Letter, the inclusion of charge-conjugated processes is implied and the symbol χ_{cJ} is used to denote the χ_{c1} and χ_{c2} states collectively. All Λ_b^0 decay modes considered here proceed via the same quark-level process, whose dominant contribution is shown in Fig. 1. A measurement of the Λ_b^0 baryon mass is also presented.

Previous measurements of the branching fractions of $B \rightarrow \chi_{cJ}K$ decays [18–20] have shown that the χ_{c2} mode is suppressed relative to the χ_{c1} mode, in agreement with the predictions from the factorisation approach [21], although the suppression appears to be lessened when additional particles are present in the final state [22]. Studying the production of χ_{cJ} mesons in Λ_b^0 baryon decays will help to further test the factorisation approach, as the additional spectator quark in the baryon decay may play an important role in modifying final-state interactions.

The measurements described in this Letter are based on a data sample corresponding to 1.0 fb^{-1} of integrated luminosity collected by the LHCb experiment in pp collisions at a centre-of-mass energy of 7 TeV in 2011, and 2.0 fb^{-1} at 8 TeV in 2012. The LHCb detector [23,24] is a single-arm forward spectrometer covering the pseudorapidity range $2 < \eta < 5$, designed for the study of particles containing b or c quarks. The detector includes a high-precision tracking system consisting of a silicon-strip vertex detector surrounding the pp interaction region, a large-area silicon-strip detector located upstream of a dipole magnet with a bending power of about 4 Tm, and three stations of silicon-strip detectors and straw drift tubes placed downstream of the magnet. The tracking system provides a measurement of momentum, p , of charged particles with a relative




Figure 1: Feynman diagram of $\Lambda_b^0 \rightarrow J/\psi \Lambda^*$ and $\Lambda_b^0 \rightarrow \chi_{cJ} \Lambda^*$ decays, where Λ^* refers to an excited Λ baryon.

uncertainty that varies from 0.5 % at low momentum to 1.0 % at 200 GeV/ c . The minimum distance of a track to a primary vertex (PV), the impact parameter (IP), is measured with a resolution of $(15 + 29/p_T)$ μm , where p_T is the component of the momentum transverse to the beam, in GeV/ c . Different types of charged hadrons are distinguished using information from two ring-imaging Cherenkov detectors. Photons, electrons and hadrons are identified by a calorimeter system consisting of scintillating-pad and preshower detectors, an electromagnetic calorimeter and a hadronic calorimeter. Muons are identified by a system composed of alternating layers of iron and multiwire proportional chambers. The online event selection is performed by a trigger, which consists of a hardware stage, based on information from the calorimeter and muon systems, followed by a software stage, which applies a full event reconstruction. The software trigger selects events that contain a pair of oppositely charged muons that form a vertex that is significantly separated from all PVs.

In the simulation, pp collisions are generated using PYTHIA 8 [25] with a specific LHCb configuration [26]. Decays of hadronic particles are described by EVTGEN [27], in which final-state radiation is generated using PHOTOS [28]. The interaction of the generated particles with the detector, and its response, are implemented using the GEANT4 toolkit [29] as described in Ref. [30]. The products of the Λ_b^0 decays are generated uniformly within the available phase space.

The $\Lambda_b^0 \rightarrow \chi_{cJ} p K^-$ and $\Lambda_b^0 \rightarrow J/\psi p K^-$ candidates are reconstructed via the decays $\chi_{cJ} \rightarrow J/\psi \gamma$ and $J/\psi \rightarrow \mu^+ \mu^-$. To separate signal from background, an offline selection is applied after the trigger, consisting of a loose preselection followed by a multivariate classifier based on a gradient-boosted decision tree (GBDT) [31].

The J/ψ candidates are formed from two oppositely charged particles with $p_T > 550$ MeV/ c , identified as muons and consistent with originating from a common vertex, but inconsistent with originating from any PV. The invariant mass of the $\mu^+ \mu^-$ pair is required to be in the range [3000, 3170] MeV/ c^2 . The χ_{cJ} candidates are formed from a J/ψ candidate and a photon with $p_T > 700$ MeV/ c . Photons that are consistent with originating from a π^0 meson when combined with any other photon in the event are removed. The invariant mass of the $\mu^+ \mu^- \gamma$ combination is required to be in the range [3400, 3700] MeV/ c^2 . In the following, the notation $[c\bar{c}]$ will be used to refer to the initial J/ψ or χ_{cJ} candidate from the Λ_b^0 baryon decay, while the notation $m(J/\psi X)$ or $m(\chi_{cJ} X)$ denotes an invariant mass that has been calculated with a mass constraint applied to the J/ψ or χ_{cJ} candidate.

The Λ_b^0 candidates are formed from a $[c\bar{c}]$ candidate and two good-quality oppositely charged tracks each with $p_T > 200$ MeV/ c , identified as a proton and kaon. Both tracks are required to be significantly displaced from any PV. A kinematic fit [32] is applied to the Λ_b^0 candidate, with the J/ψ and χ_{c1} masses constrained to their known values [33], and the Λ_b^0 candidate constrained to point back to a PV. This has the effect of producing separated peaks for the two decay modes. The mass resolution for $\Lambda_b^0 \rightarrow \chi_{c2} p K^-$ decays is lower compared to that for $\Lambda_b^0 \rightarrow \chi_{c1} p K^-$ decays due to the wrong mass hypothesis of the $[c\bar{c}]$ candidate.

Contributions from $B^0 \rightarrow [c\bar{c}] K^+ \pi^-$ ($B_s^0 \rightarrow [c\bar{c}] \phi$, $\phi \rightarrow K^+ K^-$) decays, where the π^- (K^-) is misidentified as an antiproton, are suppressed by placing tighter particle identification requirements on the misidentified hadron for candidates with an invariant mass within 30 MeV/ c^2 of the B^0 (B_s^0) mass [33] when evaluated using π^- or K^- mass assignments for the antiproton candidates. Typical mis-identification probabilities are

5 – 15 %, dependent on the particle momentum [34]. In the $\Lambda_b^0 \rightarrow \chi_{cJ} p K^-$ samples, small contributions from $B^0 \rightarrow J/\psi K^+ \pi^-$ or $B_s^0 \rightarrow J/\psi \phi$ decays, where in addition to the misidentification the J/ψ meson is combined with a random photon in the event, are removed by the requirement that $m(J/\psi K^+ \pi^-)$ or $m(J/\psi K^+ K^-)$ are within $30 \text{ MeV}/c^2$ of the B^0 or B_s^0 mass. Additionally, $\phi \rightarrow K^+ K^-$ decays are vetoed by removing all candidates where the invariant mass of the $p K^-$ combination is within $12 \text{ MeV}/c^2$ of the known ϕ meson mass when the kaon mass is used instead of the proton mass. Further misreconstructed backgrounds are studied using a fast simulation package [35] and are found to have no peaking components in the invariant mass window of interest.

The GBDT is used to further suppress the combinatorial background. It is trained on a simulated sample of $\Lambda_b^0 \rightarrow \chi_{c1} p K^-$ decays for the signal and candidates from data with $m(\chi_{c1} p K^-)$ in the range $[5700, 5800] \text{ MeV}/c^2$ for the background. Twelve variables are used as input. The first of these is the χ^2 value obtained from a kinematic fit with the Λ_b^0 candidate constrained to point back to a PV and a mass constraint applied to the J/ψ . In addition, for the signal mode, χ_{c1} or χ_{c2} mass constraints are applied, with the smaller χ^2 values being used; note that this differs from the fit used to separate the mass peaks, which does not include a χ_{c2} constraint. The remaining variables are the p_T of the Λ_b^0 , proton and kaon; the Λ_b^0 decay-length significance; the cosine of the angle between the momentum of the Λ_b^0 candidate and its displacement from the PV; the proton and kaon χ_{IP}^2 , defined as the difference in the vertex-fit χ^2 of the PV when reconstructed with and without the considered particle; and the estimated probabilities that the two muons, kaon and proton are correctly identified by the particle identification detectors.

Prior to the training, several modifications are made to the simulated samples to better match the kinematic distributions observed in data. First, the simulated $\Lambda_b^0 \rightarrow J/\psi p K^-$ events are weighted according to the six-dimensional amplitude model developed in Ref. [4]. Second, a multidimensional gradient-boosting algorithm [36] is used to weight the simulated $\Lambda_b^0 \rightarrow J/\psi p K^-$ decays such that the distributions of Λ_b^0 pseudorapidity, the number of tracks in the event and the GBDT training variables (apart from those related to particle identification) match those observed in the preselected background-subtracted $\Lambda_b^0 \rightarrow J/\psi p K^-$ data sample. These weights are also applied to the simulated $\Lambda_b^0 \rightarrow \chi_{cJ} p K^-$ samples. Finally, the simulated distributions of the particle identification variables for the muon, proton and kaon candidates are resampled from data calibration samples ($D^{*+} \rightarrow D \pi^+$, $J/\psi \rightarrow \mu^+ \mu^-$, $\Lambda \rightarrow p \pi^-$ and $\Lambda_c^+ \rightarrow p K^- \pi^+$ decays) in bins of track p , p_T and the number of tracks.

The optimal working point for the GBDT response is chosen by maximising a figure of merit, $S/\sqrt{S+B}$, where $S = S_0 \epsilon$ and B are the expected signal and background yields within $\pm 20 \text{ MeV}/c^2$ of the known Λ_b^0 baryon mass [33], S_0 is the signal yield determined from data without any cut on the GBDT response, and ϵ is the relative efficiency of the GBDT selection, evaluated using the simulated sample. The Λ_b^0 mass sidebands from the data are used to estimate B . The same GBDT and working point are used for the $\Lambda_b^0 \rightarrow J/\psi p K^-$ normalisation mode. The GBDT selection efficiencies are 78 %, 75 % and 68 % for the $\Lambda_b^0 \rightarrow \chi_{c1} p K^-$, $\Lambda_b^0 \rightarrow \chi_{c2} p K^-$ and $\Lambda_b^0 \rightarrow J/\psi p K^-$ channels, respectively.

After applying the GBDT requirement, $(2.9 \pm 0.4) \%$ of the selected events contain multiple $\Lambda_b^0 \rightarrow \chi_{cJ} p K^-$ candidates. In approximately 80 % of these cases, the same $J/\psi p K^-$ combination is combined with an additional, unrelated, photon in the event. The results reported in this Letter retain all candidates and the reported branching fractions are corrected to account for this. The correction factor is 0.993 ± 0.006 (0.986 ± 0.009)




Figure 2: Fits to the (a) $\Lambda_b^0 \rightarrow \chi_{c1} p K^-$ and (b) $\Lambda_b^0 \rightarrow J/\psi p K^-$ invariant mass distributions. Data points are shown in black and the results of the fits are shown as solid blue lines. The components are $\Lambda_b^0 \rightarrow \chi_{c1} p K^-$ and $\Lambda_b^0 \rightarrow \chi_{c2} p K^-$ signal and combinatorial background (Comb.).

for $\Lambda_b^0 \rightarrow \chi_{c1} p K^-$ ($\Lambda_b^0 \rightarrow \chi_{c2} p K^-$) decays, which is evaluated using a combination of the simulated samples and pseudoexperiments. The larger width of the $\Lambda_b^0 \rightarrow \chi_{c2} p K^-$ component leads to the larger uncertainty on the correction factor. For the selected $\Lambda_b^0 \rightarrow J/\psi p K^-$ sample, $(0.75 \pm 0.05)\%$ of the events have multiple candidates.

Extended unbinned maximum-likelihood fits are performed to the distributions of $m(\chi_{c1} p K^-)$ and $m(J/\psi p K^-)$ for the signal and normalisation modes, respectively. The fit models consist of signal components, each described by the sum of two Crystal Ball (CB) functions [37] with a common mean and power-law tails on both sides, and a linear combinatorial background component. Due to the small $\chi_{c0} \rightarrow J/\psi \gamma$ branching fraction [33] the contribution from the χ_{c0} mode is negligible. Several parameters of the signal shapes are determined from fits to simulated samples. These include the tail parameters of the CB functions, the ratio of the widths of the two CB functions, and their relative normalisations. The $\Lambda_b^0 \rightarrow \chi_{c2} p K^-$ signal component is shifted to a lower mass in $m(\chi_{c1} p K^-)$ due to the χ_{c1} mass constraint. The signal and background yields, the gradient of the background shape and the mean of each signal component are free parameters in the fit to data. In addition, the widths of the χ_{c1} and χ_{c2} components in the fit to $m(\chi_{c1} p K^-)$ are allowed to differ from simulation by a common scaling factor, while the width of the narrower CB function in the $\Lambda_b^0 \rightarrow J/\psi p K^-$ signal component is a free parameter in the fit to data. The results of these fits are shown in Fig. 2. The measured yields are 453 ± 25 , 285 ± 23 and $29\,815 \pm 178$ for the χ_{c1} , χ_{c2} and J/ψ modes, respectively. The significance of each of the signal components in the fit to $m(\chi_{c1} p K^-)$ is calculated using Wilks' theorem [38]. This gives statistical significances of 29 and 17 standard deviations for the decay modes with χ_{c1} and χ_{c2} , respectively.

Simulated samples are used to determine, for each decay mode, the reconstruction and selection efficiency as a function of the Dalitz plot coordinates [39], $m^2(p K^-)$ and $m^2([c\bar{c}]p)$. This approach focuses on the dimensions where the efficiency variation is expected to be largest whilst averaging over other dimensions in the phase space of $\Lambda_b^0 \rightarrow [c\bar{c}] p K^-$ decays. This assumption is treated as a source of systematic uncertainty. The efficiency of each mode, $\epsilon_{[c\bar{c}]}$, is due to geometric acceptance, reconstruction and selection including the GBDT. The ratios of the phase-space-averaged values of these efficiencies are 0.182 ± 0.005 for $\epsilon_{\chi_{c1}}/\epsilon_{J/\psi}$, and 0.196 ± 0.005 for $\epsilon_{\chi_{c2}}/\epsilon_{J/\psi}$, where the




Figure 3: Distributions of (a) $m(\chi_{c1} p K^-)$ versus $\Delta m + m_{J/\psi}$, where Δm is $m(\mu^+ \mu^- \gamma) - m(\mu^+ \mu^-)$ and $m_{J/\psi}$ is the known mass of the J/ψ meson [33], and (b) $\Delta m + m_{J/\psi}$ for background-subtracted $\Lambda_b^0 \rightarrow \chi_{c1} p K^-$ (green circles) and $\Lambda_b^0 \rightarrow \chi_{c2} p K^-$ (magenta triangles) candidates (cands).

uncertainties are due to the size of the simulated samples. This includes a correction factor for the χ_{cJ} decay modes to account for differences in the photon reconstruction efficiency between data and simulation [20, 40].

An efficiency-corrected, background-subtracted yield is determined for each decay mode as $N^{\text{corr}}([\bar{c}\bar{c}]pK^-) = \sum_i w_i / \epsilon_{[\bar{c}\bar{c}],i}$, where the index i runs over all candidates in the fit range. The w_i are weights determined using the *sPlot* background-subtraction technique [41], which project out the signal component from the combined signal plus background densities using the $[\bar{c}\bar{c}]pK^-$ invariant mass as discriminating variable. The corrected yields are found to be about $99\,700 \pm 5300$, $57\,800 \pm 4400$ and $1\,213\,500 \pm 7300$ for the χ_{c1} , χ_{c2} and J/ψ decay modes, respectively, where the uncertainties are determined from the sum in quadrature of the event weights. Since these weights are determined from a fit in which all shape parameters are fixed, following the *sPlot* prescription, the effect on the yield uncertainty due to the statistical uncertainties on these parameters is not included. To quantify this, the unweighted fit is performed with the shape parameters fixed and then free, and the difference in quadrature of the uncertainties is found to be 1.5% of the yield for $\Lambda_b^0 \rightarrow \chi_{c1} p K^-$ decays and 2.1% for $\Lambda_b^0 \rightarrow \chi_{c2} p K^-$. A corresponding uncertainty is added in quadrature to that on the efficiency-corrected yield. The effect is negligible for the much larger $\Lambda_b^0 \rightarrow J/\psi p K^-$ signal.

The ratios of branching fractions are determined as

$$R_{1(2)} \equiv \frac{\mathcal{B}(\Lambda_b^0 \rightarrow \chi_{c1(2)} p K^-)}{\mathcal{B}(\Lambda_b^0 \rightarrow J/\psi p K^-)} = \frac{N^{\text{corr}}(\chi_{c1(2)} p K^-)}{N^{\text{corr}}(J/\psi p K^-)} \times \frac{1}{\mathcal{B}(\chi_{c1(2)} \rightarrow J/\psi \gamma)}, \quad (1)$$

$$R_{2/1} \equiv \frac{\mathcal{B}(\Lambda_b^0 \rightarrow \chi_{c2} p K^-)}{\mathcal{B}(\Lambda_b^0 \rightarrow \chi_{c1} p K^-)} = \frac{N^{\text{corr}}(\chi_{c2} p K^-)}{N^{\text{corr}}(\chi_{c1} p K^-)} \times \frac{\mathcal{B}(\chi_{c1} \rightarrow J/\psi \gamma)}{\mathcal{B}(\chi_{c2} \rightarrow J/\psi \gamma)}, \quad (2)$$

where the branching fraction of the $\chi_{c1} \rightarrow J/\psi \gamma$ ($\chi_{c2} \rightarrow J/\psi \gamma$) decay is taken to be $(33.9 \pm 1.2)\%$ ($(19.2 \pm 0.7)\%$) [33]. Figure 3(a) shows the distribution of $\Delta m + m_{J/\psi}$, where Δm is $m(\mu^+ \mu^- \gamma) - m(\mu^+ \mu^-)$ and $m_{J/\psi}$ is the known mass of the J/ψ meson [33], while Fig. 3(b) shows $\Delta m + m_{J/\psi}$ for background-subtracted $\Lambda_b^0 \rightarrow \chi_{cJ} p K^-$ candidates. Both distributions show clear enhancements at the known masses of the χ_{cJ} mesons.

The $\Lambda_b^0 \rightarrow \chi_{cJ} p K^-$ data sample is also used to make a measurement of the Λ_b^0 mass, $m_{\Lambda_b^0}$. The momenta of the particles are scaled to account for known miscalibration of the

detector [42]. In addition, the separation between the $\Lambda_b^0 \rightarrow \chi_{c1} p K^-$ and $\Lambda_b^0 \rightarrow \chi_{c2} p K^-$ components in the $m(\chi_{c1} p K^-)$ spectrum is fixed to the known mass difference between the χ_{c1} and χ_{c2} mesons [33], to obtain a single measurement of $m_{\Lambda_b^0}$ using both decay modes. The mass fit is repeated after these changes, yielding $m_{\Lambda_b^0} = 5619.44 \pm 0.28 \text{ MeV}/c^2$.

Systematic uncertainties on the ratios of branching fractions are assigned due to imperfect knowledge of the trigger efficiency and data/simulation discrepancies in the photon reconstruction as well as for uncertainties on the corrections applied to the simulated data (kinematic reweighting, phase-space weighting and particle identification resampling), the treatment of multiple candidates, the limited size of simulated data samples, and the models for the signal and background components in the fits. The per-candidate efficiencies as a function of the Dalitz plot coordinates are also replaced by phase-space-averaged efficiencies, and the difference with the nominal result is assigned as a systematic uncertainty.

The systematic uncertainties due to the trigger and photon reconstruction are taken from previous LHCb studies [20, 40, 43]. The uncertainties assigned due to the kinematic weighting are evaluated by repeating the analysis with alternative efficiency histograms that make use of a simplified three-dimensional weighting procedure with only the Λ_b^0 pseudorapidity, $\Lambda_b^0 p_T$ and event track-multiplicity. The uncertainty on the correction for multiple candidates is assigned as a systematic uncertainty. The uncertainty from the size of the simulated samples is determined from pseudoexperiments by varying the efficiency in each bin of each efficiency histogram within its uncertainties. The signal and background components of the invariant mass fits are replaced with the sum of two Gaussian functions and an exponential function, respectively, to estimate systematic uncertainties due to the choice of models.

The largest systematic uncertainties on R_1 and R_2 come from the photon reconstruction (both 4.0%) and kinematic weighting of the simulated samples (2.2% and 2.0%, respectively). For $R_{2/1}$, the largest systematic uncertainties are due to the size of the simulated samples (1.2%) and the treatment of multiple candidates (1.1%). The total systematic uncertainties on R_1 , R_2 and $R_{2/1}$ are 5.2%, 5.4% and 2.0%, respectively.

For the Λ_b^0 mass measurement, systematic uncertainties are assigned due to the uncertainty on the momentum scale for charged-particle tracks, uncertainties on the χ_{cJ} and kaon masses, energy loss in the material, miscalibration of the electromagnetic calorimeter, and the models for the signal and background components in the fit. The total systematic uncertainty ($0.26 \text{ MeV}/c^2$) is dominated by the uncertainty on the momentum scale ($0.24 \text{ MeV}/c^2$), the effect of which is determined by repeating the analysis with the momentum scaling parameter varied up and down by one standard deviation [42]. The uncertainty from the miscalibration of the electromagnetic calorimeter [44] is found to be small due to the mass constraints that are applied, as is the uncertainty on the energy loss in material, which is taken from a previous LHCb study [45].

In conclusion, the ratios of branching fractions are found to be

$$\begin{aligned} R_1 &= 0.242 \pm 0.014 \pm 0.013 \pm 0.009, \\ R_2 &= 0.248 \pm 0.020 \pm 0.014 \pm 0.009, \\ R_{2/1} &= 1.02 \pm 0.10 \pm 0.02 \pm 0.05, \end{aligned}$$

where the first uncertainty is statistical, the second systematic and the third due to the uncertainty on the branching fractions of the $\chi_{cJ} \rightarrow J/\psi \gamma$ decays. The values of R_1 and R_2 may be combined with existing measurements of

$\mathcal{B}(\Lambda_b^0 \rightarrow J/\psi p K^-) / \mathcal{B}(B^0 \rightarrow J/\psi K^*(892)^0)$ [46] and $\mathcal{B}(B^0 \rightarrow J/\psi K^*(892)^0)$ [47] to obtain absolute branching fraction measurements. As the result in Ref. [47] assumes equal production of B^+B^- and $B^0\bar{B}^0$ pairs at the $\Upsilon(4S)$ resonance, a correction is applied using the current world average value of $\mathcal{B}(\Upsilon(4S) \rightarrow B^+B^-) / \mathcal{B}(\Upsilon(4S) \rightarrow B^0\bar{B}^0) = 1.058 \pm 0.024$ [33], yielding $\mathcal{B}(B^0 \rightarrow J/\psi K^*(892)^0) = (1.22 \pm 0.08) \times 10^{-3}$ and $\mathcal{B}(\Lambda_b^0 \rightarrow J/\psi p K^-) = (3.01 \pm 0.21^{+0.43}_{-0.26}) \times 10^{-4}$, where the second uncertainty is due to the ratio of fragmentation fractions, $f_{\Lambda_b^0}/f_d$ [48, 49], and the first incorporates all other sources. This gives

$$\begin{aligned}\mathcal{B}(\Lambda_b^0 \rightarrow \chi_{c1} p K^-) &= (7.3 \pm 0.4 \pm 0.4 \pm 0.6^{+1.0}_{-0.6}) \times 10^{-5}, \\ \mathcal{B}(\Lambda_b^0 \rightarrow \chi_{c2} p K^-) &= (7.5 \pm 0.6 \pm 0.4 \pm 0.6^{+1.1}_{-0.6}) \times 10^{-5},\end{aligned}$$

where the third uncertainty is due to uncertainties on the $\chi_{cJ} \rightarrow J/\psi \gamma$, $\Lambda_b^0 \rightarrow J/\psi p K^-$ and $B^0 \rightarrow J/\psi K^*(892)^0$ branching fractions and the fourth is due to $f_{\Lambda_b^0}/f_d$ [48, 49]. These results show no suppression of the χ_{c2} mode relative to the χ_{c1} mode in Λ_b^0 baryon decays, which is different to what is observed in $B \rightarrow \chi_{cJ} K$ decays [18–20]. These decays will be useful for future investigations into the nature of the two pentaquark candidates observed by the LHCb collaboration and provide further information on the applicability of the factorisation approach in describing b -hadron decays to final states containing charmonium.

The Λ_b^0 mass has also been measured and is found to be $5619.44 \pm 0.28 \pm 0.26 \text{ MeV}/c^2$, where the first uncertainty is statistical and the second systematic. This result is combined with previous LHCb measurements from $\Lambda_b^0 \rightarrow [c\bar{c}]X$ decays [45, 50, 51] assuming that systematic uncertainties on the momentum scale and energy loss are fully correlated between the measurements while other sources of systematic uncertainties are uncorrelated. The procedure for the combination is the same as that used in Ref. [45]. This yields a new average value of $5619.62 \pm 0.16 \pm 0.13 \text{ MeV}/c^2$, which supersedes previous combinations of these results.

Acknowledgements

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); NWO (The Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MinES and FASO (Russia); MinECo (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (The Netherlands), PIC (Spain), GridPP (United Kingdom), RRCKI and Yandex LLC (Russia), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), PL-GRID (Poland) and OSC (USA). We are indebted to the communities behind the multiple open source software packages on which we depend. Individual groups or members have received support from AvH Foundation (Germany), EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union), Conseil Général de Haute-Savoie, Labex ENIGMASS and OCEVU, Région Auvergne (France), RFBR and Yandex LLC (Russia), GVA, XuntaGal and GENCAT (Spain), Herchel Smith Fund, The

Royal Society, Royal Commission for the Exhibition of 1851 and the Leverhulme Trust (United Kingdom).

References

- [1] M. Gell-Mann, *A schematic model of baryons and mesons*, Phys. Lett. **8** (1964) 214.
- [2] G. Zweig, *An $SU(3)$ model for strong interaction symmetry and its breaking, Part 1*, CERN-TH-401 (1964).
- [3] G. Zweig, *An $SU(3)$ model for strong interaction symmetry and its breaking, Part 2*, CERN-TH-412, reprinted in *Developments in the Quark Theory of Hadrons 1* (1980) 22 (ed. D. Lichtenberg and S. Rosen).
- [4] LHCb collaboration, R. Aaij *et al.*, *Observation of $J/\psi p$ resonances consistent with pentaquark states in $\Lambda_b^0 \rightarrow J/\psi p K^-$ decays*, Phys. Rev. Lett. **115** (2015) 072001, arXiv:1507.03414.
- [5] LHCb collaboration, R. Aaij *et al.*, *Model-independent evidence for $J/\psi p$ contributions to $\Lambda_b^0 \rightarrow J/\psi p K^-$ decays*, Phys. Rev. Lett. **117** (2016) 082002, arXiv:1604.05708.
- [6] M. Karliner and J. L. Rosner, *New exotic meson and baryon resonances from doubly-heavy hadronic molecules*, Phys. Rev. Lett. **115** (2015) 122001, arXiv:1506.06386.
- [7] H.-X. Chen *et al.*, *Towards exotic hidden-charm pentaquarks in QCD*, Phys. Rev. Lett. **115** (2015) 172001, arXiv:1507.03717.
- [8] L. Roca, J. Nieves, and E. Oset, *LHCb pentaquark as a $\bar{D}^* \Sigma_c - \bar{D} \Sigma_c^*$ molecular state*, Phys. Rev. **D92** (2015) 094003, arXiv:1507.04249.
- [9] L. Maiani, A. D. Polosa, and V. Riquer, *The new pentaquarks in the diquark model*, Phys. Lett. **B749** (2015) 289, arXiv:1507.04980.
- [10] R. F. Lebed, *The pentaquark candidates in the dynamical diquark picture*, Phys. Lett. **B749** (2015) 454, arXiv:1507.05867.
- [11] G.-N. Li, X.-G. He, and M. He, *Some predictions of diquark model for hidden charm pentaquark discovered at the LHCb*, JHEP **12** (2015) 128, arXiv:1507.08252.
- [12] F.-K. Guo, U.-G. Meissner, W. Wang, and Z. Yang, *How to reveal the exotic nature of the $P_c(4450)$* , Phys. Rev. **D92** (2015) 071502, arXiv:1507.04950.
- [13] M. Mikhasenko, *A triangle singularity and the LHCb pentaquarks*, arXiv:1507.06552.
- [14] X.-H. Liu, Q. Wang, and Q. Zhao, *Understanding the newly observed heavy pentaquark candidates*, Phys. Lett. **B757** (2016) 231, arXiv:1507.05359.
- [15] U.-G. Meissner and J. A. Oller, *Testing the $\chi_{c1} p$ composite nature of the $P_c(4450)$* , Phys. Lett. **B751** (2015) 59, arXiv:1507.07478.

- [16] F.-K. Guo, U. G. Meissner, J. Nieves, and Z. Yang, *Remarks on the P_c structures and triangle singularities*, Eur. Phys. J. **A52** (2016) 318, [arXiv:1605.05113](#).
- [17] M. Bayar, F. Aceti, F.-K. Guo, and E. Oset, *A Discussion on Triangle Singularities in the $\Lambda_b \rightarrow J/\psi K^- p$ Reaction*, Phys. Rev. **D94** (2016) 074039, [arXiv:1609.04133](#).
- [18] Belle collaboration, R. Mizuk *et al.*, *Observation of two resonance-like structures in the $\pi^+ \chi_{c1}$ mass distribution in exclusive $\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}$ decays*, Phys. Rev. **D78** (2008) 072004, [arXiv:0806.4098](#).
- [19] BaBar collaboration, B. Aubert *et al.*, *Evidence for $X(3872) \rightarrow \psi(2S)\gamma$ in $B^\pm \rightarrow X_{3872} K^\pm$ decays, and a study of $B \rightarrow c\bar{c}\gamma K$* , Phys. Rev. Lett. **102** (2009) 132001, [arXiv:0809.0042](#).
- [20] LHCb collaboration, R. Aaij *et al.*, *Observation of $B_s^0 \rightarrow \chi_{c1} \phi$ decay and study of $B^0 \rightarrow \chi_{c1,2} K^{*0}$ decays*, Nucl. Phys. **B874** (2013) 663, [arXiv:1305.6511](#).
- [21] M. Beneke and L. Vernazza, *$B \rightarrow \chi_{cJ} K$ decays revisited*, Nucl. Phys. **B811** (2009) 155, [arXiv:0810.3575](#).
- [22] Belle collaboration, V. Bhardwaj *et al.*, *Inclusive and exclusive measurements of B decays to χ_{c1} and χ_{c2} at Belle*, Phys. Rev. **D93** (2016) 052016, [arXiv:1512.02672](#).
- [23] LHCb collaboration, A. A. Alves Jr. *et al.*, *The LHCb detector at the LHC*, JINST **3** (2008) S08005.
- [24] LHCb collaboration, R. Aaij *et al.*, *LHCb detector performance*, Int. J. Mod. Phys. **A30** (2015) 1530022, [arXiv:1412.6352](#).
- [25] T. Sjöstrand, S. Mrenna, and P. Skands, *PYTHIA 6.4 physics and manual*, JHEP **05** (2006) 026, [arXiv:hep-ph/0603175](#); T. Sjöstrand, S. Mrenna, and P. Skands, *A brief introduction to PYTHIA 8.1*, Comput. Phys. Commun. **178** (2008) 852, [arXiv:0710.3820](#).
- [26] I. Belyaev *et al.*, *Handling of the generation of primary events in Gauss, the LHCb simulation framework*, J. Phys. Conf. Ser. **331** (2011) 032047.
- [27] D. J. Lange, *The EvtGen particle decay simulation package*, Nucl. Instrum. Meth. **A462** (2001) 152.
- [28] P. Golonka and Z. Was, *PHOTOS Monte Carlo: A precision tool for QED corrections in Z and W decays*, Eur. Phys. J. **C45** (2006) 97, [arXiv:hep-ph/0506026](#).
- [29] Geant4 collaboration, J. Allison *et al.*, *Geant4 developments and applications*, IEEE Trans. Nucl. Sci. **53** (2006) 270; Geant4 collaboration, S. Agostinelli *et al.*, *Geant4: A simulation toolkit*, Nucl. Instrum. Meth. **A506** (2003) 250.
- [30] M. Clemencic *et al.*, *The LHCb simulation application, Gauss: Design, evolution and experience*, J. Phys. Conf. Ser. **331** (2011) 032023.
- [31] L. Breiman, J. H. Friedman, R. A. Olshen, and C. J. Stone, *Classification and regression trees*, Wadsworth international group, Belmont, California, USA, 1984.

- [32] W. D. Hulsbergen, *Decay chain fitting with a Kalman filter*, Nucl. Instrum. Meth. **A552** (2005) 566, [arXiv:physics/0503191](#).
- [33] Particle Data Group, C. Patrignani *et al.*, *Review of particle physics*, Chin. Phys. **C40** (2016) 100001.
- [34] M. Adinolfi *et al.*, *Performance of the LHCb RICH detector at the LHC*, Eur. Phys. J. **C73** (2013) 2431, [arXiv:1211.6759](#).
- [35] G. A. Cowan, D. C. Craik, and M. D. Needham, *RapidSim: An application for the fast simulation of heavy-quark hadron decays*, Comput. Phys. Commun. **214** (2017) 239, [arXiv:1612.07489](#).
- [36] A. Rogozhnikov, *Reweighting with boosted decision trees*, J. Phys. Conf. Ser. **762** (2016) 012036, [arXiv:1608.05806](#).
- [37] T. Skwarnicki, *A study of the radiative cascade transitions between the Upsilon-prime and Upsilon resonances*, PhD thesis, Institute of Nuclear Physics, Krakow, 1986, DESY-F31-86-02.
- [38] S. S. Wilks, *The large-sample distribution of the likelihood ratio for testing composite hypotheses*, Ann. Math. Stat. **9** (1938) 60.
- [39] R. H. Dalitz, *On the analysis of τ -meson data and the nature of the τ -meson*, Phil. Mag. **44** (1953) 1068.
- [40] E. Govorkova, *Study of photon reconstruction efficiency using $B^+ \rightarrow J/\psi K^{(*)+}$ decays in the LHCb experiment*, Phys. Atom. Nucl. **79** (2016) 1474, [arXiv:1505.02960](#).
- [41] M. Pivk and F. R. Le Diberder, *sPlot: A statistical tool to unfold data distributions*, Nucl. Instrum. Meth. **A555** (2005) 356, [arXiv:physics/0402083](#).
- [42] LHCb collaboration, R. Aaij *et al.*, *Precision measurement of D meson mass differences*, JHEP **06** (2013) 065, [arXiv:1304.6865](#).
- [43] LHCb collaboration, R. Aaij *et al.*, *Measurement of relative branching fractions of B decays to $\psi(2S)$ and J/ψ mesons*, Eur. Phys. J. **C72** (2012) 2118, [arXiv:1205.0918](#).
- [44] I. M. Belyaev *et al.*, *Calibration of the LHCb electromagnetic calorimeter via reconstructing the neutral-pion invariant mass*, Phys. Atom. Nucl. **78** (2015) 1026.
- [45] LHCb collaboration, R. Aaij *et al.*, *Observation of $\Lambda_b^0 \rightarrow \psi(2S)pK^-$ and $\Lambda_b^0 \rightarrow J/\psi \pi^+ \pi^- pK^-$ decays and a measurement of the Λ_b^0 baryon mass*, JHEP **05** (2016) 132, [arXiv:1603.06961](#).
- [46] LHCb collaboration, R. Aaij *et al.*, *Study of the productions of Λ_b^0 and \bar{B}^0 hadrons in pp collisions and first measurement of the $\Lambda_b^0 \rightarrow J/\psi pK^-$ branching fraction*, Chin. Phys. **C40** (2016) 011001, [arXiv:1509.00292](#).
- [47] Belle collaboration, K. Chilikin *et al.*, *Observation of a new charged charmoniumlike state in $\bar{B}^0 \rightarrow J/\psi K^- \pi^+$ decays*, Phys. Rev. **D90** (2014) 112009, [arXiv:1408.6457](#).

- [48] LHCb collaboration, R. Aaij *et al.*, *Measurement of b hadron production fractions in 7 TeV pp collisions*, Phys. Rev. **D85** (2012) 032008, [arXiv:1111.2357](#).
- [49] LHCb collaboration, R. Aaij *et al.*, *Study of the kinematic dependences of Λ_b^0 production in pp collisions and a measurement of the $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ branching fraction*, JHEP **08** (2014) 143, [arXiv:1405.6842](#).
- [50] LHCb collaboration, R. Aaij *et al.*, *Measurement of b -hadron masses*, Phys. Lett. **B708** (2012) 241, [arXiv:1112.4896](#).
- [51] LHCb collaboration, R. Aaij *et al.*, *Measurements of the Λ_b^0 , Ξ_b^- , and Ω_b^- baryon masses*, Phys. Rev. Lett. **110** (2013) 182001, [arXiv:1302.1072](#).

LHCb collaboration

R. Aaij⁴⁰, B. Adeva³⁹, M. Adinolfi⁴⁸, Z. Ajaltouni⁵, S. Akar⁵⁹, J. Albrecht¹⁰, F. Alessio⁴⁰, M. Alexander⁵³, S. Ali⁴³, G. Alkhazov³¹, P. Alvarez Cartelle⁵⁵, A.A. Alves Jr⁵⁹, S. Amato², S. Amerio²³, Y. Amhis⁷, L. An³, L. Anderlini¹⁸, G. Andreassi⁴¹, M. Andreotti^{17,g}, J.E. Andrews⁶⁰, R.B. Appleby⁵⁶, F. Archilli⁴³, P. d'Argent¹², J. Arnau Romeu⁶, A. Artamonov³⁷, M. Artuso⁶¹, E. Aslanides⁶, G. Auriemma²⁶, M. Baalouch⁵, I. Babuschkin⁵⁶, S. Bachmann¹², J.J. Back⁵⁰, A. Badalov³⁸, C. Baesso⁶², S. Baker⁵⁵, V. Balagura^{7,c}, W. Baldini¹⁷, A. Baranov³⁵, R.J. Barlow⁵⁶, C. Barschel⁴⁰, S. Barsuk⁷, W. Barter⁵⁶, F. Baryshnikov³², M. Baszczyk^{27,l}, V. Batozskaya²⁹, V. Battista⁴¹, A. Bay⁴¹, L. Beaucourt⁴, J. Beddow⁵³, F. Bedeschi²⁴, I. Bediaga¹, A. Beiter⁶¹, L.J. Bel⁴³, V. Bellee⁴¹, N. Belloli^{21,i}, K. Belous³⁷, I. Belyaev³², E. Ben-Haim⁸, G. Bencivenni¹⁹, S. Benson⁴³, S. Beranek⁹, A. Berezhnoy³³, R. Bernet⁴², A. Bertolin²³, C. Betancourt⁴², F. Betti¹⁵, M.-O. Bettler⁴⁰, M. van Beuzekom⁴³, I.A. Bezshyiko⁴², S. Bifani⁴⁷, P. Billoir⁸, A. Birnkraut¹⁰, A. Bitadze⁵⁶, A. Bizzeti^{18,u}, T. Blake⁵⁰, F. Blanc⁴¹, J. Blouw^{11,†}, S. Blusk⁶¹, V. Bocci²⁶, T. Boettcher⁵⁸, A. Bondar^{36,w}, N. Bondar³¹, W. Bonivento¹⁶, I. Bordyuzhin³², A. Borgheresi^{21,i}, S. Borghi⁵⁶, M. Borisyak³⁵, M. Borsato³⁹, F. Bossu⁷, M. Boubdir⁹, T.J.V. Bowcock⁵⁴, E. Bowen⁴², C. Bozzi^{17,40}, S. Braun¹², T. Britton⁶¹, J. Brodzicka⁵⁶, E. Buchanan⁴⁸, C. Burr⁵⁶, A. Bursche¹⁶, J. Buytaert⁴⁰, S. Cadeddu¹⁶, R. Calabrese^{17,g}, M. Calvi^{21,i}, M. Calvo Gomez^{38,m}, A. Camboni³⁸, P. Campana¹⁹, D.H. Campora Perez⁴⁰, L. Capriotti⁵⁶, A. Carbone^{15,e}, G. Carboni^{25,j}, R. Cardinale^{20,h}, A. Cardini¹⁶, P. Carniti^{21,i}, L. Carson⁵², K. Carvalho Akiba², G. Casse⁵⁴, L. Cassina^{21,i}, L. Castillo Garcia⁴¹, M. Cattaneo⁴⁰, G. Cavallero²⁰, R. Cenci^{24,t}, D. Chamont⁷, M. Charles⁸, Ph. Charpentier⁴⁰, G. Chatzikonstantinidis⁴⁷, M. Chefdeville⁴, S. Chen⁵⁶, S.F. Cheung⁵⁷, V. Chobanova³⁹, M. Chrzaszcz^{42,27}, A. Chubykin³¹, X. Cid Vidal³⁹, G. Ciezarek⁴³, P.E.L. Clarke⁵², M. Clemencic⁴⁰, H.V. Cliff⁴⁹, J. Closier⁴⁰, V. Coco⁵⁹, J. Cogan⁶, E. Cogneras⁵, V. Cogoni^{16,f}, L. Cojocariu³⁰, P. Collins⁴⁰, A. Comerma-Montells¹², A. Contu⁴⁰, A. Cook⁴⁸, G. Coombs⁴⁰, S. Coquereau³⁸, G. Corti⁴⁰, M. Corvo^{17,g}, C.M. Costa Sobral⁵⁰, B. Couturier⁴⁰, G.A. Cowan⁵², D.C. Craik⁵², A. Crocombe⁵⁰, M. Cruz Torres⁶², S. Cunliffe⁵⁵, R. Currie⁵², C. D'Ambrosio⁴⁰, F. Da Cunha Marinho², E. Dall'Occo⁴³, J. Dalseno⁴⁸, A. Davis³, O. De Aguiar Francisco⁵⁴, K. De Bruyn⁶, S. De Capua⁵⁶, M. De Cian¹², J.M. De Miranda¹, L. De Paula², M. De Serio^{14,d}, P. De Simone¹⁹, C.T. Dean⁵³, D. Decamp⁴, M. Deckenhoff¹⁰, L. Del Buono⁸, H.-P. Dembinski¹¹, M. Demmer¹⁰, A. Dendek²⁸, D. Derkach³⁵, O. Deschamps⁵, F. Dettori⁵⁴, B. Dey²², A. Di Canto⁴⁰, P. Di Nezza¹⁹, H. Dijkstra⁴⁰, F. Dordei⁴⁰, M. Dorigo⁴¹, A. Dosil Suárez³⁹, A. Dovbnya⁴⁵, K. Dreimanis⁵⁴, L. Dufour⁴³, G. Dujany⁵⁶, K. Dungs⁴⁰, P. Durante⁴⁰, R. Dzhelezhyan³⁷, M. Dziewiecki¹², A. Dziurda⁴⁰, A. Dzyuba³¹, N. Déleage⁴, S. Easo⁵¹, M. Ebert⁵², U. Egede⁵⁵, V. Egorychev³², S. Eidelman^{36,w}, S. Eisenhardt⁵², U. Eitschberger¹⁰, R. Ekelhof¹⁰, L. Eklund⁵³, S. Ely⁶¹, S. Esen¹², H.M. Evans⁴⁹, T. Evans⁵⁷, A. Falabella¹⁵, N. Farley⁴⁷, S. Farry⁵⁴, R. Fay⁵⁴, D. Fazzini^{21,i}, D. Ferguson⁵², G. Fernandez³⁸, A. Fernandez Prieto³⁹, F. Ferrari¹⁵, F. Ferreira Rodrigues², M. Ferro-Luzzi⁴⁰, S. Filippov³⁴, R.A. Fini¹⁴, M. Fiore^{17,g}, M. Fiorini^{17,g}, M. Firlej²⁸, C. Fitzpatrick⁴¹, T. Fiutowski²⁸, F. Fleuret^{7,b}, K. Fohl⁴⁰, M. Fontana^{16,40}, F. Fontanelli^{20,h}, D.C. Forshaw⁶¹, R. Forty⁴⁰, V. Franco Lima⁵⁴, M. Frank⁴⁰, C. Frei⁴⁰, J. Fu^{22,g}, W. Funk⁴⁰, E. Furfaro^{25,j}, C. Färber⁴⁰, E. Gabriel⁵², A. Gallas Torreira³⁹, D. Galli^{15,e}, S. Gallorini²³, S. Gambetta⁵², M. Gandelman², P. Gandini⁵⁷, Y. Gao³, L.M. Garcia Martin⁶⁹, J. García Pardiñas³⁹, J. Garra Tico⁴⁹, L. Garrido³⁸, P.J. Garsed⁴⁹, D. Gascon³⁸, C. Gaspar⁴⁰, L. Gavardi¹⁰, G. Gazzoni⁵, D. Gerick¹², E. Gersabeck¹², M. Gersabeck⁵⁶, T. Gershon⁵⁰, Ph. Ghez⁴, S. Giani⁴¹, V. Gibson⁴⁹, O.G. Girard⁴¹, L. Giubega³⁰, K. Gizdov⁵², V.V. Gligorov⁸, D. Golubkov³², A. Golutvin^{55,40}, A. Gomes^{1,a}, I.V. Gorelov³³, C. Gotti^{21,i}, E. Govorkova⁴³, R. Graciani Diaz³⁸, L.A. Granado Cardoso⁴⁰, E. Graugés³⁸, E. Graverini⁴², G. Graziani¹⁸, A. Grecu³⁰, R. Greim⁹, P. Griffith¹⁶, L. Grillo^{21,40,i}, L. Gruber⁴⁰, B.R. Gruber Cazon⁵⁷, O. Grünberg⁶⁷, E. Gushchin³⁴,

Yu. Guz³⁷, T. Gys⁴⁰, C. Göbel⁶², T. Hadavizadeh⁵⁷, C. Hadjivasiliou⁵, G. Haefeli⁴¹, C. Haen⁴⁰,
 S.C. Haines⁴⁹, B. Hamilton⁶⁰, X. Han¹², S. Hansmann-Menzemer¹², N. Harnew⁵⁷,
 S.T. Harnew⁴⁸, J. Harrison⁵⁶, M. Hatch⁴⁰, J. He⁶³, T. Head⁴¹, A. Heister⁹, K. Hennessy⁵⁴,
 P. Henrard⁵, L. Henry⁶⁹, E. van Herwijnen⁴⁰, M. Heß⁶⁷, A. Hicheur², D. Hill⁵⁷, C. Hombach⁵⁶,
 P.H. Hopchev⁴¹, Z.-C. Huard⁵⁹, W. Hulsbergen⁴³, T. Humair⁵⁵, M. Hushchyn³⁵,
 D. Hutchcroft⁵⁴, M. Idzik²⁸, P. Ilten⁵⁸, R. Jacobsson⁴⁰, J. Jalocha⁵⁷, E. Jans⁴³, A. Jawahery⁶⁰,
 F. Jiang³, M. John⁵⁷, D. Johnson⁴⁰, C.R. Jones⁴⁹, C. Joram⁴⁰, B. Jost⁴⁰, N. Jurik⁵⁷,
 S. Kandybei⁴⁵, M. Karacson⁴⁰, J.M. Kariuki⁴⁸, S. Karodia⁵³, M. Kecke¹², M. Kelsey⁶¹,
 M. Kenzie⁴⁹, T. Ketel⁴⁴, E. Khairullin³⁵, B. Khanji¹², C. Khurewathanakul⁴¹, T. Kirn⁹,
 S. Klaver⁵⁶, K. Klimaszewski²⁹, T. Klimkovich¹¹, S. Koliiev⁴⁶, M. Kolpin¹², I. Komarov⁴¹,
 R. Kopecna¹², P. Koppenburg⁴³, A. Kosmyntseva³², S. Kotriakhova³¹, A. Kozachuk³³,
 M. Kozeiha⁵, L. Kravchuk³⁴, M. Krepes⁵⁰, P. Krokovny^{36,w}, F. Kruse¹⁰, W. Krzemien²⁹,
 W. Kucewicz^{27,l}, M. Kucharczyk²⁷, V. Kudryavtsev^{36,w}, A.K. Kuonen⁴¹, K. Kurek²⁹,
 T. Kvaratskheliya^{32,40}, D. Lacarrere⁴⁰, G. Lafferty⁵⁶, A. Lai¹⁶, G. Lanfranchi¹⁹,
 C. Langenbruch⁹, T. Latham⁵⁰, C. Lazzeroni⁴⁷, R. Le Gac⁶, J. van Leerdam⁴³, A. Leflat^{33,40},
 J. Lefrançois⁷, R. Lefèvre⁵, F. Lemaitre⁴⁰, E. Lemos Cid³⁹, O. Leroy⁶, T. Lesiak²⁷,
 B. Leverington¹², T. Li³, Y. Li⁷, Z. Li⁶¹, T. Likhomanenko^{35,68}, R. Lindner⁴⁰, F. Lionetto⁴²,
 X. Liu³, D. Loh⁵⁰, I. Longstaff⁵³, J.H. Lopes², D. Lucchesi^{23,o}, M. Lucio Martinez³⁹, H. Luo⁵²,
 A. Lupato²³, E. Luppi^{17,g}, O. Lupton⁴⁰, A. Lusiani²⁴, X. Lyu⁶³, F. Machefert⁷, F. Maciuc³⁰,
 O. Maev³¹, K. Maguire⁵⁶, S. Malde⁵⁷, A. Malinin⁶⁸, T. Maltsev³⁶, G. Manca^{16,f}, G. Mancinelli⁶,
 P. Manning⁶¹, J. Maratas^{5,v}, J.F. Marchand⁴, U. Marconi¹⁵, C. Marin Benito³⁸,
 M. Marinangeli⁴¹, P. Marino^{24,t}, J. Marks¹², G. Martellotti²⁶, M. Martin⁶, M. Martinelli⁴¹,
 D. Martinez Santos³⁹, F. Martinez Vidal⁶⁹, D. Martins Tostes², L.M. Massacrier⁷,
 A. Massafferri¹, R. Matev⁴⁰, A. Mathad⁵⁰, Z. Mathe⁴⁰, C. Matteuzzi²¹, A. Mauri⁴²,
 E. Maurice^{7,b}, B. Maurin⁴¹, A. Mazurov⁴⁷, M. McCann^{55,40}, A. McNab⁵⁶, R. McNulty¹³,
 B. Meadows⁵⁹, F. Meier¹⁰, D. Melnychuk²⁹, M. Merk⁴³, A. Merli^{22,40,q}, E. Michielin²³,
 D.A. Milanes⁶⁶, M.-N. Minard⁴, D.S. Mitzel¹², A. Mogini⁸, J. Molina Rodriguez¹,
 I.A. Monroy⁶⁶, S. Monteil⁵, M. Morandin²³, M.J. Morello^{24,t}, O. Morgunova⁶⁸, J. Moron²⁸,
 A.B. Morris⁵², A.P. Morris⁵², R. Mountain⁶¹, F. Muheim⁵², M. Mulder⁴³, M. Mussini¹⁵,
 D. Müller⁵⁶, J. Müller¹⁰, K. Müller⁴², V. Müller¹⁰, P. Naik⁴⁸, T. Nakada⁴¹, R. Nandakumar⁵¹,
 A. Nandi⁵⁷, I. Nasteva², M. Needham⁵², N. Neri^{22,40}, S. Neubert¹², N. Neufeld⁴⁰, M. Neuner¹²,
 T.D. Nguyen⁴¹, C. Nguyen-Mau^{41,n}, S. Nieswand⁹, R. Niet¹⁰, N. Nikitin³³, T. Nikodem¹²,
 A. Nogay⁶⁸, D.P. O'Hanlon⁵⁰, A. Oblakowska-Mucha²⁸, V. Obraztsov³⁷, S. Ogilvy¹⁹,
 R. Oldeman^{16,f}, C.J.G. Onderwater⁷⁰, A. Ossowska²⁷, J.M. Otalora Goicochea², P. Owen⁴²,
 A. Oyanguren⁶⁹, P.R. Pais⁴¹, A. Palano^{14,d}, M. Palutan^{19,40}, A. Papanestis⁵¹, M. Pappagallo^{14,d},
 L.L. Pappalardo^{17,g}, C. Pappenheimer⁵⁹, W. Parker⁶⁰, C. Parkes⁵⁶, G. Passaleva¹⁸,
 A. Pastore^{14,d}, M. Patel⁵⁵, C. Patrignani^{15,e}, A. Pearce⁴⁰, A. Pellegrino⁴³, G. Penso²⁶,
 M. Pepe Altarelli⁴⁰, S. Perazzini⁴⁰, P. Perret⁵, L. Pescatore⁴¹, K. Petridis⁴⁸, A. Petrolini^{20,h},
 A. Petrov⁶⁸, M. Petruzzo^{22,q}, E. Picatoste Olloqui³⁸, B. Pietrzyk⁴, M. Pikies²⁷, D. Pinci²⁶,
 A. Pistone²⁰, A. Piucci¹², V. Placinta³⁰, S. Playfer⁵², M. Plo Casasus³⁹, T. Poikela⁴⁰, F. Polci⁸,
 M. Poli Lener¹⁹, A. Poluektov^{50,36}, I. Polyakov⁶¹, E. Polcarpo², G.J. Pomery⁴⁸, S. Ponce⁴⁰,
 A. Popov³⁷, D. Popov^{11,40}, B. Popovici³⁰, S. Poslavskii³⁷, C. Potterat², E. Price⁴⁸,
 J. Prisciandaro³⁹, C. Prouve⁴⁸, V. Pugatch⁴⁶, A. Puig Navarro⁴², G. Punzi^{24,p}, C. Qian⁶³,
 W. Qian⁵⁰, R. Quagliani^{7,48}, B. Rachwal²⁸, J.H. Rademacker⁴⁸, M. Rama²⁴, M. Ramos Pernas³⁹,
 M.S. Rangel², I. Raniuk^{45,†}, F. Ratnikov³⁵, G. Raven⁴⁴, M. Ravonel Salzgeber⁴⁰, M. Reboud⁴,
 F. Redi⁵⁵, S. Reichert¹⁰, A.C. dos Reis¹, C. Remon Alepuz⁶⁹, V. Renaudin⁷, S. Ricciardi⁵¹,
 S. Richards⁴⁸, M. Rihl⁴⁰, K. Rinnert⁵⁴, V. Rives Molina³⁸, P. Robbe⁷, A.B. Rodrigues¹,
 E. Rodrigues⁵⁹, J.A. Rodriguez Lopez⁶⁶, P. Rodriguez Perez^{56,†}, A. Rogozhnikov³⁵, S. Roiser⁴⁰,
 A. Rollings⁵⁷, V. Romanovskiy³⁷, A. Romero Vidal³⁹, J.W. Ronayne¹³, M. Rotondo¹⁹,
 M.S. Rudolph⁶¹, T. Ruf⁴⁰, P. Ruiz Valls⁶⁹, J.J. Saborido Silva³⁹, E. Sadykhov³², N. Sagidova³¹,

B. Saitta^{16,f}, V. Salustino Guimaraes¹, D. Sanchez Gonzalo³⁸, C. Sanchez Mayordomo⁶⁹,
 B. Sanmartin Sedes³⁹, R. Santacesaria²⁶, C. Santamarina Rios³⁹, M. Santimaria¹⁹,
 E. Santovetti^{25,j}, A. Sarti^{19,k}, C. Satriano^{26,s}, A. Satta²⁵, D.M. Saunders⁴⁸, D. Savrina^{32,33},
 S. Schael⁹, M. Schellenberg¹⁰, M. Schiller⁵³, H. Schindler⁴⁰, M. Schlupp¹⁰, M. Schmelling¹¹,
 T. Schmelzer¹⁰, B. Schmidt⁴⁰, O. Schneider⁴¹, A. Schopper⁴⁰, H.F. Schreiner⁵⁹, K. Schubert¹⁰,
 M. Schubiger⁴¹, M.-H. Schune⁷, R. Schwemmer⁴⁰, B. Sciascia¹⁹, A. Sciubba^{26,k},
 A. Semennikov³², A. Sergi⁴⁷, N. Serra⁴², J. Serrano⁶, L. Sestini²³, P. Seyfert²¹, M. Shapkin³⁷,
 I. Shapoval⁴⁵, Y. Shcheglov³¹, T. Shears⁵⁴, L. Shekhtman^{36,w}, V. Shevchenko⁶⁸, B.G. Siddi^{17,40},
 R. Silva Coutinho⁴², L. Silva de Oliveira², G. Simi^{23,o}, S. Simone^{14,d}, M. Sirendi⁴⁹,
 N. Skidmore⁴⁸, T. Skwarnicki⁶¹, E. Smith⁵⁵, I.T. Smith⁵², J. Smith⁴⁹, M. Smith⁵⁵,
 I. Soares Lavra¹, M.D. Sokoloff⁵⁹, F.J.P. Soler⁵³, B. Souza De Paula², B. Spaan¹⁰, P. Spradlin⁵³,
 S. Sridharan⁴⁰, F. Stagni⁴⁰, M. Stahl¹², S. Stahl⁴⁰, P. Stefko⁴¹, S. Stefkova⁵⁵, O. Steinkamp⁴²,
 S. Stemmler¹², O. Stenyakin³⁷, H. Stevens¹⁰, S. Stoica³⁰, S. Stone⁶¹, B. Storaci⁴², S. Stracka^{24,p},
 M.E. Stramaglia⁴¹, M. Straticiuc³⁰, U. Straumann⁴², L. Sun⁶⁴, W. Sutcliffe⁵⁵, K. Swientek²⁸,
 V. Syropoulos⁴⁴, M. Szczekowski²⁹, T. Szumlak²⁸, S. T'Jampens⁴, A. Tayduganov⁶,
 T. Tekampe¹⁰, G. Tellarini^{17,g}, F. Teubert⁴⁰, E. Thomas⁴⁰, J. van Tilburg⁴³, M.J. Tilley⁵⁵,
 V. Tisserand⁴, M. Tobin⁴¹, S. Tolk⁴⁹, L. Tomassetti^{17,g}, D. Tonelli²⁴, S. Topp-Joergensen⁵⁷,
 F. Toriello⁶¹, R. Tourinho Jadallah Aoude¹, E. Tournefier⁴, S. Tourneur⁴¹, K. Trabelsi⁴¹,
 M. Traill⁵³, M.T. Tran⁴¹, M. Tresch⁴², A. Trisovic⁴⁰, A. Tsaregorodtsev⁶, P. Tsopelas⁴³,
 A. Tully⁴⁹, N. Tuning⁴³, A. Ukleja²⁹, A. Ustyuzhanin³⁵, U. Uwer¹², C. Vacca^{16,f},
 V. Vagnoni^{15,40}, A. Valassi⁴⁰, S. Valat⁴⁰, G. Valenti¹⁵, R. Vazquez Gomez¹⁹,
 P. Vazquez Regueiro³⁹, S. Vecchi¹⁷, M. van Veghel⁴³, J.J. Velthuis⁴⁸, M. Veltri^{18,r},
 G. Veneziano⁵⁷, A. Venkateswaran⁶¹, T.A. Verlage⁹, M. Vernet⁵, M. Vesterinen¹²,
 J.V. Viana Barbosa⁴⁰, B. Viaud⁷, D. Vieira⁶³, M. Vieites Diaz³⁹, H. Viemann⁶⁷,
 X. Vilasis-Cardona^{38,m}, M. Vitti⁴⁹, V. Volkov³³, A. Vollhardt⁴², B. Voneki⁴⁰, A. Vorobyev³¹,
 V. Vorobyev^{36,w}, C. Voß⁹, J.A. de Vries⁴³, C. Vázquez Sierra³⁹, R. Waldi⁶⁷, C. Wallace⁵⁰,
 R. Wallace¹³, J. Walsh²⁴, J. Wang⁶¹, D.R. Ward⁴⁹, H.M. Wark⁵⁴, N.K. Watson⁴⁷,
 D. Websdale⁵⁵, A. Weiden⁴², M. Whitehead⁴⁰, J. Wicht⁵⁰, G. Wilkinson^{57,40}, M. Wilkinson⁶¹,
 M. Williams⁴⁰, M.P. Williams⁴⁷, M. Williams⁵⁸, T. Williams⁴⁷, F.F. Wilson⁵¹, J. Wimberley⁶⁰,
 M.A. Winn⁷, J. Wishahi¹⁰, W. Wislicki²⁹, M. Witek²⁷, G. Wormser⁷, S.A. Wotton⁴⁹,
 K. Wraight⁵³, K. Wyllie⁴⁰, Y. Xie⁶⁵, Z. Xu⁴, Z. Yang³, Z. Yang⁶⁰, Y. Yao⁶¹, H. Yin⁶⁵, J. Yu⁶⁵,
 X. Yuan⁶¹, O. Yushchenko³⁷, K.A. Zarebski⁴⁷, M. Zavertyaev^{11,c}, L. Zhang³, Y. Zhang⁷,
 A. Zhelezov¹², Y. Zheng⁶³, X. Zhu³, V. Zhukov³³, J.B. Zonneveld⁵², S. Zucchelli¹⁵.

¹Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

²Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

³Center for High Energy Physics, Tsinghua University, Beijing, China

⁴LAPP, Université Savoie Mont-Blanc, CNRS/IN2P3, Annecy-Le-Vieux, France

⁵Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, Clermont-Ferrand, France

⁶CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France

⁷LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France

⁸LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris, France

⁹I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany

¹⁰Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany

¹¹Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany

¹²Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

¹³School of Physics, University College Dublin, Dublin, Ireland

¹⁴Sezione INFN di Bari, Bari, Italy

¹⁵Sezione INFN di Bologna, Bologna, Italy

¹⁶Sezione INFN di Cagliari, Cagliari, Italy

¹⁷Sezione INFN di Ferrara, Ferrara, Italy

¹⁸Sezione INFN di Firenze, Firenze, Italy

- ¹⁹ *Laboratori Nazionali dell'INFN di Frascati, Frascati, Italy*
- ²⁰ *Sezione INFN di Genova, Genova, Italy*
- ²¹ *Sezione INFN di Milano Bicocca, Milano, Italy*
- ²² *Sezione INFN di Milano, Milano, Italy*
- ²³ *Sezione INFN di Padova, Padova, Italy*
- ²⁴ *Sezione INFN di Pisa, Pisa, Italy*
- ²⁵ *Sezione INFN di Roma Tor Vergata, Roma, Italy*
- ²⁶ *Sezione INFN di Roma La Sapienza, Roma, Italy*
- ²⁷ *Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*
- ²⁸ *AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland*
- ²⁹ *National Center for Nuclear Research (NCBJ), Warsaw, Poland*
- ³⁰ *Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*
- ³¹ *Petersburg Nuclear Physics Institute (PNPI), Gatchina, Russia*
- ³² *Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia*
- ³³ *Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia*
- ³⁴ *Institute for Nuclear Research of the Russian Academy of Sciences (INR RAN), Moscow, Russia*
- ³⁵ *Yandex School of Data Analysis, Moscow, Russia*
- ³⁶ *Budker Institute of Nuclear Physics (SB RAS), Novosibirsk, Russia*
- ³⁷ *Institute for High Energy Physics (IHEP), Protvino, Russia*
- ³⁸ *ICCUB, Universitat de Barcelona, Barcelona, Spain*
- ³⁹ *Universidad de Santiago de Compostela, Santiago de Compostela, Spain*
- ⁴⁰ *European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- ⁴¹ *Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- ⁴² *Physik-Institut, Universität Zürich, Zürich, Switzerland*
- ⁴³ *Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands*
- ⁴⁴ *Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, The Netherlands*
- ⁴⁵ *NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- ⁴⁶ *Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- ⁴⁷ *University of Birmingham, Birmingham, United Kingdom*
- ⁴⁸ *H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- ⁴⁹ *Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- ⁵⁰ *Department of Physics, University of Warwick, Coventry, United Kingdom*
- ⁵¹ *STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- ⁵² *School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- ⁵³ *School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- ⁵⁴ *Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- ⁵⁵ *Imperial College London, London, United Kingdom*
- ⁵⁶ *School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- ⁵⁷ *Department of Physics, University of Oxford, Oxford, United Kingdom*
- ⁵⁸ *Massachusetts Institute of Technology, Cambridge, MA, United States*
- ⁵⁹ *University of Cincinnati, Cincinnati, OH, United States*
- ⁶⁰ *University of Maryland, College Park, MD, United States*
- ⁶¹ *Syracuse University, Syracuse, NY, United States*
- ⁶² *Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to ²*
- ⁶³ *University of Chinese Academy of Sciences, Beijing, China, associated to ³*
- ⁶⁴ *School of Physics and Technology, Wuhan University, Wuhan, China, associated to ³*
- ⁶⁵ *Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China, associated to ³*
- ⁶⁶ *Departamento de Física, Universidad Nacional de Colombia, Bogota, Colombia, associated to ⁸*
- ⁶⁷ *Institut für Physik, Universität Rostock, Rostock, Germany, associated to ¹²*
- ⁶⁸ *National Research Centre Kurchatov Institute, Moscow, Russia, associated to ³²*
- ⁶⁹ *Instituto de Física Corpuscular, Centro Mixto Universidad de Valencia - CSIC, Valencia, Spain, associated to ³⁸*
- ⁷⁰ *Van Swinderen Institute, University of Groningen, Groningen, The Netherlands, associated to ⁴³*

^a *Universidade Federal do Triângulo Mineiro (UFTM), Uberaba-MG, Brazil*

- ^b*Laboratoire Leprince-Ringuet, Palaiseau, France*
- ^c*P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia*
- ^d*Università di Bari, Bari, Italy*
- ^e*Università di Bologna, Bologna, Italy*
- ^f*Università di Cagliari, Cagliari, Italy*
- ^g*Università di Ferrara, Ferrara, Italy*
- ^h*Università di Genova, Genova, Italy*
- ⁱ*Università di Milano Bicocca, Milano, Italy*
- ^j*Università di Roma Tor Vergata, Roma, Italy*
- ^k*Università di Roma La Sapienza, Roma, Italy*
- ^l*AGH - University of Science and Technology, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland*
- ^m*LIFAEELS, La Salle, Universitat Ramon Llull, Barcelona, Spain*
- ⁿ*Hanoi University of Science, Hanoi, Viet Nam*
- ^o*Università di Padova, Padova, Italy*
- ^p*Università di Pisa, Pisa, Italy*
- ^q*Università degli Studi di Milano, Milano, Italy*
- ^r*Università di Urbino, Urbino, Italy*
- ^s*Università della Basilicata, Potenza, Italy*
- ^t*Scuola Normale Superiore, Pisa, Italy*
- ^u*Università di Modena e Reggio Emilia, Modena, Italy*
- ^v*Iligan Institute of Technology (IIT), Iligan, Philippines*
- ^w*Novosibirsk State University, Novosibirsk, Russia*
- [†]*Deceased*