

# The Dynamics of Beauty & Charm Hadrons and top quarks in the Era of the LHCb & Belle II and ATLAS/CMS – Non-perturbative QCD & Many-body Final States <sup>1</sup>

I.I. Bigi<sup>a</sup>

<sup>a</sup> *Department of Physics, University of Notre Dame du Lac  
Notre Dame, IN 46556, USA  
email addresses: ibigi@nd.edu*

*This article is dedicated to Timothy O’Meara, mathematician  
& first lay Provost of the University of Notre Dame du Lac.*

## Abstract

Our community has to apply *non-perturbative* QCD on different levels of flavor dynamics in strange, charm & beauty hadrons and even for top quarks. We need *consistent* parameterization of the CKM matrix and describe weak decays of beauty hadrons with *many-body* final states. It is crucial to use the *Wilsonian* OPE as much as possible and discuss ”duality” in the worlds of quarks and hadrons. The pole mass of heavy quarks is *not* well-defined on the *non-perturbative* level – i.e., it is *not* Borel summable in total QCD. We need a novel team to combine the strengths of our tools from MEP and HEP.

## Contents

<b>1</b>	<b>Prologue</b>	<b>2</b>
<b>2</b>	<b>Introduction to the ‘Roads’</b>	<b>4</b>
<b>3</b>	<b>Anomalies: ”deep” or not so</b>	<b>6</b>
<b>4</b>	<b>Wilsonian OPE &amp; Renormalons</b>	<b>7</b>
	4.1 First Step to deal with Renormalons . . . . .	7
	4.2 Non-perturbative Renormalons . . . . .	8
<b>5</b>	<b>Describing the CKM Matrix <i>consistently</i></b>	<b>9</b>

---

<sup>1</sup>Invited talk given at the *2018 Epiphany Conference*

<b>6</b>	<b>Schemes of Quark Masses</b>	<b>9</b>
6.1	" $\overline{MS}$ ", "kinetic", "PS" . . . . .	9
6.2	'Pole mass', '1S' . . . . .	10
6.3	Short comments . . . . .	10
<b>7</b>	<b>Duality: Measuring <math> V_{qb} </math> with <math>q = c, u</math></b>	<b>11</b>
<b>8</b>	<b>Many-body Final States for <math>\Delta B \neq 0 \neq \Delta C</math> Hadrons</b>	<b>11</b>
8.1	Tools . . . . .	11
8.2	Probing Dalitz plot for $B^\pm$ . . . . .	13
8.3	<b>CP</b> asymmetries in the decays of beauty baryons . . . . .	15
<b>9</b>	<b>Top quark in the search for ND</b>	<b>16</b>
9.1	<b>CP</b> asymmetries without Higgs dynamics . . . . .	17
9.2	<b>CP</b> asymmetries with <i>on</i> -shell Higgs dynamics . . . . .	17
<b>10</b>	<b>Summary and a New Alliance for the Future</b>	<b>18</b>
<b>11</b>	<b>Personal Epilogue from my week in Krakow</b>	<b>19</b>

---

# 1 Prologue



Figure 1: Painting of 'Epiphany' in a museum in Krakow (picture taken by IIB)

I have truly enjoyed the 2018 Epiphany Conference in Krakow, learnt about fundamental dynamics – and the 'landscapes' of history & art on the true European scale. Very special event happened on January 6 long time ago (see the **Figure 1**): three sages came to meet with the Christ child.

The old center of the city of Krakow is an amazing part of the European culture. I try to show that by pictures I took on Jan. 13, 2018:

- The Barbican in Krakow is just outside of one of the gates on the north wall and very close to the Guesthouse. The Barbican was to protect the city against the ‘barbarians’, see the **Figure 2**. An analogy to use Dalitz plots to probe the impact




Figure 2: Barbican in Krakow (Model)

of New Dynamics (ND): ‘barbarians’ = ‘perturbative QCD’?

- Copernicus was a student in the Jagiellonian University of Krakow that had large impact on the understand of our Universe then; see the **Figure 3**.



Figure 3: Solar system (picture taken by IIB)

- Just south of the Main Market Square one can see a wonderful connection of Renaissance architecture & modern sculpture, see the **Figure 4** – if one can find it inside a building.



Figure 4: Renaissance architecture & modern sculpture (picture taken by IIB)

## 2 Introduction to the ‘Roads’

The Greek word ‘Epiphany’ means: ‘manifestation of a divine being an intuitive grasp of reality through something both simple and striking’ ! I have always been a fan of local Super-Symmetry & still am; however, we are in a different situation: it is *neither* simple *nor* striking. As I will discuss here in some details: best ‘fitted’ analyses of the data do *not* give us the best information about the underlying dynamics – it is crucial to use correlations with other data & judgments! Furthermore I can admire the courage of the young physicists to deal with the challenges in our world on different levels, while listen also to the talks of ‘mature’ colleague like Danish Buras and Swiss Jegerlehner. To make progress we have to discuss the disagreements. Fashion does *not* help us to go closer to our goals as my Italian colleague Augusto said at the conference.

First I will general comments including disagreements I have with some speakers here; some of obvious, while others are more subtle. If a reader finds it interesting (I hope), she/he wants details (with many references): N. Uraltsev: ”The Heavy Quark Expansion”, CRAD96 in Krakow, Acta Phys.Polon.B28(1997)755 [1]; N. Uraltsev: ”Topics in the Heavy Quark Expansion” in 2000 [2]. Somebody might think it is ‘old stuff’; however in my view it is still up-to-data of our understanding of fundamental forces.

One might think the choice of words is in the details: HQE vs. HQET. The titles are: HQE = ”Heavy Quark Expansion” vs. HQET = ”Heavy Quark Effective Theory”; in the latter item I want to mention that the applications of HQET in local QCD vs.

Lattice QCD are different, and I have less problems in the second than the first one. The differences go much deeper in their ‘meaning’.

The usual HQET papers claim to show the impact of non-perturbative physics:

$$\text{”observable”} = \text{perturbative forces} + \text{non-perturbative forces} \quad (1)$$

Instead Kolya Uraltsev (& collaborators like Shifman & me [3]) pointed out that is much deeper to describe the situations by

$$\text{”observable”} = \text{short-distance dynamics} + \text{long-distance dynamics} \quad (2)$$

Crucial statements in my view:

- It is not enough to say that OPE is an important theoretical tool: it is the *Wilsonian* OPE. The separation of short- vs. long-distances dynamics is scale *dependent* around 1 GeV for QCD. One might think it is a bad idea and gives more work without better understanding of the underlying dynamics. However, I will explain why I disagree with such a ‘feeling’.
- What the left hand does does not matter what the right hand does? No – perturbative & non-perturbative QCD effects have to be treated *simultaneously* with accuracy; furthermore we have to think about the correlations with experimental analyses.

These will be discussed with some details or some examples.

General comments:

- Anomalies – ”deep” or not so far
- Wilsonian Operator Product Expansion
- Infrared renormalon with non-perturbative QCD

Items with some details:

1. Consistent parameterization of the CKM matrix
2. Definition of quark masses: ” $\overline{MS}$ ”, ”kinetic”, ”PS”, ‘1S’, ‘pole mass’
3.  $V_{qb}$  [q=c,u]: exclusive vs. inclusive rates and duality
4. Broken U- & V-spin symmetries
5. 3- & 4-body final states in beauty & charm mesons
6. Challenges for understanding weak decays of beauty & charm baryons
7. The stage of top quarks - in a search for New Dynamics
8. Collaboration of HEP & MEP/Hadrodynamics

For some of these Sections I have very short comments, while for others I give some discussions with more references. In a talk at a conference like this one can only to ‘paint the landscape’, but not beyond. For that one has to go to a summer (or winter) schools.

### 3 Anomalies: "deep" or not so

The word ‘anomaly’ is often used in the literature – in particular, when one looks for the impact of New Dynamics (ND). It is easier to discuss exclusive semi-leptonic transitions. However, the situation is ‘complex’.

There is a "quantum anomaly": a classical symmetry is no longer conserved, once one-loop corrections are included. In this well-known case of chiral invariance: for massless quarks we have a "triangle anomaly", since it is produced by a diagram with a triangular fermion loop – or called the "Adler-Bardeen-Bell-Jackiw anomaly":

$$\partial_\mu J_\mu^{(5)} = \frac{\alpha_S}{8\pi} \tilde{G} \cdot G( + m_q \bar{\psi} \gamma_5 \psi ) ; \quad (3)$$

that is not renormalizable in  $4 - \epsilon$  dimensions. The SM ‘deals’ with that by connecting the world of quarks & charged leptons (i.e., 3 colors of quarks) <sup>2</sup>.

Our community has found ‘anomalies’ in previous & present data, namely the differences between expectations from the SM vs. measured data as a sign of the impact of ND. Even in my view it is not just a fashionable one; we have to work & think about semi-leptonic transitions in beauty hadrons with several examples like  $B \rightarrow K^* l^+ l^-$  &  $\Lambda_b^0 \rightarrow \Lambda l^+ l^-$ . One discusses (tiny) rates & the landscape in  $M_{l^+ l^-}$ . Present data show *more* events than expected with  $3 \sigma$  uncertainties. Of course, I am not surprised that our colleagues are waiting impatiently to reach  $5 \sigma$  uncertainties or more.

Allow me to give another lesson in the history: after losing the 1811 battle of Albuera in Spain Marechal Soult said: ‘I had beaten the British – it was just they did not know when they were beaten.’ He was right on both counts. To ‘battle with the British’ there is an analogue to probe the SM & its limitations: HEP theorists start with a penguin operator  $b \rightarrow s$  to describe the transitions of  $B \rightarrow l^+ l^- X_s$  as  $[b\bar{q}] \rightarrow l^+ l^- s \dots \bar{q}$ . In the worlds of hadrons one can measure the final states with  $K\pi$ 's,  $2K\bar{K}$  etc. It makes it in steps:  $K$ ,  $K^*$ , broad resonance  $\kappa$ , in general  $K\pi$ 's,  $2K\bar{K}$  etc. The question is: with which certainties can one describe the connection in the world of quarks & gluons with that of hadrons, namely the "duality". I want to pointed out that duality is *not* an additional assumption. Duality is well-defined in the deep Euclidean region thus avoiding proximity to singularities, cuts induced by hadronic thresholds etc.; then one analytically continues it into the Minkowskian domain. There is a price to be paid for this ‘prize’: in general one cannot apply local duality, but averaged one over an energy interval of around 1 - 1.5 GeV. Furthermore it is not a mathematical statement: we understand the source of the underlying dynamics; it needs some judgment where & how to apply duality in the world of current quarks & gluons:

$$[b\bar{q}] \rightarrow l^+ l^- s \dots \bar{q} \Rightarrow l^+ l^- K / K\pi' s / 2K\bar{K} \dots \quad (4)$$

Except that the branch ratios are tiny, the situations are simpler for these transitions: the underlying dynamics can be probed with  $M_{l^+ l^-}$ . The situations are much more ‘complex’,

---

<sup>2</sup>There must be a deep reason for that.

when I discuss non-leptonic weak decays below. In the future one can probe  $b \rightarrow d$ . The good side is that the SM penguin amplitudes suppressed; unfortunately the landscape has much background.

## 4 Wilsonian OPE & Renormalons

Almost all authors invoke OPE – but mostly *without* ”Wilsonian” prescription. One might think it is about bragging right. However, Shifman & collaborators [4] have a long record to emphasize that applying OPE is subtle: the Wilsonian OPE has to stop around 1 GeV, not lower. It is one thing to draw diagrams, while another thing is understand the underlying dynamics, in particular about non-perturbative QCD with some accuracy. I will come back in the next Section about infrared renormalon and later also about the definition of quark masses. Mostly I follow the ‘road’ described by Shifman in the Ref.[4] with more details now and for the future.

### 4.1 First Step to deal with Renormalons

Dyson pointed out in his famous 1952 paper ”Divergences of Perturbation Theory in QED” [5] that amplitudes *cannot* be convergent. Later it was realized perturbative series in a QFT are *factorially divergent* like  $Z = \sum_k C_k \alpha^k k^{b-1} A^{-k} k!$  with  $k \gg 1$  is the number of loops,  $C_k$ ’s are numerical coefficients of order one, and  $b$  &  $A$  are numbers. It is traced back to the factorially large number of multi-loop Feynman diagrams. The features responsible for the renormalon factorial divergence is the logarithmic running of the effective coupling constant.

Instead of *asymptotic* series one can introduce a Borel transform

$$B_Z = \sum_k C_k \alpha^k k^{b-1} A^{-k} ; \quad (5)$$

the singularity of  $B_Z(\alpha)$  closest to the origin of the  $\alpha$  plain is at a distance  $A$ , and thus  $B_Z(\alpha)$  is *convergent*. One recovers the original function  $Z$  by

$$Z(\alpha) = \int_0^\infty dt e^{-t} B_Z(\alpha t) \quad (6)$$

The integral representation is well-defined provided that  $B_Z(\alpha)$  has *no* singularities on the real positive semi-axis in the complex  $\alpha$  plane. That is not a problem for QED. For other weak couplings it is not trivial, but one can deal with that.

If  $B_Z(\alpha)$  has a singularity on the real positive semi-axis – like coefficients  $C_k$  are all positive or all negative – the integrated in the Eq.(6) become *ambiguous*. This ambiguity is of the order of  $e^{-A/\alpha}$ ; more information is needed from the underlying dynamics. The question comes from QCD with

$$\alpha_S(Q^2) \simeq \frac{\alpha_S(\mu^2)}{1 - \frac{\beta_0 \alpha_S(\mu^2)}{4\pi} \log(\mu^2/Q^2)} = \frac{\alpha_S(\mu^2)}{1 + \frac{\beta_0 \alpha_S(\mu^2)}{4\pi} \log(Q^2/\mu^2)}, \quad \beta_0 = 11 - \frac{2}{3} N_f ; \quad (7)$$

the energy scale  $\mu$  is used to calibrate  $\alpha_S(Q^2)$ . The good side is: at large scales the strong couplings go down to zero with  $Q^2/\mu^2$  (on the log scale) – i.e. ”asymptotic freedom”.

On the other hand, there is a true challenge. With  $\mu^2 \gg Q^2$   $\alpha_S(Q^2)$  gets larger and larger; thus QCD gives us true strong forces at low scales. First one might say it goes to infinite, but that is too naive. One has to stop at  $\mu \sim 1$  GeV based on perturbative QCD.

## 4.2 Non-perturbative Renormalons

It was pointed out first in 1994 that the pole mass is *not* well-defined at the non-perturbative level [6, 7]. Furthermore a rather powerful renormalon-based tool was suggested for evaluating the corresponding non-perturbative contribution [4]. Pole mass is sensitive to large distance dynamics, although this fact is not obvious in perturbative calculations. IR contributions lead to an *intrinsic uncertainty* in the pole mass of order  $\Lambda$  – i.e., a  $\Lambda/m_Q$  power correction. It comes from the factorial growth of the high order terms in the  $\alpha_S$  expansion corresponding to a singularity residing at the  $2\pi/\beta_0$  in the Borel plane. Thus one cannot say it is a correction.

Actually, there are two renormalon-based tools, namely ultraviolet (UV) and infrared (IR) dynamics <sup>3</sup>. One has to include non-perturbative QCD with IR one. Those give contribution to  $b$  quark mass numerically [3], see the **Figure 5**:

$$\begin{aligned}
 m_b^{\text{pole}} &= m_b(1 \text{ GeV}) + \delta m_{\text{pert}}(\leq 1 \text{ GeV}) \simeq \\
 &\simeq 4.55 \text{ GeV} + 0.25 \text{ GeV} + 0.22 \text{ GeV} + 0.38 \text{ GeV} + 1 \text{ GeV} + 3.3 \text{ GeV} \dots, \quad (8)
 \end{aligned}$$

where  $\delta m_{\text{pert}}(\leq 1 \text{ GeV})$  is the perturbative series taking account of the loop momenta down to zero.




Figure 5: Perturbative diagrams leading to the IR renormalon uncertainty in  $m_Q^{\text{pole}}$  of order  $\bar{\Lambda}$ . The number of bubble insertions in the gluon propagator is arbitrary. The horizontal line at the bottom is the heavy quark Green’s function.

Top quarks decays before they have produce top hadrons. Still they carry unbroken color symmetry and thus find partners with color to produce hadrons with color zero in the FS. I will come back to that below.

<sup>3</sup>”All animals are equal, but some animals are more equal than others!” G. Orwell, ‘Animal Farm’.



## 5 Describing the CKM Matrix *consistently*

Wolfenstein's parameterization was very smart, easily usable & well-known. The SM with three families of quarks describes the CKM matrix with four parameters, namely  $\lambda$ ,  $A$ ,  $\bar{\rho}$  &  $\bar{\eta}$ . One uses expansion of the Cabibbo angle  $\lambda = \sin\theta_C \simeq 0.223$ , while  $A$ ,  $\bar{\rho}$  and  $\bar{\eta}$  should be of the order of unity [8]. It is an important item (in particular about finding the impact of ND), but a subtle one: what does one mean by 'maximal' **CP** violation? In principle 100 % asymmetry is possible: I give just three example:  $\bar{\rho} \sim 1$  &  $\bar{\eta} \sim -1$ ;  $\bar{\rho} \sim -1$  &  $\bar{\eta} \sim -0.5$ ;  $\bar{\rho} \sim -0.5$  &  $\bar{\eta} \sim -0.3$ .

Measured values are  $A \simeq 0.82$  as assumed. However, *measured*  $\bar{\eta} \sim 0.35$  &  $\bar{\rho} \simeq 0.14$ , which are not close to unity; thus we have not real control over systematic uncertainties here.

The SM produces at least the leading source of **CP** violation in  $K_L \rightarrow 2\pi$  and  $B$  decays with good accuracy. Searching for ND we need even precision and to measure the correlations with other FS's. The landscape of the CKM matrix is more subtle as pointed out through  $\mathcal{O}(\lambda^6)$  consistently [9]:

$$V_{\text{CKM}} \simeq \begin{pmatrix} 1 - \frac{\lambda^2}{2} - \frac{\lambda^4}{8} - \frac{\lambda^6}{16}, & \lambda, & \bar{h}\lambda^4 e^{-i\delta_{\text{QM}}}, \\ -\lambda + \frac{\lambda^5}{2}f^2, & 1 - \frac{\lambda^2}{2} - \frac{\lambda^4}{8}(1 + 4f^2) - f\bar{h}\lambda^5 e^{i\delta_{\text{QM}}} + \frac{\lambda^6}{16}(4f^2 - 4\bar{h}^2 - 1), & f\lambda^2 + \bar{h}\lambda^3 e^{-i\delta_{\text{QM}}} - \frac{\lambda^5}{2}\bar{h}e^{-i\delta_{\text{QM}}}, \\ f\lambda^3, & -f\lambda^2 - \bar{h}\lambda^3 e^{i\delta_{\text{QM}}} + \frac{\lambda^4}{2}f + \frac{\lambda^6}{8}f, & 1 - \frac{\lambda^4}{2}f^2 - f\bar{h}\lambda^5 e^{-i\delta_{\text{QM}}} - \frac{\lambda^6}{2}\bar{h}^2 \end{pmatrix} \quad (9)$$

with  $\bar{h} \simeq 1.35$ ,  $f \simeq 0.75$  &  $\delta_{\text{QM}} \sim 90^\circ$  and only expansion in  $\lambda \simeq 0.223$ . The pattern in flavor dynamics is less obvious for **CP** violation in hadron decays as stated before [10]; the situation has changed: we have to measure the correlations between four triangles, not focus only on the 'golden triangle'. Some of the important points are emphasized:

- (a) The maximal SM value of  $S(B^0 \rightarrow J/\psi K_S)$  for indirect **CP** violation is  $\sim 0.74$ .
- (b) For  $S(B_s^0 \rightarrow J/\psi\phi)$  is  $\sim 0.03 - 0.05$ .
- (c) The SM gives basically zero **CP** value for doubly Cabibbo suppressed transitions. One has to measure accurately the correlations with several triangles.

## 6 Schemes of Quark Masses

Quark masses are *not* observables in general. Therefore I use the word of 'Schemes'.

### 6.1 "MS", "kinetic", "PS"

$\overline{\text{MS}}$  mass  $\bar{m}_Q(m_Q)$  stands for 'modified minimal subtraction scheme'. It represents a quantity of computational convenience, in particular when calculating perturbative contributions in "dimensional regularization"<sup>4</sup>. For  $\mu \geq m_Q$  it basically coincides with the

<sup>4</sup>It does not necessarily mean we understand the underlying dynamics.

running mass in the Lagrangian and is best normalized at  $\mu \sim m_Q$ . It is appropriate for describing heavy-flavor *production* like  $Z^0 \rightarrow \bar{b}b$  and now also  $H \rightarrow \bar{b}b$ . However, it diverges logarithmically for  $\mu \rightarrow 0$ .

The "kinetic" mass of the heavy quark is regular in the infrared regime including a non-leading source [7, 11, 12, 13]:  $\frac{dm_Q^{\text{kin}}(\mu)}{d\mu} = -\frac{16}{9} \frac{\alpha_S}{\pi} - \frac{4}{3} \frac{\alpha_S}{\pi} \frac{\mu}{m_Q} + \mathcal{O}(\alpha_S^2)$ . For  $b$  quarks  $\mu \sim 1$  GeV is the best scale to describe their weak decays<sup>5</sup>. Using  $\mu \sim m_b$  instead, it leads to higher-order perturbative corrections that are artificially large, for which one has *no* control [12].

"PS" = "potential-subtracted": the schemes "kinetic" and "PS" are quite different already on the conception level; technical problems of "PS" arise at  $\mathcal{O}(\alpha_S^4)$ . Still they are in the same 'division' of fundamental physics. I will come back to this point below about top quarks.

## 6.2 'Pole mass', '1S'

A pole mass for quarks is gauge independent and infrared stable in perturbative QCD; furthermore it is easy to apply pole mass in Feynman graphs. However, it is *not* infrared stable non-perturbatively. Make the same statement with different words: pole mass depends on long distance dynamics, for what we have little control.

Recent PDG reviews basically ignore the "kinetic" scheme, while focus on the '1S' scheme based on  $m_b^{1S} \simeq M_{\Upsilon(1S)}/2$ <sup>6</sup>. It claims these schemes give us the same information about underlying dynamics. However, it is incorrect, as Uraltsev pointed out [14]:  $m_b^{1S} = m_b^{\text{pole}}[1 - C_F^2(\alpha_S^2/8) + \mathcal{O}(\alpha_S^3, \beta_0\alpha_S^3\log\alpha_S)]$  - i.e., also  $m_b^{1S}$  is *not* well-defined at the *non-perturbative* level.

## 6.3 Short comments

Flavor dynamics is 'complex'. At a conference the goal is to 'paint' the landscape, but not to discuss the details. However, it is important to give short, but subtle comments. I give a reference to an important (& large) 2001 paper [15]. My main disagreements with A. Pineda: his Abstract does not mention some of his important results. However, a careful reader can find it on page 16: (a) "... it is achieved by the threshold scheme, i.e. the *kinetic*, the PS-like, the 1S ...". I would say the meaning of 'threshold' is not obvious. When one talks about  $b \rightarrow c$  [ $W_{\text{off-shell}}^-$ ], it means to get one or two charm quarks. However, the situations are quite different for  $b \rightarrow u$  [ $W_{\text{off-shell}}^-$ ]. (b) "Note also that the 1S and PS schemes depend on  $\nu_{us}$ ." At three-loops diagrams the ultrasoft scale appears in the static potential and the heavy quarkonium mass. Again, the situations are quite different for the impact of perturbative QCD vs. non-perturbative one.

---

<sup>5</sup>A reader might think, my judgment is 'biased'; however, I stay by my statement.

<sup>6</sup>due to 'par ordre du Mufti' (= no right of appeal).

## 7 Duality: Measuring $|V_{qb}|$ with $q = c, u$

The item of "duality" is referred to a very complex situations, namely the connections of the worlds of hadrons vs. quark & gluons. In this Section I give short comments at the very specific case: compare the values of  $|V_{cb}|$  and  $|V_{ub}|$  from inclusive vs. exclusive semi-leptonic amplitudes.

It seems the difference between the  $|V_{cb}|_{\text{incl.}}$  vs.  $|V_{cb}|_{\text{excl.}}$  has become smaller now based on realistic theoretical uncertainties, mostly due to LQCD analyses.

On the other hand, the difference between  $|V_{ub}|_{\text{incl.}}$  vs.  $|V_{ub}|_{\text{excl.}}$  has not changed. It has been pointed out that the values of  $|V_{ub}|_{\text{incl.}}$  based on the data from  $B \rightarrow l\nu\pi$ 's, while assuming that  $B \rightarrow l\nu\bar{K}K\dots$  are irrelevant due to a traditional understand duality. It is a good assumption – but *local* duality does not work close to thresholds. Maybe the real  $|V_{ub}|_{\text{incl.}}$  are smaller and thus solve that challenge. LHCb experiment cannot measure inclusive rates. However, it might go able to go after the rates of  $B^+ \rightarrow l^+\nu K^+K^-$  and  $B^0 \rightarrow l^+\nu K^+K^-\pi^-$  with non-zero values. Furthermore Belle II should measure values there or limits.

## 8 Many-body Final States for $\Delta B \neq 0 \neq \Delta C$ Hadrons

Indirect **CP** violation has been established in  $K_L \rightarrow 2\pi$  &  $B^0 \rightarrow J/\psi K_S$ . On the other hand, the landscapes are much more 'complex' as expected, since direct **CP** asymmetries depend on final state interactions:

$$|T(\bar{P} \rightarrow \bar{a})|^2 - |T(P \rightarrow a)|^2 = 4 \sum_{a_j, a} T_{a_j, a}^{\text{resc}} \text{Im} T_a^* T_{a_j}; \quad (10)$$

*without* non-zero re-scattering direct **CP** asymmetries cannot happen, even if there are weak phases [16, 17, 18, 19]. One expects large impact of strong re-scattering, and the LHCb data of suppressed  $B \rightarrow 3$  mesons have shown that; I will discuss below. It is obvious that the crucial information about the underlying dynamics cannot be found in two-body FS. Even so, it is a very good hunting region for the impact of ND, since they can depend only one ND amplitude.

### 8.1 Tools

One has to think about the tools that can be applied. Not surprisingly, it comes to your mind, namely symmetries broken or not.

- One can apply  $SU(3)_{\text{light flavor}}$  (not  $SU(3)_{\text{color}}$ ). The global  $SU(3)_{\text{light flavor}}$  is broken. It was pointed out by Lipkin, it helps the thinking by using 3  $SU(2)$ : one combines  $(u, d)$  quarks for I-spin, while  $s \rightleftharpoons d$  for U-spin and  $s \rightleftharpoons u$  for V-spin symmetries.

Broken U-spin symmetry *without* V-spin is usable for spectroscopy with a good record. Yet the situation is quite different for weak transitions. I give one example

from the PDG2017 data **CP** asymmetry:

$$A_{\text{CP}}(B^0 \rightarrow K^+\pi^-) = -0.082 \pm 0.006 \quad (11)$$

(In 1987 Sanda & I had given a prediction:  $A_{\text{CP}}(B^0 \rightarrow K^+\pi^-) \sim -0.1$ .) It shows the impact of Penguin diagrams – but (semi-)quantitatively. Then looks at the PDG2017 data:

$$A_{\text{CP}}(B_s^0 \rightarrow \pi^+K^-) = +0.26 \pm 0.04 . \quad (12)$$

Can we predict this connection?

It had been suggested by Lipkin in 2005 [20] to use U-spin symmetry <sup>7</sup>:

$$\Delta = \frac{A_{\text{CP}}(B^0 \rightarrow K^+\pi^-)}{A_{\text{CP}}(B_s^0 \rightarrow \pi^+K^-)} + \frac{\Gamma(B_s^0 \rightarrow \pi^+K^-)}{\Gamma(B^0 \rightarrow K^+\pi^-)} = 0 ; \quad (13)$$

The LHCb collab. had published a short 2013 paper [21]:

$$\Delta_{\text{LHCb}} = -0.02 \pm 0.05 \pm 0.04 \quad (14)$$

saying: "These results allow a stringent test of the validity of the ...". I disagree with this statement for several reasons! First examples from two-body FS:

- Indeed, the value of  $\Delta_{\text{LHCb}}$  is consistent with zero.
- Yet, it is also consistent with a value  $\sim 0.1$  expected for direct **CP** asymmetry for two-body FS.
- One has to think about correlations of U-spin symmetry with V-spin one due to re-scattering. What about  $B^0 \rightarrow K^0\pi^0/K^0\eta$  &  $B_s^0 \rightarrow \pi^0K^0/\eta K^0$ ? One has to remember that these transitions are affected by oscillations & indirect **CP** violation.
- One can look at the situation with two-body FS of  $B^+$  decays:

$$A_{\text{CP}}(B^+ \rightarrow K_S\pi^+) = -0.017 \pm 0.016 \quad (15)$$

$$A_{\text{CP}}(B^+ \rightarrow K^+\pi^0) = +0.037 \pm 0.021 \quad (16)$$

$$A_{\text{CP}}(B^+ \rightarrow K^+\eta') = +0.004 \pm 0.011 \quad (17)$$

with no sign of **CP** asymmetry, while it was found in

$$A_{\text{CP}}(B^+ \rightarrow K^+\eta) = -0.37 \pm 0.08 . \quad (18)$$

It shows the impact of the strong re-scattering. There are two lessons: difference between U- & V-spin is ‘fuzzy’ due to re-scattering – and we have to go *beyond* two-body FS.

---

<sup>7</sup>The positive sign in the Eq.(13) is not surprising in the SM.

- Probing FS in non-leptonic decays with two hadrons (including narrow resonances) is not trivial to measure **CP** violations, if one has enough data for suppressed transitions; theorists can ‘predict’ those & analyze the data. On the other hand, one gets ‘just’ numbers. We have to remember that two-body FS of suppressed non-leptonic weak decays are a small part of charm mesons & tiny ones for beauty mesons; data show that – it is not surprising. Three- & four-body FS are described by two & more dimensional plots. There is a price: lots of work for experimenters and theorists. There is also a prize: to find the existence of ND and also its features.

The situations are very different for strange hadrons with  $\Delta S = 1$  & 2 as you listened to my Danish colleague Buras (member of the Bavarian Academy!): it is produced by *local* operators and FS with only one & two pions.

## 8.2 Probing Dalitz plot for $B^\pm$

The data of CKM suppressed  $B^+$  decays show no surprising rates for  $B^+ \rightarrow K^+\pi^-\pi^+$  &  $B^+ \rightarrow K^+K^-K^+$  and  $B^+ \rightarrow \pi^+\pi^-\pi^+$  &  $B^+ \rightarrow \pi^+K^-K^+$ .

LHCb data from run-1 show averaged direct **CP** asymmetries [22]:

$$\begin{aligned} \Delta A_{CP}(B^\pm \rightarrow K^\pm\pi^+\pi^-) &= +0.032 \pm 0.008_{\text{stat}} \pm 0.004_{\text{syst}} \pm 0.007_{\psi K^\pm} \\ \Delta A_{CP}(B^\pm \rightarrow K^\pm K^+K^-) &= -0.043 \pm 0.009_{\text{stat}} \pm 0.003_{\text{syst}} \pm 0.007_{\psi K^\pm} \end{aligned} \quad (19)$$

with  $2.8 \sigma$  &  $3.7 \sigma$  from zero. Based on our experience with the impact of penguin diagrams on the best measured  $B^0 \rightarrow K^+\pi^-$ , the sizes of these averaged asymmetries are not surprising; however it does not mean that we could really predict them. It is very interesting that they come with opposite signs due to **CPT** invariance.

LHCb data show *regional* **CP** asymmetries [22]:

$$\begin{aligned} A_{CP}(B^\pm \rightarrow K^\pm\pi^+\pi^-)|_{\text{regional}} &= +0.678 \pm 0.078_{\text{stat}} \pm 0.032_{\text{syst}} \pm 0.007_{\psi K^\pm} \\ A_{CP}(B^\pm \rightarrow K^\pm K^+K^-)|_{\text{regional}} &= -0.226 \pm 0.020_{\text{stat}} \pm 0.004_{\text{syst}} \pm 0.007_{\psi K^\pm} . \end{aligned} \quad (20)$$

”Regional” **CP** asymmetries are defined by the LHCb collaboration: positive asymmetry at low  $m_{\pi^+\pi^-}$  just below  $m_{\rho^0}$ ; negative asymmetry both at low and high  $m_{K^+K^-}$  values. One should note again the opposite signs in Eqs.(20). It is not surprising that ”regional” asymmetries are very different from averaged ones. Even when one uses states only from the SM –  $SU(3)_C \times SU(2)_L \times U(1)$  – one expects that; it shows the *impact of re-scattering* due to  $SU(3)_C$  (actually  $SU(3)_C \times \text{QED}$ ) in general. Of course, our community needs more data, but that is not enough. There are important questions and/or statements:

- How do we *define* regional asymmetries and probe them on the experimental and theoretical sides?
- Can it show the impact of broad resonances like  $f_0(500)/\sigma$  and  $K^*(800)/\kappa$ ?
- Again, the best fitted analyses often do not give us the best understanding of the underlying fundamental dynamics.

LHCb data from the run-1 show *larger* averaged **CP** asymmetries as discussed above in Eqs.(19) (again with the opposite signs):

$$\begin{aligned}\Delta A_{CP}(B^\pm \rightarrow \pi^\pm \pi^+ \pi^-) &= +0.117 \pm 0.021_{\text{stat}} \pm 0.009_{\text{syst}} \pm 0.007_{\psi K^\pm} \\ \Delta A_{CP}(B^\pm \rightarrow \pi^\pm K^+ K^-) &= -0.141 \pm 0.040_{\text{stat}} \pm 0.018_{\text{syst}} \pm 0.007_{\psi K^\pm} .\end{aligned}\quad (21)$$

It is interesting already with the averaged ones, since  $b \implies d$  penguin diagrams are more suppressed than  $b \implies s$  ones. Again **CP** asymmetries focus on small regions in the Dalitz plots [22].

$$\begin{aligned}\Delta A_{CP}(B^\pm \rightarrow \pi^\pm \pi^+ \pi^-)|_{\text{regional}} &= +0.584 \pm 0.082_{\text{stat}} \pm 0.027_{\text{syst}} \pm 0.007_{\psi K^\pm} \\ \Delta A_{CP}(B^\pm \rightarrow \pi^\pm K^+ K^-)|_{\text{regional}} &= -0.648 \pm 0.070_{\text{stat}} \pm 0.013_{\text{syst}} \pm 0.007_{\psi K^\pm} .\end{aligned}\quad (22)$$

Again, it should be noted also the signs in Eqs.(21) & Eqs.(22). Does it show the impact of broad scalar resonances like  $f_0(500)/\sigma$  and/or  $K^*(800)/\kappa$ ?

First one analyzes the data using model-independent techniques [23], compares them and discuss the results – but that is not the end of our ‘traveling’. Well-known tools like dispersion relations are ‘waiting’ to apply – but we have to do it with some ‘judgement’. I had visited a museum in the north Wall of Krakow and looked at this painting, see the **Figure 6**: I was very happy to see it again – but after more looking at that, I realized



Figure 6: ‘Lady with an Ermine’ by Leonardo da Vinci (picture taken by IIB)

I did *not* see the ‘real’ paint. It has colors, but *pale* ones. The real paint with wonderful colors is still in its original part of the museum just a very steps behind this Wall, but it is closed for a year. It gives us an idea about the painting of Leonardo da Vinci, but not beyond.

Coming back to fundamental physics: one has to be prepared for analyses of Dalitz plots (& beyond); first one has to produce simulations to see both the strong & weak features for hardware & software of a detector. Yet that is *not* the final step. The best fitted analyses often do not give us the best information about the underlying dynamics. Final steps need judgment based on correlations with other data applying resonances, threshold enhancements etc. with dispersion relations & other refined tools.

### 8.3 CP asymmetries in the decays of beauty baryons

Before I had suggested to probe Dalitz plots of  $\Lambda_b^0 \rightarrow \Lambda\pi^+\pi^-/\Lambda K^+K^-/\Lambda D^-\pi^+$  and  $\Xi_b^0 \rightarrow \Lambda\pi^+\pi^-/\Lambda K^+K^-$  that do not depend on production asymmetries [19].

However, at the ICHEP2016 conference in Chicago that the LHCb collaboration showed data with evidence for **CP** asymmetry in  $\Lambda_b^0 \rightarrow p\pi^-\pi^+\pi^-$  with a novel idea. It is discussed in [24] with details. In  $pp$  collisions one gets different numbers of  $\Lambda_b^0$  vs.  $\bar{\Lambda}_b^0$  due to production asymmetries. Therefore one focuses on **T**-odd moments. LHCb measured the angle between two planes: in the rest frame of  $\Lambda_b^0$  one plane is defined by  $[\vec{p} \times \vec{\pi}_{\text{fast}}^-]$ , while the other one by  $[\vec{\pi}^+ \times \vec{\pi}_{\text{slow}}^-]$ ; likewise for  $\bar{\Lambda}_b^0$ . They found evidence for **CP** asymmetry on the level of  $3.3\sigma$  based in its run-1 of  $3\text{ fb}^{-1}$ . Actually, they found *regional CP* asymmetry  $\sim 20\%$  without saying that clearly. In principle it is not surprising due to strong dynamics with  $\Delta(1232)[\Delta(1600)/\Delta(1620)] \Rightarrow p\pi^-$ . We should keep in mind the situations should be affected by different broad resonances, thresholds etc.

Are we lucky to find this effect and its size? Of course, we need more data. Yet, the *present* data can give us more information about the underlying dynamics by measuring the angle between two different planes: one is defined by  $[\vec{p} \times \vec{\pi}_{\text{slow}}^-]$ , while the other one  $[\vec{\pi}^+ \times \vec{\pi}_{\text{fast}}^-]$ . Can we find **CP** symmetries, too? *Regional* ones, where and its size?

The data are very interesting for several reasons:

- Maybe **CP** asymmetry was found in a decay of a baryon for the first time (except ‘our existence’); it is for a beauty baryon.
- It is another example that many-body FS are *not* a background for the information our community got it from two-body FS.
- The plot given at the ICHEP2016 shows the strength of regional **T** asymmetry around  $20 \times 10^{-2}$ . Very interesting, but we cannot claim to understand the underlying dynamics – yet! Furthermore in the world of quarks & gluons one looks at CKM penguin of  $b \rightarrow d$ , where one expects less than for  $b \rightarrow s$ . LHCb data already shown similar lessons for **CP** asymmetries in  $B^+ \rightarrow \pi^+\pi^+\pi^-/\pi^+K^+K^-$  vs.  $B^+ \rightarrow K^+\pi^+\pi^-/K^+K^+K^-$ , see Eqs.(19,20,21,22) just above.
- LHCb collaboration did not get enough data from run-1 to probe  $\Lambda_b^0 \rightarrow p\pi^-K^+K^-$  &  $\bar{\Lambda}_b^0 \rightarrow \bar{p}\pi^-K^+K^-$ . It will change very ‘soon’.
- Furthermore, LHCb collab. can measure rates and **CP** ”regional” asymmetries in  $\Lambda_b^0 \rightarrow pK^-\pi^+\pi^-$  and  $\Lambda_b^0 \rightarrow pK^-K^+K^-$  ‘soon’ – and has no competition from other experiments. First we have to discuss  $\Lambda_b^0 \rightarrow p\pi^-\pi^+\pi^-$  &  $\Lambda_b^0 \rightarrow p\pi^-K^+K^-$  and  $\Lambda_b^0 \rightarrow pK^-\pi^+\pi^-$  &  $\Lambda_b^0 \rightarrow pK^-K^+K^-$ . Will they follow the same ‘landscape’ for  $B^+ \rightarrow \pi^+\pi^+\pi^-/\pi^+K^+K^-$  vs.  $B^+ \rightarrow K^+\pi^+\pi^-/K^+K^+K^-$  as discussed above qualitatively or not? So say it with different words: will they show the strengths of ‘penguin diagrams’ in  $\Lambda_b^0$  decays or not? The situations are similar for beauty mesons and beauty baryons or only on the qualitatively way?

## 9 Top quark in the search for ND

The landscape of top quark dynamics is very different from  $\Delta B \neq 0 \neq \Delta C$ , as I had ‘painted’ it, see the **Figure 6** above. To find its direct impact, power is not enough – we have to think, see the **Figure 7**.



Figure 7: Greek Goddess Athena

My suggestion is that the 2020 Epiphany Conference in Krakow can mostly focus on describing the landscape of top quark dynamics, namely the productions of a pair & single top quarks and forward vs. center regions of  $pp$  collisions; furthermore one discusses **CP** asymmetries together  $W^\pm$  &  $Z^0$  &  $H^0$ , where one hardly get background from the SM. Finally one discusses the future with new technologies for collisions & detectors. The ‘future’ is defined the time schedule of  $\sim 30$  years, which is beyond my personal ‘horizon’. I give you only three references coming from the 10<sup>th</sup> International Workshop on Top Quark Physics Braga, Portugal, 2017. Unfortunately (for me) I did not attend this Workshop. I disagree with several statements given in these papers I found on the internet; maybe our real disagreements are smaller, since I am unable to follow the discussions there. It will make progress, but it will need a lot of time. My statements below makes my point.

Now the ‘top quark community’ is hitting the ‘Systematics Wall’ in different ways, see the ‘Experimental Summary’ [25] <sup>8</sup>:

- take ratios – go differential – stop and think.
- Production rates of  $\bar{t}t$  pairs are powerful handles to constraint the parton distribution functions (PDF). It has been suggested that  $\bar{t}t$  rates may be the relative luminometer of the future for LHC and possible future hadron colliders.
- The landscapes for the cross sections of  $\bar{t}tV$  with  $V = W, Z$  have changed with the 2016 data, where statistical uncertainties are smaller than the systematic ones; likewise for  $\bar{t}tt$ : there are possible hunting regions for ND – and even more for *single* (anti-)top quarks [26].

---

<sup>8</sup>He gave a reference to: V.I. Lenin: ‘What Is To Be Done?’



First I make general statements and later examples for special situations. There is a comment about the use of the ‘words’: Nason said in the Abstract in his paper [27]: ‘... calculations interfaced to shower generators (NLO + PS) of increasing accuracy, interfaced to both Pythia8 and Herwig7 Monte Carlo generators.’. The first statement: obviously the meaning of his ‘PS’ is quite different from Beneke’s ‘word’ as I had discussed in the **Sect.6.1**.

The second statement is not so short: Nason and I talk about different worlds.

(a) He focus on the perturbative impact of QCD if only with a short comment claimed ‘the renormalon ambiguity’ is safely below the current experimental errors’, namely the ‘ambiguity’ of 110 MeV or 250 MeV. I quite disagree. One *cannot* ignore he works of Shifman [4], which has an excellent record <sup>9</sup>. There is an important different between perturbative renormalon vs. non-perturbative one. Furthermore, how ‘safe’ we are to depend on Monte Carlo generators?

(b) As I have said above, the ‘pole mass’ is *not* well defined. Using simulations & modeling are one thing (see the **Figure 6** above), while understanding the underlying dynamics are quite another thing. Of course, using pole masses are popular – in particular in experimental papers & analyses –, but it is only the first step, as I had said above. So far, we are not close to ‘precision’ or even ‘accuracy’.

It was said top quarks decay before they can produce top hadrons [28]. Still they carry ”color” based on a local unbroken QFT; thus they can evolve with other ”color” states in connection to produce hadrons without ”color” states in the end. It means that the ‘world’ of simulations is *less* complex than the FS in the real word.

## 9.1 CP asymmetries without Higgs dynamics

One can measure  $pp$  collisions with a pair of  $[\bar{b}W^-]_t[bW^+]_t\dots$  in the center region with  $gg$ ; or forward(backward) region  $qg \rightarrow q\bar{t}t$  with  $q = u, d$ . It is unlikely to find **CP** asymmetries there; on the other hand, we might learn new lessons about very heavy resonances.

Another ‘road’:  $pp \rightarrow [\bar{b}W^-]_t\dots[q'W^+]_t/[\bar{q}'W^-]_t\dots[bW^+]_t$  with  $q' = s, d$ . One might find **CP** asymmetries there; a possible source is an asymmetric Dark Matter.

One can probe **CP** asymmetry with a *single* top quark:  $bg \rightarrow W^-tg \rightarrow W^-[bW^+]_tg$  vs.  $\bar{b}g \rightarrow W^+\bar{t}g \rightarrow W^+[\bar{b}W^-]_tg$ . Again a possible source is an asymmetric Dark Matter.

## 9.2 CP asymmetries with *on-shell* Higgs dynamics

Collisions at LHC have enough energies to produce very often  $pp \rightarrow \bar{t}H^0tX$ ; to use different words: one talks about short distance forces like  $gg \rightarrow \bar{t}H^0t$ . However, can one find these events with a huge background? While I disagree with some statements in these articles, I have to say first I admire the courage of these experimenters that enter this challenge.

---

<sup>9</sup>Of course, I am ‘biased’.

## 10 Summary and a New Alliance for the Future

The ruler of a Greek city in southern Italy once approached the resident sage (Pythagoras) with the request to be educated in mathematics, but in a ‘royal way’, since he was busy with many obligations. Whereupon Pythagoras replied with admirable candor: ‘There is no royal way to mathematics.’

Likewise is there *no* ‘royal insights’ into the inner working of ‘our’ Nature as I try to show first with pictures: power is not enough – we have to think as the **Figure 7** shows.

The painting of Piero della Francesca shows the dream before the crucial battle outside of Rome between Constantine and Maxentius on different dimensions, see **Figure 8**<sup>10</sup>. Kolya Uraltsev & I had looked at this painting in person and realized that it is symbol



Figure 8: The dream in the night *before* the battle by the Milvian Bridge over the Tiber of a true collaboration.

Our community proceeds in steps: first one uses models to describe the data and then model-independent analyses. However, those should not be the final step(s). Often best fitted analyses do not give us the best information about the underlying dynamics. How to do that? We have theoretical tools with a good record like dispersion relations & other refined tools. They are ‘waiting’ – it ‘only’ needs to work with *judgements* and tests it with correlations with other data! Yes, the data are the referees, but in the end – theorists should not be the slaves of the data.

In the previous century we had talked about fundamental physics: Nuclear Physics at low energies, while HEP at high energies; flavor dynamics are part of HEP. In this century one thinks (or should) about Nuclear Physics and MEP and HEP. Probing jets, Higgs & top quarks dynamics and direct SUSY is the ‘job’ for HEP still again. However, the landscape is more complex with many interconnected parts: decays of strange/beauty/charm hadrons, where tools applied to Dalitz plots with dispersion relations etc. We have to go

---

<sup>10</sup>Another example of divine manifestation in the old history?

for accuracy and even precision to find the impact of ND. To make progress, it is crucial to connect the world of hadrons, where MEP applies – or with a better choice of word, namely "hadro-dynamics" – with the world of quarks & gluons, where HEP works; it is highly non-trivial.

## 11 Personal Epilogue from my week in Krakow

In a museum of Krakow that is inside of the north Wall of the old center I have seen a very good Roman sculpture to show the goddess 'Minerva/Athena'. I saw a group of pairs of ladies, where one was blind and the other was a guide: the blind one was allowed to *touch* this sculpture in some details – a wonderful experience!

**Acknowledgments:** This work was supported by the NSF under the grant number PHY-1520966. I truly thank the organizers of the 2018 Epiphany Conference! If a reader of my article has seen the center of Krakow, she/he should understand why I used pictures, not only diagrams.

## References

- [1] N. Uraltsev: "The Heavy Quark Expansion", Symposium on Radiative Corrections (CRAD96) in Krakow, arXiv:hep-ph/9612349, *Acta Phys.Polon.* **B28** (1997) 755.
- [2] N. Uraltsev: "Topics in the Heavy Quark Expansion", arXiv:hep-ph/0010328.
- [3] I. Bigi, M. Shifman, N. Uraltsev, "Aspects of Heavy-Quark Theory", *Annu.Rev.Nucl.Sci.* **47** (1997) 591; arXiv: hep-ph/9703290.
- [4] M. Shifman, "New and Old About Renormalons", arXiv:1310.1966 [hep-th]; contribution to the book: "QCD and Heavy Quarks – In Memoriam Nikolai Uraltsev", World Scientific Publishing Co., 2015; it shows the progress since 1994; M. Shifman, "Quark-hadron duality", Boris Ioffe Festschrift, At the Frontier of Particle Physics, Singapore, World Scientific Publishing Co., 2001; arXiv: hep-ph/0009131.
- [5] F.J. Dyson, *Phys.Rev.* **85** (1952) 631.
- [6] I.I. Bigi, M. Shifman, N.G. Uraltsev, A. Vainshtein, *Phys.Rev.* **D50** (1994) 2234; arXiv: hep-ph/9402360.
- [7] M. Beneke, V. Braun, *Nucl.Phys.* **B426** (1994) 301; arXiv:hep-ph/9402364.
- [8] L. Wolfenstein, *Phys.Rev.Lett.* **51** (1983) 1945.
- [9] Y.H. Ahn, H-Y. Cheng, S. Oh, *Phys.Lett.* **B703** (2011) 571; arXiv: 1106.0935.

- [10] I.I. Bigi, arXiv:1306.6014 [hep-ph], talk given at FPCP2013 "Flavor Physics and CP Violation 2013".
- [11] M. Voloshin, *Phys.Rev.***D46**(1992)3062; N. Uraltsev, *Int.J.Mod.Phys.***A10**(1995)515.
- [12] I. Bigi, M. Shifman, N. Uraltsev, A. Vainshtein, *Phys.Rev.* **D56** (1997) 4017.
- [13] S. Bianco, F.L. Fabbri, D. Benson, I. Bigi, "A Cicerone for the physics of charm", *Riv.Nuovo Cim.***26N7**(2003)1 [arXiv:hep-ex/0309021].
- [14] N. Uraltsev, "Heavy quark expansion in beauty: recent successes and problems", Invited talk at *Continuous Advances in QCD 2004*, Minneapolis; arXiv:hep-ph/0409125.
- [15] A. Pineda, "Determination of the bottom quark mass from the  $\Upsilon(1S)$  system", JHEP 0106 (2001) 022; hep-ph/0105008.
- [16] I.I. Bigi, V.A. Khoze, N.G. Uraltsev, A.I. Sanda, p. 175-248 in "CP Violation", C. Jarlskog (Editor), World Scientific (1988).
- [17] L. Wolfenstein, *Phys.Rev.* **D43** (1991) 151.
- [18] N.G. Uraltsev, Proceedings of DPF-92, arXiv:hep-ph/9212233.
- [19] I.I. Bigi, *Frontiers of Physics* **10**, #3 (June 2015) 101203; arXiv:1503.07719 [hep-ph].
- [20] H. Lipkin, *Phys.Lett.* **B62** (2005) 126; arXiv:hep-ph/0503022.
- [21] R. Aaij *et al.*, LHCb collab., *Phys.Rev.Lett.* **110** (2013) 221601.
- [22] R. Aaij *et al.*, LHCb collab., *Phys.Rev.Lett.* **111** (2013) 101801; R. Aaij *et al.*, LHCb collab., *Phys.Rev.Lett.* **112** (2014) 011801.
- [23] I. Bediaga *et al.*, *Phys.Rev.* **D89** (2014) 074024; arXiv:1401.3310 [hep-ph].
- [24] R. Aaij *et al.*, LHCb collaboration, nature physics, published online: 30 January 2017; arXiv: 1609.05216 [hep-ex].
- [25] A. Giammanco, "TOP2017: Experimental Summary", arXiv:1712.02177 [hep-ex].
- [26] M. Komm,"Measurements of single top quark cross sections at 13 TeV with the CMS experiment", arXiv:1711.11445 [hep-ex].
- [27] P. Nason, "The Top Quark Mass at the LHC", arXiv:1801.04826 [hep-ph].
- [28] I.I. Bigi, Y.L. Dokshitzer, V. A. Khoze, J. H. Kühn, P. M. Zerwas, "Production and Decay Properties of Ultraheavy Quarks", *Phys.Lett.* **B181** (1986) 157.