Hyperfine spectroscopy of hydrogen and antihydrogen in ASACUSA

E. Widmann · C. Amsler · S. Arguedas Cuendis · H. Breuker · M. Diermaier · P. Dupré C. Evans · M. Fleck · A. Gligorova · H. Higaki · Y. Kanai · B. Kolbinger · N. Kuroda · M. Leali A.M.M. Leite · V. Mäckel · C. Malbrunot · V. Mascagna · O. Massiczek · Y. Matsuda D.J. Murtagh · Y. Nagata · A. Nanda · D. Phan · C. Sauerzopf · M.C. Simon · M. Tajima · H. Spitzer · M. Strube · S. Ulmer · L. Venturelli · M. Wiesinger · Y. Yamazaki · J. Zmeskal

Received: date / Accepted: date

Abstract The ASACUSA collaboration at the Antiproton Decelerator of CERN aims at a precise measurement of the antihydrogen ground-state hyperfine structure as a test of the fundamental CPT symmetry. A beam of antihydrogen atoms is formed in a CUSP trap, undergoes Rabi-type spectroscopy and is detected downstream in a dedicated antihydrogen detector. In parallel measurements using a polarized hydrogen beam are being performed to commission the spectroscopy apparatus and to perform measurements of parameters of

C. Malbrunot

CERN, Geneva Switzerland

H. Breuker, P. Dupré, Y. Kanai, V. Mäckel, M. Tajima, S. Ulmer, Y. Yamazaki RIKEN, 351-0198 Saitama, Japan

N. Kuroda, M. Fleck, Y. Matsuda University of Tokyo, 153-8902 Tokyo, Japan *Present address:* of M. Tajima: RIKEN, 351-0198 Saitama, Japan

H. Higaki

Hiroshima University, 739-8530 Hiroshima, Japan

C. Evans, M. Leali, V. Mascagna, L. Venturelli Dipartimento di Ingegneria dell'Informazione, Universita' degli Studi di Brescia, Brescia, Italy and Istituto ` Nazionale di Fisica Nucleare (INFN), sez. Pavia, Italy *Present address:* of V. Mascagna: Dipartimento di Scienza e Alta Tecnologia, Univerita' dell'Insubria and ` INFN sez. di Pavia, Italy

Y. Nagata

Tokyo University of Science, 162-8601 Tokyo, Japan

E. Widmann, C. Amsler, S. Arguedas Cuendis, M. Diermaier, A. Gligorova, B. Kolbinger, A.M.M. Leite, C. Malbrunot, O.Massiczek, D.J. Murtagh, A. Nanda, D. Phan, C. Sauerzopf, M.C. Simon, H. Spitzer, M. Strube, M. Wiesinger, J. Zmeskal

Stefan Meyer Institute for Subatomic Physics, Austrian Academy of Sciences, 1090 Vienna, Austria E-mail: eberhard.widmann@oeaw.ac.at

Present address: of S. Arguedas Cuendis: CERN, Geneva, Switzerland

Present address: of M. Wiesinger: Max Planck Institute for Nuclear Physics, Heidelberg, Germany

the Standard Model Extension (SME). The current status of antihydrogen spectroscopy is reviewed and progress of ASACUSA is presented.

Keywords Antihydrogen · CPT · hyperfine spectroscopy

1 Antihydrogen and CPT symmetry

CPT symmetry is a cornerstone of the Standard Model (SM) of particle physics due the existence of the CPT theorem which states that in any Lorentz-invariant, unitary, local field theory of point-like particles is invariant under the combined operation of charge conjugation C, parity P and time reversal T. The observed matter - antimatter asymmetry in the Universe as well as theories beyond the SM like string theory, where some of the prerequisites of the mathematical proof of the CPT theorem do not hold any more, have encouraged experimental tests of CPT symmetry.

One of the consequences of the CPT theorem is that particles and antiparticles have exactly the same (mass, total lifetime) or exactly the opposite (charge, magnetic moment) properties. An overview of existing measurements is available at the Particle Data Group [\[1\]](#page-5-0), some of the standard measurements of masses of elementary particles and antiparticles are shown in Fig. [1](#page-1-0) (a more elaborate comparison including e.g. *g*-factor measurements as performed by the BASE collaboration to ppb precision [\[6\]](#page-5-1) is possible within the Standard Model Extension (SME) framework [\[7,](#page-5-2) [8,](#page-5-3) [9\]](#page-5-4)).

From Fig. [1](#page-1-0) it becomes evident that atomic physics measurements like the cyclotron frequency of protons and antiprotons in a trap yielding the charge-to-mass ratio *q*/*m* and determinations of the internal structure of antihydrogen offer the most sensitive tests of CPT symmetry on an absolute scale. The potential of antihydrogen, the simplest anti-atom whose matter equivalent hydrogen is one of the best studied systems in physics, for precision CPT tests has been realized early [\[10\]](#page-5-5), the most difficult challenge being the formation of cold

Fig. 1 Comparison of several tests of CPT symmetry on an energy scale. Bar's right hand side: measured quantity, length of bar: relative precision of CPT test, left hand side: sensitivity on an absolute energy scale. Blue: existing test. Orange: predicted sensitivity if existing precision for hydrogen is achieved. In the case of HFS: orange: first goal for in-beam measurement, paler orange: line width for fountain, yellow: hydrogen maser result. In case of Lamb shift: orange: estimated achievable accuracy, paler orange: accuracy for hydrogen. Values are from PDG [\[1\]](#page-5-0) except for \overline{H} results for HFS [\[2\]](#page-5-6), 1S–2S [\[3\]](#page-5-7), and Lamb shift [\[4\]](#page-5-8). \overline{p} chargeto-mass ratio [\[5\]](#page-5-9), note that in this case only the left hand side is well defined, while the right hand side is less precise since the cyclotron frequency is proportional to the magnetic field in which the measurement is taken.

antihydrogen atoms suitable for spectroscopy. At the Antiproton Decelerator of CERN now several experiments are routinely producing antihydrogen [\[11\]](#page-5-10), with the ALPHA collaboration having been able to perform first spectroscopy measurements recently [\[2,](#page-5-6) [3,](#page-5-7) [12\]](#page-5-11).

ASACUSA is aiming at a measurement of the antihydrogen ground-state hyperfine structure GS-HFS [\[13,](#page-5-12) [14\]](#page-5-13), transporting the \overline{H} atoms to a region far away from the stray magnetic fields at the formation region, like it was done for hydrogen by Rabi [\[15\]](#page-5-14) and others. This will ultimately allow for precision exceeding the one of in-trap hyperfine spectroscopy as done in [\[3\]](#page-5-7).

2 Antihydrogen beam formation and hyperfine spectroscopy

In ASACUSA antihydrogen atoms are formed from their ingredients [\[16\]](#page-6-0). Antiprotons from the CERN Antiproton Decelerator are trapped and cooled in a Penning trap MUSASHI [\[17\]](#page-6-1), Positrons are produced by a 22 Na source in combination with a solid Ne ice moderator, they are accumulated in a Surko type buffer gas trap. Both species are brought together and mixed in a magnetic CUSP configuration [\[18\]](#page-6-2) using a nested Penning trap configuration [\[19\]](#page-6-3) which is widely used at the AD.

Fig. [2](#page-2-0) (left) shows a schematic view of the Rabi-type beam spectroscopy experiment. Fig. [2](#page-2-0) (right) depicts the behavior of the four hyperfine states characterized by the quantum number *F* connected to the total spin $\mathbf{F} = \mathbf{S_1} + \mathbf{S_2}$ and its projection *M* to the quantization axis. States with increasing (decreasing) energy in an external magnetic field are called lowfield seekers (high-field seekers). Two possible transitions induced by microwaves (σ_1 , π_1) that are accessible in this experiment are denoted by arrows.

After the initial success of \overline{H} formation [\[20\]](#page-6-4), a beam of cold antihydrogen atoms was observed 2.7 m downstream of the formation region [\[21\]](#page-6-5). In 2017 for the first time the principle quantum number distribution could be measured using the antihydrogen detector [\[22\]](#page-6-6) and a field ionization stage, which – as expected from three-body formation – contains mainly Rydberg atoms [\[23\]](#page-6-7). \overline{H} with $n < 14$ were detected with 4.5 σ significance, which could within their life time of $\sim 100 \mu s$ decay to the ground state before reaching the cavity. The overall observed rate of such events of ~ 0.001 s⁻¹ is however far too small for spec-

Fig. 2 Left: schematic antihydrogen formation and hyperfine measurement beam line. Antihydrogen is created by mixing antiprotons and positrons in the CUSP trap that polarizes the outgoing beam by focusing low-field seekers (red) and defocussing high-field seekers (green). The beam then passes a microwave cavity where spin-flips are induced under the presence of an external static magnetic field B_{ext} , a superconducting sextupole which again selects low-field seekers, and an antihydrogen detector. Right: Breit-Rabi diagram showing the magnetic field dependence of the four hyperfine states of antihydrogen.

troscopy measurements. The primary task is to increase the production rate of ground-state antihydrogen by 1–2 orders of magnitude.

3 Hydrogen hyperfine spectroscopy results

Due to the scarce availability of antihydrogen, ASACUSA decided to develop a hydrogen beam with the initial goal to commission the hyperfine spectroscopy apparatus. A source of cold polarized hydrogen with a temperature of \sim 50 K as expected for the H formed in the CUSP was connected to the microwave cavity and superconducting sextupole magnet to be used for antihydrogen spectroscopy. For hydrogen detection a commercial Q-mass spectrometer was used. With this setup the σ_1 transition in hydrogen could be determined to a few Hertz precision at $v \sim 1.42$ GHz, resulting in a 2.7 ppb measurement of the hydrogen GS-HFS [\[24\]](#page-6-8). This constitutes the most precise determination of this quantity in a beam so far, the highest precision of 2 mHz having been obtained in a hydrogen maser [\[25,](#page-6-9) [26\]](#page-6-10)), which cannot be used for antimatter. The result using the σ_1 transition also allows us to estimate that with this method about 8000 \overline{H} atoms in the ground state are needed to obtain a precision of ∼ 1 ppm, which is the initial goal of ASACUSA.

While many coefficients of the minimal SME have been constrained by hydrogen maser measurements, in the newly developed non-minimal SME a set of coefficients is present that has not yet been investigated experimentally [\[27\]](#page-6-11), especially those depending on the orientation of the static magnetic field with respect to the Earth rotation axis [\[23\]](#page-6-7). Within the SME, the π_1 transition is sensitive to CPT violations while the σ_1 transition is not. Since the π_1 transition is more sensitive to magnetic field inhomogeneities, we constructed new McKeehan-like coils [\[28\]](#page-6-12) and a three-layer cylindrical shielding [\[29\]](#page-6-13). Additionally we changed the beam optics so to produce the same rate of hydrogen atoms in the different quantum states at the detector [\[30\]](#page-6-14), using permanent sextupole magnets and ring apertures. With this setup, first measurements were done determining alternatingly the transition frequencies v_{σ} and v_{π} at each setting of the external constant magnetic field B_{ext} , which is controlled by the McKeehan-like coils. Fig. [3](#page-3-0) shows preliminary measurements of v_{σ} and v_π at several B_{ext} . The *B*-field dependence of the two transitions follows from the Breit-Rabi formula [\[31\]](#page-6-15), yielding a hyperbolic dependence of $v_{\sigma}(B_{ext})$ and a dominantly linear one

Fig. 3 Preliminary results of the first simultaneously taken extrapolations of v_π (a) and v_σ (c) as a function of the external static magnetic field for hydrogen. (b) and (d) show the deviations from the fit as standard scores. The dashed red line in panel (a) shows v_{σ} to illustrate the much smaller *B*-field dependence compared to v_π .

for $v_\pi(B_{ext})$. Table [1](#page-4-0) gives the values of the hyperfine transition frequency v_0 obtained by extrapolating each type of transition to zero external field, yielding typical precisions of a 10–30 ppb in a first test run lasting five days.

Instead of extrapolations, the measurement of v_{σ} and v_{π} at the same B_{ext} can be used to directly determine the zero-field GS-HFS frequency v_0 [\[32\]](#page-7-0). This method, using the same data sets, leads to a factor 2 smaller errors than the individual extrapolations and is of interest to antihydrogen, as it includes the more interesting π_1 transition and requires the acquisition of only two resonances to predict the zero-field value (thereby reducing the amount of required \overline{H} events to 4000 for a ~ 1 ppm result). Further tests of this method are ongoing and systematic measurements of both v_{σ} and v_{π} are planned in the near future.

Table 1 Hydrogen ground-state hyperfine splitting results using the ASACUSA hydrogen beam (preliminary). The results agree within 1 standard deviation to the literature value v_{lit} obtained by a hydrogen maser [\[25\]](#page-6-9). From [\[29\]](#page-6-13).

	v_0 [Hz]	Relative error	$v_0 - v_{\text{lit}}$ [Hz]
σ_1 extrapolation	1 420 405 767(15)	1.04×10^{-8}	15
π_1 extrapolation	1 420 405 760(34)	2.38×10^{-8}	8
Mean value of the two extrapolations	1 420 405 766(14)	9.96×10^{-9}	14
v_{σ} and v_{π} determined at same static magnetic field	1 420 405 753(8)	5.60×10^{-9}	

4 Summary and outlook

The ground-state hyperfine splitting of antihydrogen offers one of the most sensitive tests of CPT symmetry. ASACUSA is preparing an antihydrogen beam for an in-flight measurement of this quantity using a Rabi-type spectroscopy method which has the advantage of being performed in a field-free environment. The rate of antihydrogen formation in the CUSP needs further improvement of 1–2 orders of magnitude to allow for spectroscopy measurements.

In parallel in-beam hyperfine spectroscopy of ordinary hydrogen is pursued. A first measurement using the same cavity and sextupole as will be employed for antihydrogen resulted in a 2.7 ppb precision showing that the spectroscopy method is fully functional. Using the hydrogen beam, measurements are under way to determine coefficients of the non-minimal Standard Model Extension that have so far not been experimentally constrained.

Acknowledgements This work has been supported by the European Research Council under European Union's Seventh Framework Programme (FP7/2007-2013)/ERC Grant agreement (291242), the Austrian Ministry of Science and Research, the Austrian Science Fund (FWF): W1252-N27, a Grant-in-Aid for Specially Promoted Research (24000008) of MEXT and the RIKEN Pioneering Project. We express our gratitude towards the AD group of CERN.

References

- 1. Tanabashi, M., et al., (Particle Data Group): Review of particle properties. Phys. Rev. D 98, 010001 (2018)
- 2. Ahmadi, M., Alves, B.X.R., Baker, C.J., Bertsche, W., Butler, E., Capra, A., Carruth, C., Cesar, C.L., Charlton, M., Cohen, S., Collister, R., Eriksson, S., Evans, A., Evetts, N., Fajans, J., Friesen, T., Fujiwara, M.C., Gill, D.R., Gutierrez, A., Hangst, J.S., Hardy, W.N., Hayden, M.E., Isaac, C.A., Ishida, A., Johnson, M.A., Jones, S.A., Jonsell, S., Kurchaninov, L., Madsen, N., Mathers, M., Maxwell, D., McKenna, J.T.K., Menary, S., Michan, J.M., Momose, T., Munich, J.J., Nolan, P., Olchanski, K., Olin, A., Pusa, P., Rasmussen, C.., Robicheaux, F., Sacramento, R.L., Sameed, M., Sarid, E., Silveira, D.M., Stracka, S., Stutter, G., So, C., Tharp, T.D., Thompson, J.E., Thompson, R.I., van der Werf, D.P., Wurtele, J.S.: Observation of the hyperfine spectrum of antihydrogen. Nature 548, 66 (2017). URL <http://dx.doi.org/10.1038/nature23446>
- 3. Ahmadi, M., Alves, B.X.R., Baker, C.J., Bertsche, W., Capra, A., Carruth, C., Cesar, C.L., Charlton, M., Cohen, S., Collister, R., Eriksson, S., Evans, A., Evetts, N., Fajans, J., Friesen, T., Fujiwara, M.C., Gill, D.R., Hangst, J.S., Hardy, W.N., Hayden, M.E., Isaac, C.A., Johnson, M.A., Jones, J.M., Jones, S.A., Jonsell, S., Khramov, A., Knapp, P., Kurchaninov, L., Madsen, N., Maxwell, D., McKenna, J.T.K., Menary, S., Momose, T., Munich, J.J., Olchanski, K., Olin, A., Pusa, P., Rasmussen, C.., Robicheaux, F., Sacramento, R.L., Sameed, M., Sarid, E., Silveira, D.M., Stutter, G., So, C., Tharp, T.D., Thompson, R.I., van der Werf, D.P., Wurtele, J.S.: Characterization of the 1s-2s transition in antihydrogen. Nature 557(7703), 71–75 (2018). URL <https://doi.org/10.1038/s41586-018-0017-2>
- 4. Crivelli, P., Cooke, D., Heiss, M.W.: Antiproton charge radius. Phys. Rev. D 94, 052008 (2016). DOI 10. 1103/PhysRevD.94.052008. URL <https://link.aps.org/doi/10.1103/PhysRevD.94.052008>
- 5. Ulmer, S., Smorra, C., Mooser, A., Franke, K., Nagahama, H., Schneider, G., Higuchi, T., Van Gorp, S., Blaum, K., Matsuda, Y., et al.: High-precision comparison of the antiproton-to-proton charge-to-mass ratio. Nature 524(7564), 196–199 (2015)
- 6. Smorra, C., Sellner, S., Borchert, M.J., Harrington, J.A., Higuchi, T., Nagahama, H., Tanaka, T., Mooser, A., Schneider, G., Bohman, M., Blaum, K., Matsuda, Y., Ospelkaus, C., Quint, W., Walz, J., Yamazaki, Y., Ulmer, S.: A parts-per-billion measurement of the antiproton magnetic moment. Nature 550, 371 (2017). URL <http://dx.doi.org/10.1038/nature24048>
- 7. Colladay, D., Kostelecký, V.A.: CPT violation and the standard model. Physical Review D 55, 6760– 6774 (1997)
- 8. Kostelecký, V., Russell, N.: Data tables for Lorentz and CPT violation. Review of Modern Physics $83(1)$, 11–32 (2011)
- 9. Kostelecky, A., Russell, N.: Data Tables for Lorentz and CPT Violation. [arXiv:0801.0287](http://arxiv.org/abs/0801.0287) (2018 edition) 10. Charlton, M., Eades, J., Horvath, D., Hughes, R., Zimmermann, C.: Antihydrogen physics. Physics Reports 241(2), 65–117 (1994)
- 11. Hori, M., Walz, J.: Physics at CERN's antiproton decelerator. Progress in Particle and Nuclear Physics 72, 206 – 253 (2013). DOI https://doi.org/10.1016/j.ppnp.2013.02.004. URL [http://www.](http://www.sciencedirect.com/science/article/pii/S0146641013000069) [sciencedirect.com/science/article/pii/S0146641013000069](http://www.sciencedirect.com/science/article/pii/S0146641013000069)
- 12. Ahmadi, M., Alves, B.X.R., Baker, C.J., Bertsche, W., Capra, A., Carruth, C., Cesar, C.L., Charlton, M., Cohen, S., Collister, R., Eriksson, S., Evans, A., Evetts, N., Fajans, J., Friesen, T., Fujiwara, M.C., Gill, D.R., Hangst, J.S., Hardy, W.N., Hayden, M.E., Hunter, E.D., Isaac, C.A., Johnson, M.A., Jones, J.M., Jones, S.A., Jonsell, S., Khramov, A., Knapp, P., Kurchaninov, L., Madsen, N., Maxwell, D., McKenna, J.T.K., Menary, S., Michan, J.M., Momose, T., Munich, J.J., Olchanski, K., Olin, A., Pusa, P., Rasmussen, C.., Robicheaux, F., Sacramento, R.L., Sameed, M., Sarid, E., Silveira, D.M., Starko, D.M., Stutter, G., So, C., Tharp, T.D., Thompson, R.I., van der Werf, D.P., Wurtele, J.S.: Observation of the 1s-2p Lyman- α transition in antihydrogen. Nature 561, 211 (2018). URL [https://doi.org/10.1038/](https://doi.org/10.1038/s41586-018-0435-1) [s41586-018-0435-1](https://doi.org/10.1038/s41586-018-0435-1)
- 13. Widmann, E., Eades, J., Hayano, R.S., Hori, M., Horváth, D., Ishikawa, T., Juhász, B., Sakaguchi, J., Torii, H.A., Yamaguchi, H., Yamazaki, T.: Hyperfine structure measurements of antiprotonic helium and antihydrogen. In: S.G. Karshenboim, F.S. Pavone, F. Bassani, M. Inguscio, T.W. Hansch (eds.) The Hydrogen Atom: Precision Physics of Simple Atomic Systems, *Lecture Notes in Physics*, vol. 570, pp. 528–542. Springer-Verlag Berlin Heidelberg (2001). *[arXiv:nucl-ex/0102002](http://arxiv.org/abs/nucl-ex/0102002)*
- 14. Widmann, E., Diermaier, M., Juhasz, B., Malbrunot, C., Massiczek, O., Sauerzopf, C., Suzuki, K., ´ Wünschek, B., Zmeskal, J., Federmann, S., Kuroda, N., Ulmer, S., Yamazaki, Y.: Measurement of the hyperfine structure of antihydrogen in a beam. Hyperfine Interactions 215(1-3), 1–8 (2013). DOI 10.1007/ s10751-013-0809-6. URL <http://dx.doi.org/10.1007/s10751-013-0809-6>. arXiv:1301.4670
- 15. Rabi, I.I., Zacharias, J.R., Millman, S., Kusch, P.: A New Method of Measuring Nuclear Magnetic Moment. Physical Review 53, 318 (1938)
- 16. Kuroda, N., Tajima, M., Radics, B., Dupre, P., Nagata, Y., Kaga, C., Kanai, Y., Leali, M., Rizzini, E.L., ´ Mascagna, V., Matsudate, T., Breuker, H., Higaki, H., Matsuda, Y., Ulmer, S., Venturelli, L., Yamazaki, Y.: Antihydrogen Synthesis in a Double-Cusp Trap (2017). DOI 10.7566/JPSCP.18.011009. URL <https://journals.jps.jp/doi/abs/10.7566/JPSCP.18.011009>
- 17. Kuroda, N., Torii, H., Franzen, K., Wang, Z., Yoneda, S., Inoue, M., Hori, M., Juhász, B., Horváth, D., Higaki, H., Mohri, A., Eades, J., Komaki, K., Yamazaki, Y.: Confinement of a Large Number of Antiprotons and Production of an Ultraslow Antiproton Beam. Physical Review Letters 94(2), 023401 (2005)
- 18. Mohri, A., Yamazaki, Y.: A possible new scheme to synthesize antihydrogen and to prepare a polarised antihydrogen beam. Europhysics Letters 63, 207 (2003)
- 19. Gabrielse, G., Rolston, S., Haarsma, L., Kells, W.: Antihydrogen production using trapped plasmas. Physics Letters A 129(1), 38 – 42 (1988). DOI https://doi.org/10.1016/0375-9601(88)90470-7. URL <http://www.sciencedirect.com/science/article/pii/0375960188904707>
- 20. Enomoto, Y., Kuroda, N., Michishio, K., Kim, C.H., Higaki, H., Nagata, Y., Kanai, Y., Torii, H.A., Corradini, M., Leali, M., Lodi-Rizzini, E., Mascagna, V., Venturelli, L., Zurlo, N., Fujii, K., Ohtsuka, M., Tanaka, K., Imao, H., Nagashima, Y., Matsuda, Y., Juhasz, B., Mohri, A., Yamazaki, Y.: Synthesis of ´ cold antihydrogen in a cusp trap. Phys. Rev. Lett. 105(24), 243401 (2010). DOI 10.1103/PhysRevLett. 105.243401
- 21. Kuroda, N., Ulmer, S., Murtagh, D.J., Van Gorp, S., Nagata, Y., Diermaier, M., Federmann, S., Leali, M., Malbrunot, C., Mascagna, V., Massiczek, O., Michishio, K., Mizutani, T., Mohri, A., Nagahama, H., Ohtsuka, M., Radics, B., Sakurai, S., Sauerzopf, C., Suzuki, K., Tajima, M., Torii, H.A., Venturelli, L., Wunschek, B., Zmeskal, J., Zurlo, N., Higaki, H., Kanai, Y., Lodi Rizzini, E., Nagashima, Y., Matsuda, ¨ Y., Widmann, E., Yamazaki, Y.: A source of antihydrogen for in-flight hyperfine spectroscopy. Nat Commun 5, 3089 (2014). URL <http://dx.doi.org/10.1038/ncomms4089>
- 22. Kolbinger, B., Amsler, C., Breuker, H., Diermaier, M., Dupre, P., Fleck, M., Gligorova, A., Higaki, H., ´ Kanai, Y., Kobayashi, T., Leali, M., Mackel, V., Malbrunot, C., Mascagna, V., Massiczek, O., Matsuda, ¨ Y., Murtagh, D., Nagata, Y., Sauerzopf, C., Simon, M.C., Tajima, M., Ulmer, S., Kuroda, N., Venturelli, L., Widmann, E., Yamazaki, Y., Zmeskal, J.: Recent developments from ASACUSA on antihydrogen detection. EPJ Web of Conferences 181, 01003 (2018). URL [https://doi.org/10.1051/epjconf/](https://doi.org/10.1051/epjconf/201818101003) [201818101003](https://doi.org/10.1051/epjconf/201818101003)
- 23. Malbrunot, C., Amsler, C., Arguedas Cuendis, S., Breuker, H., Dupre, P., Fleck, M., Higaki, H., ´ Kanai, Y., Kolbinger, B., Kuroda, N., Leali, M., Mäckel, V., Mascagna, V., Massiczek, O., Matsuda, Y., Nagata, Y., Simon, M.C., Spitzer, H., Tajima, M., Ulmer, S., Venturelli, L., Widmann, E., Wiesinger, M., Yamazaki, Y., Zmeskal, J.: The ASACUSA antihydrogen and hydrogen program: results and prospects. Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences 376(2116), 20170273 (2018). DOI 10.1098/rsta.2017.0273. URL <http://rsta.royalsocietypublishing.org/content/376/2116/20170273>
- 24. Diermaier, M., Jepsen, C.B., Kolbinger, B., Malbrunot, C., Massiczek, O., Sauerzopf, C., Simon, M.C., Zmeskal, J., Widmann, E.: In-beam measurement of the hydrogen hyperfine splitting and prospects for antihydrogen spectroscopy. Nature Communications 8, 15749 (2017). URL [http://dx.doi.org/10.](http://dx.doi.org/10.1038/ncomms15749) [1038/ncomms15749](http://dx.doi.org/10.1038/ncomms15749)
- 25. Hellwig, H., Vessot, R.F., Levine, M.W., Zitzewitz, P.W., Allan, D.W., Glaze, D.J.: Measurement of the unperturbed hydrogen hyperfine transition frequency. Instrumentation and Measurement, IEEE Transactions on 19(4), 200–209 (1970)
- 26. Karshenboim, S.G.: Some possibilities for laboratory searches for variations of fundamental constants. Canadian Journal of Physics 78(7), 639–678 (2000)
- 27. Kostelecky, V.A., Vargas, A.J.: Lorentz and CPT tests with hydrogen, antihydrogen, and related systems. ´ Phys. Rev. D 92, 056002 (2015). DOI 10.1103/PhysRevD.92.056002. URL [http://link.aps.org/](http://link.aps.org/doi/10.1103/PhysRevD.92.056002) [doi/10.1103/PhysRevD.92.056002](http://link.aps.org/doi/10.1103/PhysRevD.92.056002)
- 28. McKeehan, L.W.: Combinations of circular currents for producing uniform magnetic field gradients. Review of Scientific Instruments 7(4), 178–179 (1936). DOI 10.1063/1.1752110. URL [https://doi.](https://doi.org/10.1063/1.1752110) [org/10.1063/1.1752110](https://doi.org/10.1063/1.1752110)
- 29. Arguedas Cuendis, S.: Measuring the hydrogen ground-state hyperfine splitting through the π_1 and σ_1 transitions. M. thesis, Universität Wien. Fakultät für Physik (2017). URL [http://othes.univie.ac.](http://othes.univie.ac.at/22584/) [at/22584/](http://othes.univie.ac.at/22584/)
- 30. Wiesinger, M.: Design and implementation of new optics for the atomic hydrogen beam of ASACUSA's antihydrogen hyperfine spectrscopy experiment. M. thesis, Technische Universitat Wien (2017). URL ¨ <http://othes.univie.ac.at/22584/>
- 31. Breit, G., Rabi, I.I.: Measurement of nuclear spin. Phys. Rev. 38, 2082–2083 (1931). DOI 10.1103/ PhysRev.38.2082.2. URL <https://link.aps.org/doi/10.1103/PhysRev.38.2082.2>

32. Kolbinger, B., Capon, A., Diermaier, M., Lehner, S., Malbrunot, C., Massiczek, O., Sauerzopf, C., Simon, M.C., Widmann, E.: Numerical simulations of hyperfine transitions of antihydrogen. Hyperfine Interactions 233(1), 47–51 (2015). DOI 10.1007/s10751-015-1130-3. URL [http://dx.doi.org/](http://dx.doi.org/10.1007/s10751-015-1130-3) [10.1007/s10751-015-1130-3](http://dx.doi.org/10.1007/s10751-015-1130-3)