



CERN-EP-2019-241
28 October 2019

Probing the effects of strong electromagnetic fields with charge-dependent directed flow in Pb–Pb collisions at the LHC

ALICE Collaboration*

Abstract

The first measurement at the LHC of charge-dependent directed flow (v_1) relative to the spectator plane is presented for Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV. Results are reported for charged hadrons and D^0 mesons for the transverse momentum intervals $p_T > 0.2$ GeV/ c and $3 < p_T < 6$ GeV/ c in the 5–40% and 10–40% centrality classes, respectively. The difference between the positively and negatively charged hadron v_1 has a positive slope as a function of pseudorapidity η , $d\Delta v_1/d\eta = [1.68 \pm 0.49$ (stat.) ± 0.41 (syst.)] $\times 10^{-4}$. The same measurement for D^0 and \bar{D}^0 mesons yields a positive value $d\Delta v_1/d\eta = [4.9 \pm 1.7$ (stat.) ± 0.6 (syst.)] $\times 10^{-1}$, which is about three orders of magnitude larger than the one of the charged hadrons. These measurements can provide new insights into the effects of the strong electromagnetic field and the initial tilt of matter created in non-central heavy-ion collisions on the dynamics of light (u, d, and s) and heavy (c) quarks. The large difference between the observed Δv_1 of charged hadrons and D^0 mesons may reflect different sensitivity of the charm and light quarks to the early time dynamics of a heavy-ion collision. These observations challenge some recent theoretical calculations, which predicted a negative and an order of magnitude smaller value of $d\Delta v_1/d\eta$ for both light-flavour and charmed hadrons.

© 2019 CERN for the benefit of the ALICE Collaboration.

Reproduction of this article or parts of it is allowed as specified in the CC-BY-4.0 license.

*See Appendix A for the list of collaboration members

Quantum Chromo-dynamic (QCD) calculations on the lattice [1–6] predict at high temperatures the existence of a deconfined state of quarks and gluons, known as the quark–gluon plasma (QGP). Characterizing the QGP properties is among the main goals of the experimental program with ultra-relativistic heavy-ion collisions at the Large Hadron Collider (LHC). Measurements of the anisotropic transverse flow [7–11], quantified by the second (v_2) and higher order ($n > 2$) harmonic coefficients v_n , allowed one to characterize the different phases of a heavy-ion collision and to constrain the properties of the QGP [12–16].

The directed flow, v_1 , has a special role due to its sensitivity to the three-dimensional spatial profile of the initial conditions and the pre-equilibrium early time dynamics in the evolution of the collision. The space-momentum correlations in particle production from a longitudinally tilted source result in a non-zero v_1 . The tilt arises from the asymmetries in the number of forward and backward moving participant nucleons at different positions in the transverse plane [17–19]. The directed flow of charged hadrons at the LHC [20] has significantly smaller magnitude compared to that at lower RHIC energies [21], which can be interpreted as a smaller initial tilt at the LHC [22–24].

Charm quarks are produced early in the collision via hard scattering processes. Their emission region is not tilted in the longitudinal direction [19] unlike the one of light quarks, which are predominantly produced in soft processes at later stages of the collision [18, 25]. Consequently, the region of charm quark production in the transverse plane is shifted with respect to that of light quarks and gluons, resulting in an enhanced dipole asymmetry in the charm quark distribution [19]. During the system expansion, charm quarks would be dragged by the flow of the light quarks in the transverse direction of the shift, which is predicted to result in a larger v_1 of charm hadrons compared to light-flavour hadrons [19, 26]. Consequently, the measurements of the charge-integrated directed flow of hadrons containing light (u, d, and s) and heavy (c) quarks together with their difference in magnitude are of great interest and allow one to probe the three-dimensional space-time evolution of the produced matter.

Heavy-ion collisions are also characterized by extremely strong electromagnetic fields primarily induced by spectator protons, which do not undergo inelastic collisions. There is strong interest in characterizing the time evolution of these fields, which are estimated to reach $10^{18} - 10^{19}$ Gauss in the early stages (< 0.5 fm) of Pb–Pb collisions at LHC energies [27, 28]. Phenomena predicted to occur in the presence of this strong electromagnetic field include the chiral magnetic effect (CME), which is driven by the generation of an electric current along the magnetic field in a medium with chiral imbalance [29–32]. While experimental results for charge-dependent correlations are in qualitative agreement with theoretical expectations for the CME [33–35], possible background contributions, such as effects of local charge conservation coupled with the anisotropic flow, prevent their unambiguous interpretation [36] and have led to upper limits on the CME at LHC energies. Thus it is fundamental to use other observables with direct sensitivity to the electromagnetic fields in order to constrain their magnitudes and time evolution in heavy-ion collisions.

The charge dependence of the produced particle directed flow relative to the spectator plane is directly sensitive to the presence of electromagnetic fields. The spectator plane is defined by the deflection direction of the collision spectators. On average its orientation is perpendicular to the direction of the magnetic field generated by the positively charged spectators. The charge dependence of v_1 comes from two competing effects, acting in opposite directions, i.e. the Lorentz force and the Coulomb force, both induced by the rapidly decreasing magnetic field. In an electrically conducting plasma the induced charged currents might slow down the decay of the magnetic field [27]. The measurement of charge-dependent v_1 can therefore constrain the QGP electric conductivity.

First estimates of these effects on the v_1 of charged particles were presented in [37, 38]. Using the electric conductivity from lattice QCD calculations [39, 40], the difference between the v_1 of positively and negatively charged pions, $\Delta v_1(\pi) = v_1(\pi^+) - v_1(\pi^-)$, in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV

was estimated to be not larger than 10^{-5} for $|\eta| < 1$ [38]. Charm quarks, which are produced in the early stages when the magnetic field is maximal, should be more strongly affected by the electromagnetic fields than light quarks [26, 41]. The difference Δv_1 between the v_1 of D^0 ($c\bar{u}$) and \bar{D}^0 ($\bar{c}u$) mesons should therefore provide better sensitivity to this initial magnetic field. A calculation of v_1 of charmed mesons in the rapidity interval $|y| < 1$ gives a value $\Delta v_1(D) \sim 10^{-2}$ [41], which is three orders of magnitude larger than the one expected for pions. Recently, the STAR Collaboration published a measurement of directed flow of D^0 and \bar{D}^0 mesons in 10–80% central Au–Au collisions at $\sqrt{s_{NN}} = 200$ GeV [42]. The slope of v_1 of D^0 and \bar{D}^0 mesons as a function of rapidity is negative and about a factor of 25 times larger than that of charged kaons.

This Letter reports the first measurements at the LHC of the charge dependence of v_1 relative to the spectator plane for charged hadrons and D^0 mesons as a function of pseudorapidity in mid-central Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV.

About 23 (19) million Pb–Pb collisions in the 5–40% (10–40%) centrality interval are used for the charged hadron (D^0 and \bar{D}^0 meson) v_1 measurements. Only events with a primary vertex reconstructed within ± 10 cm from the detector centre along the beam direction are analysed. Two forward scintillator arrays (V0A and V0C) [43] are used to determine the collision centrality. For the most central (0–5%) collisions, the small number of spectators prevents an accurate reconstruction of their deflection. In the 5–10% centrality interval the large combinatorial background does not allow the measurement of the D^0 and \bar{D}^0 v_1 .

The deflection direction of the collision spectators is reconstructed from spectator neutrons detected using two Zero Degree Calorimeters (ZDC) [44, 45]. The ZDCs have a 2×2 segmentation in the plane transverse to the beam direction and are installed at 112.5 m distance from the detector centre on both sides of the interaction point, covering the "projectile" ($\eta > 8.78$) and the "target" ($\eta < -8.78$) spectator regions. For each ZDC a flow vector is constructed following the procedure described in [20]

$$\mathbf{Q}^{t,p} \equiv (Q_x^{t,p}, Q_y^{t,p}) = \frac{\sum_{i=1}^4 \mathbf{n}_i E_i^{t,p}}{\sum_{i=1}^4 E_i^{t,p}}, \quad (1)$$

where p and t denote the ZDC on the projectile and target side, E_i is the measured signal and $\mathbf{n}_i = (x_i, y_i)$ are the coordinates of the center of the i -th ZDC segment.

The deflection direction of the spectator neutrons is estimated event-by-event with the $\mathbf{Q}^{t,p}$ vectors corrected for the run-dependent variation of the LHC beam crossing position [46]. In mid-central collisions this deflection direction is strongly correlated with the magnetic field orientation. The deflection is expected to be opposite (anti-correlated) for the projectile and the target sides, i.e. $\langle Q_x^p Q_x^t \rangle = \langle Q_y^p Q_y^t \rangle < 0$ and $\langle Q_y^p Q_x^t \rangle$ and $\langle Q_x^p Q_y^t \rangle = 0$. A deviation from these expectations, mostly for peripheral collisions with centrality above 40%, is observed even after applying the flow-vector correction. These residual variations are used in the estimation of the systematic uncertainty as described in [20] and discussed below.

The directed flow is measured using the scalar-product method [47]

$$v_1^{t,p} = \frac{\langle \mathbf{u} \mathbf{Q}^{t,p} \rangle}{\sqrt{|\langle \mathbf{Q}^t \mathbf{Q}^p \rangle|}} = \frac{\langle u_x Q_x^{t,p} + u_y Q_y^{t,p} \rangle}{\sqrt{|\langle Q_x^t Q_x^p + Q_y^t Q_y^p \rangle|}}, \quad (2)$$

where $\mathbf{u} = (\cos \varphi, \sin \varphi)$ is the unit flow vector of the charged hadron or D^0 meson candidate with azimuthal angle φ . The directed flow is calculated as: $v_1 = (v_1^p - v_1^t)/2$. The sign of v_1 is defined relative to the deflection of the projectile spectators, corresponding to the rapidity-odd component of the v_1 discussed in [20]. The measurement of v_1 using spectators does not require any treatment of the momentum

conservation unlike the measurements based on correlations between particles produced at midrapidity [48]. This is justified by the observation of a vanishing relative momentum shift along the spectator plane at $\eta = 0$ [20].

The charged-hadron v_1 is measured from tracks reconstructed with the Inner Tracking System (ITS) [49] and the Time Projection Chamber (TPC) [50] and selected requiring $p_T > 0.2$ GeV/c, $|\eta| < 0.8$, at least 70 (out of a maximum of 159) TPC space points and $\chi^2/\text{ndf} < 2$ for the momentum fit in the TPC. In order to reduce the contamination from secondary particles only tracks with a maximum distance of closest approach (DCA) to the reconstructed primary vertex in both the transverse ($\text{DCA}_{xy} < 2.4$ cm) and the longitudinal direction ($\text{DCA}_z < 3.2$ cm) are accepted.

The D^0 and \bar{D}^0 mesons are reconstructed using the decay channel $D^0 \rightarrow K^- \pi^+$ and its charge conjugate for $3 < p_T < 6$ GeV/c. Pions and kaons are reconstructed in the TPC and ITS detectors. Tracks are selected requiring $|\eta| < 0.8$, $p_T > 0.4$ GeV/c, at least 70 hits in TPC and at least 2 hits (out of a maximum of 6) in the ITS, out of which at least one has to be in the two innermost layers. Particle identification is based on measurements of the specific ionisation energy loss dE/dx in the TPC and the flight time from the interaction point to the Time-Of-Flight (TOF) detector [51]. The charge of the identified pions and kaons allows one to distinguish between the $D^0 \rightarrow K^- \pi^+$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ candidates. Geometrical selections on the displaced decay vertex topology are applied to reduce the combinatorial background [52].

The v_1^D is extracted separately for D^0 and \bar{D}^0 mesons via a simultaneous fit to the number $N(M)$ of $K^\mp \pi^\pm$ pairs and their $v_1(M)$ as a function of the invariant mass, M :

$$N(M) = N_D(M) + N_{\text{bg}}(M), \quad (3)$$

$$v_1(M) = [v_1^D N_D(M) + v_1^{\text{bg}}(M) N_{\text{bg}}(M)] / [N_D(M) + N_{\text{bg}}(M)]. \quad (4)$$

An example of the simultaneous fit is shown in Fig. 1. The invariant mass distribution is fitted with the sum of a Gaussian function $N_D(M)$ for the D^0 and \bar{D}^0 signal and an exponential function $N_{\text{bg}}(M)$ for the background. The invariant mass dependence of the directed flow of background candidates $v_1^{\text{bg}}(M)$ is parametrized by a linear function.

Candidates, which satisfy both the $K^- \pi^+$ and $K^+ \pi^-$ hypotheses (reflected kinematics) and therefore can not be tagged uniquely as D^0 or \bar{D}^0 , are rejected. This removes about 35% of the signal and increases the signal-to-background ratio by about 30–40%, with a net result of a negligible reduction of the statistical significance of the D^0 and \bar{D}^0 yield. The extracted v_1^D includes contributions from both prompt D^0 mesons and feed-down D^0 mesons from beauty-hadron decays. The fraction of prompt D^0 meson is about 85% for the analysed centrality class and p_T interval [53].

Common sources of systematic uncertainty between charged hadrons and D mesons are related to the resolution of the spectator plane and to the dependence on the ALICE magnet polarity. The absolute systematic uncertainty related to the residual asymmetry in the spectator plane estimation is given by the difference between the v_1 obtained separately from $\langle u_x Q_x \rangle$ and $\langle u_y Q_y \rangle$ correlations with the ZDCs in Eq. 2. It is about 3.5×10^{-5} (2×10^{-2}) for charged hadrons (D^0 and \bar{D}^0 mesons). Effects related to track reconstruction and geometrical alignment of the detectors, which could influence positive and negative tracks differently, are estimated by comparing the v_1 results obtained using data taken with opposite magnet polarity. This comparison also probes the bias in the spectator plane estimation due to the non-zero beam crossing angle in the vertical plane, which had opposite values (± 60 μrad) for the opposite magnet polarities. The absolute difference between the v_1 values obtained with the two field polarities is 2.5×10^{-5} (2×10^{-2}) for charged hadrons (D^0 and \bar{D}^0). These systematic uncertainties are correlated in pseudorapidity for charged hadrons, while for D^0 and \bar{D}^0 mesons no significant correlation, beyond statistical uncertainties, is observed.




Figure 1: (color online) Illustration of the D^0 meson v_1 extraction procedure via a simultaneous fit to the candidate invariant mass distribution (upper panel) and v_1 (lower panel) for $3 < p_T < 6$ GeV/c and $-0.4 < \eta < 0$ in the 10–40% centrality interval. The blue solid lines correspond to the combined signal and background fit functions, while the red dashed lines represent the background contribution.

For charged hadrons, the track quality selections are varied and an absolute systematic uncertainty of 2.5×10^{-5} is assigned. The contribution from secondaries is varied by changing the maximum DCA_{xy}, which resulted in a negligible variation of v_1 . The contamination due to TPC tracks originating from pile-up collisions during the readout time of the TPC is estimated by varying the selections on the correlations between the event multiplicity (centrality) estimated with detectors with different readout times. The resulting systematic uncertainty is 10^{-5} for the charged hadrons. No systematic uncertainty is assigned for D^0 and \bar{D}^0 because the topological selections effectively remove tracks from pile-up. The uncertainty due to the D^0 and \bar{D}^0 signal extraction is estimated by varying (i) the fit functions in Eq. (3) and (4) for $N(M)$ and $v_1(M)$, (ii) fixing the Gaussian width and mean to the values extracted from Monte Carlo simulations, and (iii) varying the invariant mass fit range. The absolute systematic uncertainty assigned to v_1 due to the D^0 and \bar{D}^0 yield extraction is 2×10^{-2} . The possible bias due to the p_T -dependent efficiency in the D^0 and \bar{D}^0 v_1 analysis is tested by re-weighting both signal and background with the inverse value of the signal reconstruction efficiency as a function of p_T . The assigned absolute systematic uncertainty is 10^{-2} .

The total systematic uncertainty on v_1 is obtained by adding in quadrature the contributions described above. In the calculation of $\Delta v_1(D)$ all individual systematic uncertainties are propagated as fully uncorrelated between D^0 and \bar{D}^0 . For charged hadrons, the systematic uncertainties due to the asymmetry in the spectator plane estimation and the magnet polarity are correlated between positive and negative tracks and largely cancel in $\Delta v_1(h)$.

The pseudorapidity dependence of the directed flow of positively and negatively charged hadrons for the 5–40% centrality class in Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV is shown in the upper left panel of Fig. 2. The negative slope of v_1 is usually attributed to the effect of the initial tilt [18] or rotation [25] of the particle-emitting source. The charge-integrated v_1 at $\sqrt{s_{NN}} = 5.02$ TeV agrees within uncertainties with the results at $\sqrt{s_{NN}} = 2.76$ TeV [20].

The difference $\Delta v_1(h)$ between the v_1 of positively and negatively charged hadrons as a function of pseu-

rapidity is shown in the lower left panel of Fig. 2. The rapidity slope $d\Delta v_1/d\eta$, extracted with a linear fit (constrained to $v_1 = 0$ at $\eta = 0$), is $d\Delta v_1/d\eta = [1.68 \pm 0.49$ (stat.) ± 0.41 (syst.)] $\times 10^{-4}$ with a significance of 2.6σ for having a positive value. The $d\Delta v_1/d\eta$ is expected to reflect different effects, including those originating from the early-time magnetic field dynamics [19, 26, 41], the Coulomb interaction with charged spectators [54], as well as the transport to mid-rapidity via the baryon stopping mechanism [17] of the positive charge carried by the protons from the colliding nuclei. The importance of baryon stopping for the charge dependence of unidentified hadron v_1 is supported by the observed difference, even at top RHIC energy, between proton and anti-proton v_1 [22, 55, 56]. The baryon stopping effects are expected to decrease with increasing collision energy, as supported by the observation of a smaller magnitude of v_1 [20] and of a proton-to-antiproton ratio closer to unity at the LHC as compared to RHIC [57]. Despite the overall decrease, the baryon stopping can contribute significantly to the proton and anti-proton v_1 difference, and as such to the charge dependence of the inclusive hadron v_1 .

The charged-hadron $d\Delta v_1/d\eta$ at $\sqrt{s_{NN}} = 5.02$ TeV is one order of magnitude larger and has an opposite sign with respect to calculations for charged pions at $\sqrt{s_{NN}} = 2.76$ TeV [38] based on the analytic solution of relativistic hydrodynamics [58] with a constant electrical conductivity of the QGP. More recent calculations [54], using viscous hydrodynamic calculations [59], yield an absolute value of $d\Delta v_1/d\eta$ of similar magnitude as the one measured for charged hadrons, but with opposite sign.




Figure 2: (color online) Upper left: v_1 of positively (red) and negatively (blue) charged hadrons for the 5–40% centrality interval. Upper right: v_1 of D^0 (red) and \bar{D}^0 (blue) for the 10–40% centrality interval. Lower left and right: $\Delta v_1(h) = v_1(h^+) - v_1(h^-)$ and $\Delta v_1(D) = v_1(D^0) - v_1(\bar{D}^0)$, respectively. Dashed lines represent fits with a linear function.

The D^0 and \bar{D}^0 v_1 as a function of pseudorapidity is shown in the upper right panel of Fig. 2. The data

suggest a positive slope for the rapidity dependence of the v_1 of D^0 and a negative slope for \bar{D}^0 , with a significance of about 2σ in both cases. The slopes are different from the measurements in Au–Au collisions at $\sqrt{s_{NN}} = 200$ GeV [42], where a negative value is observed for both D^0 and \bar{D}^0 . Additionally, the v_1 for D^0 and \bar{D}^0 mesons with $3 < p_T < 6$ GeV/c ($\langle p_T \rangle \approx 4.2$ GeV/c) in the 10–40% centrality interval is about three orders of magnitude larger than that of charged hadrons with $p_T > 0.2$ GeV/c ($\langle p_T \rangle \approx 0.7$ GeV/c) in the 5–40% centrality class. The different p_T intervals used for the charged-hadron and D^0 -meson v_1 measurements are imposed by the statistical precision of the data, which simultaneously limits the yield of high- p_T charged hadrons and results in low significance of the D^0 and \bar{D}^0 meson yield at low p_T . The charged hadron v_1 at the LHC has a weak centrality dependence and changes sign around $p_T \approx 1.5$ GeV/c [20]. The differences in centrality and transverse momentum intervals should not be responsible for the observed difference between the magnitude of the v_1 of charged hadrons and D^0 and \bar{D}^0 mesons. The D^0 and \bar{D}^0 v_1 is an order of magnitude larger than the predictions from the transport [41] and hydrodynamic [19, 26] model calculations. The difference between the v_1 values of D^0 and \bar{D}^0 mesons $\Delta v_1(D)$ is shown in the lower right panel of Fig. 2. The value of $d\Delta v_1/d\eta = [4.9 \pm 1.7$ (stat.) ± 0.6 (syst.)] $\times 10^{-1}$ corresponds to a significance of 2.7σ to have a positive slope. A negative value for $d\Delta v_1/d\eta$ was predicted in [41] and is observed in Au–Au collisions at $\sqrt{s_{NN}} = 200$ GeV [42]. The opposite sign of the measured D^0 -meson and charged-hadron Δv_1 slope with respect to model calculations might indicate a stronger effect of the Lorentz force relative to the Coulomb one. These results demonstrate the sensitivity of the v_1 to the interplay among the effects of the rapidly decreasing magnetic field and the initial tilt of the source.

In summary, first measurements of the charge dependence of v_1 relative to the spectator plane in mid-central Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV are presented. The v_1 and the difference Δv_1 between positively and negatively charged hadrons and D^0 mesons are sensitive to the effects of the electromagnetic fields induced by spectator protons, baryon number transport, and the initial tilt or rotation of the particle-emitting source for non-central collisions. An indication of a positive slope $d\Delta v_1/d\eta$ of the charge-dependent v_1 at midrapidity for both charged hadrons and D^0 and \bar{D}^0 mesons is observed. The slope $d\Delta v_1/d\eta$ is $[1.68 \pm 0.49$ (stat.) ± 0.41 (syst.)] $\times 10^{-4}$ for charged hadrons with $p_T > 0.2$ GeV/c and $[4.9 \pm 1.7$ (stat.) ± 0.6 (syst.)] $\times 10^{-1}$ for D^0 and \bar{D}^0 mesons with $3 < p_T < 6$ GeV/c, with significance of 2.6σ and 2.7σ for having a positive value, respectively. The measured values of v_1 for D^0 and \bar{D}^0 mesons are about three orders of magnitude larger than the one of charged hadrons. These measurements together with those at RHIC [42] provide new insights and can constrain the theoretical modelling [38, 41] of electromagnetic effects. Further constraints will be set by future higher precision measurements at the LHC [60, 61].

Acknowledgements

The ALICE Collaboration would like to thank all its engineers and technicians for their invaluable contributions to the construction of the experiment and the CERN accelerator teams for the outstanding performance of the LHC complex. The ALICE Collaboration gratefully acknowledges the resources and support provided by all Grid centres and the Worldwide LHC Computing Grid (WLCG) collaboration. The ALICE Collaboration acknowledges the following funding agencies for their support in building and running the ALICE detector: A. I. Alikhanyan National Science Laboratory (Yerevan Physics Institute) Foundation (ANSL), State Committee of Science and World Federation of Scientists (WFS), Armenia; Austrian Academy of Sciences, Austrian Science Fund (FWF): [M 2467-N36] and Nationalstiftung für Forschung, Technologie und Entwicklung, Austria; Ministry of Communications and High Technologies, National Nuclear Research Center, Azerbaijan; Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Financiadora de Estudos e Projetos (Finep), Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) and Universidade Federal do Rio Grande do Sul (UFRGS), Brazil; Ministry of Education of China (MOEC), Ministry of Science & Technology of China (MSTC)

and National Natural Science Foundation of China (NSFC), China; Ministry of Science and Education and Croatian Science Foundation, Croatia; Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Cubaenergía, Cuba; Ministry of Education, Youth and Sports of the Czech Republic, Czech Republic; The Danish Council for Independent Research | Natural Sciences, the VILLUM FONDEN and Danish National Research Foundation (DNRF), Denmark; Helsinki Institute of Physics (HIP), Finland; Commissariat à l’Energie Atomique (CEA), Institut National de Physique Nucléaire et de Physique des Particules (IN2P3) and Centre National de la Recherche Scientifique (CNRS) and Région des Pays de la Loire, France; Bundesministerium für Bildung und Forschung (BMBF) and GSI Helmholtzzentrum für Schwerionenforschung GmbH, Germany; General Secretariat for Research and Technology, Ministry of Education, Research and Religions, Greece; National Research, Development and Innovation Office, Hungary; Department of Atomic Energy Government of India (DAE), Department of Science and Technology, Government of India (DST), University Grants Commission, Government of India (UGC) and Council of Scientific and Industrial Research (CSIR), India; Indonesian Institute of Science, Indonesia; Centro Fermi - Museo Storico della Fisica e Centro Studi e Ricerche Enrico Fermi and Istituto Nazionale di Fisica Nucleare (INFN), Italy; Institute for Innovative Science and Technology, Nagasaki Institute of Applied Science (IIST), Japanese Ministry of Education, Culture, Sports, Science and Technology (MEXT) and Japan Society for the Promotion of Science (JSPS) KAKENHI, Japan; Consejo Nacional de Ciencia (CONACYT) y Tecnología, through Fondo de Cooperación Internacional en Ciencia y Tecnología (FONCICYT) and Dirección General de Asuntos del Personal Académico (DGAPA), Mexico; Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO), Netherlands; The Research Council of Norway, Norway; Commission on Science and Technology for Sustainable Development in the South (COMSATS), Pakistan; Pontificia Universidad Católica del Perú, Peru; Ministry of Science and Higher Education and National Science Centre, Poland; Korea Institute of Science and Technology Information and National Research Foundation of Korea (NRF), Republic of Korea; Ministry of Education and Scientific Research, Institute of Atomic Physics and Ministry of Research and Innovation and Institute of Atomic Physics, Romania; Joint Institute for Nuclear Research (JINR), Ministry of Education and Science of the Russian Federation, National Research Centre Kurchatov Institute, Russian Science Foundation and Russian Foundation for Basic Research, Russia; Ministry of Education, Science, Research and Sport of the Slovak Republic, Slovakia; National Research Foundation of South Africa, South Africa; Swedish Research Council (VR) and Knut & Alice Wallenberg Foundation (KAW), Sweden; European Organization for Nuclear Research, Switzerland; Suranaree University of Technology (SUT), National Science and Technology Development Agency (NSDTA) and Office of the Higher Education Commission under NRU project of Thailand, Thailand; Turkish Atomic Energy Agency (TAEK), Turkey; National Academy of Sciences of Ukraine, Ukraine; Science and Technology Facilities Council (STFC), United Kingdom; National Science Foundation of the United States of America (NSF) and United States Department of Energy, Office of Nuclear Physics (DOE NP), United States of America.

References

- [1] **HotQCD** Collaboration, A. Bazavov *et al.*, “Chiral crossover in QCD at zero and non-zero chemical potentials”, *Phys. Lett.* **B795** (2019) 15–21, arXiv:1812.08235 [hep-lat].
- [2] A. Bazavov *et al.*, “The QCD Equation of State to $\mathcal{O}(\mu_B^6)$ from Lattice QCD”, *Phys. Rev.* **D95** no. 5, (2017) 054504, arXiv:1701.04325 [hep-lat].
- [3] **HotQCD** Collaboration, A. Bazavov *et al.*, “Equation of state in (2+1)-flavor QCD”, *Phys. Rev.* **D90** (2014) 094503, arXiv:1407.6387 [hep-lat].
- [4] A. Bazavov *et al.*, “The chiral and deconfinement aspects of the QCD transition”, *Phys. Rev.* **D85** (2012) 054503, arXiv:1111.1710 [hep-lat].

- [5] **Wuppertal-Budapest** Collaboration, S. Borsanyi, Z. Fodor, C. Hoelbling, S. D. Katz, S. Krieg, C. Ratti, and K. K. Szabo, “Is there still any T_c mystery in lattice QCD? Results with physical masses in the continuum limit III”, *JHEP* **09** (2010) 073, arXiv:1005.3508 [hep-lat].
- [6] **Wuppertal-Budapest** Collaboration, S. Borsanyi, G. Endrodi, Z. Fodor, A. Jakovac, S. D. Katz, S. Krieg, C. Ratti, and K. K. Szabo, “The QCD equation of state with dynamical quarks”, *JHEP* **11** (2010) 077, arXiv:1007.2580 [hep-lat].
- [7] **ALICE** Collaboration, K. Aamodt *et al.*, “Elliptic flow of charged particles in Pb-Pb collisions at 2.76 TeV”, *Phys. Rev. Lett.* **105** (2010) 252302, arXiv:1011.3914 [nucl-ex].
- [8] **ALICE** Collaboration, K. Aamodt *et al.*, “Higher harmonic anisotropic flow measurements of charged particles in Pb-Pb collisions at $\sqrt{s_{NN}}=2.76$ TeV”, *Phys. Rev. Lett.* **107** (2011) 032301, arXiv:1105.3865 [nucl-ex].
- [9] **ATLAS** Collaboration, G. Aad *et al.*, “Measurement of the azimuthal anisotropy for charged particle production in $\sqrt{s_{NN}} = 2.76$ TeV lead-lead collisions with the ATLAS detector”, *Phys. Rev.* **C86** (2012) 014907, arXiv:1203.3087 [hep-ex].
- [10] **CMS** Collaboration, S. Chatrchyan *et al.*, “Measurement of the elliptic anisotropy of charged particles produced in PbPb collisions at $\sqrt{s_{NN}}=2.76$ TeV”, *Phys. Rev.* **C87** no. 1, (2013) 014902, arXiv:1204.1409 [nucl-ex].
- [11] **ALICE** Collaboration, B. Abelev *et al.*, “Elliptic flow of identified hadrons in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV”, *JHEP* **06** (2015) 190, arXiv:1405.4632 [nucl-ex].
- [12] U. Heinz and R. Snellings, “Collective flow and viscosity in relativistic heavy-ion collisions”, *Ann. Rev. Nucl. Part. Sci.* **63** (2013) 123–151, arXiv:1301.2826 [nucl-th].
- [13] C. Gale, S. Jeon, and B. Schenke, “Hydrodynamic Modeling of Heavy-Ion Collisions”, *Int. J. Mod. Phys.* **A28** (2013) 1340011, arXiv:1301.5893 [nucl-th].
- [14] H. Niemi, K. J. Eskola, and R. Paatelainen, “Event-by-event fluctuations in a perturbative QCD + saturation + hydrodynamics model: Determining QCD matter shear viscosity in ultrarelativistic heavy-ion collisions”, *Phys. Rev.* **C93** no. 2, (2016) 024907, arXiv:1505.02677 [hep-ph].
- [15] J. E. Bernhard, J. S. Moreland, S. A. Bass, J. Liu, and U. Heinz, “Applying Bayesian parameter estimation to relativistic heavy-ion collisions: simultaneous characterization of the initial state and quark-gluon plasma medium”, *Phys. Rev.* **C94** no. 2, (2016) 024907, arXiv:1605.03954 [nucl-th].
- [16] A. Dubla, S. Masciocchi, J. M. Pawłowski, B. Schenke, C. Shen, and J. Stachel, “Towards QCD-assisted hydrodynamics for heavy-ion collision phenomenology”, *Nucl. Phys.* **A979** (2018) 251–264, arXiv:1805.02985 [nucl-th].
- [17] R. J. M. Snellings, H. Sorge, S. A. Voloshin, F. Q. Wang, and N. Xu, “Novel rapidity dependence of directed flow in high-energy heavy ion collisions”, *Phys. Rev. Lett.* **84** (2000) 2803–2805, arXiv:nucl-ex/9908001 [nucl-ex].
- [18] P. Bozek and I. Wiskiel, “Directed flow in ultrarelativistic heavy-ion collisions”, *Phys. Rev.* **C81** (2010) 054902, arXiv:1002.4999 [nucl-th].
- [19] S. Chatterjee and P. Bozek, “Large directed flow of open charm mesons probes the three dimensional distribution of matter in heavy ion collisions”, *Phys. Rev. Lett.* **120** no. 19, (2018) 192301, arXiv:1712.01189 [nucl-th].

- [20] **ALICE** Collaboration, B. Abelev *et al.*, “Directed Flow of Charged Particles at Midrapidity Relative to the Spectator Plane in Pb-Pb Collisions at $\sqrt{s_{NN}}=2.76$ TeV”, *Phys. Rev. Lett.* **111** no. 23, (2013) 232302, arXiv:1306.4145 [nucl-ex].
- [21] **STAR** Collaboration, B. I. Abelev *et al.*, “System-size independence of directed flow at the Relativistic Heavy-Ion Collider”, *Phys. Rev. Lett.* **101** (2008) 252301, arXiv:0807.1518 [nucl-ex].
- [22] **STAR** Collaboration, L. Adamczyk *et al.*, “Charge-dependent directed flow in Cu+Au collisions at $\sqrt{s_{NN}} = 200$ GeV”, *Phys. Rev. Lett.* **118** no. 1, (2017) 012301, arXiv:1608.04100 [nucl-ex].
- [23] A. Adil, M. Gyulassy, and T. Hirano, “3D jet tomography of the twisted color glass condensate”, *Phys. Rev.* **D73** (2006) 074006, arXiv:nucl-th/0509064 [nucl-th].
- [24] A. Adil and M. Gyulassy, “3D jet tomography of twisted strongly coupled quark gluon plasmas”, *Phys. Rev.* **C72** (2005) 034907, arXiv:nucl-th/0505004 [nucl-th].
- [25] L. P. Csernai, V. K. Magas, H. Stocker, and D. D. Strottman, “Fluid Dynamical Prediction of Changed v_1 -flow at LHC”, *Phys. Rev.* **C84** (2011) 024914, arXiv:1101.3451 [nucl-th].
- [26] S. Chatterjee and P. Bozek, “Interplay of drag by hot matter and electromagnetic force on the directed flow of heavy quarks”, *Phys. Lett.* **B798** (2019) 134955, arXiv:1804.04893 [nucl-th].
- [27] K. Tuchin, “Particle production in strong electromagnetic fields in relativistic heavy-ion collisions”, *Adv. High Energy Phys.* **2013** (2013) 490495, arXiv:1301.0099 [hep-ph].
- [28] V. Skokov, A. Yu. Illarionov, and V. Toneev, “Estimate of the magnetic field strength in heavy-ion collisions”, *Int. J. Mod. Phys.* **A24** (2009) 5925–5932, arXiv:0907.1396 [nucl-th].
- [29] K. Fukushima, D. E. Kharzeev, and H. J. Warringa, “The Chiral Magnetic Effect”, *Phys. Rev.* **D78** (2008) 074033, arXiv:0808.3382 [hep-ph].
- [30] D. E. Kharzeev, L. D. McLerran, and H. J. Warringa, “The Effects of topological charge change in heavy ion collisions: ‘Event by event P and CP violation’”, *Nucl. Phys.* **A803** (2008) 227–253, arXiv:0711.0950 [hep-ph].
- [31] K. Tuchin, “Synchrotron radiation by fast fermions in heavy-ion collisions”, *Phys. Rev.* **C82** (2010) 034904, arXiv:1006.3051 [nucl-th]. [Erratum: *Phys. Rev.* **C83**,039903(2011)].
- [32] V. Voronyuk, V. D. Toneev, W. Cassing, E. L. Bratkovskaya, V. P. Konchakovski, and S. A. Voloshin, “(Electro-)Magnetic field evolution in relativistic heavy-ion collisions”, *Phys. Rev.* **C83** (2011) 054911, arXiv:1103.4239 [nucl-th].
- [33] **CMS** Collaboration, V. Khachatryan *et al.*, “Observation of charge-dependent azimuthal correlations in p -Pb collisions and its implication for the search for the chiral magnetic effect”, *Phys. Rev. Lett.* **118** no. 12, (2017) 122301, arXiv:1610.00263 [nucl-ex].
- [34] **ALICE** Collaboration, B. Abelev *et al.*, “Charge separation relative to the reaction plane in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV”, *Phys. Rev. Lett.* **110** no. 1, (2013) 012301, arXiv:1207.0900 [nucl-ex].
- [35] **STAR** Collaboration, B. I. Abelev *et al.*, “Azimuthal Charged-Particle Correlations and Possible Local Strong Parity Violation”, *Phys. Rev. Lett.* **103** (2009) 251601, arXiv:0909.1739 [nucl-ex].

- [36] S. Schlichting and S. Pratt, “Charge conservation at energies available at the BNL Relativistic Heavy Ion Collider and contributions to local parity violation observables”, *Phys. Rev.* **C83** (2011) 014913, arXiv:1009.4283 [nucl-th].
- [37] A. Rybicki and A. Szczyrek, “Spectator induced electromagnetic effect on directed flow in heavy ion collisions”, *Phys. Rev.* **C87** no. 5, (2013) 054909, arXiv:1303.7354 [nucl-th].
- [38] U. Gursoy, D. Kharzeev, and K. Rajagopal, “Magnetohydrodynamics, charged currents and directed flow in heavy ion collisions”, *Phys. Rev.* **C89** no. 5, (2014) 054905, arXiv:1401.3805 [hep-ph].
- [39] H. T. Ding, A. Francis, O. Kaczmarek, F. Karsch, E. Laermann, and W. Soeldner, “Thermal dilepton rate and electrical conductivity: An analysis of vector current correlation functions in quenched lattice QCD”, *Phys. Rev.* **D83** (2011) 034504, arXiv:1012.4963 [hep-lat].
- [40] A. Amato, G. Aarts, C. Allton, P. Giudice, S. Hands, and J.-I. Skullerud, “Electrical conductivity of the quark-gluon plasma across the deconfinement transition”, *Phys. Rev. Lett.* **111** no. 17, (2013) 172001, arXiv:1307.6763 [hep-lat].
- [41] S. K. Das, S. Plumari, S. Chatterjee, J. Alam, F. Scardina, and V. Greco, “Directed Flow of Charm Quarks as a Witness of the Initial Strong Magnetic Field in Ultra-Relativistic Heavy Ion Collisions”, *Phys. Lett.* **B768** (2017) 260–264, arXiv:1608.02231 [nucl-th].
- [42] **STAR** Collaboration, J. Adam *et al.*, “First observation of the directed flow of D^0 and \bar{D}^0 in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV”, *Phys. Rev. Lett.* **123** (2019) 162301, arXiv:1905.02052 [nucl-ex].
- [43] **ALICE** Collaboration, E. Abbas *et al.*, “Performance of the ALICE VZERO system”, *JINST* **8** (2013) P10016, arXiv:1306.3130 [nucl-ex].
- [44] **ALICE** Collaboration, K. Aamodt *et al.*, “The ALICE experiment at the CERN LHC”, *JINST* **3** (2008) S08002.
- [45] **ALICE** Collaboration, B. Abelev *et al.*, “Performance of the ALICE Experiment at the CERN LHC”, *Int.J.Mod.Phys.* **A29** (2014) 1430044, arXiv:1402.4476 [nucl-ex].
- [46] I. Selyuzhenkov and S. Voloshin, “Effects of non-uniform acceptance in anisotropic flow measurement”, *Phys. Rev.* **C77** (2008) 034904, arXiv:0707.4672 [nucl-th].
- [47] M. Luzum and J.-Y. Ollitrault, “Eliminating experimental bias in anisotropic-flow measurements of high-energy nuclear collisions”, *Phys.Rev.* **C87** no. 4, (2013) 044907, arXiv:1209.2323 [nucl-ex].
- [48] E. Retinskaya, M. Luzum, and J.-Y. Ollitrault, “Directed flow at midrapidity in $\sqrt{s_{NN}} = 2.76$ TeV Pb+Pb collisions”, *Phys. Rev. Lett.* **108** (2012) 252302, arXiv:1203.0931 [nucl-th].
- [49] **ALICE** Collaboration, K. Aamodt *et al.*, “Alignment of the ALICE Inner Tracking System with cosmic-ray tracks”, *JINST* **5** (2010) P03003, arXiv:1001.0502 [physics.ins-det].
- [50] J. Alme *et al.*, “The ALICE TPC, a large 3-dimensional tracking device with fast readout for ultra-high multiplicity events”, *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment* **622** no. 1, (2010) 316 – 367, arXiv:1001.1950 [physics.ins-det].
- [51] **ALICE** Collaboration, F. Carnesecchi, “Performance of the ALICE Time-Of-Flight detector at the LHC”, *JINST* **14** no. 06, (2019) C06023, arXiv:1806.03825 [physics.ins-det].

- [52] **ALICE** Collaboration, S. Acharya *et al.*, “D-meson azimuthal anisotropy in midcentral Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV”, *Phys. Rev. Lett.* **120** no. 10, (2018) 102301, arXiv:1707.01005 [nucl-ex].
- [53] **ALICE** Collaboration, S. Acharya *et al.*, “Measurement of D^0 , D^+ , D^{*+} and D_s^+ production in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV”, *JHEP* **10** (2018) 174, arXiv:1804.09083 [nucl-ex].
- [54] U. Gursoy, D. Kharzeev, E. Marcus, K. Rajagopal, and C. Shen, “Charge-dependent Flow Induced by Magnetic and Electric Fields in Heavy Ion Collisions”, *Phys. Rev.* **C98** no. 5, (2018) 055201, arXiv:1806.05288 [hep-ph].
- [55] **STAR** Collaboration, L. Adamczyk *et al.*, “Directed Flow of Identified Particles in Au + Au Collisions at $\sqrt{s_{NN}} = 200$ GeV at RHIC”, *Phys. Rev. Lett.* **108** (2012) 202301, arXiv:1112.3930 [nucl-ex].
- [56] **STAR** Collaboration, L. Adamczyk *et al.*, “Beam-Energy Dependence of the Directed Flow of Protons, Antiprotons, and Pions in Au+Au Collisions”, *Phys. Rev. Lett.* **112** no. 16, (2014) 162301, arXiv:1401.3043 [nucl-ex].
- [57] **ALICE** Collaboration, B. Abelev *et al.*, “Centrality dependence of π , K, p production in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV”, *Phys. Rev.* **C88** (2013) 044910, arXiv:1303.0737 [hep-ex].
- [58] S. S. Gubser, “Symmetry constraints on generalizations of Bjorken flow”, *Phys. Rev.* **D82** (2010) 085027, arXiv:1006.0006 [hep-th].
- [59] C. Shen, Z. Qiu, H. Song, J. Bernhard, S. Bass, and U. Heinz, “The iEBE-VISHNU code package for relativistic heavy-ion collisions”, *Comput. Phys. Commun.* **199** (2016) 61–85, arXiv:1409.8164 [nucl-th].
- [60] Z. Citron *et al.*, “Future physics opportunities for high-density QCD at the LHC with heavy-ion and proton beams”, in *HL/HE-LHC Workshop: Workshop on the Physics of HL-LHC, and Perspectives at HE-LHC Geneva, Switzerland, June 18-20, 2018*. 2018. arXiv:1812.06772 [hep-ph].
- [61] **ALICE** Collaboration, S. Acharya *et al.*, “ALICE upgrade physics performance studies for 2018 Report on HL/HE-LHC physics”, *ALICE-PUBLIC-2019-001* (Feb, 2019). <https://cds.cern.ch/record/2661798>.

A The ALICE Collaboration

S. Acharya¹⁴¹, D. Adamová⁹⁴, A. Adler⁷⁴, J. Adolfsson⁸⁰, M.M. Aggarwal⁹⁹, G. Aglieri Rinella³³, M. Agnello³⁰, N. Agrawal^{10,53}, Z. Ahammed¹⁴¹, S. Ahmad¹⁶, S.U. Ahn⁷⁶, A. Akindinov⁹¹, M. Al-Turany¹⁰⁶, S.N. Alam¹⁴¹, D.S.D. Albuquerque¹²², D. Aleksandrov⁸⁷, B. Alessandro⁵⁸, H.M. Alfanda⁶, R. Alfaro Molina⁷¹, B. Ali¹⁶, Y. Ali¹⁴, A. Alici^{10,26,53}, A. Alkin², J. Alme²¹, T. Alt⁶⁸, L. Altenkamper²¹, I. Altsybeev¹¹², M.N. Anaam⁶, C. Andrei⁴⁷, D. Andreou³³, H.A. Andrews¹¹⁰, A. Andronic¹⁴⁴, M. Angeletti³³, V. Anguelov¹⁰³, C. Anson¹⁵, T. Antičić¹⁰⁷, F. Antinori⁵⁶, P. Antonioli⁵³, R. Anwar¹²⁵, N. Apadula⁷⁹, L. Aphecetche¹¹⁴, H. Appelshäuser⁶⁸, S. Arcelli²⁶, R. Arnaldi⁵⁸, M. Arratia⁷⁹, I.C. Arsene²⁰, M. Arslanok¹⁰³, A. Augustinus³³, R. Averbek¹⁰⁶, S. Aziz⁶¹, M.D. Azmi¹⁶, A. Badalà⁵⁵, Y.W. Baek⁴⁰, S. Bagnasco⁵⁸, X. Bai¹⁰⁶, R. Bailhache⁶⁸, R. Bala¹⁰⁰, A. Baldisseri¹³⁷, M. Ball⁴², S. Balouza¹⁰⁴, R. Barbera²⁷, L. Barioglio²⁵, G.G. Barnaföldi¹⁴⁵, L.S. Barnby⁹³, V. Barret¹³⁴, P. Bartalini⁶, K. Barth³³, E. Bartsch⁶⁸, F. Baruffaldi²⁸, N. Bastid¹³⁴, S. Basu¹⁴³, G. Batigne¹¹⁴, B. Batyunya⁷⁵, D. Bauri⁴⁸, J.L. Bazo Alba¹¹¹, I.G. Bearden⁸⁸, C. Bedda⁶³, N.K. Behera⁶⁰, I. Belikov¹³⁶, A.D.C. Bell Hechavarria¹⁴⁴, F. Bellini³³, R. Bellwied¹²⁵, V. Belyaev⁹², G. Bencedi¹⁴⁵, S. Beole²⁵, A. Bercuci⁴⁷, Y. Berdnikov⁹⁷, D. Berenyi¹⁴⁵, R.A. Bertens¹³⁰, D. Berzano⁵⁸, M.G. Besoiu⁶⁷, L. Betev³³, A. Bhasin¹⁰⁰, I.R. Bhat¹⁰⁰, M.A. Bhat³, H. Bhatt⁴⁸, B. Bhattacharjee⁴¹, A. Bianchi²⁵, L. Bianchi²⁵, N. Bianchi⁵¹, J. Bielčik³⁶, J. Bielčíková⁹⁴, A. Bilandzic^{104,117}, G. Biro¹⁴⁵, R. Biswas³, S. Biswas³, J.T. Blair¹¹⁹, D. Blau⁸⁷, C. Blume⁶⁸, G. Boca¹³⁹, F. Bock^{33,95}, A. Bogdanov⁹², S. Boi²³, L. Boldizsár¹⁴⁵, A. Bolozdynya⁹², M. Bombara³⁷, G. Bonomi¹⁴⁰, H. Borel¹³⁷, A. Borisso^{92,144}, H. Bossi¹⁴⁶, E. Botta²⁵, L. Bratrud⁶⁸, P. Braun-Munzinger¹⁰⁶, M. Bregant¹²¹, M. Broz³⁶, E.J. Brucken⁴³, E. Bruna⁵⁸, G.E. Bruno¹⁰⁵, M.D. Buckland¹²⁷, D. Budnikov¹⁰⁸, H. Buesching⁶⁸, S. Bufalino³⁰, O. Bugnon¹¹⁴, P. Buhler¹¹³, P. Buncic³³, Z. Buthelezi^{72,131}, J.B. Butt¹⁴, J.T. Buxton⁹⁶, S.A. Bysiak¹¹⁸, D. Caffarri⁸⁹, A. Caliva¹⁰⁶, E. Calvo Villar¹¹¹, R.S. Camacho⁴⁴, P. Camerini²⁴, A.A. Capon¹¹³, F. Carnesecchi^{10,26}, R. Caron¹³⁷, J. Castillo Castellanos¹³⁷, A.J. Castro¹³⁰, E.A.R. Casula⁵⁴, F. Catalano³⁰, C. Ceballos Sanchez⁵², P. Chakraborty⁴⁸, S. Chandra¹⁴¹, W. Chang⁶, S. Chapeland³³, M. Chartier¹²⁷, S. Chattopadhyay¹⁴¹, S. Chattopadhyay¹⁰⁹, A. Chauvin²³, C. Cheshkov¹³⁵, B. Cheynis¹³⁵, V. Chibante Barroso³³, D.D. Chinellato¹²², S. Cho⁶⁰, P. Chochula³³, T. Chowdhury¹³⁴, P. Christakoglou⁸⁹, C.H. Christensen⁸⁸, P. Christiansen⁸⁰, T. Chujo¹³³, C. Cicalo⁵⁴, L. Cifarelli^{10,26}, F. Cindolo⁵³, J. Cleymans¹²⁴, F. Colamaria⁵², D. Colella⁵², A. Collu⁷⁹, M. Colocci²⁶, M. Concas^{58,ii}, G. Conesa Balbastre⁷⁸, Z. Conesa del Valle⁶¹, G. Contin^{24,127}, J.G. Contreras³⁶, T.M. Cormier⁹⁵, Y. Corrales Morales²⁵, P. Cortese³¹, M.R. Cosentino¹²³, F. Costa³³, S. Costanza¹³⁹, P. Crochet¹³⁴, E. Cuautle⁶⁹, P. Cui⁶, L. Cunqueiro⁹⁵, D. Dabrowski¹⁴², T. Dahms^{104,117}, A. Dainese⁵⁶, F.P.A. Damas^{114,137}, M.C. Danisch¹⁰³, A. Danu⁶⁷, D. Das¹⁰⁹, I. Das¹⁰⁹, P. Das⁸⁵, P. Das³, S. Das³, A. Dash⁸⁵, S. Dash⁴⁸, S. De⁸⁵, A. De Caro²⁹, G. de Cataldo⁵², J. de Cuveland³⁸, A. De Falco²³, D. De Gruttola¹⁰, N. De Marco⁵⁸, S. De Pasquale²⁹, S. Deb⁴⁹, B. Debjani³, H.F. Degenhardt¹²¹, K.R. Deja¹⁴², A. Deloff⁸⁴, S. Delsanto^{25,131}, D. Devetak¹⁰⁶, P. Dhankher⁴⁸, D. Di Bari³², A. Di Mauro³³, R.A. Diaz⁸, T. Dietel¹²⁴, P. Dillenseger⁶⁸, Y. Ding⁶, R. Divia³³, D.U. Dixit¹⁹, Ø. Djuvsland²¹, U. Dmitrieva⁶², A. Dobrin^{33,67}, B. Dönigus⁶⁸, O. Dordic²⁰, A.K. Dubey¹⁴¹, A. Dubla¹⁰⁶, S. Dudi⁹⁹, M. Dukhishyam⁸⁵, P. Dupieux¹³⁴, R.J. Ehlers¹⁴⁶, V.N. Eikeland²¹, D. Elia⁵², H. Engel⁷⁴, E. Epple¹⁴⁶, B. Erazmus¹¹⁴, F. Erhardt⁹⁸, A. Erokhin¹¹², M.R. Ersdal²¹, B. Espagnon⁶¹, S. Esumi¹³³, G. Eulisse³³, D. Evans¹¹⁰, S. Evdokimov⁹⁰, L. Fabbietti^{104,117}, M. Faggin²⁸, J. Faivre⁷⁸, F. Fan⁶, A. Fantoni⁵¹, M. Fasel⁹⁵, P. Fecchio³⁰, A. Feliciello⁵⁸, G. Feofilov¹¹², A. Fernández Téllez⁴⁴, A. Ferrero¹³⁷, A. Ferretti²⁵, A. Festanti³³, V.J.G. Feuillard¹⁰³, J. Figiel¹¹⁸, S. Filchagin¹⁰⁸, D. Finogeev⁶², F.M. Fiorda²¹, G. Fiorenza⁵², F. Flor¹²⁵, S. Foertsch⁷², P. Foka¹⁰⁶, S. Fokin⁸⁷, E. Fragiaco⁵⁹, U. Frankfeld¹⁰⁶, U. Fuchs³³, C. Furget⁷⁸, A. Furs⁶², M. Fusco Girard²⁹, J.J. Gaardhøje⁸⁸, M. Gagliardi²⁵, A.M. Gago¹¹¹, A. Gal¹³⁶, C.D. Galvan¹²⁰, P. Ganoti⁸³, C. Garabatos¹⁰⁶, E. Garcia-Solis¹¹, K. Garg²⁷, C. Gargiulo³³, A. Garibli⁸⁶, K. Garner¹⁴⁴, P. Gasik^{104,117}, E.F. Gauger¹¹⁹, M.B. Gay Ducati⁷⁰, M. Germain¹¹⁴, J. Ghosh¹⁰⁹, P. Ghosh¹⁴¹, S.K. Ghosh³, P. Gianotti⁵¹, P. Giubellino^{58,106}, P. Giubilato²⁸, P. Gläsel¹⁰³, D.M. Gomez Coral⁷¹, A. Gomez Ramirez⁷⁴, V. Gonzalez¹⁰⁶, P. González-Zamora⁴⁴, S. Gorbunov³⁸, L. Görlich¹¹⁸, S. Gotovac³⁴, V. Grabski⁷¹, L.K. Graczykowski¹⁴², K.L. Graham¹¹⁰, L. Greiner⁷⁹, A. Grelli⁶³, C. Grigoras³³, V. Grigoriev⁹², A. Grigoryan¹, S. Grigoryan⁷⁵, O.S. Groettvik²¹, F. Grosa³⁰, J.F. Grosse-Oetringhaus³³, R. Grosso¹⁰⁶, R. Guernane⁷⁸, M. Guittiere¹¹⁴, K. Gulbrandsen⁸⁸, T. Gunji¹³², A. Gupta¹⁰⁰, R. Gupta¹⁰⁰, I.B. Guzman⁴⁴, R. Haake¹⁴⁶, M.K. Habib¹⁰⁶, C. Hadjidakis⁶¹, H. Hamagaki⁸¹, G. Hamar¹⁴⁵, M. Hamid⁶, R. Hannigan¹¹⁹, M.R. Haque^{63,85}, A. Harlanderova¹⁰⁶, J.W. Harris¹⁴⁶, A. Harton¹¹, J.A. Hasenbichler³³, H. Hassan⁹⁵, D. Hatzifotiadou^{10,53}, P. Hauer⁴², S. Hayashi¹³², S.T. Heckel^{68,104}, E. Hellbär⁶⁸, H. Helstrup³⁵, A. Hergelegiu⁴⁷, T. Herman³⁶, E.G. Hernandez⁴⁴, G. Herrera Corral⁹, F. Herrmann¹⁴⁴, K.F. Hetland³⁵, T.E. Hilden⁴³, H. Hillemanns³³, C. Hills¹²⁷, B. Hippolyte¹³⁶, B. Hohlweger¹⁰⁴, D. Horak³⁶, A. Hornung⁶⁸, S. Hornung¹⁰⁶, R. Hosokawa^{15,133}, P. Hristov³³, C. Huang⁶¹, C. Hughes¹³⁰, P. Huhn⁶⁸, T.J. Humanic⁹⁶,

H. Hushnud¹⁰⁹, L.A. Husova¹⁴⁴, N. Hussain⁴¹, S.A. Hussain¹⁴, D. Hutter³⁸, J.P. Iddon^{33,127}, R. Ilkaev¹⁰⁸, M. Inaba¹³³, G.M. Innocenti³³, M. Ippolitov⁸⁷, A. Isakov⁹⁴, M.S. Islam¹⁰⁹, M. Ivanov¹⁰⁶, V. Ivanov⁹⁷, V. Izucheev⁹⁰, B. Jacak⁷⁹, N. Jacazio⁵³, P.M. Jacobs⁷⁹, S. Jadlovská¹¹⁶, J. Jadlovsky¹¹⁶, S. Jaelani⁶³, C. Jahnke¹²¹, M.J. Jakubowska¹⁴², M.A. Janik¹⁴², T. Janson⁷⁴, M. Jercic⁹⁸, O. Jevons¹¹⁰, M. Jin¹²⁵, F. Jonas^{95,144}, P.G. Jones¹¹⁰, J. Jung⁶⁸, M. Jung⁶⁸, A. Jusko¹¹⁰, P. Kalinak⁶⁴, A. Kalweit³³, V. Kaplin⁹², S. Kar⁶, A. Karasu Uysal⁷⁷, O. Karavichev⁶², T. Karavicheva⁶², P. Karczmarczyk³³, E. Karpechev⁶², A. Kazantsev⁸⁷, U. Kebschull⁷⁴, R. Keidel⁴⁶, M. Keil³³, B. Ketzer⁴², Z. Khabanova⁸⁹, A.M. Khan⁶, S. Khan¹⁶, S.A. Khan¹⁴¹, A. Khanzadeev⁹⁷, Y. Kharlov⁹⁰, A. Khatun¹⁶, A. Khuntia¹¹⁸, B. Kileng³⁵, B. Kim⁶⁰, B. Kim¹³³, D. Kim¹⁴⁷, D.J. Kim¹²⁶, E.J. Kim⁷³, H. Kim^{17,147}, J. Kim¹⁴⁷, J.S. Kim⁴⁰, J. Kim¹⁰³, J. Kim¹⁴⁷, J. Kim⁷³, M. Kim¹⁰³, S. Kim¹⁸, T. Kim¹⁴⁷, T. Kim¹⁴⁷, S. Kirsch^{38,68}, I. Kisel³⁸, S. Kiselev⁹¹, A. Kisiel¹⁴², J.L. Klay⁵, C. Klein⁶⁸, J. Klein⁵⁸, S. Klein⁷⁹, C. Klein-Bösing¹⁴⁴, M. Kleiner⁶⁸, A. Kluge³³, M.L. Knichel³³, A.G. Knospe¹²⁵, C. Kobdaj¹¹⁵, M.K. Köhler¹⁰³, T. Kollegger¹⁰⁶, A. Kondratyev⁷⁵, N. Kondratyeva⁹², E. Kondratyuk⁹⁰, J. König⁶⁸, P.J. Konopka³³, L. Koska¹¹⁶, O. Kovalenko⁸⁴, V. Kovalenko¹¹², M. Kowalski¹¹⁸, I. Králik⁶⁴, A. Kravčáková³⁷, L. Kreis¹⁰⁶, M. Krivda^{64,110}, F. Krizek⁹⁴, K. Krizkova Gajdosova³⁶, M. Krüger⁶⁸, E. Kryshen⁹⁷, M. Krzewicki³⁸, A.M. Kubera⁹⁶, V. Kučera⁶⁰, C. Kuhn¹³⁶, P.G. Kuijjer⁸⁹, L. Kumar⁹⁹, S. Kumar⁴⁸, S. Kundu⁸⁵, P. Kurashvili⁸⁴, A. Kurepin⁶², A.B. Kurepin⁶², A. Kuryakin¹⁰⁸, S. Kushpil⁹⁴, J. Kvapil¹¹⁰, M.J. Kweon⁶⁰, J.Y. Kwon⁶⁰, Y. Kwon¹⁴⁷, S.L. La Pointe³⁸, P. La Rocca²⁷, Y.S. Lai⁷⁹, R. Langoy¹²⁹, K. Lapidus³³, A. Lardeux²⁰, P. Larionov⁵¹, E. Laudi³³, R. Lavicka³⁶, T. Lazareva¹¹², R. Lea²⁴, L. Leardini¹⁰³, J. Lee¹³³, S. Lee¹⁴⁷, F. Lehas⁸⁹, S. Lehner¹¹³, J. Lehrbach³⁸, R.C. Lemmon⁹³, I. León Monzón¹²⁰, E.D. Lesser¹⁹, M. Lettrich³³, P. Lévai¹⁴⁵, X. Li¹², X.L. Li⁶, J. Lien¹²⁹, R. Lietava¹¹⁰, B. Lim¹⁷, V. Lindenstruth³⁸, S.W. Lindsay¹²⁷, C. Lippmann¹⁰⁶, M.A. Lisa⁹⁶, V. Litichevskiy⁴³, A. Liu¹⁹, S. Liu⁹⁶, W.J. Llope¹⁴³, I.M. Lofnes²¹, V. Loginov⁹², C. Loizides⁹⁵, P. Loncar³⁴, X. Lopez¹³⁴, E. López Torres⁸, J.R. Luhder¹⁴⁴, M. Lunardon²⁸, G. Luparello⁵⁹, Y. Ma³⁹, A. Maevskaya⁶², M. Mager³³, S.M. Mahmood²⁰, T. Mahmoud⁴², A. Maire¹³⁶, R.D. Majka¹⁴⁶, M. Malaev⁹⁷, Q.W. Malik²⁰, L. Malinina^{75,iii}, D. Mal'Kevich⁹¹, P. Malzacher¹⁰⁶, G. Mandaglio⁵⁵, V. Manko⁸⁷, F. Manso¹³⁴, V. Manzari⁵², Y. Mao⁶, M. Marchisone¹³⁵, J. Mareš⁶⁶, G.V. Margagliotti²⁴, A. Margotti⁵³, J. Margutti⁶³, A. Marín¹⁰⁶, C. Markert¹¹⁹, M. Marquard⁶⁸, N.A. Martin¹⁰³, P. Martinengo³³, J.L. Martinez¹²⁵, M.I. Martínez⁴⁴, G. Martínez García¹¹⁴, M. Martinez Pedreira³³, S. Masciocchi¹⁰⁶, M. Masera²⁵, A. Masoni⁵⁴, L. Massacrier⁶¹, E. Masson¹¹⁴, A. Mastroserio^{52,138}, A.M. Mathis^{104,117}, O. Matonoha⁸⁰, P.F.T. Matuoka¹²¹, A. Matyja¹¹⁸, C. Mayer¹¹⁸, M. Mazzilli⁵², M.A. Mazzoni⁵⁷, A.F. Mechler⁶⁸, F. Meddi²², Y. Melikyan^{62,92}, A. Menchaca-Rocha⁷¹, C. Mengke⁶, E. Meninno^{29,113}, M. Meres¹³, S. Mhlanga¹²⁴, Y. Miake¹³³, L. Micheletti²⁵, D.L. Mihaylov¹⁰⁴, K. Mikhaylov^{75,91}, A. Mischke^{63,i}, A.N. Mishra⁶⁹, D. Miśkowiec¹⁰⁶, A. Modak³, N. Mohammadi³³, A.P. Mohanty⁶³, B. Mohanty⁸⁵, M. Mohisin Khan^{16,iv}, C. Mordasini¹⁰⁴, D.A. Moreira De Godoy¹⁴⁴, L.A.P. Moreno⁴⁴, I. Morozov⁶², A. Morsch³³, T. Mrnjavac³³, V. Muccifora⁵¹, E. Mudnic³⁴, D. Mühlheim¹⁴⁴, S. Muhuri¹⁴¹, J.D. Mulligan⁷⁹, M.G. Munhoz¹²¹, R.H. Munzer⁶⁸, H. Murakami¹³², S. Murray¹²⁴, L. Musa³³, J. Musinsky⁶⁴, C.J. Myers¹²⁵, J.W. Myrcha¹⁴², B. Naik⁴⁸, R. Nair⁸⁴, B.K. Nandi⁴⁸, R. Nania^{10,53}, E. Nappi⁵², M.U. Naru¹⁴, A.F. Nassirpour⁸⁰, C. Nattrass¹³⁰, R. Nayak⁴⁸, T.K. Nayak⁸⁵, S. Nazarenko¹⁰⁸, A. Neagu²⁰, R.A. Negrao De Oliveira⁶⁸, L. Nellen⁶⁹, S.V. Nesbo³⁵, G. Neskovic³⁸, D. Nesterov¹¹², L.T. Neumann¹⁴², B.S. Nielsen⁸⁸, S. Nikolaev⁸⁷, S. Nikulin⁸⁷, V. Nikulin⁹⁷, F. Noferini^{10,53}, P. Nomokonov⁷⁵, J. Norman^{78,127}, N. Novitzky¹³³, P. Nowakowski¹⁴², A. Nyanin⁸⁷, J. Nystrand²¹, M. Ogino⁸¹, A. Ohlson^{80,103}, J. Oleniacz¹⁴², A.C. Oliveira Da Silva^{121,130}, M.H. Oliver¹⁴⁶, C. Oppedisano⁵⁸, R. Orava⁴³, A. Ortiz Velasquez⁶⁹, A. Oskarsson⁸⁰, J. Otwinowski¹¹⁸, K. Oyama⁸¹, Y. Pachmayer¹⁰³, V. Pacik⁸⁸, D. Pagano¹⁴⁰, G. Paic⁶⁹, J. Pan¹⁴³, A.K. Pandey⁴⁸, S. Panebianco¹³⁷, P. Pareek^{49,141}, J. Park⁶⁰, J.E. Parkkila¹²⁶, S. Parmar⁹⁹, S.P. Pathak¹²⁵, R.N. Patra¹⁴¹, B. Paul^{23,58}, H. Pei⁶, T. Peitzmann⁶³, X. Peng⁶, L.G. Pereira⁷⁰, H. Pereira Da Costa¹³⁷, D. Peresunko⁸⁷, G.M. Perez⁸, E. Perez Lezama⁶⁸, V. Peskov⁶⁸, Y. Pestov⁴, V. Petráček³⁶, M. Petrovici⁴⁷, R.P. Pezzi⁷⁰, S. Piano⁵⁹, M. Pikna¹³, P. Pillot¹¹⁴, O. Pinazza^{33,53}, L. Pinsky¹²⁵, C. Pinto²⁷, S. Pisano^{10,51}, D. Pistone⁵⁵, M. Płoskoń⁷⁹, M. Planinic⁹⁸, F. Pliquett⁶⁸, J. Pluta¹⁴², S. Pochybova^{145,i}, M.G. Poghosyan⁹⁵, B. Polichtchouk⁹⁰, N. Poljak⁹⁸, A. Pop⁴⁷, H. Poppenborg¹⁴⁴, S. Porteboeuf-Houssais¹³⁴, V. Pozdniakov⁷⁵, S.K. Prasad³, R. Preghenella⁵³, F. Prino⁵⁸, C.A. Pruneau¹⁴³, I. Pshenichnov⁶², M. Puccio^{25,33}, J. Putschke¹⁴³, R.E. Quishpe¹²⁵, S. Ragoni¹¹⁰, S. Raha³, S. Rajput¹⁰⁰, J. Rak¹²⁶, A. Rakotozafindrabe¹³⁷, L. Ramello³¹, F. Rami¹³⁶, R. Raniwala¹⁰¹, S. Raniwala¹⁰¹, S.S. Räsänen⁴³, R. Rath⁴⁹, V. Ratza⁴², I. Ravasenga^{30,89}, K.F. Read^{95,130}, K. Redlich^{84,v}, A. Rehman²¹, P. Reichelt⁶⁸, F. Reidt³³, X. Ren⁶, R. Renfordt⁶⁸, Z. Rescakova³⁷, J.-P. Revol¹⁰, K. Reygers¹⁰³, V. Riabov⁹⁷, T. Richert^{80,88}, M. Richter²⁰, P. Riedler³³, W. Riegler³³, F. Riggi²⁷, C. Ristea⁶⁷, S.P. Rode⁴⁹, M. Rodríguez Cahuantzi⁴⁴, K. Røed²⁰, R. Rogalev⁹⁰, E. Rogochaya⁷⁵, D. Rohr³³, D. Röhrich²¹, P.S. Rokita¹⁴², F. Ronchetti⁵¹, E.D. Rosas⁶⁹, K. Roslon¹⁴², A. Rossi^{28,56}, A. Rotondi¹³⁹, A. Roy⁴⁹, P. Roy¹⁰⁹, O.V. Rueda⁸⁰, R. Rui²⁴,

B. Rumyantsev⁷⁵, A. Rustamov⁸⁶, E. Ryabinkin⁸⁷, Y. Ryabov⁹⁷, A. Rybicki¹¹⁸, H. Rytkonen¹²⁶, O.A.M. Saari⁴³, S. Sadhu¹⁴¹, S. Sadovsky⁹⁰, K. Šafařík³⁶, S.K. Saha¹⁴¹, B. Sahoo⁴⁸, P. Sahoo^{48,49}, R. Sahoo⁴⁹, S. Sahoo⁶⁵, P.K. Sahu⁶⁵, J. Saini¹⁴¹, S. Sakai¹³³, S. Sambyal¹⁰⁰, V. Samsonov^{92,97}, D. Sarkar¹⁴³, N. Sarkar¹⁴¹, P. Sarma⁴¹, V.M. Sarti¹⁰⁴, M.H.P. Sas⁶³, E. Scapparone⁵³, B. Schaefer⁹⁵, J. Schambach¹¹⁹, H.S. Scheid⁶⁸, C. Schiaua⁴⁷, R. Schicker¹⁰³, A. Schmah¹⁰³, C. Schmidt¹⁰⁶, H.R. Schmidt¹⁰², M.O. Schmidt¹⁰³, M. Schmidt¹⁰², N.V. Schmidt^{68,95}, A.R. Schmier¹³⁰, J. Schukraft⁸⁸, Y. Schutz^{33,136}, K. Schwarz¹⁰⁶, K. Schweda¹⁰⁶, G. Scioli²⁶, E. Scomparin⁵⁸, M. Šefčík³⁷, J.E. Seger¹⁵, Y. Sekiguchi¹³², D. Sekihata¹³², I. Selyuzhenkov^{92,106}, S. Senyukov¹³⁶, D. Serebryakov⁶², E. Serradilla⁷¹, A. Sevcenco⁶⁷, A. Shabanov⁶², A. Shabetai¹¹⁴, R. Shahoyan³³, W. Shaikh¹⁰⁹, A. Shangaraev⁹⁰, A. Sharma⁹⁹, A. Sharma¹⁰⁰, H. Sharma¹¹⁸, M. Sharma¹⁰⁰, N. Sharma⁹⁹, A.I. Sheikh¹⁴¹, K. Shigaki⁴⁵, M. Shimomura⁸², S. Shirinkin⁹¹, Q. Shou³⁹, Y. Sibiriak⁸⁷, S. Siddhanta⁵⁴, T. Siemarczuk⁸⁴, D. Silvermyr⁸⁰, G. Simatovic⁸⁹, G. Simonetti^{33,104}, R. Singh⁸⁵, R. Singh¹⁰⁰, R. Singh⁴⁹, V.K. Singh¹⁴¹, V. Singhal¹⁴¹, T. Sinha¹⁰⁹, B. Sitar¹³, M. Sitta³¹, T.B. Skaali²⁰, M. Slupecki¹²⁶, N. Smirnov¹⁴⁶, R.J.M. Snellings⁶³, T.W. Snellman^{43,126}, C. Soncco¹¹¹, J. Song^{60,125}, A. Songmoolnak¹¹⁵, F. Soramel²⁸, S. Sorensen¹³⁰, I. Sputowska¹¹⁸, J. Stachel¹⁰³, I. Stan⁶⁷, P. Stankus⁹⁵, P.J. Steffanic¹³⁰, E. Stenlund⁸⁰, D. Stocco¹¹⁴, M.M. Stortvedt³⁵, L.D. Stritto²⁹, A.A.P. Suaide¹²¹, T. Sugitate⁴⁵, C. Suire⁶¹, M. Suleymanov¹⁴, M. Suljic³³, R. Sultanov⁹¹, M. Šumbera⁹⁴, S. Sumowidagdo⁵⁰, S. Swain⁶⁵, A. Szabo¹³, I. Szarka¹³, U. Tabassam¹⁴, G. Taillepied¹³⁴, J. Takahashi¹²², G.J. Tambave²¹, S. Tang^{6,134}, M. Tarhini¹¹⁴, M.G. Tarzila⁴⁷, A. Tauro³³, G. Tejeda Muñoz⁴⁴, A. Telesca³³, C. Terrevoli¹²⁵, D. Thakur⁴⁹, S. Thakur¹⁴¹, D. Thomas¹¹⁹, F. Thoresen⁸⁸, R. Tieulent¹³⁵, A. Tikhonov⁶², A.R. Timmins¹²⁵, A. Toia⁶⁸, N. Topilskaya⁶², M. Toppi⁵¹, F. Torres-Acosta¹⁹, S.R. Torres^{9,120}, A. Trifiro⁵⁵, S. Tripathy⁴⁹, T. Tripathy⁴⁸, S. Trogolo²⁸, G. Trombetta³², L. Tropp³⁷, V. Trubnikov², W.H. Trzaska¹²⁶, T.P. Trzcinski¹⁴², B.A. Trzeciak⁶³, T. Tsuji¹³², A. Tumkin¹⁰⁸, R. Turrisi⁵⁶, T.S. Tveter²⁰, K. Ullaland²¹, E.N. Umaka¹²⁵, A. Uras¹³⁵, G.L. Usai²³, A. Utrobicic⁹⁸, M. Vala³⁷, N. Valle¹³⁹, S. Vallero⁵⁸, N. van der Kolk⁶³, L.V.R. van Doremalen⁶³, M. van Leeuwen⁶³, P. Vande Vyvre³³, D. Varga¹⁴⁵, Z. Varga¹⁴⁵, M. Varga-Kofarago¹⁴⁵, A. Vargas⁴⁴, M. Vasileiou⁸³, A. Vasiliev⁸⁷, O. Vázquez Doce^{104,117}, V. Vechernin¹¹², A.M. Veen⁶³, E. Vercellin²⁵, S. Vergara Limón⁴⁴, L. Vermunt⁶³, R. Vernet⁷, R. Vértesi¹⁴⁵, L. Vickovic³⁴, Z. Vilakazi¹³¹, O. Villalobos Baillie¹¹⁰, A. Villatoro Tello⁴⁴, G. Vito⁵², A. Vinogradov⁸⁷, T. Virgili²⁹, V. Vislavicius⁸⁸, A. Vodopyanov⁷⁵, B. Volkel³³, M.A. Völkl¹⁰², K. Voloshin⁹¹, S.A. Voloshin¹⁴³, G. Volpe³², B. von Haller³³, I. Vorobyev¹⁰⁴, D. Voscek¹¹⁶, J. Vrláková³⁷, B. Wagner²¹, M. Weber¹¹³, S.G. Weber¹⁴⁴, A. Wegrzynek³³, D.F. Weiser¹⁰³, S.C. Wenzel³³, J.P. Wessels¹⁴⁴, J. Wiechula⁶⁸, J. Wikne²⁰, G. Wilk⁸⁴, J. Wilkinson^{10,53}, G.A. Willems³³, E. Willsher¹¹⁰, B. Windelband¹⁰³, M. Winn¹³⁷, W.E. Witt¹³⁰, Y. Wu¹²⁸, R. Xu⁶, S. Yalcin⁷⁷, K. Yamakawa⁴⁵, S. Yang²¹, S. Yano¹³⁷, Z. Yin⁶, H. Yokoyama⁶³, I.-K. Yoo¹⁷, J.H. Yoon⁶⁰, S. Yuan²¹, A. Yuncu¹⁰³, V. Yurchenko², V. Zaccaro²⁴, A. Zaman¹⁴, C. Zampolli³³, H.J.C. Zanoli⁶³, N. Zardoshti³³, A. Zarochentsev¹¹², P. Závada⁶⁶, N. Zaviyalov¹⁰⁸, H. Zbroszczyk¹⁴², M. Zhalov⁹⁷, S. Zhang³⁹, X. Zhang⁶, Z. Zhang⁶, V. Zhrebchevskii¹¹², D. Zhou⁶, Y. Zhou⁸⁸, Z. Zhou²¹, J. Zhu^{6,106}, Y. Zhu⁶, A. Zichichi^{10,26}, M.B. Zimmermann³³, G. Zinovjev², N. Zurlo¹⁴⁰,

Affiliation notes

ⁱ Deceased

ⁱⁱ Dipartimento DET del Politecnico di Torino, Turin, Italy

ⁱⁱⁱ M.V. Lomonosov Moscow State University, D.V. Skobeltsyn Institute of Nuclear Physics, Moscow, Russia

^{iv} Department of Applied Physics, Aligarh Muslim University, Aligarh, India

^v Institute of Theoretical Physics, University of Wrocław, Poland

Collaboration Institutes

¹ A.I. Alikhanyan National Science Laboratory (Yerevan Physics Institute) Foundation, Yerevan, Armenia

² Bogolyubov Institute for Theoretical Physics, National Academy of Sciences of Ukraine, Kiev, Ukraine

³ Bose Institute, Department of Physics and Centre for Astroparticle Physics and Space Science (CAPSS), Kolkata, India

⁴ Budker Institute for Nuclear Physics, Novosibirsk, Russia

⁵ California Polytechnic State University, San Luis Obispo, California, United States

⁶ Central China Normal University, Wuhan, China

⁷ Centre de Calcul de l'IN2P3, Villeurbanne, Lyon, France

⁸ Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Havana, Cuba

⁹ Centro de Investigación y de Estudios Avanzados (CINVESTAV), Mexico City and Mérida, Mexico

- 10 Centro Fermi - Museo Storico della Fisica e Centro Studi e Ricerche “Enrico Fermi”, Rome, Italy
- 11 Chicago State University, Chicago, Illinois, United States
- 12 China Institute of Atomic Energy, Beijing, China
- 13 Comenius University Bratislava, Faculty of Mathematics, Physics and Informatics, Bratislava, Slovakia
- 14 COMSATS University Islamabad, Islamabad, Pakistan
- 15 Creighton University, Omaha, Nebraska, United States
- 16 Department of Physics, Aligarh Muslim University, Aligarh, India
- 17 Department of Physics, Pusan National University, Pusan, Republic of Korea
- 18 Department of Physics, Sejong University, Seoul, Republic of Korea
- 19 Department of Physics, University of California, Berkeley, California, United States
- 20 Department of Physics, University of Oslo, Oslo, Norway
- 21 Department of Physics and Technology, University of Bergen, Bergen, Norway
- 22 Dipartimento di Fisica dell’Università ‘La Sapienza’ and Sezione INFN, Rome, Italy
- 23 Dipartimento di Fisica dell’Università and Sezione INFN, Cagliari, Italy
- 24 Dipartimento di Fisica dell’Università and Sezione INFN, Trieste, Italy
- 25 Dipartimento di Fisica dell’Università and Sezione INFN, Turin, Italy
- 26 Dipartimento di Fisica e Astronomia dell’Università and Sezione INFN, Bologna, Italy
- 27 Dipartimento di Fisica e Astronomia dell’Università and Sezione INFN, Catania, Italy
- 28 Dipartimento di Fisica e Astronomia dell’Università and Sezione INFN, Padova, Italy
- 29 Dipartimento di Fisica ‘E.R. Caianiello’ dell’Università and Gruppo Collegato INFN, Salerno, Italy
- 30 Dipartimento DISAT del Politecnico and Sezione INFN, Turin, Italy
- 31 Dipartimento di Scienze e Innovazione Tecnologica dell’Università del Piemonte Orientale and INFN Sezione di Torino, Alessandria, Italy
- 32 Dipartimento Interateneo di Fisica ‘M. Merlin’ and Sezione INFN, Bari, Italy
- 33 European Organization for Nuclear Research (CERN), Geneva, Switzerland
- 34 Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, University of Split, Split, Croatia
- 35 Faculty of Engineering and Science, Western Norway University of Applied Sciences, Bergen, Norway
- 36 Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Prague, Czech Republic
- 37 Faculty of Science, P.J. Šafárik University, Košice, Slovakia
- 38 Frankfurt Institute for Advanced Studies, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany
- 39 Fudan University, Shanghai, China
- 40 Gangneung-Wonju National University, Gangneung, Republic of Korea
- 41 Gauhati University, Department of Physics, Guwahati, India
- 42 Helmholtz-Institut für Strahlen- und Kernphysik, Rheinische Friedrich-Wilhelms-Universität Bonn, Bonn, Germany
- 43 Helsinki Institute of Physics (HIP), Helsinki, Finland
- 44 High Energy Physics Group, Universidad Autónoma de Puebla, Puebla, Mexico
- 45 Hiroshima University, Hiroshima, Japan
- 46 Hochschule Worms, Zentrum für Technologietransfer und Telekommunikation (ZTT), Worms, Germany
- 47 Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest, Romania
- 48 Indian Institute of Technology Bombay (IIT), Mumbai, India
- 49 Indian Institute of Technology Indore, Indore, India
- 50 Indonesian Institute of Sciences, Jakarta, Indonesia
- 51 INFN, Laboratori Nazionali di Frascati, Frascati, Italy
- 52 INFN, Sezione di Bari, Bari, Italy
- 53 INFN, Sezione di Bologna, Bologna, Italy
- 54 INFN, Sezione di Cagliari, Cagliari, Italy
- 55 INFN, Sezione di Catania, Catania, Italy
- 56 INFN, Sezione di Padova, Padova, Italy
- 57 INFN, Sezione di Roma, Rome, Italy
- 58 INFN, Sezione di Torino, Turin, Italy
- 59 INFN, Sezione di Trieste, Trieste, Italy
- 60 Inha University, Incheon, Republic of Korea

- 61 Institut de Physique Nucléaire d'Orsay (IPNO), Institut National de Physique Nucléaire et de Physique des Particules (IN2P3/CNRS), Université de Paris-Sud, Université Paris-Saclay, Orsay, France
- 62 Institute for Nuclear Research, Academy of Sciences, Moscow, Russia
- 63 Institute for Subatomic Physics, Utrecht University/Nikhef, Utrecht, Netherlands
- 64 Institute of Experimental Physics, Slovak Academy of Sciences, Košice, Slovakia
- 65 Institute of Physics, Homi Bhabha National Institute, Bhubaneswar, India
- 66 Institute of Physics of the Czech Academy of Sciences, Prague, Czech Republic
- 67 Institute of Space Science (ISS), Bucharest, Romania
- 68 Institut für Kernphysik, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany
- 69 Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Mexico City, Mexico
- 70 Instituto de Física, Universidade Federal do Rio Grande do Sul (UFRGS), Porto Alegre, Brazil
- 71 Instituto de Física, Universidad Nacional Autónoma de México, Mexico City, Mexico
- 72 iThemba LABS, National Research Foundation, Somerset West, South Africa
- 73 Jeonbuk National University, Jeonju, Republic of Korea
- 74 Johann-Wolfgang-Goethe Universität Frankfurt Institut für Informatik, Fachbereich Informatik und Mathematik, Frankfurt, Germany
- 75 Joint Institute for Nuclear Research (JINR), Dubna, Russia
- 76 Korea Institute of Science and Technology Information, Daejeon, Republic of Korea
- 77 KTO Karatay University, Konya, Turkey
- 78 Laboratoire de Physique Subatomique et de Cosmologie, Université Grenoble-Alpes, CNRS-IN2P3, Grenoble, France
- 79 Lawrence Berkeley National Laboratory, Berkeley, California, United States
- 80 Lund University Department of Physics, Division of Particle Physics, Lund, Sweden
- 81 Nagasaki Institute of Applied Science, Nagasaki, Japan
- 82 Nara Women's University (NWU), Nara, Japan
- 83 National and Kapodistrian University of Athens, School of Science, Department of Physics, Athens, Greece
- 84 National Centre for Nuclear Research, Warsaw, Poland
- 85 National Institute of Science Education and Research, Homi Bhabha National Institute, Jatni, India
- 86 National Nuclear Research Center, Baku, Azerbaijan
- 87 National Research Centre Kurchatov Institute, Moscow, Russia
- 88 Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark
- 89 Nikhef, National institute for subatomic physics, Amsterdam, Netherlands
- 90 NRC Kurchatov Institute IHEP, Protvino, Russia
- 91 NRC «Kurchatov Institute» - ITEP, Moscow, Russia
- 92 NRNU Moscow Engineering Physics Institute, Moscow, Russia
- 93 Nuclear Physics Group, STFC Daresbury Laboratory, Daresbury, United Kingdom
- 94 Nuclear Physics Institute of the Czech Academy of Sciences, Řež u Prahy, Czech Republic
- 95 Oak Ridge National Laboratory, Oak Ridge, Tennessee, United States
- 96 Ohio State University, Columbus, Ohio, United States
- 97 Petersburg Nuclear Physics Institute, Gatchina, Russia
- 98 Physics department, Faculty of science, University of Zagreb, Zagreb, Croatia
- 99 Physics Department, Panjab University, Chandigarh, India
- 100 Physics Department, University of Jammu, Jammu, India
- 101 Physics Department, University of Rajasthan, Jaipur, India
- 102 Physikalisches Institut, Eberhard-Karls-Universität Tübingen, Tübingen, Germany
- 103 Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
- 104 Physik Department, Technische Universität München, Munich, Germany
- 105 Politecnico di Bari, Bari, Italy
- 106 Research Division and ExtreMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung GmbH, Darmstadt, Germany
- 107 Rudjer Bošković Institute, Zagreb, Croatia
- 108 Russian Federal Nuclear Center (VNIIEF), Sarov, Russia
- 109 Saha Institute of Nuclear Physics, Homi Bhabha National Institute, Kolkata, India
- 110 School of Physics and Astronomy, University of Birmingham, Birmingham, United Kingdom
- 111 Sección Física, Departamento de Ciencias, Pontificia Universidad Católica del Perú, Lima, Peru

- 112 St. Petersburg State University, St. Petersburg, Russia
- 113 Stefan Meyer Institut für Subatomare Physik (SMI), Vienna, Austria
- 114 SUBATECH, IMT Atlantique, Université de Nantes, CNRS-IN2P3, Nantes, France
- 115 Suranaree University of Technology, Nakhon Ratchasima, Thailand
- 116 Technical University of Košice, Košice, Slovakia
- 117 Technische Universität München, Excellence Cluster 'Universe', Munich, Germany
- 118 The Henryk Niewodniczanski Institute of Nuclear Physics, Polish Academy of Sciences, Cracow, Poland
- 119 The University of Texas at Austin, Austin, Texas, United States
- 120 Universidad Autónoma de Sinaloa, Culiacán, Mexico
- 121 Universidade de São Paulo (USP), São Paulo, Brazil
- 122 Universidade Estadual de Campinas (UNICAMP), Campinas, Brazil
- 123 Universidade Federal do ABC, Santo Andre, Brazil
- 124 University of Cape Town, Cape Town, South Africa
- 125 University of Houston, Houston, Texas, United States
- 126 University of Jyväskylä, Jyväskylä, Finland
- 127 University of Liverpool, Liverpool, United Kingdom
- 128 University of Science and Technology of China, Hefei, China
- 129 University of South-Eastern Norway, Tonsberg, Norway
- 130 University of Tennessee, Knoxville, Tennessee, United States
- 131 University of the Witwatersrand, Johannesburg, South Africa
- 132 University of Tokyo, Tokyo, Japan
- 133 University of Tsukuba, Tsukuba, Japan
- 134 Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France
- 135 Université de Lyon, Université Lyon 1, CNRS/IN2P3, IPN-Lyon, Villeurbanne, Lyon, France
- 136 Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France, Strasbourg, France
- 137 Université Paris-Saclay Centre d'Etudes de Saclay (CEA), IRFU, Département de Physique Nucléaire (DPhN), Saclay, France
- 138 Università degli Studi di Foggia, Foggia, Italy
- 139 Università degli Studi di Pavia, Pavia, Italy
- 140 Università di Brescia, Brescia, Italy
- 141 Variable Energy Cyclotron Centre, Homi Bhabha National Institute, Kolkata, India
- 142 Warsaw University of Technology, Warsaw, Poland
- 143 Wayne State University, Detroit, Michigan, United States
- 144 Westfälische Wilhelms-Universität Münster, Institut für Kernphysik, Münster, Germany
- 145 Wigner Research Centre for Physics, Budapest, Hungary
- 146 Yale University, New Haven, Connecticut, United States
- 147 Yonsei University, Seoul, Republic of Korea