



# Measurement of the $B_c^-$ meson production fraction and asymmetry in 7 and 13 TeV $pp$ collisions

LHCb collaboration<sup>†</sup>

## Abstract

The production fraction of the  $B_c^-$  meson with respect to the sum of  $B^-$  and  $\bar{B}^0$  mesons is measured in both 7 and 13 TeV center-of-mass energy  $pp$  collisions produced by the Large Hadron Collider (LHC), using the LHCb detector. The rate, approximately 3.7 per mille, does not change with energy, but shows a transverse momentum dependence. The  $B_c^- - B_c^+$  production asymmetry is also measured, and is consistent with zero within the determined statistical and systematic uncertainties of a few percent.

Published in Phys. Rev. D100 (2019) 112006

© 2019 CERN for the benefit of the LHCb collaboration. CC-BY-4.0 licence.

---

<sup>†</sup>Authors are listed at the end of this paper.



# 1 Introduction

The  $B_c^-$  meson is a bound state containing a  $b$  quark with a  $\bar{c}$  quark.<sup>1</sup> It has the largest mass of any two differently flavored quarks in a mesonic ground state. Studies of its production or determination of individual decay widths require measurements of its branching fractions to exclusive final states. Since the branching fractions of some decay modes of  $B^-$  and  $\bar{B}^0$  mesons are accurately known, we determine the ratio of  $B_c^-$  meson production relative to the sum of  $B^-$  and  $\bar{B}^0$  mesons. Here we use techniques similar to those employed for the measurement of  $\bar{B}_s^0$  meson and  $\Lambda_b^0$  baryon fractions [1]. In that paper use is made of the fact that the semileptonic widths of all  $b$ -flavored hadrons with light and strange quarks are equal. However, both the  $b$  and  $c$  quarks can decay, rendering that concept inapplicable. Instead we rely on theoretical predictions of the semileptonic decay branching fraction  $\mathcal{B}(B_c^- \rightarrow J/\psi \mu^- \bar{\nu})$ . Currently, only the relative production cross-section times the branching fraction of either the  $B_c^- \rightarrow J/\psi \mu^- \bar{\nu}$  or  $B_c^- \rightarrow J/\psi \pi^-$  modes have been measured by the CDF [2, 3], LHCb [4, 5] and CMS [6, 7] experiments.

The  $B_c^-$  meson production fraction ( $f_c$ ) relative to the sum of  $\bar{B}^0$  ( $f_d$ ) and  $B^-$  ( $f_u$ ) mesons is defined as

$$R_c = \frac{f_c}{f_u + f_d} \equiv \frac{n_{\text{cor}}(B_c^- \rightarrow J/\psi \mu^- \bar{\nu})}{n_{\text{cor}}(B \rightarrow D^0 X \mu^- \bar{\nu}) + n_{\text{cor}}(B \rightarrow D^+ X \mu^- \bar{\nu})} \cdot \frac{\langle \mathcal{B}_{\text{sl}} \rangle}{\mathcal{B}(B_c^- \rightarrow J/\psi \mu^- \bar{\nu})}, \quad (1)$$

where  $n_{\text{cor}}$  refers to the efficiency and branching fraction corrected number of signal events. The modes containing  $D^0$  and  $D^+$  mesons are also corrected for cross-feeds with  $\bar{B}_s^0$  and  $\Lambda_b^0$  decays. The determination of the corrected yields of the  $B \rightarrow DX \mu^- \bar{\nu}$  decays follows our previous measurement strategy in Ref. [1] where the equations relating the fractions to the corrected yields, including cross-feed contributions, are given. We also correct for the 0.4% effect of doubly-Cabibbo-suppressed decays and  $D^0$  mixing. The relevant hadron branching fractions are listed in Table 1. The average semileptonic branching fractions of  $\bar{B}^0$  and  $B^-$ ,  $\langle \mathcal{B}_{\text{sl}} \rangle = (10.70 \pm 0.19)\%$  is found by averaging measurements from the CLEO [8], BaBar [9] and Belle [10] experiments, detailed in Ref. [11]. Since only  $b \rightarrow c \mu^- \bar{\nu}_\mu$  modes are detected in this analysis, a correction for the small  $b \rightarrow u \mu^- \bar{\nu}_\mu$  rate of 1% is applied to the denominator of Eq. 1.

Table 1: Charm and charmonium branching fractions for the decay modes used in this analysis.

Particle and decay	$\mathcal{B}(\%)$	Source
$D^0 \rightarrow K^- \pi^+$	$3.93 \pm 0.03$	PDG average [12]
$D^+ \rightarrow K^- \pi^+ \pi^+$	$9.22 \pm 0.17$	CLEO III [13]
$J/\psi \rightarrow \mu^+ \mu^-$	$5.96 \pm 0.03$	PDG average [12]

The dominant production mechanism for  $B_c^-$  mesons is gluon-gluon fusion,  $gg \rightarrow B_c^- + \bar{b} + c$ . Non-relativistic quantum chromodynamics is used along with fragmentation functions to predict cross-sections as functions of transverse momentum ( $p_T$ ) and pseudorapidity ( $\eta$ ). The literature is nicely summarized in Ref. [14]. We define  $H_b$  to refer to  $B_c$ ,  $\bar{B}^0$ , and  $B^-$  mesons, while  $H_c$  refers to  $D^0$  and  $D^+$  mesons.

<sup>1</sup>The mention of a particular state implies the use of the charge-conjugate state as well, except when discussing production asymmetries.

In this analysis  $\eta$  is determined by measuring the angle of the  $B$  meson with respect to the beam direction by using the positions of the primary  $pp$  interaction vertex (PV) and the  $B$  meson decay point into either  $J/\psi \mu^-$ ,  $D^0 \mu^-$ , or  $D^+ \mu^-$ . The transverse momentum initially refers to the vector sum of the charmed-hadron and  $\mu^-$  momentum transverse to the proton beams. However, the results are re-interpreted in terms of the  $H_b$  meson  $p_T(H_b)$  by simulating and correcting the effects of the missing momenta.

The production asymmetry between  $B_c^-$  and  $B_c^+$  mesons, which should be small, is defined as

$$a_{\text{prod}} \equiv \frac{\sigma(B_c^-) - \sigma(B_c^+)}{\sigma(B_c^-) + \sigma(B_c^+)} = a_{\text{raw}} - a_{\text{det}}, \quad (2)$$

where  $a_{\text{raw}}$  and  $a_{\text{det}}$  are the asymmetries in the signal yields and the efficiencies of  $B_c^-$  and  $B_c^+$  detection, respectively. The  $CP$  asymmetry in the  $B_c^- \rightarrow J/\psi \mu^- \bar{\nu}$  decay is assumed to be zero in this analysis.

The branching fraction predictions from various models of semileptonic  $B_c^-$  decays are listed in Table 2. For  $\mathcal{B}(B_c^- \rightarrow J/\psi \mu^- \bar{\nu})$  they range from 1.4 to 7.5%, which is quite a large interval. Branching fractions for other modes are also listed where available. We use the  $Z$  expansion fit results from Ref. [32], and the method II results for Ref. [33].

Table 2: Branching fractions predictions (%). The  $B_c^-$  lifetime is taken as 0.507 ps [12]. The value for the semileptonic decays of the  $B_c^-$  meson,  $\mathcal{B}_{\text{sl}}^c$ , is derived by summing the  $J/\psi \mu^- \bar{\nu}$  and  $\eta_c \mu^- \bar{\nu}$  individual predictions with the average predictions of 0.1% for  $\psi(2S) \mu^- \bar{\nu}$ , the sum of  $\chi_{c0,1,2} \mu^- \bar{\nu}$  as 0.6%, and 0.3% for  $h_c \mu^- \bar{\nu}$ . In the one case where  $\eta_c \mu^- \bar{\nu}$  was not predicted averages from other measurements are used.

Ref. \ Mode	$J/\psi \mu^- \bar{\nu}$	$\eta_c \mu^- \bar{\nu}$	$\psi(2S) \mu^- \bar{\nu}$	$\chi_{c0,1,2} \mu^- \bar{\nu}$	$h_c \mu^- \bar{\nu}$	$\mathcal{B}_{\text{sl}}^c$
[15]	6.4	5.0	1.3			13.6
[16]				0.5		
[17]	1.4	0.5				2.9
[18]	7.5	2.4				10.9
[19]	1.9	0.6	0.1			3.5
[20]	2.3	0.9		0.8		4.2
[21]	2.7	1.8				5.5
[22]	1.6	0.8				3.4
[23]	1.7	0.5		0.6		3.3
[24]	1.7	0.2				2.9
[25]	1.9	0.8	0.1			3.7
[26]	2.3	0.9				4.2
[27]	2.2	0.8	0.1			4.0
[28]	2.6		0.1	1.1		4.2
[29]	2.5	1.1				4.6
[30]	1.3	0.8	0.2			3.1
[31]	1.4	0.7				3.1
[32]	1.5	0.7		0.5	0.3	3.2
[33]	1.9	0.6	0.1	0.3	0.3	3.5
[34]	2.2	0.8				4.0

Some restrictions on models are possible by comparing to lighter  $B$  meson decays. Since the inclusive semileptonic branching fraction for these decays,  $\mathcal{B}_{\text{sl}}$ , is about 10.5% and the  $B_c^-$  lifetime,  $\tau_{B_c}$ , is 1/3 that of the  $\bar{B}^0$ ,<sup>2</sup> we disregard models that predict 10% or larger values for  $\mathcal{B}_{\text{sl}}^c$  of the  $B_c^-$ . This excludes from consideration the models of Refs. [15] and [18]. The average model prediction is then  $\mathcal{B}(B_c^- \rightarrow J/\psi \mu^- \bar{\nu}) = 1.95\%$ . The standard deviation is 0.46%, which we use to estimate the systematic uncertainty on the model variation. Results of our measurement without using this branching fraction are also quoted.

## 2 Detector, trigger and simulation

The LHCb detector [35,36] is a single-arm forward spectrometer covering the pseudorapidity range  $2 < \eta < 5$ , designed for the study of particles containing  $b$  or  $c$  quarks. The detector elements that are particularly relevant to this analysis are: a silicon-strip vertex detector surrounding the  $pp$  interaction region that allows  $c$  and  $b$  hadrons to be identified from their characteristically long flight distance; a tracking system that provides a measurement of the momentum,  $p$ , of charged particles; two ring-imaging Cherenkov detectors that are able to discriminate between different species of charged hadrons; and a downstream system of iron interspersed with chambers is used to identify muons.

The magnetic field deflects positively and negatively charged particles in opposite directions and this can lead to detection asymmetries. Periodically reversing the magnetic field polarity throughout the data taking almost cancels the effect. The configuration with the magnetic field pointing upwards (downwards) bends positively (negatively) charged particles in the horizontal plane towards the centre of the LHC ring. This analysis uses data collected in 2011 (7 TeV) and 2016 (13 TeV) where appropriate triggers are available. The data taking was split between magnetic field up and down configurations. In the 2011 data  $0.6 \text{ fb}^{-1}$  ( $0.4 \text{ fb}^{-1}$ ) were collected with the field pointing up (down), while in 2016 the split was  $0.9 \text{ fb}^{-1}$  with field up and  $0.8 \text{ fb}^{-1}$  with field down.

The trigger [37] consists of a hardware stage, based on information from the calorimeter and muon systems, followed by a software stage, in which all charged particles with  $p_{\text{T}} > 500$  (300) MeV are reconstructed for 2011 (2016) data.

Separate hardware triggers are used for the  $J/\psi \mu^-$  and  $H_c$  samples. For the former we require a  $\mu^+ \mu^-$  pair. For the latter, we require a single muon with large  $p_{\text{T}}$  for the 7 TeV data as used in Ref. [38]. For the 13 TeV data, the single muon trigger was not available, therefore at the hardware trigger stage, events are required to have a hadron, photon or electron transverse energy greater than approximately 3.5 GeV in the calorimeters. The software trigger requires a two-, three- or four-track secondary vertex with a significant displacement from any primary  $pp$  interaction vertex as described in Ref. [1]. At least one charged particle must have  $p_{\text{T}} > 1.6$  GeV and be inconsistent with originating from a PV. A multivariate algorithm [39] is used for the identification of secondary vertices consistent with the decay of a  $b$  hadron.

Simulation is required to model the effects of the detector acceptance and the imposed selection requirements. In the simulation,  $pp$  collisions are generated using PYTHIA [40] with a specific LHCb configuration [41]. Decays of unstable particles are described

<sup>2</sup>This is evident since  $\mathcal{B}_{\text{sl}}^c = \Gamma_{\text{sl}} \cdot \tau_{B_c}$ , and  $\Gamma_{\text{sl}}$  is approximately the same for all  $b$ -hadron species. We use natural units where  $c = \hbar = 1$ .

by EVTGEN [42], in which final-state radiation is generated using PHOTOS [43]. The interaction of the generated particles with the detector, and its response, are implemented using the GEANT4 toolkit [44] as described in Ref. [45].

### 3 Event selection, signal efficiencies and yields

#### 3.1 Selection of $B_c^- \rightarrow J/\psi \mu^- \bar{\nu}$ candidates

The analysis is done separately for the light  $B$  meson modes and the  $B_c^- \rightarrow J/\psi \mu^- \bar{\nu}$  decay. In each case the triggered events are subject to further filtering requirements. In addition, the  $J/\psi \mu^-$  sample is subjected to a boosted decision tree (BDT), a multivariate classification method, using the TMVA toolkit [46]. This is not necessary for the  $D^0$  or  $D^+$  modes because they have large signals and are relatively free from backgrounds [1].

For the  $J/\psi \mu^- \bar{\nu}$  final state the initial selection requires that muons that satisfy the  $J/\psi$  candidate trigger each have minimum  $p_T > 550$  MeV, have large impact parameters with the PV, form a good quality vertex, have a reasonable flight distance significance from the PV, and have a summed  $p_T > 2$  GeV. The ‘‘companion’’ muon that is not part of the  $J/\psi$  decay must be well identified and form a good quality vertex with the  $J/\psi$  candidate, which must be downstream of the PV.

To suppress muon tracks that are reconstructed more than once, we require a small minimum opening angle between the muons from the  $J/\psi$  decay and the companion muon momentum measured in the plane transverse to the beam line. Specifically, this opening angle must be greater than  $0.8^\circ$ . The invariant mass of the companion muon and the oppositely charged muon from  $J/\psi$  must differ from the known value of the  $J/\psi$  mass by more than 50 MeV [12], while the invariant mass with the same charged muon is required to be larger than 400 MeV.

Since we are dealing with an exclusive final state, we define

$$m_{\text{cor}} \equiv \sqrt{m(J/\psi \mu^-)^2 + p_\perp^2} + p_\perp, \quad (3)$$

where  $p_\perp$  is the magnitude of the combination’s momentum component transverse to the  $b$ -hadron flight direction. Figure 1 shows the distributions of  $m_{\text{cor}}$  versus the invariant  $J/\psi \mu^-$  mass,  $m(J/\psi \mu^-)$ , for both data and simulation. To remove background, a requirement of  $m(J/\psi \mu^-) > 4.5$  GeV is applied, as indicated by the (red) dashed line.

Since we also measure the production asymmetry between  $B_c^+$  and  $B_c^-$  mesons, we restrict the angular acceptance of the companion muon to make it more uniform by removing muons close to the edge of the detector, in the bending direction ( $x$ -direction), where large acceptance-induced asymmetries can occur. Thus, we require that the  $x$ -component of the momentum satisfies

$$|p_x| \leq 0.294(p_z - 2 \text{ GeV}), \quad (4)$$

where  $p_z$  is the muon momentum along the direction of the proton beam downstream of the PV, as is done in Refs. [47, 48].

After these initial restrictions, we turn to the multivariate selection, forming the classifier denoted BDT in the following. The discriminating variables used are: (a) the  $\chi^2$  of the vertex fit of the  $J/\psi$  with the  $\mu^-$ ; (b) the  $\ln \chi_{\text{IP}}^2$ , where  $\chi_{\text{IP}}^2$  is defined as the  $\chi^2$




Figure 1: Distributions of corrected mass  $m_{\text{cor}}$  and  $m(J/\psi \mu^-)$  for (top) 7 TeV and (bottom) 13 TeV data, where (a) and (c) are data and (b) and (d) simulated signal. The (red) dashed line indicates the  $m(J/\psi \mu^-) > 4.5$  GeV requirement.

of the impact parameter with respect to the PV, of the  $J/\psi$ ,  $\mu^-$  and their combination; (c) the  $p_T$  of the  $J/\psi$  and the  $\mu^-$ ; and (d) the cosine of the angle between the  $\mu^-$  and the  $J/\psi$  meson in the plane perpendicular to the beam direction. The training sample for signal is simulated  $B_c^- \rightarrow J/\psi \mu^- \bar{\nu}$  events, and for background is inclusive  $b \rightarrow J/\psi X$  simulated events.

We then optimize the BDT output threshold by maximizing  $S/\sqrt{S+B}$ , where  $S$  and  $B$  are the number of the signal and background yields in the signal region defined as  $m_{\text{cor}} \in (4.8, 10.8)$  GeV. The sum,  $S+B$ , is the total number of events within these limits, and  $S$  is taken from a fit to the  $m_{\text{cor}}$  distribution. The optimal BDT output threshold results in a BDT signal efficiency of 89% with a background rejection of 63%, as determined by observing the resulting samples of input signal simulation events and background candidates.

The  $m_{\text{cor}}$  distribution is shown in Fig. 2. It consists not only of signal  $B_c^-$  events, but also of  $B_c^- \rightarrow J/\psi \tau^- \bar{\nu}$  decays, where  $\tau^- \rightarrow \mu^- \nu \bar{\nu}$ , and other  $c\bar{c}$  final states, most importantly  $B_c^- \rightarrow \psi(2S)\mu^- \bar{\nu}$  and  $B_c^- \rightarrow \chi_{c1}\mu^- \bar{\nu}$ . We find shapes for these final states using simulation. The signal shape is a sum of a double Crystal Ball and a bifurcated Gaussian functions. The sum of the combinatorial and misidentification backgrounds are represented by a Gaussian kernel shape [49]. For the other background modes, we use histograms directly. These shapes are fitted to the  $m_{\text{cor}}$  distributions in Fig. 2 in order to determine the  $B_c^- \rightarrow J/\psi \mu^- \bar{\nu}$  yields. The ratio of the  $J/\psi \tau^- \bar{\nu}$  yield to the  $J/\psi \mu^- \bar{\nu}$  yield is fixed, after accounting for the relative detection efficiencies, from the LHCb measurement of  $0.71 \pm 0.17 \pm 0.18$ , where the first uncertainty is statistical and the second




Figure 2: Fitted  $m_{\text{cor}}$  distributions in (a) 7 TeV and (b) 13 TeV samples. The signal and the backgrounds are shown as the dark (orange) and lighter (gray) areas. The dashed (cyan) curves show the  $B_c^- \rightarrow J/\psi \tau^- \nu_\tau$  components, while the dotted (blue) curves show the  $B_c^- \rightarrow \chi_{c0,1,2} \mu^- \bar{\nu}$  components. The  $B_c^- \rightarrow \psi(2S) \mu^- \bar{\nu}$  contribution is also in the fit but is too small to be seen. The total fit is shown by the solid (red) curve.

systematic [50]; this convention is used throughout this paper. The other components of the fit are allowed to vary. We find  $4010 \pm 200$  and  $15170 \pm 710$  signal  $B_c^- \rightarrow J/\psi \mu^- \bar{\nu}$  events at 7 and 13 TeV, respectively, while the backgrounds sum to 950 and 5170 events at the same energies. These signal yields need to be corrected for the small background from candidates with a correctly reconstructed  $J/\psi$  meson that is paired with a hadron mis-identified as a muon.

### 3.2 Efficiency for $B_c^- \rightarrow J/\psi \mu^- \bar{\nu}$

Efficiencies are determined using both data [51, 52] and simulation of  $B_c^- \rightarrow J/\psi \mu^- \bar{\nu}$ , with the generated events weighted to match the  $p_T(H_b)$ , and  $\eta$  distributions observed in data. In addition, we weight accordingly the  $\chi_{\text{IP}}^2$  distribution of the muon associated with the  $J/\psi$ . Weighting the simulation is important since the total efficiencies are functions of these variables. Efficiencies using data include trigger, and muon identification. Efficiencies using simulation include detector acceptance, reconstruction and event selection, and removal of beam crossings with an excess number of hits in the detector. Total efficiencies as a function of  $p_T(B_c^-)$  for different  $\eta$  intervals are shown in Fig. 3.

### 3.3 $H_c X \mu^- \bar{\nu}$ selection criteria

Selection criteria for  $H_b \rightarrow H_c X \mu^- \bar{\nu}$  final states differ from those containing a  $J/\psi$ . The transverse momentum of each hadron must be greater than 0.3 GeV, and that of the muon larger than 1.3 GeV. We require  $\chi_{\text{IP}}^2 > 9$  with respect to any PV, ensuring that tracks do not originate from primary  $pp$  interactions. All final state particles are required to be positively identified using information from the RICH detectors. Particles from  $H_c$  decay candidates must have a good fit to a common vertex with  $\chi^2/\text{ndof} < 9$ , where ndof is the number of degrees of freedom. They must also be well separated from the nearest PV, with the flight distance divided by its uncertainty greater than 5.

Candidate  $b$  hadrons are formed by combining  $H_c$  and muon candidates originating from a common vertex with  $\chi^2/\text{ndof} < 9$  and an  $H_c \mu^-$  invariant mass in the range



Table 3: Yields of  $B \rightarrow DX\mu^- \bar{\nu}$  decays.

Mode	7 TeV Yields		13 TeV yields	
	Signal	fake muons	Signal	fake muons
$D^0 X\mu^- \bar{\nu}$	$789\,800 \pm 940$	$5500 \pm 160$	$12\,285\,000 \pm 3700$	$115\,155 \pm 580$
$D^+ X\mu^- \bar{\nu}$	$263\,190 \pm 570$	$990 \pm 70$	$3\,686\,240 \pm 2130$	$21\,370 \pm 240$

3.0–5.0 GeV.

Background from prompt  $H_c$  production at the PV needs to be considered. We use the natural logarithm of the  $H_c$  impact parameter, IP, with respect to the PV in units of mm. Requiring  $\ln(\text{IP}/\text{mm}) > -3$  is found to reduce the prompt component to be below 0.1%, while preserving 97% of all signals. This restriction allows us to perform fits only to the  $H_c$  candidate mass spectra to find the  $b$ -hadron decay yields.

The  $H_c$  candidate mass distributions integrated over  $p_T(H_b)$  and  $\eta$  are shown in Fig. 4 and consist of a prominent peak resulting from signal, and a small contribution due to combinatorial background from random combinations of particles that pass the selection. They are fit with a signal component comprised of two Gaussian functions, and a combinatorial background component modeled as a linear function. The fitted yields are listed in Table 3. These numbers must be corrected for hadrons that are mis-identified as muons, and for semileptonic decays of  $\bar{B}_s^0$  and  $\Lambda_b^0$  hadrons that produce  $D^0$  and  $D^+$  mesons.

In Table 3 the column labeled “fake muons” shows the yields of wrong-sign  $D^0 X\mu^+$  and  $D^+ X\mu^+$  combinations that pass the selections. These yields provide good estimates of the fake muon contributions in the signal samples, which are very small. Following the procedure in Ref. [1], we find the cross-feed corrections of  $\bar{B}_s^0 \rightarrow (D^0 + D^+)X\mu^- \bar{\nu}$  and  $\Lambda_b^0 \rightarrow (D^0 + D^+)X\mu^- \bar{\nu}$  to be twice the measured yields for  $\bar{B}_s^0 \rightarrow D^0 K^+ X\mu^- \bar{\nu}$ , which are  $8500 \pm 340$  (7 TeV) and  $69\,390 \pm 1130$  (13 TeV), and for  $\Lambda_b^0 \rightarrow D^0 p X\mu^- \bar{\nu}$ , which are  $2330 \pm 140$  (7 TeV) and  $33\,050 \pm 460$  (13 TeV). Relative efficiencies for detecting final states with a single extra hadron are taken into account when subtracting these yields.




Figure 3: The total efficiency for  $B_c^- \rightarrow J/\psi \mu^- \bar{\nu}$ , as a function of  $p_T(B_c^-)$  in different intervals of  $\eta$  in (a) 7 TeV and (b) 13 TeV samples.

### 3.4 Efficiencies for $B \rightarrow D^0 X \mu^- \bar{\nu}$ and $B \rightarrow D^+ X \mu^- \bar{\nu}$

Similar methods based on data, as implemented for the  $B_c^-$  decay, are used to evaluate the efficiencies for trigger and particle identification. Simulation is also used to determine the efficiencies of event selection and reconstruction of these modes. The total efficiencies for  $B$  meson decays into  $D^0 X \mu^- \bar{\nu}$  and  $D^+ X \mu^- \bar{\nu}$  are shown in Fig. 5.

## 4 Results

### 4.1 Corrections to the $p_T(H_b)$ distributions due to the missing neutrino

Since the production kinematics of  $B$  and  $B_c^-$  mesons can differ as functions of  $p_T(H_b)$  and  $\eta$ , we need to measure  $f_c/(f_u + f_d)$  as functions of these variables. The measurement of  $\eta$  is straightforward, however, we do not measure directly the  $p_T(H_b)$  of the  $b$ -flavored hadron because of the missing neutrino, and in the case of the  $B$  meson possible missing extra particles. Following a procedure similar to the one used in Ref. [1], we determine a correction factor,  $k$ , that is the ratio of the average reconstructed to true  $p_T(H_b)$  as a function of the invariant mass of the charmed hadron plus muon. The ratio distribution as a function of hadron-muon invariant mass are shown in Fig. 6. The average correction, the  $k$ -factor, is shown on the figure. For the  $B$  meson it varies from 0.75 to unity over




Figure 4: Invariant-mass distributions of (a)  $K^- \pi^+$  and (b)  $K^- \pi^+ \pi^+$  for 7 TeV, and (c) and (d) for 13 TeV collisions. The data are shown by solid points. The (red) dashed lines represent the signal components. The combinatorial backgrounds are shown as the dotted (magenta) line, and the solid (blue) line shows the total fit.




Figure 5: Total efficiencies for the (a)  $D^0 X \mu^- \bar{\nu}$  and (b)  $D^+ X \mu^- \bar{\nu}$  signals in 7 TeV and (c) and (d) in 13 TeV samples as functions of  $p_T$  in  $\eta$  intervals.

Table 4: Results of the fits to Eq. 5

Energy	$p_1$	$p_2 \cdot 10^{-2} \text{ (GeV}^{-1}\text{)}$
7 TeV	$3.82 \pm 0.09 \pm 0.05$	$-6.2 \pm 1.7 \pm 1.1$
13 TeV	$4.13 \pm 0.05 \pm 0.04$	$-9.7 \pm 0.8 \pm 1.0$

the interval from 3 GeV to the  $B$  mass, and for the  $B_c^-$  meson it varies from 0.85 to unity over the interval from 4 GeV to the  $B_c^-$  mass.

## 4.2 $B_c^-$ fraction results

The ratio of production fractions,  $f_c/(f_u + f_d)$ , are shown as functions of  $p_T(H_b)$  and  $\eta$  in Fig. 7. There is little dependence on  $\eta$ , but the decrease as a function of  $p_T(H_b)$  is noticeable.

To describe the  $p_T(H_b)$  dependence we use an equation of the form

$$\frac{f_c}{f_u + f_d}(p_T) = A [p_1 + p_2 (p_T(H_b) - \langle p_T \rangle)], \quad (5)$$

where  $A$  represents the overall normalization and contains the total global systematic uncertainty, thus,  $A = 1 \pm 0.24$ ,<sup>3</sup>  $\langle p_T \rangle$  is taken as 7.2 GeV, close to the average  $p_T$  of the  $B_c^-$ . The slopes,  $p_2$ , are similar in size to those measured for the  $B_s$  meson fraction ratio as a function of  $p_T$  [1, 53]. Results of fits to the data using Eq. 5 are listed in Table 4.

<sup>3</sup>See Section 5 for the discussion of the systematic uncertainties.

The average fractions in the interval  $4 < p_{\text{T}}(H_b) < 25$  GeV are found by integrating over  $p_{\text{T}}(H_b)$ . To allow for facile changes to our results due to improved theoretical predictions, we provide the results for

$$\frac{f_c}{f_u + f_d} \cdot \mathcal{B}(B_c^- \rightarrow J/\psi \mu^- \bar{\nu}) = (7.07 \pm 0.15 \pm 0.24) \cdot 10^{-5} \text{ for 7 TeV,}$$

$$\frac{f_c}{f_u + f_d} \cdot \mathcal{B}(B_c^- \rightarrow J/\psi \mu^- \bar{\nu}) = (7.36 \pm 0.08 \pm 0.30) \cdot 10^{-5} \text{ for 13 TeV.}$$

Next we give the result on the fractions ratio

$$\frac{f_c}{f_u + f_d} = (3.63 \pm 0.08 \pm 0.12 \pm 0.86) \cdot 10^{-3} \text{ for 7 TeV,}$$

$$\frac{f_c}{f_u + f_d} = (3.78 \pm 0.04 \pm 0.15 \pm 0.89) \cdot 10^{-3} \text{ for 13 TeV,}$$

where the third uncertainty is due to the theoretical prediction of  $\mathcal{B}(B_c^- \rightarrow J/\psi \mu^- \bar{\nu})$ . To find  $f_c/f_u$  just double these numbers.

We also measure the ratio of the  $B_c^-$  production fraction at 7 TeV to that at 13 TeV. Figure 8 shows the ratio as functions of  $p_{\text{T}}$  and  $\eta$ . Here most of the systematic uncertainties cancel. The integrated value of the ratio of 13 TeV and 7 TeV is measured as  $1.02 \pm 0.02 \pm$




Figure 6: The  $k$ -factor corrections as a function of invariant mass of (a)  $m(J/\psi \mu^-)$ , (b)  $m(D^0 \mu^-)$ , and (c)  $m(D^+ \mu^-)$  for the 13 TeV simulation samples. (The 7 TeV results are almost identical.) The points (magenta) are the average  $k$ -factor corrections, and the (green) dashed line shows a second-order polynomial fit to the average data.

0.04, consistent with no increase in the  $B_c^-$  fraction ratio as a function of center-of-mass energy.

The  $B_c^-$  fraction with respect to inclusive  $b$ -hadron production can be derived from the information in previous LHCb  $b$ -hadron fraction papers Ref. [1, 38, 53]. There the measured values of the ratios of  $b$ -hadron fractions over the same  $p_T$  range in terms of the  $b$ -hadron  $p_T$  are for  $\bar{B}_s^0$  mesons ( $f_s$ ) and  $\Lambda_b^0$  baryons

$$\frac{f_s}{f_u + f_d} = \begin{cases} 0.124 \pm 0.010 & (7 \text{ TeV}) [53] \\ 0.122 \pm 0.006 & (13 \text{ TeV}) [1], \end{cases} \quad (6)$$

$$\frac{f_{\Lambda_b^0}}{f_u + f_d} = \begin{cases} 0.223 \pm 0.036 & (7 \text{ TeV}) [38] \\ 0.259 \pm 0.018 & (13 \text{ TeV}) [1], \end{cases} \quad (7)$$

where the uncertainties contain both statistical and systematic components added in quadrature. For the measurement of the  $f_{\Lambda_b^0}$  fraction at 7 TeV, the dominant systematic uncertainty is from the lack of the knowledge of  $\mathcal{B}(\Lambda_c^+ \rightarrow pK^-\pi^+)$  at that time [38]; here the value and uncertainty have been recalculated according to the latest value of  $\mathcal{B}(\Lambda_c^+ \rightarrow pK^-\pi^+)$  from the PDG [12].

Taking the sum of all the  $b$ -hadron fractions to be unity, and ignoring  $f_c$  here because it is so small,

$$f_u + f_d + f_s + f_{\Lambda_b^0}(1 + \delta) = 1, \quad (8)$$




Figure 7: Ratio of production fractions after the  $k$ -factor correction as a function of (a)  $p_T(H_b)$  and (c)  $\eta$  in 7 TeV data and (b) and (d) in 13 TeV data. The smaller error bars show the statistical uncertainties and the larger ones include the statistical and systematic uncertainties. The horizontal (green) dashed-lines show the average values.

where  $\delta = 0.25 \pm 0.10$  is a correction factor derived in Ref. [11] that accounts for heavier  $b$ -baryons, mainly the  $\Xi_b$ . Solving for  $f_u + f_d$  yields

$$f_u + f_d = \left( 1 + \frac{f_s}{f_u + f_d} + \frac{f_{\Lambda_b^0}}{f_u + f_d} (1 + \delta) \right)^{-1},$$

$$= \begin{cases} 0.713 \pm 0.026 & (7 \text{ TeV}) \\ 0.692 \pm 0.015 & (13 \text{ TeV}). \end{cases} \quad (9)$$

We find that

$$f_c \cdot \mathcal{B}(B_c^- \rightarrow J/\psi \mu^- \bar{\nu}) = \begin{cases} (5.04 \pm 0.11 \pm 0.17 \pm 0.18) \cdot 10^{-5} & (7 \text{ TeV}) \\ (5.09 \pm 0.06 \pm 0.21 \pm 0.11) \cdot 10^{-5} & (13 \text{ TeV}) \end{cases},$$

where the first uncertainty is statistical, the second is systematic, and the third is from the fractions of the  $B_s^0$  and  $\Lambda_b^0$  given in Eq. 9. We also provide the result for  $f_c$ ,

$$f_c = \begin{cases} (2.58 \pm 0.05 \pm 0.62 \pm 0.09) \cdot 10^{-3} & (7 \text{ TeV}) \\ (2.61 \pm 0.03 \pm 0.62 \pm 0.06) \cdot 10^{-3} & (13 \text{ TeV}) \end{cases},$$

where the first uncertainty is statistical, the second is systematic including that from  $\mathcal{B}(B_c^- \rightarrow J/\psi \mu^- \bar{\nu})$  and the third is from the fractions of the  $B_s^0$  and  $\Lambda_b^0$  given in Eq. 9.

### 4.3 The $B_c^- - B_c^+$ production asymmetry

The production asymmetries are measured in two different magnetic field configurations and then averaged. No significant asymmetry is observed in any intervals of  $p_T(H_b)$  or  $\eta$ . The results are summarized in Table 5.

Averaging the  $B_c^- - B_c^+$  production asymmetries over  $p_T(H_b)$  and  $\eta$ , we find  $(-2.5 \pm 2.1 \pm 0.5)\%$ , and  $(-0.5 \pm 1.1 \pm 0.4)\%$  at center-of-mass energies of 7 and 13 TeV, respectively.




Figure 8: Ratio of the  $B_c^-$  production fractions at 13 TeV to 7 TeV as a function of (a)  $p_T(H_b)$  and (b)  $\eta$ . The smaller error bars show the statistical uncertainties and the larger ones include the statistical and systematic uncertainties added in quadrature.

Table 5: The  $B_c^- - B_c^+$  production asymmetry ( $\times 10^{-2}$ ) as a function of  $p_T(H_b)$  and  $\eta$  at 7 TeV and 13 TeV.

7 TeV production asymmetry		
$p_T(\text{GeV}) \setminus \eta$	2.5 – 3.5	3.5 – 4.5
4 – 6	$7.91 \pm 7.00 \pm 1.03$	$-6.44 \pm 6.44 \pm 2.10$
6 – 8	$-4.34 \pm 5.43 \pm 1.62$	$-6.66 \pm 6.65 \pm 2.03$
8 – 10	$-1.13 \pm 6.31 \pm 1.56$	$-9.63 \pm 7.23 \pm 0.81$
10 – 25	$0.24 \pm 4.13 \pm 0.98$	$-4.87 \pm 8.63 \pm 1.44$
13 TeV production asymmetry		
$p_T(\text{GeV}) \setminus \eta$	2.5 – 3.5	3.5 – 4.5
4 – 6	$3.13 \pm 3.33 \pm 1.16$	$1.76 \pm 3.23 \pm 0.91$
6 – 8	$-0.34 \pm 2.79 \pm 1.26$	$-5.03 \pm 3.61 \pm 1.06$
8 – 10	$2.03 \pm 2.73 \pm 0.94$	$-2.48 \pm 4.29 \pm 1.78$
10 – 25	$1.50 \pm 2.05 \pm 0.73$	$-1.47 \pm 4.20 \pm 2.18$

## 5 Systematic uncertainties

Systematic uncertainties are separated into two categories: “global”, which apply across the phase space, and “local”, which are calculated in each two-dimensional  $p_T(H_b) - \eta$  bin. These uncertainties are listed in Table 6.

First let us consider the  $B_c^- \rightarrow J/\psi \mu^- \bar{\nu}$  decay. The uncertainty due to the signal shape used to fit the  $m_{\text{cor}}$  distribution is determined by changing the baseline signal shape, the sum of a double sided Crystal Ball function and a bifurcated Gaussian, to a kernel estimation. To find the shape of the combinatorial and misidentification backgrounds we use simulated inclusive samples of  $b \rightarrow J/\psi X$  events not including  $B_c^-$  decays. A total of 500 samples are generated and different fits to the samples are performed to determine the possible uncertainty. This procedure is also used for the  $a_{\text{prod}}$  measurement. We call contributions to the  $J/\psi \mu^-$  mass spectrum “feed-down” contributions, occurring from other  $B_c^-$  decay channels including  $J/\psi \tau \bar{\nu}$ ,  $\psi(2S) \mu^- \bar{\nu}$ , and  $\chi_{c\ell} \mu^- \bar{\nu}$ . The systematic uncertainty results from the uncertainties in their branching fractions. Different decay models for  $B_c^- \rightarrow J/\psi \mu^- \bar{\nu}$  decays can change the  $m_{\text{cor}}$  shape. We use the model of Ebert *et al.* [16] for our baseline prediction. Then we also use the model by Kiselev [27] to find the efficiencies and take half the difference as the systematic uncertainty. We also estimate the uncertainty due to the sensitivity to various selection requirements and simulation statistics. The muon identification efficiencies are determined from data using inclusive samples of  $J/\psi$  decay where one of the muon candidates is not identified. The trigger efficiency is determined by using three independent samples of events, those that trigger on a  $J/\psi$ , those that triggered on something else in the event, and those that trigger on both the  $J/\psi$  and something else. These samples are then used to compute the trigger efficiencies in two-dimensional  $p_T(H_b)$  and  $\eta$  bins.

Next, we turn to the  $B \rightarrow DX\mu\nu$  modes. The efficiencies and their uncertainties for identifying pions and kaons are determined by using almost background free samples of  $D^{*+} \rightarrow \pi^+ D^0$ ,  $D^0 \rightarrow K^- \pi^+$  decays. The trigger and muon identification efficiencies, and

Table 6: Summary of the relative systematic uncertainties for  $f_c/(f_u + f_d)$ (%) and the absolute production asymmetries  $a_{\text{prod}}$ (%). For local uncertainties, the ranges correspond to the minimum and maximum uncertainties evaluated in the  $p_T(H_b)$  and  $\eta$  ranges.

	$f_c/(f_u + f_d)$		$a_{\text{prod}}$	
	7 TeV	13 TeV	7 TeV	13 TeV
<i>Local uncertainties</i>				
Signal shape	0.12–9.56	0.14–2.80	0.04–1.80	0.01–0.78
Background shape	0.34–6.16	0.02–5.80	0.06–3.05	0.05–2.45
Feed-down channels	0.12–5.00	0.43–2.27	0.01–1.11	0.03–0.65
Decay models	0.00–2.00	0.01–3.84	0.02–0.28	0.02–0.61
Muon ID in $J/\psi \mu^-$	0.06–5.79	0.03–2.92	0.02–0.37	0.01–0.18
Trigger for $J/\psi$	0.00–0.23	0.00–0.34	0.05–2.34	0.07–4.24
Simulation decay model	0.00–2.00	0.01–3.84	0.02–0.28	0.02–0.61
Hadron ID in $DX\mu^-\bar{\nu}$	0.04–1.81	0.01–2.01	–	–
Muon trigger & ID in $DX\mu^-\bar{\nu}$	0.02–1.34	0.00–0.21	–	–
Simulation sample size	1.5–11.5	2.1–10.7	0.5–1.1	0.5–1.2
$k$ -factor	0.02–0.95	0.05–0.70	0.01–0.10	0.00–0.10
Tracking asymmetry	–	–	0.00–0.28	0.00–0.09
<i>Global uncertainties</i>				
$\mathcal{B}(J/\psi \rightarrow \mu^+ \mu^-)$	0.55	0.55	–	–
$\mathcal{B}(D^+ \rightarrow K^- \pi^+ \pi^+)$ or $\mathcal{B}(D^0 \rightarrow K^- \pi^+)$	1.0	1.0	–	–
$\mathcal{B}(B \rightarrow H_c X \mu^- \bar{\nu})$	1.8	1.8	–	–
Cross-feed contribution	0.2	0.2	–	–
Multiplicity cut	1.2	2.7	–	–
Tracking efficiency	1.8	1.8	–	–
<i>Uncertainty sum</i>	4.3–21.3	5.1–17.4	1.0–3.5	1.0–4.8
$\mathcal{B}(B_c^- \rightarrow J/\psi \mu^- \bar{\nu})$	23.6	23.6	–	–
<i>Overall uncertainty</i>	24.0–31.8	24.1–29.3	1.0–3.5	1.0–4.8

their uncertainties, are obtained in the same manner as for the  $B_c^- \rightarrow J/\psi \mu^- \bar{\nu}$  mode. There are small systematic uncertainties related to efficiency estimates and the assumed  $D^*$  to  $D$  mixtures, as well as simulation statistics. Global systematic uncertainties include the hadron branching fractions listed in Table 1, cross-feed corrections arising from  $\bar{B}_s^0$  and  $A_b^0$  decays into  $DX\mu^-\bar{\nu}$  events, and a global hadron plus photon multiplicity requirement. The latter is evaluated with data.

## 6 Conclusions

In 7 and 13 TeV  $pp$  collisions the product of  $\mathcal{B}(B_c^- \rightarrow J/\psi \mu^- \bar{\nu})$  with the relative fraction of  $B_c^-$  mesons with respect to the sum of  $B^0$  and  $B^+$  mesons in the ranges  $2.5 < \eta < 4.5$



and  $4 < p_T(H_b) < 25$  GeV is found to be

$$\begin{aligned}\frac{f_c}{f_u + f_d} \cdot \mathcal{B}(B_c^- \rightarrow J/\psi \mu^- \bar{\nu}) &= (7.07 \pm 0.15 \pm 0.24) \cdot 10^{-5} \text{ for 7 TeV,} \\ \frac{f_c}{f_u + f_d} \cdot \mathcal{B}(B_c^- \rightarrow J/\psi \mu^- \bar{\nu}) &= (7.36 \pm 0.08 \pm 0.30) \cdot 10^{-5} \text{ for 13 TeV.}\end{aligned}$$

We derive the product of  $f_c \cdot \mathcal{B}(B_c^- \rightarrow J/\psi \mu^- \bar{\nu})$  at the two energies as

$$f_c \cdot \mathcal{B}(B_c^- \rightarrow J/\psi \mu^- \bar{\nu}) = \begin{cases} (5.04 \pm 0.11 \pm 0.17 \pm 0.18) \cdot 10^{-5} & (7 \text{ TeV}) \\ (5.09 \pm 0.06 \pm 0.21 \pm 0.11) \cdot 10^{-5} & (13 \text{ TeV}) \end{cases}$$

Using the average of the theoretical prediction  $\mathcal{B}(B_c^- \rightarrow J/\psi \mu^- \bar{\nu}) = (1.95 \pm 0.46)\%$ , where the uncertainty is given by the standard deviation derived from the distribution of the models, we determine

$$\begin{aligned}\frac{f_c}{f_u + f_d} &= (3.63 \pm 0.08 \pm 0.12 \pm 0.86) \cdot 10^{-3} \text{ for 7 TeV,} \\ \frac{f_c}{f_u + f_d} &= (3.78 \pm 0.04 \pm 0.15 \pm 0.89) \cdot 10^{-3} \text{ for 13 TeV,}\end{aligned}$$

where the first uncertainties are statistical, the second systematic, and the third due to the theoretical prediction of  $\mathcal{B}(B_c^- \rightarrow J/\psi \mu^- \bar{\nu})$ . There is a small dependence on the transverse momentum of the  $B_c^+$  meson, but no dependence on its pseudorapidity is observed. We also report

$$f_c = \begin{cases} (2.58 \pm 0.05 \pm 0.62 \pm 0.09) \cdot 10^{-3} & (7 \text{ TeV}) \\ (2.61 \pm 0.03 \pm 0.62 \pm 0.06) \cdot 10^{-3} & (13 \text{ TeV}) \end{cases},$$

where the first uncertainty is statistical, the second is systematic including that from  $\mathcal{B}(B_c^- \rightarrow J/\psi \mu^- \bar{\nu})$  and the third is from the fractions of the  $B_s^0$  and  $A_b^0$  given in Eq. 9.

The ratio of fractions,  $1.02 \pm 0.02 \pm 0.04$ , for 13 TeV/7 TeV is consistent with no increase in the  $B_c^-$  fraction. Furthermore, using the assumption of no  $CP$  violation in the  $B_c^- \rightarrow J/\psi \mu^- \bar{\nu}$  decay, we find that the average asymmetry in  $B_c^- - B_c^+$  production is consistent with zero. The measurements are  $(-2.5 \pm 2.1 \pm 0.5)\%$ , and  $(-0.5 \pm 1.1 \pm 0.4)\%$  at center-of-mass energies of 7 and 13 TeV, respectively.

These results are useful to extract absolute branching fractions for  $B_c^-$  measurements, albeit with a relatively large uncertainty. They also challenge QCD calculations to predict the measured  $B_c^-$  fractions and explain the consistency between the fractions measured at 7 and 13 TeV [14, 54].

## Acknowledgements

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3

(France); BMBF, DFG and MPG (Germany); INFN (Italy); NWO (Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MSHE (Russia); MinECo (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); DOE NP and NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (Netherlands), PIC (Spain), GridPP (United Kingdom), RRCKI and Yandex LLC (Russia), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), PL-GRID (Poland) and OSC (USA). We are indebted to the communities behind the multiple open-source software packages on which we depend. Individual groups or members have received support from AvH Foundation (Germany); EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union); ANR, Labex P2IO and OCEVU, and Région Auvergne-Rhône-Alpes (France); Key Research Program of Frontier Sciences of CAS, CAS PIFI, and the Thousand Talents Program (China); RFBR, RSF and Yandex LLC (Russia); GVA, XuntaGal and GENCAT (Spain); the Royal Society and the Leverhulme Trust (United Kingdom).

## References

- [1] LHCb collaboration, R. Aaij *et al.*, *Measurement of  $b$  hadron fractions in 13 TeV  $pp$  collisions*, Phys. Rev. **D100** (2019) 031102, [arXiv:1902.06794](#).
- [2] CDF collaboration, F. Abe *et al.*, *Observation of the  $B_c$  meson in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8$  TeV*, Phys. Rev. Lett. **81** (1998) 2432, [arXiv:hep-ex/9805034](#).
- [3] CDF collaboration, F. Abe *et al.*, *Observation of  $B_c$  mesons in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8$  TeV*, Phys. Rev. **D58** (1998) 112004, [arXiv:hep-ex/9804014](#).
- [4] LHCb collaboration, R. Aaij *et al.*, *Measurements of  $B_c^+$  production and mass with the  $B_c^+ \rightarrow J/\psi \pi^+$  decay*, Phys. Rev. Lett. **109** (2012) 232001, [arXiv:1209.5634](#).
- [5] LHCb collaboration, R. Aaij *et al.*, *Measurement of  $B_c^+$  production in proton-proton collisions at  $\sqrt{s} = 8$  TeV*, Phys. Rev. Lett. **114** (2015) 132001, [arXiv:1411.2943](#).
- [6] CMS collaboration, *Measurement of  $\frac{\sigma(B_c^+) \times \mathcal{B}(B_c^+ \rightarrow J/\psi \pi^+)}{\sigma(B^+) \times \mathcal{B}(B^+ \rightarrow J/\psi K^+)}$  and  $\frac{\sigma(B_c^+) \times \mathcal{B}(B_c^+ \rightarrow J/\psi \pi^+ \pi^+ \pi^-)}{\sigma(B_c^+) \times \mathcal{B}(B^+ \rightarrow J/\psi \pi^+)}$  at  $\sqrt{s} = 7$  TeV*, CMS-PAS-BPH-12-011.
- [7] CMS collaboration, *Measurement of production cross sections times branching fraction of  $B_c^+ \rightarrow J/\psi \pi^+$  and  $B^+ \rightarrow J/\psi K^+$  in  $pp$  collisions at  $\sqrt{s} = 7$  TeV at CMS*, CMS-PAS-BPH-13-002.
- [8] CLEO collaboration, A. H. Mahmood *et al.*, *Measurement of the  $B$ -meson inclusive semileptonic branching fraction and electron energy moments*, Phys. Rev. **D70** (2004) 032003, [arXiv:hep-ex/0403053](#).
- [9] BaBar collaboration, B. Aubert *et al.*, *Measurement of the ratio  $\mathcal{B}(B^+ \rightarrow X e \nu) / \mathcal{B}(B^0 \rightarrow X e \nu)$* , Phys. Rev. **D74** (2006) 091105, [arXiv:hep-ex/0607111](#).
- [10] Belle collaboration, P. Urquijo *et al.*, *Moments of the electron energy spectrum and partial branching fraction of  $B \rightarrow X_c e \nu$  decays at Belle*, Phys. Rev. **D75** (2007) 032001, [arXiv:hep-ex/0610012](#).

- [11] LHCb collaboration, R. Aaij *et al.*, *Measurement of the  $b$ -quark production cross-section in 7 and 13 TeV  $pp$  collisions*, Phys. Rev. Lett. **118** (2017) 052002, Erratum *ibid.* **119** (2017) 169901, [arXiv:1612.05140](#).
- [12] Particle Data Group, M. Tanabashi *et al.*, *Review of particle physics*, Phys. Rev. **D98** (2018) 030001.
- [13] CLEO collaboration, G. Bonvicini *et al.*, *Updated measurements of absolute  $D^+$  and  $D^0$  hadronic branching fractions and  $\sigma(e^+e^- \rightarrow D\bar{D})$  at  $E_{\text{cm}} = 3774$  MeV*, Phys. Rev. **D89** (2014) 072002, Erratum *ibid.* **D91** (2015) 019903, [arXiv:1312.6775](#).
- [14] N. Brambilla *et al.*, *Heavy quarkonium: progress, puzzles, and opportunities*, Eur. Phys. J. **C71** (2011) 1534, [arXiv:1010.5827](#).
- [15] Z. Rui, H. Li, G.-x. Wang, and Y. Xiao, *Semileptonic decays of  $B_c$  meson to  $S$ -wave charmonium states in the perturbative QCD approach*, Eur. Phys. J. **C76** (2016) 564, [arXiv:1602.08918](#).
- [16] D. Ebert, R. N. Faustov, and V. O. Galkin, *Semileptonic and nonleptonic decays of  $B_c$  mesons to orbitally excited heavy mesons in the relativistic quark model*, Phys. Rev. **D82** (2010) 034019, [arXiv:1007.1369](#).
- [17] D. Ebert, R. N. Faustov, and V. O. Galkin, *Weak decays of the  $B_c$  meson to charmonium and  $D$  mesons in the relativistic quark model*, Phys. Rev. **D68** (2003) 094020, [arXiv:hep-ph/0306306](#).
- [18] C.-F. Qiao and R.-L. Zhu, *Estimation of semileptonic decays of  $B_c$  meson to  $S$ -wave charmonia with nonrelativistic QCD*, Phys. Rev. **D87** (2013) 014009, [arXiv:1208.5916](#).
- [19] C.-H. Chang, H.-F. Fu, G.-L. Wang, and J.-M. Zhang, *Some of semileptonic and nonleptonic decays of  $B_c$  meson in a Bethe-Salpeter relativistic quark model*, Sci. China Phys. Mech. Astron. **58** (2015) 071001, [arXiv:1411.3428](#).
- [20] M. A. Ivanov, J. G. Korner, and P. Santorelli, *Exclusive semileptonic and nonleptonic decays of the  $B_c$  meson*, Phys. Rev. **D73** (2006) 054024, [arXiv:hep-ph/0602050](#).
- [21] T. Huang and F. Zuo, *Semileptonic  $B_c$  decays and charmonium distribution amplitude*, Eur. Phys. J. **C51** (2007) 833, [arXiv:hep-ph/0702147](#).
- [22] W. Wang, Y.-L. Shen, and C.-D. Lu, *Covariant light-front approach for  $B_c$  transition form factors*, Phys. Rev. **D79** (2009) 054012, [arXiv:0811.3748](#).
- [23] E. Hernandez, J. Nieves, and J. M. Verde-Velasco, *Study of exclusive semileptonic and non-leptonic decays of  $B_c^-$  in a nonrelativistic quark model*, Phys. Rev. **D74** (2006) 074008, [arXiv:hep-ph/0607150](#).
- [24] P. Colangelo and F. De Fazio, *Using heavy quark spin symmetry in semileptonic  $B_c$  decays*, Phys. Rev. **D61** (2000) 034012, [arXiv:hep-ph/9909423](#).
- [25] I. P. Gouz *et al.*, *Prospects for the  $B_c$  studies at LHCb*, Phys. Atom. Nucl. **67** (2004) 1559, [arXiv:hep-ph/0211432](#), [*Yad. Fiz.*67,1581(2004)].

- [26] A. Abd El-Hady, J. H. Munoz, and J. P. Vary, *Semileptonic and nonleptonic  $B_c$  decays*, Phys. Rev. **D62** (2000) 014019, [arXiv:hep-ph/9909406](#).
- [27] V. V. Kiselev, *Exclusive decays and lifetime of  $B_c$  meson in QCD sum rules*, [arXiv:hep-ph/0211021](#).
- [28] C.-H. Chang and Y.-Q. Chen, *The decays of  $B_c$  meson*, Phys. Rev. **D49** (1994) 3399.
- [29] M. A. Ivanov, J. G. Korner, and P. Santorelli, *The semileptonic decays of the  $B_c$  meson*, Phys. Rev. **D63** (2001) 074010, [arXiv:hep-ph/0007169](#).
- [30] D. Scora and N. Isgur, *Semileptonic meson decays in the quark model: An update*, Phys. Rev. **D52** (1995) 2783, [arXiv:hep-ph/9503486](#).
- [31] A. Yu. Anisimov, P. Yu. Kulikov, I. M. Narodetsky, and K. A. Ter-Martirosian, *Exclusive and inclusive decays of the  $B_c$  meson in the light front ISGW model*, Phys. Atom. Nucl. **62** (1999) 1739, [arXiv:hep-ph/9809249](#), [*Yad. Fiz.*62,1868(1999)].
- [32] W. Wang and R. Zhu, *Model independent investigation of the  $R_{J/\psi, \eta_c}$  and ratios of decay widths of semileptonic  $B_c$  decays into a  $P$ -wave charmonium*, [arXiv:1808.10830](#).
- [33] Z.-K. Geng *et al.*, *Relativistic effects in the semileptonic  $B_c$  decays to charmonium with the Bethe-Salpeter method*, [arXiv:1809.02968](#).
- [34] D. Leljak, B. Melic, and M. Patra, *On lepton flavour universality in semileptonic  $B_c \rightarrow \eta_c, J/\psi$  decays*, [arXiv:1901.08368](#).
- [35] LHCb collaboration, A. A. Alves Jr. *et al.*, *The LHCb detector at the LHC*, JINST **3** (2008) S08005.
- [36] LHCb collaboration, R. Aaij *et al.*, *LHCb detector performance*, Int. J. Mod. Phys. **A30** (2015) 1530022, [arXiv:1412.6352](#).
- [37] R. Aaij *et al.*, *The LHCb trigger and its performance in 2011*, JINST **8** (2013) P04022, [arXiv:1211.3055](#).
- [38] LHCb collaboration, R. Aaij *et al.*, *Measurement of  $b$  hadron production fractions in 7 TeVpp collisions*, Phys. Rev. **D85** (2012) 032008, [arXiv:1111.2357](#).
- [39] V. V. Gligorov and M. Williams, *Efficient, reliable and fast high-level triggering using a bonsai boosted decision tree*, JINST **8** (2013) P02013, [arXiv:1210.6861](#).
- [40] T. Sjöstrand, S. Mrenna, and P. Skands, *PYTHIA 6.4 physics and manual*, JHEP **05** (2006) 026, [arXiv:hep-ph/0603175](#); T. Sjöstrand, S. Mrenna, and P. Skands, *A brief introduction to PYTHIA 8.1*, Comput. Phys. Commun. **178** (2008) 852, [arXiv:0710.3820](#).
- [41] I. Belyaev *et al.*, *Handling of the generation of primary events in Gauss, the LHCb simulation framework*, J. Phys. Conf. Ser. **331** (2011) 032047.
- [42] D. J. Lange, *The EvtGen particle decay simulation package*, Nucl. Instrum. Meth. **A462** (2001) 152.

- [43] P. Golonka and Z. Was, *PHOTOS Monte Carlo: A precision tool for QED corrections in Z and W decays*, Eur. Phys. J. **C45** (2006) 97, [arXiv:hep-ph/0506026](#).
- [44] Geant4 collaboration, J. Allison *et al.*, *Geant4 developments and applications*, IEEE Trans. Nucl. Sci. **53** (2006) 270; Geant4 collaboration, S. Agostinelli *et al.*, *Geant4: A simulation toolkit*, Nucl. Instrum. Meth. **A506** (2003) 250.
- [45] M. Clemencic *et al.*, *The LHCb simulation application, Gauss: Design, evolution and experience*, J. Phys. Conf. Ser. **331** (2011) 032023.
- [46] H. Voss, A. Hoecker, J. Stelzer, and F. Tegenfeldt, *TMVA - Toolkit for Multivariate Data Analysis with ROOT*, PoS **ACAT** (2007) 040; A. Hoecker *et al.*, *TMVA 4 — Toolkit for Multivariate Data Analysis with ROOT. Users Guide.*, [arXiv:physics/0703039](#).
- [47] LHCb collaboration, R. Aaij *et al.*, *Measurement of the ratio of branching fractions and difference in CP asymmetries of the decays  $B^+ \rightarrow J/\psi \pi^+$  and  $B^+ \rightarrow J/\psi K^+$* , JHEP **03** (2017) 036, [arXiv:1612.06116](#).
- [48] LHCb collaboration, R. Aaij *et al.*, *Measurement of the  $B^\pm$  production asymmetry and the CP asymmetry in  $B^\pm \rightarrow J/\psi K^\pm$  decays*, Phys. Rev. **D95** (2017) 052005, [arXiv:1701.05501](#).
- [49] K. S. Cranmer, *Kernel estimation in high-energy physics*, Comput. Phys. Commun. **136** (2001) 198, [arXiv:hep-ex/0011057](#).
- [50] LHCb collaboration, R. Aaij *et al.*, *Measurement of the ratio of branching fractions  $\mathcal{B}(B_c^+ \rightarrow J/\psi \tau^+ \nu_\tau)/\mathcal{B}(B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu)$* , Phys. Rev. Lett. **120** (2018) 121801, [arXiv:1711.05623](#).
- [51] R. Aaij *et al.*, *Selection and processing of calibration samples to measure the particle identification performance of the LHCb experiment in Run 2*, Eur. Phys. J. Tech. Instr. **6** (2018) 1, [arXiv:1803.00824](#).
- [52] LHCb collaboration, R. Aaij *et al.*, *Measurement of the track reconstruction efficiency at LHCb*, JINST **10** (2015) P02007, [arXiv:1408.1251](#).
- [53] LHCb collaboration, R. Aaij *et al.*, *Measurement of the fragmentation fraction ratio  $f_s/f_d$  and its dependence on B meson kinematics*, JHEP **04** (2013) 001, [arXiv:1301.5286](#).
- [54] A. Ali, Q. Qin, and W. Wang, *Discovery potential of stable and near-threshold doubly heavy tetraquarks at the LHC*, Phys. Lett. **B785** (2018) 605, [arXiv:1806.09288](#).

## LHCb collaboration

R. Aaij<sup>31</sup>, C. Abellán Beteta<sup>49</sup>, T. Ackernley<sup>59</sup>, B. Adeva<sup>45</sup>, M. Adinolfi<sup>53</sup>, H. Afsharnia<sup>9</sup>, C.A. Aidala<sup>79</sup>, S. Aiola<sup>25</sup>, Z. Ajaltouni<sup>9</sup>, S. Akar<sup>64</sup>, P. Albicocco<sup>22</sup>, J. Albrecht<sup>14</sup>, F. Alessio<sup>47</sup>, M. Alexander<sup>58</sup>, A. Alfonso Alberro<sup>44</sup>, G. Alkhazov<sup>37</sup>, P. Alvarez Cartelle<sup>60</sup>, A.A. Alves Jr<sup>45</sup>, S. Amato<sup>2</sup>, Y. Amhis<sup>11</sup>, L. An<sup>21</sup>, L. Anderlini<sup>21</sup>, G. Andreassi<sup>48</sup>, M. Andreotti<sup>20</sup>, F. Archilli<sup>16</sup>, J. Arnau Romeu<sup>10</sup>, A. Artamonov<sup>43</sup>, M. Artuso<sup>67</sup>, K. Arzymatov<sup>41</sup>, E. Aslanides<sup>10</sup>, M. Atzeni<sup>49</sup>, B. Audurier<sup>26</sup>, S. Bachmann<sup>16</sup>, J.J. Back<sup>55</sup>, S. Baker<sup>60</sup>, V. Balagura<sup>11,b</sup>, W. Baldini<sup>20,47</sup>, A. Baranov<sup>41</sup>, R.J. Barlow<sup>61</sup>, S. Barsuk<sup>11</sup>, W. Barter<sup>60</sup>, M. Bartolini<sup>23,47,h</sup>, F. Baryshnikov<sup>76</sup>, G. Bassi<sup>28</sup>, V. Batozskaya<sup>35</sup>, B. Batsukh<sup>67</sup>, A. Battig<sup>14</sup>, A. Bay<sup>48</sup>, M. Becker<sup>14</sup>, F. Bedeschi<sup>28</sup>, I. Bediaga<sup>1</sup>, A. Beiter<sup>67</sup>, L.J. Bel<sup>31</sup>, V. Belavin<sup>41</sup>, S. Belin<sup>26</sup>, N. Belyi<sup>5</sup>, V. Bellec<sup>48</sup>, K. Belous<sup>43</sup>, I. Belyaev<sup>38</sup>, G. Bencivenni<sup>22</sup>, E. Ben-Haim<sup>12</sup>, S. Benson<sup>31</sup>, S. Beranek<sup>13</sup>, A. Berezhnoy<sup>39</sup>, R. Bernet<sup>49</sup>, D. Berninghoff<sup>16</sup>, H.C. Bernstein<sup>67</sup>, E. Bertholet<sup>12</sup>, A. Bertolin<sup>27</sup>, C. Betancourt<sup>49</sup>, F. Betti<sup>19,e</sup>, M.O. Bettler<sup>54</sup>, Ia. Bezshyiko<sup>49</sup>, S. Bhasin<sup>53</sup>, J. Bhom<sup>33</sup>, M.S. Bieker<sup>14</sup>, S. Bifani<sup>52</sup>, P. Billoir<sup>12</sup>, A. Bizzeti<sup>21,u</sup>, M. Bjørn<sup>62</sup>, M.P. Blago<sup>47</sup>, T. Blake<sup>55</sup>, F. Blanc<sup>48</sup>, S. Blusk<sup>67</sup>, D. Bobulska<sup>58</sup>, V. Bocci<sup>30</sup>, O. Boente Garcia<sup>45</sup>, T. Boettcher<sup>63</sup>, A. Boldyrev<sup>77</sup>, A. Bondar<sup>42,x</sup>, N. Bondar<sup>37</sup>, S. Borghi<sup>61,47</sup>, M. Borisyak<sup>41</sup>, M. Borsato<sup>16</sup>, J.T. Borsuk<sup>33</sup>, T.J.V. Bowcock<sup>59</sup>, C. Bozzi<sup>20</sup>, M.J. Bradley<sup>60</sup>, S. Braun<sup>16</sup>, A. Brea Rodriguez<sup>45</sup>, M. Brodski<sup>47</sup>, J. Brodzicka<sup>33</sup>, A. Brossa Gonzalo<sup>55</sup>, D. Brundu<sup>26</sup>, E. Buchanan<sup>53</sup>, A. Buonauro<sup>49</sup>, C. Burr<sup>47</sup>, A. Bursche<sup>26</sup>, J.S. Butter<sup>31</sup>, J. Buytaert<sup>47</sup>, W. Byczynski<sup>47</sup>, S. Cadeddu<sup>26</sup>, H. Cai<sup>71</sup>, R. Calabrese<sup>20,g</sup>, L. Calero Diaz<sup>22</sup>, S. Cali<sup>22</sup>, R. Calladine<sup>52</sup>, M. Calvi<sup>24,i</sup>, M. Calvo Gomez<sup>44,m</sup>, A. Camboni<sup>44</sup>, P. Campana<sup>22</sup>, D.H. Campora Perez<sup>47</sup>, L. Capriotti<sup>19,e</sup>, A. Carbone<sup>19,e</sup>, G. Carboni<sup>29</sup>, R. Cardinale<sup>23,h</sup>, A. Cardini<sup>26</sup>, P. Carniti<sup>24,i</sup>, K. Carvalho Akiba<sup>31</sup>, A. Casais Vidal<sup>45</sup>, G. Casse<sup>59</sup>, M. Cattaneo<sup>47</sup>, G. Cavallero<sup>47</sup>, R. Cenci<sup>28,p</sup>, J. Cerasoli<sup>10</sup>, M.G. Chapman<sup>53</sup>, M. Charles<sup>12,47</sup>, Ph. Charpentier<sup>47</sup>, G. Chatzikonstantinidis<sup>52</sup>, M. Chefdeville<sup>8</sup>, V. Chekalina<sup>41</sup>, C. Chen<sup>3</sup>, S. Chen<sup>26</sup>, A. Chernov<sup>33</sup>, S.-G. Chitic<sup>47</sup>, V. Chobanova<sup>45</sup>, M. Chruszcz<sup>33</sup>, A. Chubykin<sup>37</sup>, P. Ciambrone<sup>22</sup>, M.F. Cicala<sup>55</sup>, X. Cid Vidal<sup>45</sup>, G. Ciezarek<sup>47</sup>, F. Cindolo<sup>19</sup>, P.E.L. Clarke<sup>57</sup>, M. Clemencic<sup>47</sup>, H.V. Cliff<sup>54</sup>, J. Closier<sup>47</sup>, J.L. Cobbedick<sup>61</sup>, V. Coco<sup>47</sup>, J.A.B. Coelho<sup>11</sup>, J. Cogan<sup>10</sup>, E. Cogneras<sup>9</sup>, L. Cojocariu<sup>36</sup>, P. Collins<sup>47</sup>, T. Colombo<sup>47</sup>, A. Comerma-Montells<sup>16</sup>, A. Contu<sup>26</sup>, N. Cooke<sup>52</sup>, G. Coombs<sup>58</sup>, S. Coquereau<sup>44</sup>, G. Corti<sup>47</sup>, C.M. Costa Sobral<sup>55</sup>, B. Couturier<sup>47</sup>, D.C. Craik<sup>63</sup>, J. Crkovska<sup>66</sup>, A. Crocombe<sup>55</sup>, M. Cruz Torres<sup>1</sup>, R. Currie<sup>57</sup>, C.L. Da Silva<sup>66</sup>, E. Dall'Occo<sup>14</sup>, J. Dalseno<sup>45,53</sup>, C. D'Ambrosio<sup>47</sup>, A. Danilina<sup>38</sup>, P. d'Argent<sup>16</sup>, A. Davis<sup>61</sup>, O. De Aguiar Francisco<sup>47</sup>, K. De Bruyn<sup>47</sup>, S. De Capua<sup>61</sup>, M. De Cian<sup>48</sup>, J.M. De Miranda<sup>1</sup>, L. De Paula<sup>2</sup>, M. De Serio<sup>18,d</sup>, P. De Simone<sup>22</sup>, J.A. de Vries<sup>31</sup>, C.T. Dean<sup>66</sup>, W. Dean<sup>79</sup>, D. Decamp<sup>8</sup>, L. Del Buono<sup>12</sup>, B. Delaney<sup>54</sup>, H.-P. Dembinski<sup>15</sup>, M. Demmer<sup>14</sup>, A. Dendek<sup>34</sup>, V. Denysenko<sup>49</sup>, D. Derkach<sup>77</sup>, O. Deschamps<sup>9</sup>, F. Desse<sup>11</sup>, F. Dettori<sup>26</sup>, B. Dey<sup>7</sup>, A. Di Canto<sup>47</sup>, P. Di Nezza<sup>22</sup>, S. Didenko<sup>76</sup>, H. Dijkstra<sup>47</sup>, V. Dobishuk<sup>51</sup>, F. Dordei<sup>26</sup>, M. Dorigo<sup>28,y</sup>, A.C. dos Reis<sup>1</sup>, L. Douglas<sup>58</sup>, A. Dovbnya<sup>50</sup>, K. Dreimanis<sup>59</sup>, M.W. Dudek<sup>33</sup>, L. Dufour<sup>47</sup>, G. Dujany<sup>12</sup>, P. Durante<sup>47</sup>, J.M. Durham<sup>66</sup>, D. Dutta<sup>61</sup>, R. Dzhelyadin<sup>43,†</sup>, M. Dziewiecki<sup>16</sup>, A. Dziurda<sup>33</sup>, A. Dzyuba<sup>37</sup>, S. Easo<sup>56</sup>, U. Egede<sup>60</sup>, V. Egorychev<sup>38</sup>, S. Eidelman<sup>42,x</sup>, S. Eisenhardt<sup>57</sup>, R. Ekelhof<sup>14</sup>, S. Ek-In<sup>48</sup>, L. Eklund<sup>58</sup>, S. Ely<sup>67</sup>, A. Ene<sup>36</sup>, S. Escher<sup>13</sup>, S. Esen<sup>31</sup>, T. Evans<sup>47</sup>, A. Falabella<sup>19</sup>, J. Fan<sup>3</sup>, N. Farley<sup>52</sup>, S. Farry<sup>59</sup>, D. Fazzini<sup>11</sup>, M. Féo<sup>47</sup>, P. Fernandez Declara<sup>47</sup>, A. Fernandez Prieto<sup>45</sup>, F. Ferrari<sup>19,e</sup>, L. Ferreira Lopes<sup>48</sup>, F. Ferreira Rodrigues<sup>2</sup>, S. Ferreres Sole<sup>31</sup>, M. Ferrillo<sup>49</sup>, M. Ferro-Luzzi<sup>47</sup>, S. Filippov<sup>40</sup>, R.A. Fini<sup>18</sup>, M. Fiorini<sup>20,g</sup>, M. Firlej<sup>34</sup>, K.M. Fischer<sup>62</sup>, C. Fitzpatrick<sup>47</sup>, T. Fiutowski<sup>34</sup>, F. Fleuret<sup>11,b</sup>, M. Fontana<sup>47</sup>, F. Fontanelli<sup>23,h</sup>, R. Forty<sup>47</sup>, V. Franco Lima<sup>59</sup>, M. Franco Sevilla<sup>65</sup>, M. Frank<sup>47</sup>, C. Frei<sup>47</sup>, D.A. Friday<sup>58</sup>, J. Fu<sup>25,q</sup>, M. Fuehring<sup>14</sup>, W. Funk<sup>47</sup>, E. Gabriel<sup>57</sup>, A. Gallas Torreira<sup>45</sup>, D. Galli<sup>19,e</sup>, S. Gallorini<sup>27</sup>, S. Gambetta<sup>57</sup>, Y. Gan<sup>3</sup>, M. Gandelman<sup>2</sup>, P. Gandini<sup>25</sup>, Y. Gao<sup>4</sup>, L.M. Garcia Martin<sup>46</sup>, J. García Pardiñas<sup>49</sup>, B. Garcia Plana<sup>45</sup>, F.A. Garcia Rosales<sup>11</sup>, J. Garra Tico<sup>54</sup>, L. Garrido<sup>44</sup>,

D. Gascon<sup>44</sup>, C. Gaspar<sup>47</sup>, D. Gerick<sup>16</sup>, E. Gersabeck<sup>61</sup>, M. Gersabeck<sup>61</sup>, T. Gershon<sup>55</sup>,  
 D. Gerstel<sup>10</sup>, Ph. Ghez<sup>8</sup>, V. Gibson<sup>54</sup>, A. Gioventù<sup>45</sup>, O.G. Girard<sup>48</sup>, P. Gironella Gironell<sup>44</sup>,  
 L. Giubega<sup>36</sup>, C. Giugliano<sup>20</sup>, K. Gizdov<sup>57</sup>, V.V. Gligorov<sup>12</sup>, C. Göbel<sup>69</sup>, D. Golubkov<sup>38</sup>,  
 A. Golutvin<sup>60,76</sup>, A. Gomes<sup>1,a</sup>, P. Gorbounov<sup>38,6</sup>, I.V. Gorelov<sup>39</sup>, C. Gotti<sup>24,i</sup>, E. Govorkova<sup>31</sup>,  
 J.P. Grabowski<sup>16</sup>, R. Graciani Diaz<sup>44</sup>, T. Grammatico<sup>12</sup>, L.A. Granado Cardoso<sup>47</sup>,  
 E. Graugés<sup>44</sup>, E. Graverini<sup>48</sup>, G. Graziani<sup>21</sup>, A. Greco<sup>36</sup>, R. Greim<sup>31</sup>, P. Griffith<sup>20</sup>, L. Grillo<sup>61</sup>,  
 L. Gruber<sup>47</sup>, B.R. Gruberg Cazon<sup>62</sup>, C. Gu<sup>3</sup>, E. Gushchin<sup>40</sup>, A. Guth<sup>13</sup>, Yu. Guz<sup>43,47</sup>, T. Gys<sup>47</sup>,  
 T. Hadavizadeh<sup>62</sup>, G. Haefeli<sup>48</sup>, C. Haen<sup>47</sup>, S.C. Haines<sup>54</sup>, P.M. Hamilton<sup>65</sup>, Q. Han<sup>7</sup>, X. Han<sup>16</sup>,  
 T.H. Hancock<sup>62</sup>, S. Hansmann-Menzemer<sup>16</sup>, N. Harnew<sup>62</sup>, T. Harrison<sup>59</sup>, R. Hart<sup>31</sup>, C. Hasse<sup>47</sup>,  
 M. Hatch<sup>47</sup>, J. He<sup>5</sup>, M. Hecker<sup>60</sup>, K. Heijhoff<sup>31</sup>, K. Heinicke<sup>14</sup>, A. Heister<sup>14</sup>, A.M. Hennequin<sup>47</sup>,  
 K. Hennessy<sup>59</sup>, L. Henry<sup>46</sup>, J. Heuel<sup>13</sup>, A. Hicheur<sup>68</sup>, R. Hidalgo Charman<sup>61</sup>, D. Hill<sup>62</sup>,  
 M. Hilton<sup>61</sup>, P.H. Hopchev<sup>48</sup>, J. Hu<sup>16</sup>, W. Hu<sup>7</sup>, W. Huang<sup>5</sup>, W. Hulsbergen<sup>31</sup>, T. Humair<sup>60</sup>,  
 R.J. Hunter<sup>55</sup>, M. Hushchyn<sup>77</sup>, D. Hutchcroft<sup>59</sup>, D. Hynds<sup>31</sup>, P. Ibis<sup>14</sup>, M. Idzik<sup>34</sup>, P. Ilten<sup>52</sup>,  
 A. Inglessi<sup>37</sup>, A. Inyakin<sup>43</sup>, K. Ivshin<sup>37</sup>, R. Jacobsson<sup>47</sup>, S. Jakobsen<sup>47</sup>, J. Jalocha<sup>62</sup>, E. Jans<sup>31</sup>,  
 B.K. Jashal<sup>46</sup>, A. Jawahery<sup>65</sup>, V. Jevtic<sup>14</sup>, F. Jiang<sup>3</sup>, M. John<sup>62</sup>, D. Johnson<sup>47</sup>, C.R. Jones<sup>54</sup>,  
 B. Jost<sup>47</sup>, N. Jurik<sup>62</sup>, S. Kandybei<sup>50</sup>, M. Karacson<sup>47</sup>, J.M. Kariuki<sup>53</sup>, N. Kazeev<sup>77</sup>, M. Kecke<sup>16</sup>,  
 F. Keizer<sup>54,54</sup>, M. Kelsey<sup>67</sup>, M. Kenzie<sup>54</sup>, T. Ketel<sup>32</sup>, B. Khanji<sup>47</sup>, A. Kharisova<sup>78</sup>, K.E. Kim<sup>67</sup>,  
 T. Kirn<sup>13</sup>, V.S. Kirsebom<sup>48</sup>, S. Klaver<sup>22</sup>, K. Klimaszewski<sup>35</sup>, S. Koliiev<sup>51</sup>, A. Kondybayeva<sup>76</sup>,  
 A. Konoplyannikov<sup>38</sup>, P. Kopciwicz<sup>34</sup>, R. Kopecna<sup>16</sup>, P. Koppenburg<sup>31</sup>, I. Kostiuk<sup>31,51</sup>,  
 O. Kot<sup>51</sup>, S. Kotriakhova<sup>37</sup>, L. Kravchuk<sup>40</sup>, R.D. Krawczyk<sup>47</sup>, M. Kreps<sup>55</sup>, F. Kress<sup>60</sup>,  
 S. Kretschmar<sup>13</sup>, P. Krokovny<sup>42,x</sup>, W. Krupa<sup>34</sup>, W. Krzemien<sup>35</sup>, W. Kucewicz<sup>33,l</sup>,  
 M. Kucharczyk<sup>33</sup>, V. Kudryavtsev<sup>42,x</sup>, H.S. Kuindersma<sup>31</sup>, G.J. Kunde<sup>66</sup>, T. Kvaratskheliya<sup>38</sup>,  
 D. Lacarrere<sup>47</sup>, G. Lafferty<sup>61</sup>, A. Lai<sup>26</sup>, D. Lancierini<sup>49</sup>, J.J. Lane<sup>61</sup>, G. Lanfranchi<sup>22</sup>,  
 C. Langenbruch<sup>13</sup>, T. Latham<sup>55</sup>, F. Lazzari<sup>28,v</sup>, C. Lazzeroni<sup>52</sup>, R. Le Gac<sup>10</sup>, R. Lefèvre<sup>9</sup>,  
 A. Leflat<sup>39</sup>, F. Lemaitre<sup>47</sup>, O. Leroy<sup>10</sup>, T. Lesiak<sup>33</sup>, B. Leverington<sup>16</sup>, H. Li<sup>70</sup>, X. Li<sup>66</sup>, Y. Li<sup>6</sup>,  
 Z. Li<sup>67</sup>, X. Liang<sup>67</sup>, R. Lindner<sup>47</sup>, V. Lisovskyi<sup>11</sup>, G. Liu<sup>70</sup>, X. Liu<sup>3</sup>, D. Loh<sup>55</sup>, A. Loi<sup>26</sup>,  
 J. Lomba Castro<sup>45</sup>, I. Longstaff<sup>58</sup>, J.H. Lopes<sup>2</sup>, G. Loustau<sup>49</sup>, G.H. Lovell<sup>54</sup>, Y. Lu<sup>6</sup>,  
 D. Lucchesi<sup>27,o</sup>, M. Lucio Martinez<sup>31</sup>, Y. Luo<sup>3</sup>, A. Lupato<sup>27</sup>, E. Luppi<sup>20,g</sup>, O. Lupton<sup>55</sup>,  
 A. Lusiani<sup>28</sup>, X. Lyu<sup>5</sup>, S. Maccolini<sup>19,e</sup>, F. Machefert<sup>11</sup>, F. Maciuc<sup>36</sup>, V. Macko<sup>48</sup>,  
 P. Mackowiak<sup>14</sup>, S. Maddrell-Mander<sup>53</sup>, L.R. Madhan Mohan<sup>53</sup>, O. Maev<sup>37,47</sup>, A. Maevskiy<sup>77</sup>,  
 D. Maisuzenko<sup>37</sup>, M.W. Majewski<sup>34</sup>, S. Malde<sup>62</sup>, B. Malecki<sup>47</sup>, A. Malinin<sup>75</sup>, T. Maltsev<sup>42,x</sup>,  
 H. Malygina<sup>16</sup>, G. Manca<sup>26,f</sup>, G. Mancinelli<sup>10</sup>, R. Manera Escalero<sup>44</sup>, D. Manuzzi<sup>19,e</sup>,  
 D. Marangotto<sup>25,q</sup>, J. Maratas<sup>9,w</sup>, J.F. Marchand<sup>8</sup>, U. Marconi<sup>19</sup>, S. Mariani<sup>21</sup>,  
 C. Marin Benito<sup>11</sup>, M. Marinangeli<sup>48</sup>, P. Marino<sup>48</sup>, J. Marks<sup>16</sup>, P.J. Marshall<sup>59</sup>, G. Martellotti<sup>30</sup>,  
 L. Martinazzoli<sup>47</sup>, M. Martinelli<sup>24</sup>, D. Martinez Santos<sup>45</sup>, F. Martinez Vidal<sup>46</sup>, A. Massafferri<sup>1</sup>,  
 M. Materok<sup>13</sup>, R. Matev<sup>47</sup>, A. Mathad<sup>49</sup>, Z. Mathe<sup>47</sup>, V. Matiunin<sup>38</sup>, C. Matteuzzi<sup>24</sup>,  
 K.R. Mattioli<sup>79</sup>, A. Mauri<sup>49</sup>, E. Maurice<sup>11,b</sup>, M. McCann<sup>60,47</sup>, L. McConnell<sup>17</sup>, A. McNab<sup>61</sup>,  
 R. McNulty<sup>17</sup>, J.V. Mead<sup>59</sup>, B. Meadows<sup>64</sup>, C. Meaux<sup>10</sup>, G. Meier<sup>14</sup>, N. Meinert<sup>73</sup>,  
 D. Melnychuk<sup>35</sup>, S. Meloni<sup>24,i</sup>, M. Merk<sup>31</sup>, A. Merli<sup>25</sup>, M. Mikhasenko<sup>47</sup>, D.A. Milanese<sup>72</sup>,  
 E. Millard<sup>55</sup>, M.-N. Minard<sup>8</sup>, O. Mineev<sup>38</sup>, L. Minzoni<sup>20,g</sup>, S.E. Mitchell<sup>57</sup>, B. Mitreska<sup>61</sup>,  
 D.S. Mitzel<sup>47</sup>, A. Mödden<sup>14</sup>, A. Mogini<sup>12</sup>, R.D. Moise<sup>60</sup>, T. Mombächer<sup>14</sup>, I.A. Monroy<sup>72</sup>,  
 S. Monteil<sup>9</sup>, M. Morandin<sup>27</sup>, G. Morello<sup>22</sup>, M.J. Morello<sup>28,t</sup>, J. Moron<sup>34</sup>, A.B. Morris<sup>10</sup>,  
 A.G. Morris<sup>55</sup>, R. Mountain<sup>67</sup>, H. Mu<sup>3</sup>, F. Muheim<sup>57</sup>, M. Mukherjee<sup>7</sup>, M. Mulder<sup>31</sup>,  
 D. Müller<sup>47</sup>, K. Müller<sup>49</sup>, V. Müller<sup>14</sup>, C.H. Murphy<sup>62</sup>, D. Murray<sup>61</sup>, P. Muzzetto<sup>26</sup>, P. Naik<sup>53</sup>,  
 T. Nakada<sup>48</sup>, R. Nandakumar<sup>56</sup>, A. Nandi<sup>62</sup>, T. Nanut<sup>48</sup>, I. Nasteva<sup>2</sup>, M. Needham<sup>57</sup>,  
 N. Neri<sup>25,q</sup>, S. Neubert<sup>16</sup>, N. Neufeld<sup>47</sup>, R. Newcombe<sup>60</sup>, T.D. Nguyen<sup>48</sup>, C. Nguyen-Mau<sup>48,n</sup>,  
 E.M. Niel<sup>11</sup>, S. Nieswand<sup>13</sup>, N. Nikitin<sup>39</sup>, N.S. Nolte<sup>47</sup>, C. Nunez<sup>79</sup>, A. Oblakowska-Mucha<sup>34</sup>,  
 V. Obraztsov<sup>43</sup>, S. Ogilvy<sup>58</sup>, D.P. O'Hanlon<sup>19</sup>, R. Oldeman<sup>26,f</sup>, C.J.G. Onderwater<sup>74</sup>, J.  
 D. Osborn<sup>79</sup>, A. Ossowska<sup>33</sup>, J.M. Otalora Goicochea<sup>2</sup>, T. Ovsianikova<sup>38</sup>, P. Owen<sup>49</sup>,  
 A. Oyanguren<sup>46</sup>, P.R. Pais<sup>48</sup>, T. Pajero<sup>28,t</sup>, A. Palano<sup>18</sup>, M. Palutan<sup>22</sup>, G. Panshin<sup>78</sup>,

A. Papanestis<sup>56</sup>, M. Pappagallo<sup>57</sup>, L.L. Pappalardo<sup>20,g</sup>, C. Pappenheimer<sup>64</sup>, W. Parker<sup>65</sup>,  
 C. Parkes<sup>61</sup>, G. Passaleva<sup>21,47</sup>, A. Pastore<sup>18</sup>, M. Patel<sup>60</sup>, C. Patrignani<sup>19,e</sup>, A. Pearce<sup>47</sup>,  
 A. Pellegrino<sup>31</sup>, M. Pepe Altarelli<sup>47</sup>, S. Perazzini<sup>19</sup>, D. Pereima<sup>38</sup>, P. Perret<sup>9</sup>, L. Pescatore<sup>48</sup>,  
 K. Petridis<sup>53</sup>, A. Petrolini<sup>23,h</sup>, A. Petrov<sup>75</sup>, S. Petrucci<sup>57</sup>, M. Petruzzo<sup>25,q</sup>, B. Pietrzyk<sup>8</sup>,  
 G. Pietrzyk<sup>48</sup>, M. Pikiés<sup>33</sup>, M. Pili<sup>62</sup>, D. Pinci<sup>30</sup>, J. Pinzino<sup>47</sup>, F. Pisani<sup>47</sup>, A. Piucci<sup>16</sup>,  
 V. Placinta<sup>36</sup>, S. Playfer<sup>57</sup>, J. Plews<sup>52</sup>, M. Plo Casasus<sup>45</sup>, F. Polci<sup>12</sup>, M. Poli Lener<sup>22</sup>,  
 M. Poliakov<sup>67</sup>, A. Poluektov<sup>10</sup>, N. Polukhina<sup>76,c</sup>, I. Polyakov<sup>67</sup>, E. Polycarpo<sup>2</sup>, G.J. Pomery<sup>53</sup>,  
 S. Ponce<sup>47</sup>, A. Popov<sup>43</sup>, D. Popov<sup>52</sup>, S. Poslavskii<sup>43</sup>, K. Prasanth<sup>33</sup>, L. Promberger<sup>47</sup>,  
 C. Prouve<sup>45</sup>, V. Pugatch<sup>51</sup>, A. Puig Navarro<sup>49</sup>, H. Pullen<sup>62</sup>, G. Punzi<sup>28,p</sup>, W. Qian<sup>5</sup>, J. Qin<sup>5</sup>,  
 R. Quagliani<sup>12</sup>, B. Quintana<sup>9</sup>, N.V. Raab<sup>17</sup>, R.I. Rabadan Trejo<sup>10</sup>, B. Rachwal<sup>34</sup>,  
 J.H. Rademacker<sup>53</sup>, M. Rama<sup>28</sup>, M. Ramos Pernas<sup>45</sup>, M.S. Rangel<sup>2</sup>, F. Ratnikov<sup>41,77</sup>,  
 G. Raven<sup>32</sup>, M. Reboud<sup>8</sup>, F. Redi<sup>48</sup>, F. Reiss<sup>12</sup>, C. Remon Alepuz<sup>46</sup>, Z. Ren<sup>3</sup>, V. Renaudin<sup>62</sup>,  
 S. Ricciardi<sup>56</sup>, S. Richards<sup>53</sup>, K. Rinnert<sup>59</sup>, P. Robbe<sup>11</sup>, A. Robert<sup>12</sup>, A.B. Rodrigues<sup>48</sup>,  
 E. Rodrigues<sup>64</sup>, J.A. Rodriguez Lopez<sup>72</sup>, M. Roehrken<sup>47</sup>, S. Roiser<sup>47</sup>, A. Rollings<sup>62</sup>,  
 V. Romanovskiy<sup>43</sup>, M. Romero Lamas<sup>45</sup>, A. Romero Vidal<sup>45</sup>, J.D. Roth<sup>79</sup>, M. Rotondo<sup>22</sup>,  
 M.S. Rudolph<sup>67</sup>, T. Ruf<sup>47</sup>, J. Ruiz Vidal<sup>46</sup>, J. Ryzka<sup>34</sup>, J.J. Saborido Silva<sup>45</sup>, N. Sagidova<sup>37</sup>,  
 B. Saitta<sup>26,f</sup>, C. Sanchez Gras<sup>31</sup>, C. Sanchez Mayordomo<sup>46</sup>, B. Sanmartin Sedes<sup>45</sup>,  
 R. Santacesaria<sup>30</sup>, C. Santamarina Rios<sup>45</sup>, M. Santimaria<sup>22</sup>, E. Santovetti<sup>29,j</sup>, G. Sarpis<sup>61</sup>,  
 A. Sarti<sup>30</sup>, C. Satriano<sup>30,s</sup>, A. Satta<sup>29</sup>, M. Saur<sup>5</sup>, D. Savrina<sup>38,39</sup>, L.G. Scantlebury Smead<sup>62</sup>,  
 S. Schael<sup>13</sup>, M. Schellenberg<sup>14</sup>, M. Schiller<sup>58</sup>, H. Schindler<sup>47</sup>, M. Schmelling<sup>15</sup>, T. Schmelzer<sup>14</sup>,  
 B. Schmidt<sup>47</sup>, O. Schneider<sup>48</sup>, A. Schopper<sup>47</sup>, H.F. Schreiner<sup>64</sup>, M. Schubiger<sup>31</sup>, S. Schulte<sup>48</sup>,  
 M.H. Schune<sup>11</sup>, R. Schwemmer<sup>47</sup>, B. Sciascia<sup>22</sup>, A. Sciubba<sup>30,k</sup>, S. Sellam<sup>68</sup>, A. Semennikov<sup>38</sup>,  
 A. Sergi<sup>52,47</sup>, N. Serra<sup>49</sup>, J. Serrano<sup>10</sup>, L. Sestini<sup>27</sup>, A. Seuthe<sup>14</sup>, P. Seyfert<sup>47</sup>, D.M. Shangase<sup>79</sup>,  
 M. Shapkin<sup>43</sup>, T. Shears<sup>59</sup>, L. Shekhtman<sup>42,x</sup>, V. Shevchenko<sup>75,76</sup>, E. Shmanin<sup>76</sup>,  
 J.D. Shupperd<sup>67</sup>, B.G. Siddi<sup>20</sup>, R. Silva Coutinho<sup>49</sup>, L. Silva de Oliveira<sup>2</sup>, G. Simi<sup>27,o</sup>,  
 S. Simone<sup>18,d</sup>, I. Skiba<sup>20</sup>, N. Skidmore<sup>16</sup>, T. Skwarnicki<sup>67</sup>, M.W. Slater<sup>52</sup>, J.G. Smeaton<sup>54</sup>,  
 A. Smetkina<sup>38</sup>, E. Smith<sup>13</sup>, I.T. Smith<sup>57</sup>, M. Smith<sup>60</sup>, A. Snoch<sup>31</sup>, M. Soares<sup>19</sup>,  
 L. Soares Lavra<sup>1</sup>, M.D. Sokoloff<sup>64</sup>, F.J.P. Soler<sup>58</sup>, B. Souza De Paula<sup>2</sup>, B. Spaan<sup>14</sup>,  
 E. Spadaro Norella<sup>25,q</sup>, P. Spradlin<sup>58</sup>, F. Stagni<sup>47</sup>, M. Stahl<sup>64</sup>, S. Stahl<sup>47</sup>, P. Stefko<sup>48</sup>,  
 S. Stefkova<sup>60</sup>, O. Steinkamp<sup>49</sup>, S. Stemmlé<sup>16</sup>, O. Stenyakin<sup>43</sup>, M. Stepanova<sup>37</sup>, H. Stevens<sup>14</sup>,  
 S. Stone<sup>67</sup>, S. Stracka<sup>28</sup>, M.E. Stramaglia<sup>48</sup>, M. Straticiu<sup>36</sup>, S. Strokov<sup>78</sup>, J. Sun<sup>3</sup>, L. Sun<sup>71</sup>,  
 Y. Sun<sup>65</sup>, P. Svihra<sup>61</sup>, K. Swientek<sup>34</sup>, A. Szabelski<sup>35</sup>, T. Szumlak<sup>34</sup>, M. Szymanski<sup>5</sup>, S. Taneja<sup>61</sup>,  
 Z. Tang<sup>3</sup>, T. Tekampe<sup>14</sup>, G. Tellarini<sup>20</sup>, F. Teubert<sup>47</sup>, E. Thomas<sup>47</sup>, K.A. Thomson<sup>59</sup>,  
 M.J. Tilley<sup>60</sup>, V. Tisserand<sup>9</sup>, S. T'Jampens<sup>8</sup>, M. Tobin<sup>6</sup>, S. Tolk<sup>47</sup>, L. Tomassetti<sup>20,g</sup>,  
 D. Tonelli<sup>28</sup>, D.Y. Tou<sup>12</sup>, E. Tournefier<sup>8</sup>, M. Traill<sup>58</sup>, M.T. Tran<sup>48</sup>, C. Trippl<sup>48</sup>, A. Trisovic<sup>54</sup>,  
 A. Tsaregorodtsev<sup>10</sup>, G. Tuci<sup>28,47,p</sup>, A. Tully<sup>48</sup>, N. Tuning<sup>31</sup>, A. Ukleja<sup>35</sup>, A. Usachov<sup>11</sup>,  
 A. Ustyuzhanin<sup>41,77</sup>, U. Uwer<sup>16</sup>, A. Vagner<sup>78</sup>, V. Vagnoni<sup>19</sup>, A. Valassi<sup>47</sup>, G. Valenti<sup>19</sup>,  
 M. van Beuzekom<sup>31</sup>, H. Van Hecke<sup>66</sup>, E. van Herwijnen<sup>47</sup>, C.B. Van Hulse<sup>17</sup>, M. van Veghel<sup>74</sup>,  
 R. Vazquez Gomez<sup>44</sup>, P. Vazquez Regueiro<sup>45</sup>, C. Vázquez Sierra<sup>31</sup>, S. Vecchi<sup>20</sup>, J.J. Velthuis<sup>53</sup>,  
 M. Veltri<sup>21,r</sup>, A. Venkateswaran<sup>67</sup>, M. Vernet<sup>9</sup>, M. Veronesi<sup>31</sup>, M. Vesterinen<sup>55</sup>,  
 J.V. Viana Barbosa<sup>47</sup>, D. Vieira<sup>5</sup>, M. Vieites Diaz<sup>48</sup>, H. Viemann<sup>73</sup>, X. Vilasis-Cardona<sup>44,m</sup>,  
 A. Vitkovskiy<sup>31</sup>, V. Volkov<sup>39</sup>, A. Vollhardt<sup>49</sup>, D. Vom Bruch<sup>12</sup>, A. Vorobyev<sup>37</sup>, V. Vorobyev<sup>42,x</sup>,  
 N. Voropaev<sup>37</sup>, R. Waldi<sup>73</sup>, J. Walsh<sup>28</sup>, J. Wang<sup>3</sup>, J. Wang<sup>71</sup>, J. Wang<sup>6</sup>, M. Wang<sup>3</sup>, Y. Wang<sup>7</sup>,  
 Z. Wang<sup>49</sup>, D.R. Ward<sup>54</sup>, H.M. Wark<sup>59</sup>, N.K. Watson<sup>52</sup>, D. Websdale<sup>60</sup>, A. Weiden<sup>49</sup>,  
 C. Weisser<sup>63</sup>, B.D.C. Westhenry<sup>53</sup>, D.J. White<sup>61</sup>, M. Whitehead<sup>13</sup>, D. Wiedner<sup>14</sup>,  
 G. Wilkinson<sup>62</sup>, M. Wilkinson<sup>67</sup>, I. Williams<sup>54</sup>, M. Williams<sup>63</sup>, M.R.J. Williams<sup>61</sup>,  
 T. Williams<sup>52</sup>, F.F. Wilson<sup>56</sup>, M. Winn<sup>11</sup>, W. Wislicki<sup>35</sup>, M. Witek<sup>33</sup>, G. Wormser<sup>11</sup>,  
 S.A. Wotton<sup>54</sup>, H. Wu<sup>67</sup>, K. Wyllie<sup>47</sup>, Z. Xiang<sup>5</sup>, D. Xiao<sup>7</sup>, Y. Xie<sup>7</sup>, H. Xing<sup>70</sup>, A. Xu<sup>3</sup>, L. Xu<sup>3</sup>,  
 M. Xu<sup>7</sup>, Q. Xu<sup>5</sup>, Z. Xu<sup>8</sup>, Z. Xu<sup>3</sup>, Z. Yang<sup>3</sup>, Z. Yang<sup>65</sup>, Y. Yao<sup>67</sup>, L.E. Yeomans<sup>59</sup>, H. Yin<sup>7</sup>,  
 J. Yu<sup>7,aa</sup>, X. Yuan<sup>67</sup>, O. Yushchenko<sup>43</sup>, K.A. Zarebski<sup>52</sup>, M. Zavertyaev<sup>15,c</sup>, M. Zdybal<sup>33</sup>,



M. Zeng<sup>3</sup>, D. Zhang<sup>7</sup>, L. Zhang<sup>3</sup>, S. Zhang<sup>3</sup>, W.C. Zhang<sup>3,z</sup>, Y. Zhang<sup>47</sup>, A. Zhelezov<sup>16</sup>,  
Y. Zheng<sup>5</sup>, X. Zhou<sup>5</sup>, Y. Zhou<sup>5</sup>, X. Zhu<sup>3</sup>, V. Zhukov<sup>13,39</sup>, J.B. Zonneveld<sup>57</sup>, S. Zucchelli<sup>19,e</sup>.

<sup>1</sup>*Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil*

<sup>2</sup>*Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil*

<sup>3</sup>*Center for High Energy Physics, Tsinghua University, Beijing, China*

<sup>4</sup>*School of Physics State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China*

<sup>5</sup>*University of Chinese Academy of Sciences, Beijing, China*

<sup>6</sup>*Institute Of High Energy Physics (IHEP), Beijing, China*

<sup>7</sup>*Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China*

<sup>8</sup>*Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, IN2P3-LAPP, Annecy, France*

<sup>9</sup>*Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France*

<sup>10</sup>*Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France*

<sup>11</sup>*LAL, Univ. Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, Orsay, France*

<sup>12</sup>*LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France*

<sup>13</sup>*I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany*

<sup>14</sup>*Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany*

<sup>15</sup>*Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany*

<sup>16</sup>*Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany*

<sup>17</sup>*School of Physics, University College Dublin, Dublin, Ireland*

<sup>18</sup>*INFN Sezione di Bari, Bari, Italy*

<sup>19</sup>*INFN Sezione di Bologna, Bologna, Italy*

<sup>20</sup>*INFN Sezione di Ferrara, Ferrara, Italy*

<sup>21</sup>*INFN Sezione di Firenze, Firenze, Italy*

<sup>22</sup>*INFN Laboratori Nazionali di Frascati, Frascati, Italy*

<sup>23</sup>*INFN Sezione di Genova, Genova, Italy*

<sup>24</sup>*INFN Sezione di Milano-Bicocca, Milano, Italy*

<sup>25</sup>*INFN Sezione di Milano, Milano, Italy*

<sup>26</sup>*INFN Sezione di Cagliari, Monserrato, Italy*

<sup>27</sup>*INFN Sezione di Padova, Padova, Italy*

<sup>28</sup>*INFN Sezione di Pisa, Pisa, Italy*

<sup>29</sup>*INFN Sezione di Roma Tor Vergata, Roma, Italy*

<sup>30</sup>*INFN Sezione di Roma La Sapienza, Roma, Italy*

<sup>31</sup>*Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands*

<sup>32</sup>*Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, Netherlands*

<sup>33</sup>*Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*

<sup>34</sup>*AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland*

<sup>35</sup>*National Center for Nuclear Research (NCBJ), Warsaw, Poland*

<sup>36</sup>*Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*

<sup>37</sup>*Petersburg Nuclear Physics Institute NRC Kurchatov Institute (PNPI NRC KI), Gatchina, Russia*

<sup>38</sup>*Institute of Theoretical and Experimental Physics NRC Kurchatov Institute (ITEP NRC KI), Moscow, Russia, Moscow, Russia*

<sup>39</sup>*Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia*

<sup>40</sup>*Institute for Nuclear Research of the Russian Academy of Sciences (INR RAS), Moscow, Russia*

<sup>41</sup>*Yandex School of Data Analysis, Moscow, Russia*

<sup>42</sup>*Budker Institute of Nuclear Physics (SB RAS), Novosibirsk, Russia*

<sup>43</sup>*Institute for High Energy Physics NRC Kurchatov Institute (IHEP NRC KI), Protvino, Russia, Protvino, Russia*

<sup>44</sup>*ICCUB, Universitat de Barcelona, Barcelona, Spain*

<sup>45</sup>*Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain*

<sup>46</sup>*Instituto de Física Corpuscular, Centro Mixto Universidad de Valencia - CSIC, Valencia, Spain*

<sup>47</sup>*European Organization for Nuclear Research (CERN), Geneva, Switzerland*

<sup>48</sup>*Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*

- <sup>49</sup> *Physik-Institut, Universität Zürich, Zürich, Switzerland*
- <sup>50</sup> *NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- <sup>51</sup> *Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- <sup>52</sup> *University of Birmingham, Birmingham, United Kingdom*
- <sup>53</sup> *H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- <sup>54</sup> *Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- <sup>55</sup> *Department of Physics, University of Warwick, Coventry, United Kingdom*
- <sup>56</sup> *STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- <sup>57</sup> *School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- <sup>58</sup> *School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- <sup>59</sup> *Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- <sup>60</sup> *Imperial College London, London, United Kingdom*
- <sup>61</sup> *Department of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- <sup>62</sup> *Department of Physics, University of Oxford, Oxford, United Kingdom*
- <sup>63</sup> *Massachusetts Institute of Technology, Cambridge, MA, United States*
- <sup>64</sup> *University of Cincinnati, Cincinnati, OH, United States*
- <sup>65</sup> *University of Maryland, College Park, MD, United States*
- <sup>66</sup> *Los Alamos National Laboratory (LANL), Los Alamos, United States*
- <sup>67</sup> *Syracuse University, Syracuse, NY, United States*
- <sup>68</sup> *Laboratory of Mathematical and Subatomic Physics , Constantine, Algeria, associated to <sup>2</sup>*
- <sup>69</sup> *Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to <sup>2</sup>*
- <sup>70</sup> *South China Normal University, Guangzhou, China, associated to <sup>3</sup>*
- <sup>71</sup> *School of Physics and Technology, Wuhan University, Wuhan, China, associated to <sup>3</sup>*
- <sup>72</sup> *Departamento de Física , Universidad Nacional de Colombia, Bogota, Colombia, associated to <sup>12</sup>*
- <sup>73</sup> *Institut für Physik, Universität Rostock, Rostock, Germany, associated to <sup>16</sup>*
- <sup>74</sup> *Van Swinderen Institute, University of Groningen, Groningen, Netherlands, associated to <sup>31</sup>*
- <sup>75</sup> *National Research Centre Kurchatov Institute, Moscow, Russia, associated to <sup>38</sup>*
- <sup>76</sup> *National University of Science and Technology “MISIS”, Moscow, Russia, associated to <sup>38</sup>*
- <sup>77</sup> *National Research University Higher School of Economics, Moscow, Russia, associated to <sup>41</sup>*
- <sup>78</sup> *National Research Tomsk Polytechnic University, Tomsk, Russia, associated to <sup>38</sup>*
- <sup>79</sup> *University of Michigan, Ann Arbor, United States, associated to <sup>67</sup>*

<sup>a</sup> *Universidade Federal do Triângulo Mineiro (UFMT), Uberaba-MG, Brazil*

<sup>b</sup> *Laboratoire Leprince-Ringuet, Palaiseau, France*

<sup>c</sup> *P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia*

<sup>d</sup> *Università di Bari, Bari, Italy*

<sup>e</sup> *Università di Bologna, Bologna, Italy*

<sup>f</sup> *Università di Cagliari, Cagliari, Italy*

<sup>g</sup> *Università di Ferrara, Ferrara, Italy*

<sup>h</sup> *Università di Genova, Genova, Italy*

<sup>i</sup> *Università di Milano Bicocca, Milano, Italy*

<sup>j</sup> *Università di Roma Tor Vergata, Roma, Italy*

<sup>k</sup> *Università di Roma La Sapienza, Roma, Italy*

<sup>l</sup> *AGH - University of Science and Technology, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland*

<sup>m</sup> *LIFAELS, La Salle, Universitat Ramon Llull, Barcelona, Spain*

<sup>n</sup> *Hanoi University of Science, Hanoi, Vietnam*

<sup>o</sup> *Università di Padova, Padova, Italy*

<sup>p</sup> *Università di Pisa, Pisa, Italy*

<sup>q</sup> *Università degli Studi di Milano, Milano, Italy*

<sup>r</sup> *Università di Urbino, Urbino, Italy*

<sup>s</sup> *Università della Basilicata, Potenza, Italy*

<sup>t</sup> *Scuola Normale Superiore, Pisa, Italy*

<sup>u</sup> *Università di Modena e Reggio Emilia, Modena, Italy*

<sup>v</sup> *Università di Siena, Siena, Italy*

<sup>w</sup> *MSU - Iligan Institute of Technology (MSU-IIT), Iligan, Philippines*

<sup>x</sup> *Novosibirsk State University, Novosibirsk, Russia*

<sup>y</sup>*INFN Sezione di Trieste, Trieste, Italy*

<sup>z</sup>*School of Physics and Information Technology, Shaanxi Normal University (SNNU), Xi'an, China*

<sup>aa</sup>*Physics and Micro Electronic College, Hunan University, Changsha City, China*

<sup>†</sup>*Deceased*