



The Compact Muon Solenoid Experiment
Conference Report

Mailing address: CMS CERN, CH-1211 GENEVA 23, Switzerland



20 April 2020 (v5, 26 June 2020)

Ultra-peripheral vector meson photoproduction in heavy-ion collisions in CMS

Marek Bohdan Walczak for the CMS Collaboration

Abstract

In this presentation an introduction to ultra-peripheral collisions (UPCs) is given. Recent results on UPCs obtained by the Compact Muon Solenoid collaboration are presented. They are also shown in conjunction with theoretical predictions. Finally, the analysis of ultra-peripheral Upsilon photoproduction in the new PbPb data collected at the end of 2018 by CMS is discussed. This analysis is still in progress.

Presented at *Epiphany XXVI Cracow Epiphany Conference on LHC Physics Standard Model and Beyond*

Ultra-peripheral vector meson photoproduction in heavy-ion collisions in CMS*

MAREK BOHDAN WALCZAK, ON BEHALF OF THE CMS COLLABORATION

University of Warsaw

In this document an introduction to ultra-peripheral collisions (UPCs) is given. The recent results on UPCs obtained by Compact Muon Solenoid collaboration are presented. They are also shown in conjunction with theoretical predictions. Finally, the analysis of ultra-peripheral Υ photoproduction in the new PbPb data collected at the end of 2018 by the CMS is discussed. This analysis is still in progress.

PACS numbers: ?

1. Introduction

Hadronic collisions at the Large Hadron Collider (LHC) are a very powerful source of γ -hadron and γ - γ interactions. Protons and ions possess an electric charge and, when colliding in the LHC, are surrounded by high energy, quasi-real photons, which can interact with another photon or with a parton inside the second hadron in a process called photoproduction. This process can be observed in a clean way during ultra-peripheral collisions (UPC), where the impact parameter b for the two hadrons is larger than the sum of their radii, $R_A + R_B$ (see Figure 1, panel (a)). A variety of final state particles can be produced in these processes. For example ultra-peripheral exclusive vector meson (VM) photoproduction has received great theoretical interest ([1], [2]). In this process both hadrons (h) stay intact after the collision, and there is a VM in the final state, $\gamma h \rightarrow \text{VM} + h$ (see Figure 1, panel (b)). The VM interacts with the hadron by the exchange of the pomeron carrying vacuum quantum numbers.

2. Exclusive $\rho(770)^0$ photoproduction in pPb data

During pPb collisions, the large electric charge of the lead nucleus enhances photon-proton interactions. Because these interactions are asym-

* Presented at the EIPHANY Conference, Cracow, Poland, 9 Jan 2020




Fig. 1: Panel (a) – the UPC: the impact parameter b for the two hadrons is larger than the sum of their radii $R_A + R_B$ / Panel (b) – Feynman diagram of the ultra-peripheral exclusive VM photoproduction. Both hadrons stay intact after the collision.

metrical it is possible to determine photon direction unambiguously – an advantage that is not present in case of symmetric PbPb or pp collisions. This feature enables determination of the centre-of-mass energy, $W_{\gamma p}$. In exclusive VM photoproduction, the dependence of the cross-section on the squared four-momentum transfer at the proton vertex (t) has been considered as a very promising way to find the onset of gluon saturation [6]. Furthermore, the Fourier transform of the t distribution is related to the two-dimensional, spatial distribution of the partons transverse to the beam direction.

In the presented analysis [4] the exclusive photoproduction of $\rho(770)^0$ meson in ultra-peripheral pPb collisions is measured in the $\pi^+\pi^-$ decay channel. The data have been collected by the Compact Muon Solenoid (CMS) detector in 2013. A full description of CMS and its subdetectors is given in ref. [3]. The center-of-mass energy per nucleon pair was $\sqrt{s_{NN}} = 5.02$ TeV, and the integrated luminosity of the analysed sample was $L = 7.4 \mu\text{b}^{-1}$ for pPb and $L = 9.6 \mu\text{b}^{-1}$ for Pbp, where pPb and Pbp mean opposite directions of beams in the LHC. The STARLIGHT Monte Carlo event generator [5] has been used for simulations of the exclusive resonant and nonresonant $\pi^+\pi^-$ production, exclusive $\rho(1700)$ events, acceptance and efficiency studies and for photon flux calculation.

The event selection consists of number a of requirements: events with




Fig. 2: Panel (a) shows the distribution of the $\pi^+\pi^-$ transverse momentum together with fits to signal and background components. Panel (b) shows the unfolded $\pi^+\pi^-$ invariant mass distribution in the $\pi^+\pi^-$ rapidity interval $|y^{\pi^+\pi^-}| < 2.0$ [4].




Fig. 3: Exclusive $\rho(770)^0$ photoproduction cross-section is shown as a function of $W_{\gamma p}$. The fit of the CMS and HERA data yields $\delta = 0.24 \pm 0.13(\text{stat}) \pm 0.04(\text{syst})$ [4].

an energy deposition in any tower in Hadronic Forward (HF) calorimeter below 3.0 GeV (consistent with only noise in HF) are selected to reject hadronic interactions. It is required that there are exactly two tracks with a pseudorapidity less than 2.0, and their transverse momenta greater than 0.4 and 0.2 GeV. Additional requirements to ensure exclusivity consist of a selection of events with energy deposited in CASTOR to be below 9 GeV

and for Zero Degree Calorimeter (ZDC) to be below 500 GeV on the positive side and below 2000 GeV on the negative side. The lack of symmetry for the ZDC requirement is due to the difference in radiation damage between them. For detailed selection requirements with numbers of events after each step see Table 1 in [4]. The final signal extraction is done in two steps, by fitting background distributions to $p_T^{\pi^+\pi^-}$ and $M^{\pi^+\pi^-}$ distributions as shown in Figure 2.




Fig. 4: Differential cross sections for exclusive $\rho(770)^0$ photoproduction as a function of $|t|$. The four panels present different corresponding center-of-mass energies, $W_{\gamma p}$, from the lowest value of 35.6 GeV on the top-left to the highest, 176 GeV on the bottom-right [6].

In Figure 3 the photon-proton cross section, $\sigma(\gamma p \rightarrow \rho(770)^0 p)$, for four bins in $W_{\gamma p}$ is shown. The total obtained cross-section for $29 < W_{\gamma p} < 213$ GeV is found to be $11.0 \pm 1.4(\text{stat}) \pm 1.0(\text{syst}) \mu\text{b}$. Figure 3 also shows results from fixed-target and HERA results (for details see [4]). The CMS results are consistent with those from the and ZEUS Collaborations at HERA, which shows that ion-proton collisions can be used similarly to electron-proton collisions.

Thanks to the large integrated luminosity, the energy dependence ($d\sigma/dt$) for the four bins in $W_{\gamma p}$ are provided for the first time. Figure 4 from a recent theoretical publication [6] compares the CMS results with different




Fig. 5: Panel (a) shows the EPPS16 nuclear gluon modifications for lead nucleus. The green dotted lines show all the contributions to the uncertainties. The largest uncertainty occurs at $x < 10^{-2}$ because of lack of data in this region [7]. Panel (b) shows the cross-section as a function of dimuon rapidity for coherent J/ψ photoproduction in ultra-peripheral PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, measured by CMS [8] and ALICE [9]. The data are consistent with the leading twist approximation calculations, which include nuclear gluon shadowing.

theoretical models. The location of the diffractive dips in exclusive UPC VMs in gamma-proton interactions is sensitive to the gluon saturation effect, which makes these measurements very promising. As can be seen, for the highest energy point, $W_{\gamma p} = 176$ GeV (lower-right panel in Figure 4) the data for low $|t|$ values are not well described by the bCGS model with only linear contribution. This can be a first indication that the model of the spatial distribution of gluons in protons will have to be reevaluated for the purpose of the study of the gluon saturation effects.

3. Exclusive Υ photoproduction in Run 2 PbPb data

The initial state of the nuclei is described by the Parton Distribution Functions (PDFs), describing the probability of finding a parton (gluons and quarks) with a longitudinal momentum fraction of the hadron x at the energy transfer to the hadron Q^2 . As can be seen in Figure 5, panel (a), the nuclear gluon PDFs for lead are poorly known, especially in the low x ($x <$




Fig. 6: The event displays show two of the candidate events for ultra-peripheral, exclusive Υ photoproduction. In an otherwise empty detector, there are two muons going into forward (panel (a)) and central (panel (b)) direction. The reconstructed invariant mass of the dimuon objects is consistent with the $\Upsilon(1S)$ mass. Events were recorded by CMS during the Run 2 PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV.

10^{-2}) region. It can be probed with exclusive J/ψ and Υ photoproduction measurements [1] [2]. CMS and ALICE have performed this analysis for the J/ψ meson at $\sqrt{s_{NN}} = 2.76$ TeV with PbPb data [8] [9] (see Figure 5, panel (b)). An analogous analysis with the Υ meson decaying to $\mu^+\mu^-$, at a higher center-of-mass energy of $\sqrt{s_{NN}} = 5.02$ TeV with 2018 Run 2 PbPb data is still in progress. The PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV took place during Run 2 in 2015 and 2018. In the later year CMS collected 3 times more data than in 2015, about 1.7 nb^{-1} . This allows to perform the measurement for $\Upsilon(1S)$, $\Upsilon(2S)$ and $\Upsilon(3S)$. The event selection consists of the requirement that there is no energy deposited in the forward calorimeters and that there are exactly two muons with kinematic requirements in an otherwise empty detector. Example event displays passing those requirements are shown in Figure 6.

4. Conclusions

The CMS detector, because of its good coverage in the forward region and excellent muon subdetectors, is perfect for UPC studies. The ultra-peripheral VM photoproduction process is a very useful tool for measuring the properties of protons and nuclei. The cross-sections obtained by CMS are in agreement with the power law dependence of $W_{\Upsilon\text{p}}$ observed at HERA, indicating that ion-proton interactions can supplement electron-proton data. UPC studies allow to set constraints on a variety of theoretical models, in particular the UPC photoproduction $\Upsilon(nS)$, which probes the kinematic range of $x < 10^{-2}$ with few experimental results. The ongoing

analysis of the Run 2 PbPb data could be the first measurement of this process. Future measurements with more data on exclusive VM photoproduction will be very useful to improve our understanding of the QCD dynamics at high energies.

Acknowledgement

This work was supported by National Science Centre (Poland) grant 2018/28/T/ST2/00199.

REFERENCES

- [1] A.J. Baltz, G. Baur, D. d'Enterria, L. Frankfurt, F. Gelis, *et al.*, “The Physics of Ultraperipheral Collisions at the LHC”, *Phys. Rept.*, 458:1–171, 2008
- [2] J. G. Contreras and J. D. Tapia Takaki, “Ultra-peripheral heavy-ion collisions at the LHC”, *Int. J. Mod. Phys. A* 30 1542012, 2015
- [3] CMS Collaboration, “The CMS experiment at the CERN LHC”, *JINST* 3 S08004, 2008
- [4] CMS Collaboration, “Exclusive $\rho(770)^0$ photoproduction in ultra-peripheral pPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV with the CMS experiment”, *Eur. Phys. J. C* 79 702, 2019
- [5] S. R. Klein, J. Nystrand, J. Seger, Y. Gorbunov, J. Butterworth, “STARlight: A Monte Carlo simulation program for ultra-peripheral collisions of relativistic ions”, *Comp. Phys. Comm.* 212-258, 2017
- [6] V. P. Gonçalves, F. S. Navarra, D. Spiering, “Exclusive ρ and J/ψ photoproduction in ultraperipheral pA collisions: Predictions of the gluon saturation models for the momentum transfer distributions”, *Phys. Lett. B* 791 299-304, 2019
- [7] K. J. Eskola, P. Paakkinen, H. Paukkunen, C. A. Salgado, “EPPS16: Nuclear parton distributions with LHC data”, *Eur. Phys. J. C* 77 163, 2017
- [8] CMS Collaboration, “Coherent J/ψ photoproduction in ultra-peripheral PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the CMS experiment”, *Phys. Lett. B* 772 489, 2017
- [9] ALICE Collaboration, “Coherent J/ψ photoproduction in ultra-peripheral Pb-Pb collisions at $\sqrt{s_{NN}}=2.76$ TeV”, *Phys. Lett.*, B718:1273–1283, 2013