



Higgs-mass prediction in the NMSSM with heavy BSM particles

Emanuele Bagnaschi^{1,a}, Mark Goodsell^{2,b}, Pietro Slavich^{2,c}

¹ CERN, Theory Division, 1211 Geneva 23, Switzerland

² Sorbonne Université, CNRS, Laboratoire de Physique Théorique et Hautes Energies, LPTHE, 75005 Paris, France

Received: 17 June 2022 / Accepted: 14 September 2022
© The Author(s) 2022

Abstract We address the prediction for the mass of the SM-like Higgs boson in NMSSM scenarios where all BSM particles, including the singlets, have masses at the TeV scale. We provide a full one-loop computation of the matching condition for the quartic Higgs coupling in the NMSSM, supplemented with the two-loop contributions that involve the strong gauge coupling. We discuss the impact of the one- and two-loop corrections that are specific to the NMSSM on the prediction for the Higgs mass, and propose a method to estimate of the uncertainty associated with the uncomputed higher-order terms. Finally, we illustrate how the measured value of the Higgs mass can be used to constrain some yet-unmeasured parameters of the NMSSM.

1 Introduction

The discovery of a Higgs boson with mass around 125 GeV and properties compatible with the predictions of the Standard Model (SM) [1–4], combined with the negative (so far) results of the searches for additional new particles at the LHC, point to scenarios with at least a mild hierarchy between the electroweak (EW) scale and the scale of beyond-the-SM (BSM) physics. In this case, the SM plays the role of an effective field theory (EFT) valid between the two scales. The requirement that a given BSM model include a state that can be identified with the observed Higgs boson can translate into important constraints on the model's parameter space.

One of the prime candidates for BSM physics is supersymmetry (SUSY), which predicts scalar partners for all SM fermions, as well as fermionic partners for all bosons. A remarkable feature of SUSY extensions of the SM is the requirement of an extended Higgs sector, with additional

neutral and charged bosons. In contrast to the case of the SM, the masses of the Higgs bosons are not free parameters, as SUSY requires all quartic scalar couplings to be related to the gauge and Yukawa couplings. Moreover, radiative corrections to the tree-level predictions for the quartic scalar couplings introduce a dependence on all of the SUSY-particle masses and couplings. In a hierarchical scenario such as the one described above, the lightest scalar of the SUSY model plays the role of the SM Higgs boson, and the prediction for its quartic self-coupling at the SUSY scale must coincide with the SM coupling extracted at the EW scale from the measured value of the Higgs mass and evolved up to the SUSY scale with appropriate renormalization group equations (RGEs). This “matching” condition can be used to constrain some yet-unmeasured parameters of the SUSY model, such as, e.g., the masses of the scalar partners of the top quarks, the stops.

In the next-to-minimal SUSY extension of the SM, or NMSSM, the Higgs sector includes two $SU(2)$ doublets H_1 and H_2 – as in the case of the minimal extension, the MSSM – plus a complex scalar S , singlet with respect to the SM gauge group.¹ The vacuum expectation value (vev) of the singlet, induced by the mass and interaction terms in the soft SUSY-breaking Lagrangian, generates a superpotential mass term for the doublets. This provides a solution to the so-called “ μ problem” of the MSSM, i.e., the question of why the supersymmetric Higgs-mass parameter μ should be at the same scale as the soft SUSY-breaking parameters. The doublet–singlet interactions of the NMSSM also induce new contributions to the prediction for the quartic self-coupling of the SM-like Higgs boson. Depending on the considered region of the NMSSM parameter space, these additional contributions can either increase or decrease the prediction for

^a e-mail: Emanuele.Bagnaschi@cern.ch

^b e-mail: goodsell@lpthe.jussieu.fr

^c e-mail: slavich@lpthe.jussieu.fr (corresponding author)

¹ For reviews of the MSSM and of the NMSSM, we point the reader to Ref. [5] and Refs. [6, 7], respectively.

the quartic Higgs coupling (and hence the Higgs mass) with respect to the case of the MSSM.

The fixed-order (FO) calculation of the NMSSM Higgs boson masses – in which the Higgs self-energies are computed up to a given order in the perturbative expansion, considering the full content of heavy and light fields of the theory – is quite advanced by now, albeit not yet at the level of the corresponding calculation in the MSSM.² After early studies of the NMSSM Higgs sector at the tree level [9, 10] and partial calculations of the dominant one-loop corrections [11–20], full calculations of the one-loop corrections – for increasingly general versions of the NMSSM and using a variety of renormalization schemes for the NMSSM parameters – were made available in Refs. [21–29]. At the two-loop level, the corrections involving the strong gauge coupling were computed in Refs. [21, 30–32], those involving only the top Yukawa coupling were computed in Ref. [33], and those involving also the remaining superpotential couplings of the NMSSM were computed in Refs. [31, 34–36]. It is worth pointing out that, in all of the two-loop calculations listed above, the two-loop part of the Higgs self-energies was computed under the approximation of vanishing external momentum and in the so-called “gaugeless limit” of vanishing EW gauge couplings. These full one-loop and partial two-loop calculations of the NMSSM Higgs masses were implemented in various public codes, such as `NMSSMTools` [37, 38], `SARAH/SPheno` [39–45], `NMSSMCalc` [46], `SOFTSUSY` [47, 48], `FlexibleSUSY` [49, 50] and `FeynHiggs` [51–53].³ Detailed comparisons among the predictions of these codes for the NMSSM Higgs masses were also presented in Refs. [54, 55].

Compared with the case of the FO calculation, much less attention has been devoted so far to the calculation of the NMSSM Higgs boson masses in the EFT approach, in which the heavy fields are “integrated out” of the theory at a scale comparable to their mass, leaving behind matching conditions for the couplings of the low-energy theory. This approach is the most appropriate to scenarios characterized by a hierarchy between the SUSY scale and the EW scale, where the FO calculation is plagued by large logarithms of the ratio of the two scales. In Ref. [56] a method was proposed to numerically obtain the boundary condition for the quartic Higgs coupling at the SUSY scale, by matching the FO calculation of the pole mass of the lightest Higgs boson in the NMSSM, as implemented in `FlexibleSUSY`, with the corresponding calculation in the SM. A similar method was later implemented in `SARAH/SPheno` in Ref. [57]. The “hybrid”

approach to the Higgs-mass calculation in Refs. [56, 57] accounts for terms suppressed by powers of v^2/M_S^2 – where v stands for the EW scale and M_S for the SUSY scale – that would be neglected in a pure EFT calculation. However, this approach required successive adjustments [50, 58] to avoid the introduction of spurious large-logarithmic effects at higher orders, and it does not provide explicit analytic results for the matching conditions. The latter would come in useful, e.g., to assess the relevance of the various contributions, to check and compare existing calculations, or as building blocks for further calculations.

A sensible approach to obtain explicit analytic results for the matching conditions consists in adapting to the NMSSM the formulas that were computed independently in Refs. [59, 60] for the one-loop matching of a general high-energy theory (without heavy gauge bosons) on a general renormalizable EFT. Indeed, in Ref. [60] analytic results for the one-loop matching condition for the quartic Higgs coupling were obtained in an extremely constrained NMSSM scenario in which all of the BSM-particle masses depend on just one parameter. Subsequently, the general formulas of Ref. [60] were employed in Ref. [61] to study a “Split-SUSY” scenario in which the NMSSM is matched to an EFT that, beyond the SM fields, includes also the complex singlet as well as the gauginos, the higgsinos and the singlino (i.e., the SUSY partners of the gauge bosons, the Higgs doublets and the singlet, respectively).

In this paper we aim to improve the accuracy of the EFT calculation of the SM-like Higgs mass in the NMSSM, and to illustrate how the measured value of the Higgs mass can constrain the parameter space of the model even in scenarios where all of the BSM particles are heavy. In Sect. 2 we adapt to the NMSSM the general formulas of Ref. [59], and obtain the full one-loop matching condition for the quartic Higgs coupling, for arbitrary values of all of the relevant parameters, in the EFT setup in which the NMSSM is matched directly to the SM. We compare our results with those of Ref. [60], in the constrained scenario considered in that paper, and find a discrepancy. We also compute directly the full two-loop-QCD contribution to the matching condition, i.e., the contribution of all two-loop terms that involve the strong gauge coupling. In Sect. 3 we discuss the effect of the corrections computed in Sect. 2 on the prediction for the mass of the SM-like Higgs boson, including an estimate of the uncertainty associated with uncomputed higher-order terms. We also discuss the constraints on the NMSSM parameters that arise from the requirement that the theory prediction for the Higgs mass correspond to the measured value. Finally, Sect. 4 contains our conclusions.

² For details on the Higgs-mass calculation in the MSSM, as well as for an overview of the different approaches to the Higgs-mass calculation in SUSY models, we point the reader to Ref. [8].

³ In fact, the version of `FeynHiggs` that implements the NMSSM calculations of Refs. [26–28] is not public yet.

2 Matching condition for the quartic Higgs coupling

In this section we describe our calculation of the full one-loop and two-loop-QCD contributions to the matching condition for the quartic Higgs coupling in the NMSSM. To fix our notation, which follows the SLHA2 conventions [62], we list here the terms in the superpotential W and in the soft SUSY-breaking Lagrangian $\mathcal{L}_{\text{soft}}$ that determine the Higgs potential:

$$W \supset -\lambda \hat{S} \hat{H}_1 \hat{H}_2 + \frac{\kappa}{3} \hat{S}^3, \tag{1}$$

$$-\mathcal{L}_{\text{soft}} \supset m_{H_1}^2 H_1^\dagger H_1 + m_{H_2}^2 H_2^\dagger H_2 + m_S^2 S^* S - \left(\lambda A_\lambda S H_1 H_2 - \frac{\kappa}{3} A_\kappa S^3 + \text{h.c.} \right), \tag{2}$$

where the hats denote superfields, and the $SU(2)$ indices of the Higgs doublets are contracted by the antisymmetric tensor ϵ_{ab} , with $\epsilon_{12} = 1$. For simplicity, we take the parameters in Eqs. (1) and (2) to be all real, and we enforce the Z_3 symmetry that forbids tadpole and mass terms in W and the corresponding SUSY-breaking terms in $\mathcal{L}_{\text{soft}}$. We assume that the singlet develops a vev $v_s \equiv \langle S \rangle$, providing effective μ and B_μ terms for the Higgs doublets. It is also convenient to rotate the two doublets to the so-called Higgs basis, in which they have the same hypercharge and only one of them develops a vev

$$\begin{pmatrix} H \\ A \end{pmatrix} = \begin{pmatrix} \cos \beta & \sin \beta \\ -\sin \beta & \cos \beta \end{pmatrix} \begin{pmatrix} -\epsilon H_1^* \\ H_2 \end{pmatrix}, \tag{3}$$

where the rotation angle is defined by $\tan \beta \equiv v_2/v_1$, with $v_i \equiv \langle H_i^0 \rangle$. In this basis the neutral component of H has the vev $v = (v_1^2 + v_2^2)^{1/2}$, while A has no vev.

We consider the hierarchical scenario in which all of the SUSY particles, as well as the Higgs doublet A and the scalar and pseudoscalar components of the singlet, are significantly heavier than the EW scale. We then adopt an EFT setup in which all of the heavy particles of the NMSSM are integrated out at a common renormalization scale $Q \approx M_S$, below which the field content of the theory is just the one of the SM, and in particular the Higgs doublet H plays the role of the SM Higgs. In the matching of the NMSSM to the SM we work in the limit of unbroken EW symmetry, $v \rightarrow 0$. This amounts to neglecting corrections suppressed by powers of v^2/M_S^2 , which can be mapped to the effect of non-renormalizable, higher-dimensional operators in the EFT Lagrangian. In this limit we can neglect the mixing among gauginos, higgsinos and singlino, as well as the mixing between the “left” and “right” sfermions (i.e., the SUSY partners of the left- and right-handed fermions of the SM).

2.1 Tree-level masses and couplings

We now discuss the tree-level masses and couplings in the Higgs/higgsino sector. As mentioned above, we work in the limit $v \rightarrow 0$, as appropriate to the calculation of the matching conditions in the EFT approach. The masses of the scalar and pseudoscalar components of the singlet, which we decompose as $S = v_s + (s + i a)/\sqrt{2}$, are

$$m_s^2 = \kappa v_s (A_\kappa + 4\kappa v_s), \quad m_a^2 = -3\kappa v_s A_\kappa, \tag{4}$$

where one of the minimum conditions of the tree-level scalar potential has been used to replace the soft SUSY-breaking mass for the singlet with a combination of the other parameters,

$$m_S^2 = -\kappa v_s (A_\kappa + 2\kappa v_s), \tag{5}$$

and the requirement that $\langle S \rangle = v_s$ be a deeper minimum than $\langle S \rangle = 0$ implies $3\kappa v_s/A_\kappa < -1$. The mass of the heavy Higgs doublet A is

$$m_A^2 = \frac{\lambda v_s (A_\lambda + \kappa v_s)}{\sin \beta \cos \beta}, \tag{6}$$

where again the minimum conditions have been used to remove the dependence on $m_{H_1}^2$ and $m_{H_2}^2$. Finally, the higgsinos \tilde{h}_1 and \tilde{h}_2 combine into a Dirac fermion with mass $\mu = \lambda v_s$, and the singlino acquires a mass $m_{\tilde{g}} = 2\kappa v_s$.

In the EFT approach, the calculation of the mass of the SM-like Higgs boson H – or, alternatively, the determination of the constraints on the NMSSM parameters that arise from the measured value of the Higgs mass – require the calculation of the NMSSM prediction for the quartic Higgs coupling at the matching scale.⁴ At the tree level, the latter reads

$$\lambda_{\text{SM}}^{\text{tree}} = \frac{1}{4}(g^2 + g'^2) \cos^2 2\beta + \frac{1}{2} \lambda^2 \sin^2 2\beta - \frac{a_{hhs}^2}{m_s^2}, \tag{7}$$

where the first term on the r.h.s. is the D-term contribution analogous to the one in the MSSM, the second term is the F-term contribution specific to the NMSSM, and the third term originates from the decoupling of the singlet scalar. The trilinear coupling a_{hhs} enters the NMSSM Lagrangian as $\mathcal{L} \supset -(a_{hhs}/2) h^2 s$, where h is the neutral scalar component of H , and at the tree level it reads

$$a_{hhs} = \frac{\lambda}{\sqrt{2}} [2\lambda v_s - (A_\lambda + 2\kappa v_s) \sin 2\beta]. \tag{8}$$

Combining Eqs. (4), (7) and (8), we remark that the contribution to $\lambda_{\text{SM}}^{\text{tree}}$ that arises from the decoupling of the singlet is always negative, and contains a piece that does not depend on $\tan \beta$.

⁴ We denote the quartic Higgs coupling as λ_{SM} to distinguish it from the doublet–singlet superpotential coupling λ . In our conventions the SM Lagrangian contains the quartic interaction term $-\frac{1}{2} \lambda_{\text{SM}} |H|^4$.

2.2 One-loop matching

We now describe our calculation of the full one-loop contribution to the matching condition for the quartic Higgs coupling. This contribution can be decomposed as:

$$\Delta\lambda_{SM}^{1\ell} = \Delta\lambda_{SM}^{1\ell,1PI} + 2\lambda_{SM}^{tree} \Delta Z_h^{1\ell} - 2\frac{a_{hhs}}{m_s^2} \Delta a_{hhs}^{1\ell} + \frac{a_{hhs}^2}{m_s^4} \Delta m_s^{2,1\ell} + \Delta\lambda_{SM}^{1\ell,RS}. \tag{9}$$

The first term on the r.h.s. of Eq. (9) originates from one-particle-irreducible (1PI) diagrams with heavy particles in the loop and four external Higgs legs. The second term arises from diagrams with a wave-function renormalization (WFR) insertion on one of the external legs, and it involves the derivative with respect to the external momentum of the heavy-particle (HP) contribution to the renormalized self-energy of the Higgs field:

$$\Delta Z_h^{1\ell} = - \left. \frac{d\hat{\Pi}_{hh}^{1\ell,HP}}{dp^2} \right|_{p^2=0}. \tag{10}$$

The third term on the r.h.s. of Eq. (9) arises from singlet-exchange diagrams with a HP loop insertion on one of the trilinear Higgs–singlet vertices, while the fourth term arises from singlet-exchange diagrams with a HP loop insertion on the singlet propagator. The exact form of the latter term depends on the choices made for the definitions of the singlet vev v_s and of the singlet mass m_s^2 entering the tree-level part of the matching condition for λ_{SM} , see Eq. (7). Assuming that v_s corresponds to the minimum of the loop-corrected scalar potential, and using the expression in Eq. (4) for the singlet mass, we get

$$\Delta m_s^{2,1\ell} = - \left. \hat{\Pi}_{ss}^{1\ell,HP} \right|_{p^2=0} + \frac{\hat{T}_s^{1\ell,HP}}{\sqrt{2} v_s}, \tag{11}$$

involving the HP contributions to the renormalized self-energy and tadpole of the singlet. We remark that the tadpole term in Eq. (11) originates from the fact that, with our choice for v_s , the minimum condition of the scalar potential used to remove the explicit dependence of the singlet mass on the soft SUSY-breaking parameter m_s^2 includes a loop correction,

$$m_s^2 = -\kappa v_s (A_\kappa + 2\kappa v_s) + \frac{\hat{T}_s^{1\ell,HP}}{\sqrt{2} v_s}. \tag{12}$$

By expressing the singlet mass as in Eq. (4), we are effectively moving the tadpole term from the singlet mass to the one-loop part of the matching condition for λ_{SM} . If we were instead to express the singlet mass entering the tree-level part of the matching condition as $m_s^2 = m_S^2 + 2\kappa v_s (A_\kappa + 3\kappa v_s)$, while still considering v_s as the minimum of the loop-corrected potential, there would be no tadpole term in Eq. (11). If, on the other hand, we were to consider v_s as the

minimum of the tree-level potential, there would still be no tadpole term in Eq. (11), and the two expressions for the singlet mass discussed above would be equivalent to each other, i.e. $m_s^2 = \kappa v_s (A_\kappa + 4\kappa v_s) = m_S^2 + 2\kappa v_s (A_\kappa + 3\kappa v_s)$. However, the one-loop contribution to the matching condition for λ_{SM} in Eq. (9) would receive direct contributions from non-1PI diagrams with tadpole insertions (see Ref. [63] for a related discussion)

$$\Delta\lambda_{SM}^{1\ell,tad} = \left[\sqrt{2}\kappa (A_\kappa + 6\kappa v_s) \frac{a_{hhs}^2}{m_s^6} - 2\lambda (\lambda - \kappa \sin 2\beta) \frac{a_{hhs}}{m_s^4} \right] \hat{T}_s^{1\ell,HP}, \tag{13}$$

where the first term within square brackets arises from a diagram with a tadpole insertion on the singlet propagator, and the second from diagrams with tadpole insertions on the singlet-doublet vertices.

Finally, the last term on the r.h.s. of Eq. (9) includes contributions arising from differences in the renormalization scheme (RS) used for the couplings of the NMSSM and for those of the EFT valid below the matching scale (i.e., the SM). In particular, the calculation of the matching condition for λ_{SM} is performed in the \overline{DR} scheme assuming the field content of the NMSSM, whereas in the EFT λ_{SM} is interpreted as an \overline{MS} parameter, and we also choose to interpret the EW gauge couplings entering Eq. (7) as \overline{MS} parameters of the SM. We remark, however, that the presence of the singlet superfield does not affect these contributions at one loop, thus they can be taken directly from the MSSM calculation of Refs. [64,65].

To obtain the quartic- and trilinear-vertex corrections and the self-energies entering the various contributions to Eq. (9) we use the formulas in appendix B of Ref. [59], which discusses the one-loop matching of a general high-energy theory (without heavy gauge bosons) on a general renormalizable EFT. To obtain the singlet tadpole we use analogous formulas from Ref. [66]. This saves us the trouble of actually calculating one-loop Feynman diagrams, but requires that we adapt to the case of the NMSSM the notation of Refs. [59,66] for masses and interactions of scalars and fermions in a general renormalizable theory.⁵ We find that, once the higgsino mass and the Higgs–sfermion interaction parameters are expressed in terms of $\mu = \lambda v_s$, the threshold correction $\Delta\lambda_{SM}^{1\ell}$ splits neatly into a part that does not depend explicitly on λ and coincides with the corresponding correction in the MSSM, see Refs. [64,65], plus an NMSSM-specific, λ -dependent part that vanishes in the limit $\lambda \rightarrow 0$.

⁵ Note that our conventions for the signs of tadpoles and self-energies are opposite to those in Refs. [59,66]. Also, Refs. [59,66] define $v_s \equiv \sqrt{2} \langle S \rangle$, whereas we define $v_s \equiv \langle S \rangle$ as mentioned after Eq. (2).

Our full formulas for $\Delta\lambda_{SM}^{1\ell}$, valid for generic values of all of the relevant NMSSM parameters, are lengthy and not particularly illuminating, therefore we make them available on request in electronic form. For later convenience, we provide here the contribution to the NMSSM-specific part of $\Delta\lambda_{SM}^{1\ell}$ from diagrams involving stop squarks. The sfermion contribution to $\Delta\lambda_{SM}^{1\ell, \text{1PI}}$ is the same as in the MSSM once we set $\mu = \lambda v_s$, and in the limit $v \rightarrow 0$ there are no one-loop sfermion contributions to the self-energy and tadpole of the singlet. The NMSSM-specific stop contribution to the one-loop matching condition for λ_{SM} thus reduces to

$$\left(\Delta\lambda_{SM}^{1\ell, \tilde{t}}\right)_\lambda = 2(\lambda_{SM}^{\text{tree}})_\lambda \Delta Z_h^{1\ell, \tilde{t}} - 2 \frac{a_{hhs}}{m_s^2} \Delta a_{hhs}^{1\ell, \tilde{t}}. \tag{14}$$

In the first term on the r.h.s. of Eq. (14) above, $(\lambda_{SM}^{\text{tree}})_\lambda = \lambda^2 \sin^2 2\beta/2 - a_{hhs}^2/m_s^2$ is the λ -dependent part of the tree-level matching condition for λ_{SM} , see Eq. (7), and $\Delta Z_h^{1\ell, \tilde{t}}$ is the stop contribution to the Higgs WFR:

$$\Delta Z_h^{1\ell, \tilde{t}} = -\frac{g_t^2 N_c}{(4\pi)^2} \frac{X_t^2}{6 m_{Q_3} m_{U_3}} \tilde{F}_5\left(\frac{m_{Q_3}}{m_{U_3}}\right), \tag{15}$$

where $N_c = 3$ is a color factor, $X_t = A_t - \mu \cot \beta$ is the trilinear Higgs-stop interaction parameter (with A_t being the corresponding soft SUSY-breaking coupling), m_{Q_3} and m_{U_3} are the soft SUSY-breaking masses for the left and right stop, respectively, and the loop function $\tilde{F}_5(x)$ is defined in the appendix A of Ref. [64]. In the second term on the r.h.s. of Eq. (14) above, the tree-level quantities m_s^2 and a_{hhs} are given in Eqs. (4) and (8), respectively, while $\Delta a_{hhs}^{1\ell, \tilde{t}}$ is the one-loop stop contribution to the Higgs–singlet coupling in the limit $v \rightarrow 0$,

$$\Delta a_{hhs}^{1\ell, \tilde{t}} = \sqrt{2} N_c \frac{\lambda g_t^2}{(4\pi)^2} \frac{X_t \cot \beta}{m_{Q_3}^2 - m_{U_3}^2} \left[m_{Q_3}^2 \left(1 - \ln \frac{m_{Q_3}^2}{Q^2}\right) - m_{U_3}^2 \left(1 - \ln \frac{m_{U_3}^2}{Q^2}\right) \right]. \tag{16}$$

The NMSSM-specific sbottom contribution $\left(\Delta\lambda_{SM}^{1\ell, \tilde{b}}\right)_\lambda$ can be obtained from Eqs. (14)–(16) with the replacements $g_t \rightarrow g_b$, $X_t \rightarrow X_b$, $\cot \beta \rightarrow \tan \beta$ and $m_{U_3} \rightarrow m_{D_3}$, where g_b is the bottom Yukawa coupling, $X_b = A_b - \mu \tan \beta$, and m_{D_3} is the soft SUSY-breaking mass for the right sbottom. Analogous replacements (with $N_c = 1$) yield also the stau contribution. For simplicity, we set the tiny Yukawa couplings of the first two generations to zero, hence there are no NMSSM-specific contributions to $\Delta\lambda_{SM}^{1\ell}$ from the corresponding sfermions.

As a non-trivial check of our full one-loop calculation, we verified that by taking the derivative of the matching condition for λ_{SM} with respect to $\ln Q^2$ we recover the corresponding one-loop RGE of the SM, i.e.,

$$(4\pi)^2 \frac{d}{d \ln Q^2} \left(\lambda_{SM}^{\text{tree}} + \Delta\lambda_{SM}^{1\ell}\right) = \lambda_{SM}^{\text{tree}} \left(6\lambda_{SM}^{\text{tree}} + 6g_t^2 + 6g_b^2 + 2g_\tau^2 - \frac{9}{2}g^2 - \frac{3}{2}g'^2\right) - 6g_t^4 - 6g_b^4 - 6g_\tau^4 + \frac{9}{8}g^4 + \frac{3}{8}g'^4 + \frac{3}{4}g^2g'^2. \tag{17}$$

To this effect, we must combine the explicit scale dependence of our result for $\Delta\lambda_{SM}^{1\ell}$ with the implicit scale dependence of all of the parameters entering $\lambda_{SM}^{\text{tree}}$. For the latter we use the RGEs of the NMSSM as listed in Ref. [7], with the exception of the EW gauge couplings for which our definitions require that we use the RGEs of the SM.

As mentioned in Sect. 1, the authors of Ref. [60] obtained the matching condition for λ_{SM} in an extremely constrained NMSSM scenario in which all of the masses and couplings in the singlet/singlino sector depend only on λ , $\tan \beta$ and a lone mass parameter m_0 . After adapting our formulas for $\Delta\lambda_{SM}^{1\ell}$ to this constrained scenario, we compared them with those of Ref. [60],⁶ but we found a discrepancy. Indeed, it appears that in Ref. [60] the one-loop matching condition for λ_{SM} misses a tadpole contribution analogous to the one in our Eq. (11). This is inconsistent with the fact that the authors of Ref. [60] appear to have defined the singlet mass entering $\lambda_{SM}^{\text{tree}}$ as in our Eq. (4) – thus obtaining $m_s^2 = 2/3 m_0^2$ in the constrained scenario – and v_s as the vev of the loop-corrected potential, as can be inferred from the absence of direct contributions from tadpole-insertion diagrams in their results.

2.3 Two-loop-QCD matching

As mentioned earlier, the one-loop squark contribution to the matching condition for λ_{SM} splits into a λ -independent part that coincides with the analogous MSSM result and a λ -dependent part that is specific to the NMSSM. This structure allows for a relatively economical calculation of the two-loop contribution that involves the strong gauge coupling. Indeed, once we identify $\mu = \lambda v_s$, the contribution from two-loop 1PI diagrams that involve the strong interactions of the squarks is the same as in the MSSM, and has already been computed in Refs. [64,67]. Furthermore, the fact that there are no squark contributions to the singlet self-energy and tadpole at one loop implies that there are no two-loop contributions involving the strong interactions either. In analogy with Eqs. (9) and (14), the NMSSM-specific, λ -dependent two-loop-QCD contribution to the matching condition for λ_{SM} can thus be decomposed as

⁶ We obtained the formulas of Ref. [60] for $\Delta\lambda_{SM}^{1\ell}$ in the constrained NMSSM scenario from their implementation in SARAH/SPheno. We also remark that we cannot reproduce Fig. 12 of Ref. [60] beyond the qualitative level.

$$\begin{aligned}
 \left(\Delta\lambda_{\text{SM}}^{2\ell,\text{QCD}}\right)_\lambda &= 2 \left(\lambda_{\text{SM}}^{\text{tree}}\right)_\lambda \Delta Z_h^{2\ell,\text{QCD}} \\
 &\quad - 2 \frac{a_{hhs}}{m_s^2} \Delta a_{hhs}^{2\ell,\text{QCD}} + \left(\Delta\lambda_{\text{SM}}^{2\ell,\text{RS}}\right)_\lambda.
 \end{aligned}
 \tag{18}$$

The quantity denoted as $\Delta Z_h^{2\ell,\text{QCD}}$ in the first term on the r.h.s. of Eq. (18) above is the two-loop contribution to the Higgs WFR from diagrams that involve the strong interactions of the squarks. Again, once we identify $\mu = \lambda v_s$, this contribution is the same as in the MSSM and has already been computed in Ref. [67]. Explicit formulas for $\Delta Z_h^{2\ell,\text{QCD}}$ in the limit of degenerate squark and gluino masses can be gleaned from Eqs. (13)–(17) in Sect. 2.2 of that paper, and the formulas with full dependence on all of the relevant parameters are available on request in electronic form. It is worth noting that, in the EFT calculation of the SM-like Higgs mass, the WFR contribution in Eq. (18) accounts for effects that, in the corresponding FO calculation, arise from the external-momentum dependence of the two-loop self-energies; in particular, it is related to the first-order term in the p^2 -expansion of $\hat{\Pi}_{hh}^{2\ell,\tilde{q}}$, the contribution to the renormalized Higgs-boson self-energy from two-loop diagrams involving the strong interactions of the squarks. Higher-order terms in the p^2 -expansion of $\hat{\Pi}_{hh}^{2\ell,\tilde{q}}$ are suppressed by powers of v^2/M_S^2 and can be neglected in our hierarchic scenario, while the momentum dependence of the two-loop quark–gluon contribution to the self-energy is fully accounted for in the calculation of the relation between λ_{SM} and M_h at the EW scale, see Sect. 3. Hence, the EFT approach allows for a straightforward inclusion of external-momentum effects that are of the same order in the relevant couplings as the other two-loop contributions computed in this section. In contrast, these effects are missed by the FO calculations of the NMSSM Higgs masses in Refs. [21, 30–32], where the two-loop part of the self-energies is computed – whether “diagrammatically” or in the effective-potential approach – under the approximation of vanishing external momentum.

The quantity denoted as $\Delta a_{hhs}^{2\ell,\text{QCD}}$ in the second term on the r.h.s. of Eq. (18) is the contribution to the trilinear Higgs–singlet coupling from two-loop, 1PI diagrams that involve the strong interactions of the squarks. This contribution is not available in the literature, but can be straightforwardly computed with the same effective-potential techniques adopted in Refs. [64, 67] for the calculation of the two-loop-QCD contributions to the matching condition for λ_{SM} . In particular, we can write

$$\Delta a_{hhs}^{2\ell,\text{QCD}} = \left. \frac{\partial^3 \Delta V^{2\ell,\tilde{q}}}{\partial^2 h \partial s} \right|_{v=0}, \tag{19}$$

where $\Delta V^{2\ell,\tilde{q}}$ is the contribution to the NMSSM scalar potential from two-loop diagrams involving the strong interactions of the stop and sbottom squarks (the first two gen-

erations of squarks do not contribute in the limit in which we neglect the corresponding Yukawa couplings). The stop contribution to $\Delta V^{2\ell,\tilde{q}}$ can be found, e.g., in Eq. (28) of Ref. [64], and the sbottom contribution can be obtained from the stop one with trivial replacements. The squark masses and mixing angles in $\Delta V^{2\ell,\tilde{q}}$ are then expressed as function of field-dependent quark masses, and Eq. (19) becomes

$$\begin{aligned}
 \Delta a_{hhs}^{2\ell,\text{QCD}} &= -\frac{\lambda g_t^2}{\sqrt{2}} \cot \beta \frac{d}{dX_t} \left[\left(\mathcal{D}_1^{tb} + m_t^2 \mathcal{D}_2^t \right) \Delta V^{2\ell,\tilde{t}} \right]_{m_t \rightarrow 0} \\
 &\quad - \frac{\lambda g_b^2}{\sqrt{2}} \tan \beta \frac{d}{dX_b} \left[\left(\mathcal{D}_1^b + m_b^2 \mathcal{D}_2^b \right) \Delta V^{2\ell,\tilde{b}} \right]_{m_b \rightarrow 0},
 \end{aligned}
 \tag{20}$$

where we define the operators

$$\mathcal{D}_i^q \equiv \left(\frac{d}{dm_q^2} \right)^i. \tag{21}$$

We use relations such as those in Eq. (32) of Ref. [64] for the derivatives of the field-dependent parameters with respect to the quark masses, then we obtain the limits of vanishing quark masses as described in that paper (note that in this calculation we do not encounter terms that diverge for $m_q \rightarrow 0$). Finally, the derivatives with respect to the parameters X_t and X_b in Eq. (20) account for the derivative with respect to the singlet field in Eq. (19). In units of $g_s^2 C_F N_c / (4\pi)^4$, where g_s is the strong gauge coupling and $C_F = 4/3$ and $N_c = 3$ are color factors, we obtain for the stop contribution

$$\begin{aligned}
 \left(\Delta a_{hhs}^{2\ell,\text{QCD}}\right)^{\tilde{t}} &= -2\sqrt{2} \lambda g_t^2 \cot \beta \left\{ X_t \left[-2 \right. \right. \\
 &\quad + \left(2 - \frac{2(1+x_Q) \ln x_Q}{x_Q - x_U} \right) \ln \frac{m_{\tilde{g}}^2}{Q^2} - \frac{1}{2} \ln^2 \frac{m_{\tilde{g}}^2}{Q^2} \\
 &\quad + \frac{2(1+2x_Q) \ln x_Q}{x_Q - x_U} - \frac{(x_Q^2 + x_Q - x_U) \ln^2 x_Q}{(x_Q - x_U)^2} \\
 &\quad + \frac{x_Q x_U \ln x_Q \ln x_U}{(x_Q - x_U)^2} - \frac{2(1-x_Q)}{x_Q - x_U} \text{Li}_2 \left(1 - \frac{1}{x_Q} \right) \left. \right] \\
 &\quad + m_{\tilde{g}} \left[\frac{5}{2} - \left(2 - \frac{2x_Q \ln x_Q}{x_Q - x_U} \right) \ln \frac{m_{\tilde{g}}^2}{Q^2} + \frac{1}{2} \ln^2 \frac{m_{\tilde{g}}^2}{Q^2} \right. \\
 &\quad - \frac{4x_Q \ln x_Q}{x_Q - x_U} \\
 &\quad - \left. \frac{(1-x_Q) \ln^2 x_Q}{x_Q - x_U} - \frac{2(1-x_Q)}{x_Q - x_U} \text{Li}_2 \left(1 - \frac{1}{x_Q} \right) \right] \\
 &\quad \left. + (x_Q \longleftrightarrow x_U) \right\},
 \end{aligned}
 \tag{22}$$

where $m_{\tilde{g}}$ is the gluino mass, and we defined the ratios $x_Q = m_{Q_3}^2/m_{\tilde{g}}^2$ and $x_U = m_{U_3}^2/m_{\tilde{g}}^2$. Again, the sbottom contribution can be obtained from the stop contribution

in Eq. (22) with the replacements $g_t \rightarrow g_b$, $X_t \rightarrow X_b$, $\cot \beta \rightarrow \tan \beta$ and $m_{U_3} \rightarrow m_{D_3}$.

The third term on the r.h.s. of Eq. (18) contains in fact two separate contributions that arise from choices of renormalization scheme for the couplings involved in the matching condition:

$$(\Delta\lambda_{SM}^{2\ell,RS})_\lambda = \frac{g_s^2 C_F N_c}{(4\pi)^4} (g_t^2 + g_b^2) (\lambda_{SM}^{tree})_\lambda + 2 \Delta g_t (\Delta\lambda_{SM}^{1\ell,\bar{t}})_\lambda. \tag{23}$$

The first term on the r.h.s. of Eq. (23) above is the NMSSM-specific completion of the contribution in Eq. (21) of Ref. [67], and stems from the fact that SUSY provides a prediction for the \overline{DR} -renormalized quartic Higgs coupling, whereas we interpret the parameter λ_{SM} in the EFT as \overline{MS} . The second term stems from the fact that we choose to express the one-loop part of the matching condition for λ_{SM} in terms of the \overline{MS} -renormalized top Yukawa coupling of the SM rather than the corresponding \overline{DR} -renormalized coupling of the NMSSM.⁷ The quantity $(\Delta\lambda_{SM}^{1\ell,\bar{t}})_\lambda$ is given in Eqs. (14)–(16), while Δg_t denotes the one-loop, $\mathcal{O}(g_s^2)$ part of the difference between the \overline{MS} coupling of the SM and the \overline{DR} coupling of the NMSSM,

$$\Delta g_t = -\frac{g_s^2 C_F}{(4\pi)^2} \left[1 + \ln \frac{m_{\tilde{g}}^2}{Q^2} + \tilde{F}_6 \left(\frac{m_{Q_3}}{m_{\tilde{g}}} \right) + \tilde{F}_6 \left(\frac{m_{U_3}}{m_{\tilde{g}}} \right) - \frac{X_t}{m_{\tilde{g}}} \tilde{F}_9 \left(\frac{m_{Q_3}}{m_{\tilde{g}}}, \frac{m_{U_3}}{m_{\tilde{g}}} \right) \right], \tag{24}$$

where the loop functions $\tilde{F}_6(x)$ and $\tilde{F}_9(x, y)$ are defined in the appendix A of Ref. [64].

Finally, we verified that, by taking the derivative with respect to $\ln Q^2$ of the NMSSM-specific part of the two-loop-QCD matching condition for λ_{SM} , we do recover the terms involving g_s^2 that we should expect from the RGE of the quartic Higgs coupling of the SM:

$$(4\pi)^4 \frac{d}{d \ln Q^2} \left[(\lambda_{SM}^{tree})_\lambda + (\Delta\lambda_{SM}^{1\ell,\bar{q}})_\lambda + (\Delta\lambda_{SM}^{2\ell,QCD})_\lambda \right] \supset 40 g_s^2 (g_t^2 + g_b^2) (\lambda_{SM}^{tree})_\lambda. \tag{25}$$

To this effect, we must combine the explicit scale dependence of our result for $(\Delta\lambda_{SM}^{2\ell,QCD})_\lambda$ with the implicit scale dependence of the parameters in $(\lambda_{SM}^{tree})_\lambda$ whose RGEs have QCD contributions at two loops (namely λ , A_λ and $\tan \beta$), and of the parameters in the one-loop stop and sbottom contributions $(\Delta\lambda_{SM}^{1\ell,\bar{q}})_\lambda$ whose RGEs have QCD contributions at

one loop (namely the soft SUSY-breaking masses and trilinear couplings of the squarks, and the top and bottom Yukawa couplings).

3 Higgs-mass prediction and constraints on the NMSSM parameters

In this section we analyze the impact of the one- and two-loop corrections to the quartic Higgs coupling computed in Sects. 2.2 and 2.3, respectively, on the prediction for the mass of the lightest Higgs scalar of the NMSSM, focusing on the scenario in which all of the SUSY particles, the heavier $SU(2)$ doublet, and the scalar and pseudoscalar components of the singlet are all significantly heavier than the EW scale. We also propose a method to estimate the so-called ‘‘theory uncertainty’’ of our calculation, i.e., the uncertainty associated with uncomputed higher-order corrections. Finally, we discuss how the prediction for the mass of the SM-like Higgs boson can constrain the parameter space of the NMSSM even in scenarios in which all of the BSM particles are heavy.

Our numerical calculations rely on an EFT setup analogous to the one developed in Ref. [67]. We use the public code `mR` [68] to extract – at full two-loop accuracy – the \overline{MS} -renormalized parameters of the SM Lagrangian from a set of physical observables, and to evolve them up to the SUSY scale using the three-loop RGEs of the SM. For the physical observables other than the Higgs mass, we use as input for the code the PDG values $G_F = 1.1663787 \times 10^{-5} \text{ GeV}^{-2}$, $M_Z = 91.1876 \text{ GeV}$, $M_W = 80.379 \text{ GeV}$, $M_t = 172.76 \text{ GeV}$, $M_b = 4.78 \text{ GeV}$ and $\alpha_s(m_Z) = 0.1179$ [69]. In order to obtain a prediction for the Higgs mass from a full set of SUSY parameters, we vary the value of the pole mass M_h that we give as input to `mR` until the value of the \overline{MS} -renormalized SM parameter $\lambda_{SM}(Q)$ returned by the code at the SUSY scale $Q = M_S$ coincides with the NMSSM prediction for the quartic coupling of the lightest scalar. In alternative, we can treat the measured value of the Higgs mass, $M_h = 125.25 \text{ GeV}$ [69], as an additional input parameter, and use the matching condition on the quartic Higgs coupling at the SUSY scale to constrain the NMSSM parameter space. In this case we vary one of the SUSY parameters until the NMSSM prediction for the quartic Higgs coupling coincides with the value of $\lambda_{SM}(M_S)$ returned by `mR`.

To obtain the NMSSM prediction for the quartic coupling of the SM-like Higgs boson at the SUSY scale, we combine the tree-level prediction in Eq. (7) with the full one-loop contribution computed in Sect. 2.2, the NMSSM-specific, λ -dependent two-loop-QCD contribution computed in Sect. 2.3, and the λ -independent two-loop contributions that are in common with the MSSM, given in Refs. [64, 65, 67]. As a result, our determination of the matching condition for λ_{SM} includes all of the two-loop contributions that

⁷ In contrast, the bottom Yukawa coupling is defined as the \overline{DR} -renormalized coupling of the NMSSM, to avoid the introduction of spurious $\tan \beta$ -enhanced effects at two loops (see Ref. [65]).

involve the strong gauge coupling, whereas the remaining two-loop contributions are included only under the approximations of vanishing EW gauge couplings and of vanishing λ . While the first approximation is generally good in view of the relative sizes of g, g', g_t and g_s , the goodness of the second approximation obviously depends on the considered values of λ . In particular – similarly to what was found in Refs. [31,34–36] in the context of the FO calculation of the NMSSM Higgs masses – when λ is of $\mathcal{O}(1)$ the two-loop, NMSSM-specific contributions that are omitted in our prediction for λ_{SM} can become as large as the dominant MSSM-like contributions that are included, i.e., those of $\mathcal{O}(g_t^4 g_s^2)$ and $\mathcal{O}(g_t^6)$. We recall that, in the EFT approach, the (next-to)^{*n*}-leading-logarithmic (N^{LL}) resummation of the large logarithmic corrections requires the combination of $(n + 1)$ -loop RGEs with n -loop matching conditions at the boundary scales. Our calculation thus provides a full NLL resummation of the large logarithmic corrections, but, in view of the incomplete determination of the two-loop boundary condition for λ_{SM} at the SUSY scale, only a partial NNLL resummation. An estimate of the impact of the omitted contributions seems desirable.

As was discussed already in Ref. [64], the EFT calculation of the Higgs mass in scenarios where all heavy particles are integrated out at a common scale M_S is subject to three distinct sources of theory uncertainty: (i) a ‘‘SM uncertainty’’ stemming from uncomputed higher-order corrections in the relations between the physical observables taken as input for the calculation and the $\overline{\text{MS}}$ -renormalized parameters of the SM Lagrangian; (ii) a ‘‘SUSY uncertainty’’ stemming from uncomputed higher-order corrections in the matching conditions for the couplings of the SM Lagrangian at the scale M_S ; (iii) an ‘‘EFT uncertainty’’ associated to the corrections suppressed by powers of v^2/M_S^2 that are omitted when the EFT Lagrangian is identified with the renormalizable Lagrangian of the SM in the unbroken phase of the EW symmetry. We neglect the third source of uncertainty in the following, because we consider scenarios in which all BSM particles are heavy enough to make the $\mathcal{O}(v^2/M_S^2)$ effects fully negligible. In contrast, we aim to simulate the effects of uncomputed higher-order corrections in the matching conditions at both the EW scale and the SUSY scale.

To obtain an estimate of the SM uncertainty, we change the accuracy of the determination of the top Yukawa coupling in $m_{\overline{\text{t}}}$, removing corrections of $\mathcal{O}(g_s^6)$ and higher that are implemented by default in the code. This simulates the effect of uncomputed $N^3\text{LL}$ corrections that involve the highest powers of the strong gauge coupling, which are expected to be the largest among those neglected in the ‘‘SM’’ part of the calculation (see, e.g., the discussion in Sect. 6.3.1 of Ref. [8]).

For what concerns the SUSY uncertainty, we combine two estimates of different classes of higher-order effects. To obtain our first estimate, which targets the uncomputed two-

loop and higher-order corrections that involve the top Yukawa coupling, we change the definition of the coupling g_t entering the matching condition for λ_{SM} from the $\overline{\text{MS}}$ -renormalized parameter of the SM to the $\overline{\text{DR}}$ -renormalized parameter of the NMSSM. The two definitions are related by

$$g_t^{\text{NMSSM}}(M_S) = \frac{g_t^{\text{SM}}(M_S)}{1 - \Delta g_t^{\text{MSSM}} - (\Delta g_t)_\lambda}, \tag{26}$$

where Δg_t^{MSSM} is the threshold correction given for the MSSM in Refs. [64,65], and $(\Delta g_t)_\lambda$ is the λ -dependent contribution that turns the MSSM coupling into the NMSSM coupling. This contribution is related to the λ -dependent part of the Higgs WFR by $(\Delta g_t)_\lambda = -(\Delta Z_h)_\lambda/2$. For the latter we find:

$$\begin{aligned} & (4\pi)^2 (\Delta Z_h)_\lambda \\ &= -\frac{a_{hhs}^2}{2m_s^2} - \frac{a_{hHs}^2}{6m_A m_s} \tilde{F}_5\left(\frac{m_s}{m_A}\right) - \frac{a_{hAa}^2}{6m_A m_a} \tilde{F}_5\left(\frac{m_a}{m_A}\right) \\ &+ \frac{\lambda^2}{3} \left[3 \ln \frac{\mu^2}{Q^2} - \sin 2\beta f\left(\frac{m_s}{\mu}\right) + g\left(\frac{m_s}{\mu}\right) \right], \tag{27} \end{aligned}$$

where the terms in the first line are the contributions of diagrams involving scalars or pseudoscalars, while those in the second line are the contributions of diagrams involving higgsinos and singlino. The loop functions $\tilde{F}_5(x)$, $f(x)$ and $g(x)$ are defined in the appendix A of Ref. [64]. The coupling a_{hhs} is given in Eq. (8), and the remaining trilinear couplings are

$$\begin{aligned} a_{hHs} &= -\frac{\lambda}{\sqrt{2}} (A_\lambda + 2\kappa v_s) \cos 2\beta, \\ a_{hAa} &= \frac{\lambda}{\sqrt{2}} (A_\lambda - 2\kappa v_s). \tag{28} \end{aligned}$$

The inclusion of this λ -dependent contribution in the redefinition of g_t , see Eq. (26), simulates the effect of two-loop contributions of $\mathcal{O}(g_t^4 \lambda^2)$ to the matching condition for λ_{SM} . As mentioned above, these can in principle compete with the known $\mathcal{O}(g_t^4 g_s^2)$ and $\mathcal{O}(g_t^6)$ contributions when λ is large enough. We remark that, to avoid the inclusion in our uncertainty estimate of effects that are in fact accounted for by our calculation, the redefinition of g_t must be accompanied by shifts in the known two-loop contributions to the matching condition for λ_{SM} . In particular, in the NMSSM-specific, λ -dependent two-loop-QCD contribution computed in Sect. 2.3 we must remove the second term on the r.h.s. of Eq. (23). The analogous shifts for the MSSM-like two-loop contributions can be trivially obtained from Refs. [64,65,67].

Our second estimate of the SUSY uncertainty targets the uncomputed two-loop and higher-order corrections that involve the highest powers of the doublet–singlet coupling λ . We replace the definitions in Eq. (4) for the scalar and pseudoscalar singlet masses with $m_s^2 = m_S^2 + 2\kappa v_s (A_\kappa + 3\kappa v_s)$ and $m_a^2 = m_S^2 + 2\kappa v_s (-A_\kappa + \kappa v_s)$, respectively.

The two sets of mass definitions are equivalent at the tree level, but they differ at one loop, because the soft SUSY-breaking singlet mass m_S is related to the other parameters as in Eq. (12) when v_s is defined as the vev of the loop-corrected potential. As discussed in Sect. 2.2, the change in the definition of the scalar singlet mass entering the tree-level part of the matching condition for λ_{SM} must be compensated for at the one-loop level by removing the tadpole term in Eq. (11). On the other hand, the change in the definition of both scalar and pseudoscalar singlet masses entering the one-loop part of the matching condition simulates the effect of two-loop λ -dependent corrections that do not involve the strong gauge coupling, and the change in the definition of the scalar singlet mass entering the two-loop-QCD part simulates the effect of three-loop λ -dependent corrections that do involve the strong gauge coupling.

To assess the impact of the different contributions to the matching condition for the quartic Higgs coupling computed in Sect. 2, we plot in Fig. 1 the prediction for the mass of the SM-like Higgs boson as a function of λ . We consider an NMSSM scenario in which all sfermions of the first and second generations have degenerate masses of 2 TeV, while those of the third generation (namely stops, sbottoms and staus) have degenerate masses $M_S = 5$ TeV. The trilinear Higgs-stop interaction parameter is fixed as $X_t = \sqrt{6} M_S$, which maximizes the one-loop stop contribution to λ_{SM} . For given values of μ and $\tan\beta$ this determines the soft SUSY-breaking coupling A_t , and the corresponding sbottom and stau couplings are fixed as $A_b = A_\tau = A_t$. For consistency with our calculation of the two-loop contributions to λ_{SM} , all sfermion masses and trilinear couplings are interpreted as \overline{DR} -renormalized parameters, at a scale that we choose as $Q = M_S$. The soft SUSY-breaking masses of bino, wino and gluino are fixed as $M_1 = 1$ TeV, $M_2 = 2$ TeV and $M_3 = 2.5$ TeV, respectively.

In contrast to the case of the MSSM, in which the tree-level masses of the heavy Higgs bosons and of the higgsinos in the limit of unbroken EW symmetry are determined by the three parameters μ , B_μ and $\tan\beta$ (with $m_A^2 = 2 B_\mu / \sin 2\beta$), the Higgs/higgsino sector of the NMSSM depends on the six parameters λ , κ , v_s , A_λ , A_κ and $\tan\beta$. In the plots of Fig. 1 we choose to vary the doublet-singlet coupling λ , because that parameter determines the size of the NMSSM-specific contributions to the quartic Higgs coupling, and in the limit $\lambda \rightarrow 0$ we recover the MSSM prediction. For the remaining parameters, we choose: $\kappa = \lambda$; a tree-level higgsino mass $\mu = \lambda v_s = 1.5$ TeV, which determines v_s for a given value of λ ; a tree-level heavy-Higgs-doublet mass $m_A = 3$ TeV, which determines A_λ via Eq. (6) for a given value of $\tan\beta$; and $A_\kappa = -2$ TeV. Finally, we fix $\tan\beta = 3$ in the left plot of Fig. 1 and $\tan\beta = 5$ in the right plot. For consistency with our calculation of the one-loop contributions to λ_{SM} , all of these six parameters – which enter the boundary

condition for λ_{SM} already at the tree level – are interpreted as \overline{DR} -renormalized parameters, also at the scale $Q = M_S$. Our choices of parameters correspond to a tree-level mass of 3 TeV for the singlino, and to tree-level masses of about 2.45 TeV and 3 TeV for the scalar and pseudoscalar components of the singlet, respectively.

In all of the lines in the plots of Fig. 1, the Higgs-mass prediction includes all of the contributions to the matching condition for λ_{SM} that are in common with the MSSM, so that the left edge of the plot where $\lambda = 0$ corresponds to our best prediction for M_h in the so-called “MSSM limit”. The three lines in each plot correspond to different accuracies for the inclusion of the NMSSM-specific contributions. The green, dot-dashed line corresponds to the inclusion of the tree-level contribution $(\lambda_{SM}^{\text{tree}})_\lambda = \lambda^2 \sin^2 2\beta/2 - a_{hhs}^2/m_s^2$ alone; the blue, dashed line corresponds to the inclusion of the full one-loop, λ -dependent contribution computed in Sect. 2.2; the red, solid line corresponds to the additional inclusion of the two-loop-QCD, λ -dependent contribution computed in Sect. 2.2. The band around the red, solid line corresponds to our total estimate of the theory uncertainty of the Higgs-mass prediction, obtained by summing linearly the absolute values of the three estimates described above. Comparing the different contributions, we find that the SM uncertainty estimate alone is generally larger than the combination of the two SUSY uncertainty estimates.

The comparison between the left and right plots in Fig. 1 shows that, in our scenario, the λ -dependent contributions increase the prediction for M_h for lower values of $\tan\beta$, and decrease it for higher values of $\tan\beta$. This behavior is driven already at the tree level by the $\tan\beta$ dependence of $(\lambda_{SM}^{\text{tree}})_\lambda$, whose first, positive-definite term is suppressed at larger $\tan\beta$, whereas the second, negative-definite term contains a $\tan\beta$ -independent piece, as remarked after Eq. (8). The comparison between the dot-dashed and dashed lines in each plot shows that the full one-loop, λ -dependent contribution to the matching condition for λ_{SM} can become substantial when $\lambda \gtrsim 0.5$, changing the prediction for M_h by several GeV. Finally, the comparison between the dashed and solid lines shows that the effect on the Higgs-mass prediction of the two-loop-QCD, λ -dependent contribution is quite modest, and it is much smaller than our estimate of the uncomputed higher-order effects. This is likely related to the fact that, with our choices of parameters, the λ -dependent stop contribution is suppressed already at the one-loop level. In particular, the second term in Eq. (14) vanishes for degenerate stop masses and $Q = M_S$, and the first term only amounts to a 2% shift of the tree-level contribution.

The knowledge of the mass of the SM-like Higgs boson can be used to constrain the parameters of the yet-undiscovered SUSY sector. For example, Fig. 3 of Ref. [67] showed the values of M_S and X_t – i.e., the two parameters that most affect the stop contribution to the matching con-




Fig. 1 Higgs-mass prediction as a function of λ , for $\tan \beta = 3$ (left) or $\tan \beta = 5$ (right), in the NMSSM scenario described in the text. The three lines in each plot correspond to different accuracies of the

NMSSM-specific contribution to the matching condition for λ_{SM} . The band around the red, solid line is our estimate of the theory uncertainty

dition for λ_{SM} – that are selected by the requirement that the prediction for M_h coincide with its measured value in an MSSM scenario with moderately large $\tan \beta$. Here we keep the parameters in the stop sector fixed, and exploit the Higgs-mass prediction to constrain two of the parameters that determine the NMSSM-specific contribution to the matching condition for λ_{SM} already at the tree level. In Fig. 2 we show the values of $\tan \beta$ and λ that yield the required prediction $M_h = 125.25$ GeV in a representative NMSSM scenario with heavy BSM particles. We adopt the same choices of SUSY parameters as in Fig. 1, except that *i*) we set M_S to 3 TeV (red lines), 5 TeV (blue lines) or 10 TeV (green lines), and *ii*) we set either $\kappa = \lambda$ (left plot) or $\kappa = 2\lambda$ (right plot).⁸ Each of the lines in Fig. 2 is obtained with our full one-loop and partial two-loop calculation of the matching condition for λ_{SM} , and is accompanied by an uncertainty estimate obtained as described earlier.⁹

The behavior of the different lines in the plots of Fig. 2 can be qualitatively understood by considering the dependence on $\tan \beta$ and λ of the three terms entering the tree-level matching condition for λ_{SM} , see Eq. (7). The first term, which is in common with the MSSM, vanishes for $\tan \beta = 1$ and increases for increasing $\tan \beta$, reaching a constant positive value at large $\tan \beta$. With our choices for the parameters that determine a_{hhs} and m_s , the remaining, NMSSM-specific contribution $(\lambda_{SM}^{tree})_\lambda = \lambda^2 \sin^2 2\beta/2 - a_{hhs}^2/m_s^2$ scales as λ^2 , with a coefficient that is positive at low values of $\tan \beta$, decreases for increasing $\tan \beta$, eventually turns negative, and

finally reaches a constant negative value at large $\tan \beta$. In each of the plots of Fig. 2, the line corresponding to a given value of M_S is split into a left branch at lower $\tan \beta$, where the NMSSM-specific contribution to the matching condition for λ_{SM} is positive, and a right branch at higher $\tan \beta$, where the NMSSM-specific contribution is negative. The point where each line meets the x axis corresponds to a value of $\tan \beta$ that we denote as $(\tan \beta)_{MSSM}$, for which the required prediction for M_h is obtained in the “MSSM limit” $\lambda \rightarrow 0$. This depends on the value of M_S , because heavier stops induce a larger positive contribution to the matching condition for λ_{SM} and thus allow for lower $\tan \beta$ in the MSSM limit.

In the left plot of Fig. 2, where we set $\kappa = \lambda$, the NMSSM-specific contribution to the matching condition for λ_{SM} turns negative for a value of $\tan \beta$ that is lower than $(\tan \beta)_{MSSM}$. At the left edge of the plot, where $\tan \beta = 1$, the MSSM prediction for the Higgs mass is too low, and the required prediction $M_h = 125.25$ GeV is obtained thanks to the positive NMSSM-specific contribution. Moving to higher values of $\tan \beta$, the value of λ that yields the required prediction for M_h decreases at first, as the tree-level, MSSM-like contribution to λ_{SM} increases. However, the coefficient of λ^2 in the tree-level,¹⁰ NMSSM-specific contribution decreases for increasing $\tan \beta$, and as it approaches zero the value of λ that yields $M_h = 125.25$ GeV shoots up. In the gap between the value of $\tan \beta$ for which the NMSSM-specific contribution turns negative and $(\tan \beta)_{MSSM}$ there is no value of λ that yields the required Higgs-mass prediction. Finally, when

⁸ For $\kappa = 2\lambda$ the tree-level masses of the singlet fields become $m_s \approx 5.5$ TeV, $m_a \approx 4.2$ TeV and $m_{\tilde{s}} = 6$ TeV.

⁹ The kinks in Fig. 2 are artifacts induced by the symmetrization of the uncertainty bands in the $\tan \beta$ - λ plane.

¹⁰ We refer only to the tree-level contribution for a qualitative interpretation of the plots. The presence of the radiative corrections, some of which scale as λ^4 in our scenario, does not alter the overall behavior described here.




Fig. 2 Regions of the $\tan \beta$ - λ plane that yield $M_h = 125.25$ GeV, for $\kappa = \lambda$ (left) or $\kappa = 2\lambda$ (right), in the NMSSM scenario described in the text. The three sets of lines in each plot correspond to different values

of the common mass term for the third-generation sfermions. The band around each line is our estimate of the theory uncertainty

$\tan \beta$ is larger than $(\tan \beta)_{\text{MSSM}}$ the MSSM prediction for M_h is too high, and the required prediction is obtained thanks to the negative NMSSM-specific contribution. As both the coefficient of λ^2 in $(\lambda_{\text{SM}}^{\text{tree}})_\lambda$ and the MSSM prediction for M_h reach a plateau at large $\tan \beta$, so does the value of λ that brings the Higgs mass down to 125.25 GeV.

In the right plot of Fig. 2, where we set $\kappa = 2\lambda$, the qualitative behavior of the red line corresponding to $M_S = 3$ TeV is the same as in the left plot. However, the blue and green lines corresponding to higher values of M_S behave differently. In this case, the value of $\tan \beta$ for which the NMSSM-specific contribution to the matching condition for λ_{SM} turns negative is higher than $(\tan \beta)_{\text{MSSM}}$. Consequently, as $\tan \beta$ approaches $(\tan \beta)_{\text{MSSM}}$ from the left, the requirement that $M_h = 125.25$ GeV drives λ to zero. On the right of $(\tan \beta)_{\text{MSSM}}$ there is a gap in which the MSSM prediction for M_h is too high but the NMSSM-specific contribution remains positive, so there is no value of λ that yields the required Higgs-mass prediction. Finally, when the NMSSM-specific contribution does turn negative, the value of λ that brings the prediction for M_h down to 125.25 GeV decreases with increasing $\tan \beta$, and eventually reaches a plateau as in the left plot.

4 Conclusions

If SUSY is realized in nature, there appears to be at least a mild hierarchy between the scale of the superparticle masses and the EW scale. In this kind of hierarchical scenario the prediction for the mass of the SM-like Higgs boson is best obtained in the EFT approach, which allows for the all-orders

resummation of potentially large corrections enhanced by powers of the logarithm of the two scales. The EFT calculation of the Higgs masses in the MSSM is by now quite advanced, with full one-loop and partial two-loop results for the matching conditions for the Higgs self-couplings under a variety of mass hierarchies (see Ref. [8] for a review). In contrast, in the case of the NMSSM analytic calculations of the matching conditions have been performed so far only at the one-loop level, in an extremely constrained scenario where all BSM particles are heavy and their masses depend on just one parameter [60], and in a Split-SUSY scenario where the low-energy EFT includes also the scalar and pseudoscalar components of the singlet plus all of the SUSY fermions [61].

In this paper we obtained a full one-loop result, valid for arbitrary values of all relevant parameters, for the matching condition for the quartic coupling of the Higgs boson, in the NMSSM scenario where all BSM particles are heavy and the EFT valid below the SUSY scale is just the SM. To this purpose, we adapted to the NMSSM the results of Ref. [59], which provides the one-loop matching of a general high-energy theory (without heavy gauge bosons) on a general renormalizable EFT. We compared our results with those of Ref. [60] – in the constrained scenario discussed in that paper – and found a discrepancy related to the definition of the singlet mass entering the tree-level part of the matching condition for λ_{SM} . In addition to the full one-loop calculation of the matching condition, we directly computed the two-loop contributions that involve the strong gauge coupling. Our result includes also terms associated with the external-momentum dependence of the two-loop Higgs self-energy that are missed by the FO calculations of the corresponding corrections in Refs. [21, 30–32]. Finally, we proposed a way

to extend to the NMSSM the estimates of the theory uncertainty associated with uncomputed higher-order effects that had previously been developed for the MSSM.

We found that, in the NMSSM scenario with heavy BSM particles, the matching condition for the quartic Higgs coupling splits neatly into a part that is in common with the MSSM, and an NMSSM-specific part which vanishes for $\lambda \rightarrow 0$. We studied the numerical impact on the Higgs-mass prediction of the different contributions to the NMSSM-specific part of the matching condition, and found that the one-loop and two-loop-QCD contributions modify only moderately, and only for quite large values of λ , the leading behavior driven by the tree-level contribution. We stress that the smallness of these effects is in fact a desirable feature of the EFT calculation of the Higgs mass, in which the logarithmically enhanced corrections are accounted for by the evolution of the parameters between the SUSY scale and the EW scale, and high-precision calculations at the EW scale can be borrowed from the SM.

Turning to the modern approach of treating the Higgs mass as an input rather than an output of the calculation, we illustrated how the requirement of a correct prediction for M_h can be used to constrain some of the yet-unmeasured parameters of the NMSSM. Focusing on the $\tan \beta$ - λ plane, we noticed how the shape of the allowed regions can change drastically depending on the choice of the remaining parameters. This is a well-known aspect of the NMSSM, in which the Higgs sector depends on a relatively large number of parameters already at the tree level. However, a systematic phenomenological study of the constraints that the Higgs-mass prediction imposes on the parameter space of the NMSSM goes beyond the scope of this paper. What we provide here is a set of fully analytic formulas for the matching condition for λ_{SM} , available on request in electronic form. Our results can be used to implement the resummation of the large logarithmic corrections in the existing public codes for the Higgs-mass calculation in the NMSSM [37–53], bringing the accuracy of those codes closer to what has already been attained for the MSSM.

Acknowledgements We thank M. Gabelmann for useful communications concerning Ref. [60]. The work of M. G. and P. S. is supported in part by French state funds managed by the Agence Nationale de la Recherche (ANR), in the context of the grant “HiggsAutomator” (ANR-15-CE31-0002). M. G. also acknowledges support from the ANR grant “DMwithLLPatLHC” (ANR-21-CE31-0013).

Data Availability Statement This manuscript has no associated data or the data will not be deposited. [Authors’ comment: Explicit formulas for the corrections computed in this paper are available on request in electronic form.]

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, pro-

vide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Funded by SCOAP³. SCOAP³ supports the goals of the International Year of Basic Sciences for Sustainable Development.

References

1. CMS Collaboration, S. Chatrchyan et al., Observation of a New Boson at a mass of 125 GeV with the CMS experiment at the LHC. *Phys. Lett. B* **716**, 30–61 (2012). [arXiv:1207.7235](https://arxiv.org/abs/1207.7235) [hep-ex]
2. ATLAS Collaboration, G. Aad et al., Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC. *Phys. Lett. B* **716**, 1–29 (2012). [arXiv:1207.7214](https://arxiv.org/abs/1207.7214) [hep-ex]
3. ATLAS, CMS Collaboration, G. Aad et al., Combined Measurement of the Higgs Boson Mass in pp Collisions at $\sqrt{s} = 7$ and 8 TeV with the ATLAS and CMS Experiments. *Phys. Rev. Lett.* **114**, 191803 (2015). [arXiv:1503.07589](https://arxiv.org/abs/1503.07589) [hep-ex]
4. ATLAS, CMS Collaboration, G. Aad et al., Measurements of the Higgs boson production and decay rates and constraints on its couplings from a combined ATLAS and CMS analysis of the LHC pp collision data at $\sqrt{s} = 7$ and 8 TeV. *JHEP* **08**, 045 (2016). [arXiv:1606.02266](https://arxiv.org/abs/1606.02266) [hep-ex]
5. S.P. Martin, A supersymmetry primer. *Adv. Ser. Direct. High Energy Phys.* **18**, 1–98 (1998). [arXiv:hep-ph/9709356](https://arxiv.org/abs/hep-ph/9709356)
6. M. Maniatis, The next-to-minimal supersymmetric extension of the standard model reviewed. *Int. J. Mod. Phys. A* **25**, 3505–3602 (2010). [arXiv:0906.0777](https://arxiv.org/abs/0906.0777) [hep-ph]
7. U. Ellwanger, C. Hugonie, A.M. Teixeira, The next-to-minimal supersymmetric standard model. *Phys. Rept.* **496**, 1–77 (2010). [arXiv:0910.1785](https://arxiv.org/abs/0910.1785) [hep-ph]
8. P. Slavich, S. Heinemeyer, et al., Higgs-mass predictions in the MSSM and beyond. *Eur. Phys. J. C* **81**(5), 450 (2021). [arXiv:2012.15629](https://arxiv.org/abs/2012.15629) [hep-ph]
9. J.R. Ellis, J. Gunion, H.E. Haber, L. Roszkowski, F. Zwirner, Higgs Bosons in a nonminimal supersymmetric model. *Phys. Rev. D* **39**, 844 (1989)
10. M. Drees, Supersymmetric models with extended higgs sector. *Int. J. Mod. Phys. A* **4**, 3635 (1989)
11. U. Ellwanger, Radiative corrections to the neutral Higgs spectrum in supersymmetry with a gauge singlet. *Phys. Lett. B* **303**, 271–276 (1993). [arXiv:hep-ph/9302224](https://arxiv.org/abs/hep-ph/9302224)
12. P. Pandita, One loop radiative corrections to the lightest Higgs scalar mass in nonminimal supersymmetric Standard Model. *Phys. Lett. B* **318**, 338–346 (1993)
13. P. Pandita, Radiative corrections to the scalar Higgs masses in a nonminimal supersymmetric Standard Model. *Z. Phys. C* **59**, 575–584 (1993)
14. T. Elliott, S. King, P. White, Squark contributions to Higgs boson masses in the next-to-minimal supersymmetric standard model. *Phys. Lett. B* **314**, 56–63 (1993). [arXiv:hep-ph/9305282](https://arxiv.org/abs/hep-ph/9305282)
15. T. Elliott, S. King, P. White, Radiative corrections to Higgs boson masses in the next-to-minimal supersymmetric Standard Model. *Phys. Rev. D* **49**, 2435–2456 (1994). [arXiv:hep-ph/9308309](https://arxiv.org/abs/hep-ph/9308309)
16. S. King, P. White, Resolving the constrained minimal and next-to-minimal supersymmetric standard models. *Phys. Rev. D* **52**, 4183–4216 (1995). [arXiv:hep-ph/9505326](https://arxiv.org/abs/hep-ph/9505326)

17. U. Ellwanger, C. Hugonie, Yukawa induced radiative corrections to the lightest Higgs boson mass in the NMSSM. *Phys. Lett. B* **623**, 93–103 (2005). [arXiv:hep-ph/0504269](#)
18. S. Ham, J. Kim, S. Oh, D. Son, The Charged Higgs boson in the next-to-minimal supersymmetric standard model with explicit CP violation. *Phys. Rev. D* **64**, 035007 (2001). [arXiv:hep-ph/0104144](#)
19. S. Ham, S. Oh, D. Son, Neutral Higgs sector of the next-to-minimal supersymmetric standard model with explicit CP violation. *Phys. Rev. D* **65**, 075004 (2002). [arXiv:hep-ph/0110052](#)
20. S. Ham, Y. Jeong, S. Oh, Radiative CP violation in the Higgs sector of the next-to-minimal supersymmetric model. [arXiv:hep-ph/0308264](#)
21. G. Degrassi, P. Slavich, On the radiative corrections to the neutral Higgs boson masses in the NMSSM. *Nucl. Phys. B* **825**, 119–150 (2010). [arXiv:0907.4682](#) [hep-ph]
22. F. Staub, W. Porod, B. Herrmann, The Electroweak sector of the NMSSM at the one-loop level. *JHEP* **10**, 040 (2010). [arXiv:1007.4049](#) [hep-ph]
23. K. Ender, T. Graf, M. Muhlleitner, H. Rzehak, Analysis of the NMSSM Higgs Boson masses at one-loop level. *Phys. Rev. D* **85**, 075024 (2012). [arXiv:1111.4952](#) [hep-ph]
24. G.G. Ross, K. Schmidt-Hoberg, F. Staub, The generalised NMSSM at one loop: Fine tuning and phenomenology. *JHEP* **08**, 074 (2012)
25. T. Graf, R. Grober, M. Muhlleitner, H. Rzehak, K. Walz, Higgs Boson masses in the complex NMSSM at one-loop level. *JHEP* **10**, 122 (2012). [arXiv:1206.6806](#) [hep-ph]
26. P. Drechsel, L. Galeta, S. Heinemeyer, G. Weiglein, Precise Predictions for the Higgs-Boson Masses in the NMSSM. *Eur. Phys. J. C* **77**(1), 42 (2017). [arXiv:1601.08100](#) [hep-ph]
27. F. Domingo, P. Drechsel, S. Paßehr, On-Shell neutral Higgs bosons in the NMSSM with complex parameters. *Eur. Phys. J. C* **77**(8), 562 (2017). [arXiv:1706.00437](#) [hep-ph]
28. W. G. Hollik, S. Liebler, G. Moortgat-Pick, S. Paßehr, G. Weiglein, Phenomenology of the inflation-inspired NMSSM at the electroweak scale. *Eur. Phys. J. C* **79**(1), 75 (2019). [arXiv:1809.07371](#) [hep-ph]
29. T. N. Dao, M. Muhlleitner, and A. V. Phan, Loop-corrected Higgs masses in the NMSSM with inverse seesaw mechanism. *Eur. Phys. J. C* **82**(8), 667 (2022). [arXiv:2108.10088](#) [hep-ph]
30. M. D. Goodsell, K. Nickel, F. Staub, Two-Loop Higgs mass calculations in supersymmetric models beyond the MSSM with SARAH and SPheno. *Eur. Phys. J. C* **75**(1), 32 (2015). [arXiv:1411.0675](#) [hep-ph]
31. M. Goodsell, K. Nickel, F. Staub, Generic two-loop Higgs mass calculation from a diagrammatic approach. *Eur. Phys. J. C* **75**(6), 290 (2015). [arXiv:1503.03098](#) [hep-ph]
32. M. Muhlleitner, D. T. Nhung, H. Rzehak, K. Walz, Two-loop contributions of the order $\mathcal{O}(\alpha_t \alpha_s)$ to the masses of the Higgs bosons in the CP-violating NMSSM. *JHEP* **05**, 128 (2015). [arXiv:1412.0918](#) [hep-ph]
33. T.N. Dao, R. Gröber, M. Krause, M. Muhlleitner, H. Rzehak, Two-loop $\mathcal{O}(\alpha_t^2)$ corrections to the neutral Higgs boson masses in the CP-violating NMSSM. *JHEP* **08**, 114 (2019). [arXiv:1903.11358](#) [hep-ph]
34. M.D. Goodsell, K. Nickel, F. Staub, Two-loop corrections to the Higgs masses in the NMSSM. *Phys. Rev. D* **91**, 035021 (2015). [arXiv:1411.4665](#) [hep-ph]
35. M.D. Goodsell, F. Staub, The Higgs mass in the CP violating MSSM, NMSSM, and beyond. *Eur. Phys. J. C* **77**(1), 46 (2017)
36. T.N. Dao, M. Gabelmann, M. Muhlleitner, H. Rzehak, Two-loop $\mathcal{O}((\alpha_t + \alpha_\lambda + \alpha_\kappa)^2)$ corrections to the Higgs boson masses in the CP-violating NMSSM. *JHEP* **09**, 193 (2021). [arXiv:2106.06990](#) [hep-ph]
37. U. Ellwanger, J.F. Gunion, C. Hugonie, NMHDECAY: A Fortran code for the Higgs masses, couplings and decay widths in the NMSSM. *JHEP* **02**, 066 (2005). [arXiv:hep-ph/0406215](#)
38. U. Ellwanger, C. Hugonie, NMHDECAY 2.0: An Updated program for sparticle masses, Higgs masses, couplings and decay widths in the NMSSM. *Comput. Phys. Commun.* **175**, 290–303 (2006). [arXiv:hep-ph/0508022](#)
39. F. Staub, SARAH. [arXiv:0806.0538](#) [hep-ph]
40. F. Staub, From superpotential to model files for FeynArts and CalcHep/CompHep. *Comput. Phys. Commun.* **181**, 1077–1086 (2010). [arXiv:0909.2863](#) [hep-ph]
41. F. Staub, Automatic calculation of supersymmetric renormalization group equations and self energies. *Comput. Phys. Commun.* **182**, 808–833 (2011). [arXiv:1002.0840](#) [hep-ph]
42. F. Staub, SARAH 3.2: Dirac Gauginos, UFO output, and more. *Comput. Phys. Commun.* **184**, 1792–1809 (2013). [arXiv:1207.0906](#) [hep-ph]
43. F. Staub, SARAH 4: A tool for (not only SUSY) model builders. *Comput. Phys. Commun.* **185**, 1773–1790 (2014). [arXiv:1309.7223](#) [hep-ph]
44. W. Porod, SPheno, a program for calculating supersymmetric spectra, SUSY particle decays and SUSY particle production at e+ e- colliders. *Comput. Phys. Commun.* **153**, 275–315 (2003). [arXiv:hep-ph/0301101](#) [hep-ph]
45. W. Porod, F. Staub, SPheno 3.1: Extensions including flavour, CP-phases and models beyond the MSSM. *Comput. Phys. Commun.* **183**, 2458–2469 (2012). [arXiv:1104.1573](#) [hep-ph]
46. J. Baglio, R. Gröber, M. Muhlleitner, D. Nhung, H. Rzehak, M. Spira, J. Streicher, K. Walz, NMSSMCALC: A program package for the calculation of loop-corrected higgs boson masses and decay widths in the (complex) NMSSM. *Comput. Phys. Commun.* **185**(12), 3372–3391 (2014). [arXiv:1312.4788](#) [hep-ph]
47. B. C. Allanach, *SOFTSUSY: a program for calculating supersymmetric spectra*. *Comput. Phys. Commun.* **143**, 305–331 (2002). [arXiv:hep-ph/0104145](#) [hep-ph]
48. B. Allanach, P. Athron, L. C. Tunstall, A. Voigt, A. Williams, Next-to-Minimal SOFTSUSY. *Comput. Phys. Commun.* **185**, 2322–2339 (2014). [arXiv:1311.7659](#) [hep-ph]. [Erratum: *Comput. Phys. Commun.* **250**, 107044 (2020)]
49. P. Athron, J.-H. Park, D. Stöckinger, A. Voigt, FlexibleSUSY-A spectrum generator for supersymmetric models. *Comput. Phys. Commun.* **190**, 139–172 (2015). [arXiv:1406.2319](#) [hep-ph]
50. P. Athron, M. Bach, D. Harries, T. Kwasnitza, J.-H. Park, D. Stöckinger, A. Voigt, J. Ziebell, FlexibleSUSY 2.0: Extensions to investigate the phenomenology of SUSY and non-SUSY models. *Comput. Phys. Commun.* **230**, 145–217 (2018). [arXiv:1710.03760](#) [hep-ph]
51. S. Heinemeyer, W. Hollik, G. Weiglein, FeynHiggs: A Program for the calculation of the masses of the neutral CP even Higgs bosons in the MSSM. *Comput. Phys. Commun.* **124**, 76–89 (2000). [arXiv:hep-ph/9812320](#) [hep-ph]
52. T. Hahn, S. Heinemeyer, W. Hollik, H. Rzehak, G. Weiglein, FeynHiggs: A program for the calculation of MSSM Higgs-boson observables - Version 2.6.5. *Comput. Phys. Commun.* **180**, 1426–1427 (2009)
53. H. Bahl, T. Hahn, S. Heinemeyer, W. Hollik, S. Paßehr, H. Rzehak, G. Weiglein, Precision calculations in the MSSM Higgs-boson sector with FeynHiggs 2.14. *Comput. Phys. Commun.* **249**, 107099 (2020)
54. F. Staub, P. Athron, U. Ellwanger, R. Gröber, M. Muhlleitner, P. Slavich, A. Voigt, Higgs mass predictions of public NMSSM spectrum generators. *Comput. Phys. Commun.* **202**, 113–130 (2016)
55. P. Drechsel, R. Gröber, S. Heinemeyer, M. M. Muhlleitner, H. Rzehak, G. Weiglein, Higgs-Boson Masses and Mixing Matrices in the NMSSM: Analysis of On-Shell Calculations. *Eur. Phys. J. C* **77**(6), 366 (2017). [arXiv:1612.07681](#) [hep-ph]
56. P. Athron, J.-H. Park, T. Stuedtner, D. Stöckinger, A. Voigt, Precise Higgs mass calculations in (non-)minimal supersymmetry at both

- high and low scales. *JHEP* **01**, 079 (2017). [arXiv:1609.00371](#) [hep-ph]
57. Florian Staub, Werner Porod, Improved predictions for intermediate and heavy supersymmetry in the MSSM and beyond. *Eur. Phys. J. C* **77**(5) (2017). [arXiv:1703.03267](#) [hep-ph]
58. T. Kwasnitza, D. Stöckinger, A. Voigt, Improved MSSM Higgs mass calculation using the 3-loop FlexibleEFTHiggs approach including xt-resummation. *J. High Energy Phys.* **2020**(7)(2020). [arXiv:2003.04639](#) [hep-ph]
59. J. Braathen, M.D. Goodsell, P. Slavich, Matching renormalisable couplings: simple schemes and a plot. *Eur. Phys. J. C* **79**(8), 669 (2019). [arXiv:1810.09388](#) [hep-ph]
60. M. Gabelmann, M. Mühlleitner, F. Staub, Automatised matching between two scalar sectors at the one-loop level. *Eur. Phys. J. C* **79**(2), 163 (2019). [arXiv:1810.12326](#) [hep-ph]
61. M. Gabelmann, M.M. Mühlleitner, F. Staub, The singlet extended standard model in the context of split supersymmetry. *Phys. Rev. D* **100**, 075026 (2019). [arXiv:1907.04338](#) [hep-ph]
62. B.C. Allanach et al., SUSY Les Houches Accord 2. *Comput. Phys. Commun.* **180**, 8–25 (2009). [arXiv:0801.0045](#) [hep-ph]
63. J. Braathen, M.D. Goodsell, S. Paßehr, E. Painsard, Expectation management. *Eur. Phys. J. C* **81**(6), 498 (2021). [arXiv:2103.06773](#) [hep-ph]
64. E. Bagnaschi, G.F. Giudice, P. Slavich, A. Strumia, Higgs mass and unnatural supersymmetry. *JHEP* **09**, 092 (2014). [arXiv:1407.4081](#) [hep-ph]
65. E. Bagnaschi, J. Pardo Vega, P. Slavich, Improved determination of the Higgs mass in the MSSM with heavy superpartners. *Eur. Phys. J. C* **77**(5), 334 (2017). [arXiv:1703.08166](#) [hep-ph]
66. J. Braathen, M. D. Goodsell, F. Staub, Supersymmetric and non-supersymmetric models without catastrophic Goldstone bosons. *Eur. Phys. J. C* **77**(11), 757 (2017). [arXiv:1706.05372](#) [hep-ph]
67. E. Bagnaschi, G. Degrandi, S. Paßehr, P. Slavich, Full two-loop QCD corrections to the Higgs mass in the MSSM with heavy superpartners. *Eur. Phys. J. C* **79**(11), 910 (2019). [arXiv:1908.01670](#) [hep-ph]
68. B.A. Kniehl, A.F. Pikelner, O.L. Veretin, mr: a C++ library for the matching and running of the Standard Model parameters. *Comput. Phys. Commun.* **206**, 84–96 (2016). [arXiv:1601.08143](#) [hep-ph]
69. Particle Data Group Collaboration, P. A. Zyla et al., Review of Particle Physics. *PTEP* **2020**(8), 083C01 (2020)