



Search for Higgs boson pair production in association with a vector boson in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector

The ATLAS Collaboration

This paper reports a search for Higgs boson pair (hh) production in association with a vector boson (W or Z) using 139 fb^{-1} of proton–proton collision data at $\sqrt{s} = 13$ TeV recorded with the ATLAS detector at the Large Hadron Collider. The search is performed in final states in which the vector boson decays leptonically ($W \rightarrow \ell\nu$, $Z \rightarrow \ell\ell, \nu\nu$ with $\ell = e, \mu$) and the Higgs bosons each decay into a pair of b -quarks. It targets Vhh signals from both non-resonant hh production, present in the Standard Model (SM), and resonant hh production, as predicted in some SM extensions. A 95% confidence-level upper limit of 183 (87) times the SM cross-section is observed (expected) for non-resonant Vhh production when assuming the kinematics are as expected in the SM. Constraints are also placed on Higgs boson coupling modifiers. For the resonant search, upper limits on the production cross-sections are derived for two specific models: one is the production of a vector boson along with a neutral heavy scalar resonance H , in the mass range 260–1000 GeV, that decays into hh , and the other is the production of a heavier neutral pseudoscalar resonance A that decays into a Z boson and H boson, where the A boson mass is 360–800 GeV and the H boson mass is 260–400 GeV. Constraints are also derived in the parameter space of two-Higgs-doublet models.

Contents

1	Introduction	3
2	ATLAS detector	5
3	Data and Monte Carlo samples	6
3.1	Non-resonant signal samples	6
3.2	Resonant signal samples	8
4	Object reconstruction and identification	8
5	Analysis	10
5.1	Signal region event selection	10
5.2	Background estimations	12
5.3	Multivariate discriminant	14
5.4	Mass requirements for resonance searches	18
6	Systematic uncertainties	19
6.1	Experimental systematic uncertainties	19
6.2	Background normalisation systematic uncertainties	21
6.3	Data-driven background modelling uncertainties	22
6.4	Theoretical uncertainties in BDT shapes and acceptances	22
6.5	Impact of systematic uncertainties	23
7	Results and interpretations	24
7.1	Search for non-resonant Vhh production	25
7.2	Searches for $VH \rightarrow Vhh$ production	26
7.3	Search for $A \rightarrow ZH$ production	26
8	Summary	34

1 Introduction

Since the discovery of the Higgs boson (h) with a mass (m_h) of approximately 125 GeV in 2012 [1, 2], its couplings to vector bosons and fermions have been found to be consistent with Standard Model (SM) predictions within the current measurement precision [3–5], providing strong evidence that the Higgs boson has SM properties. The SM also predicts the Higgs boson self-coupling (or trilinear coupling) as well as quartic couplings with itself and with massive vector bosons. These couplings are direct consequences of electroweak symmetry breaking (EWSB) [6–8] and are yet to be confirmed experimentally.

The Higgs boson self-coupling and quartic coupling to vector bosons can be probed through studies of Higgs boson pair (hh) production. In proton–proton (pp) collisions, SM production of hh is dominated by the gluon–gluon fusion (ggF) process [9, 10]. Extensive searches for this process have led to significant bounds on the Higgs boson self-coupling [11–14]. Production through the vector-boson fusion (VBF) process has the second largest cross-section [9, 10]. Searches for VBF hh production have resulted in additional constraints on the Higgs boson self-coupling and are also sensitive to the SM prediction of the Higgs boson quartic coupling to vector bosons [15–17]. While produced through non-resonant processes in the SM, hh production can also occur in resonant processes in scenarios beyond the Standard Model (BSM) through the decays of heavy resonances, such as the heavy Higgs boson predicted in two-Higgs-doublet models (2HDM) [18–20] or the spin-2 Kaluza–Klein gravitons in Randall–Sundrum models [21–23]. Searches for resonant hh production in ggF and VBF processes have also led to constraints in the parameter spaces of these models [24–26].

This paper reports a search for Higgs boson pairs produced in association with a vector boson, Vhh ($V = W, Z$), a process previously unexplored. The search targets both non-resonant hh production, which occurs in the SM, and BSM-inspired resonant hh production. It is performed on a dataset of pp collisions at a centre-of-mass energy of $\sqrt{s} = 13$ TeV collected between 2015 and 2018 with the ATLAS detector at the Large Hadron Collider (LHC), corresponding to an integrated luminosity of $139 \pm 2.4 \text{ fb}^{-1}$ [27]. The search considers vector bosons decaying into leptons ($W \rightarrow \ell\nu, Z \rightarrow \ell\ell, \nu\nu$) and Higgs bosons decaying into a pair of b -quarks ($h \rightarrow bb$), leading to three distinct leptonic channels: $Zhh \rightarrow \nu bbbbb$ (denoted by 0L), $Whh \rightarrow \ell\nu bbbbb$ (denoted by 1L), and $Zhh \rightarrow \ell\ell bbbbb$ (denoted by 2L). Here ℓ denotes either an electron (e) or a muon (μ).¹

Non-resonant hh production in association with a V boson arises in the SM from three distinct Higgs boson couplings: coupling to vector bosons, self-coupling, and quartic coupling to vector bosons. The leading-order processes are depicted in Figures 1(a), 1(b), and 1(c), respectively. In the SM, the cross-sections of these processes at the LHC are small compared with those of the ggF and VBF processes, $0.50 \pm 0.01 \text{ fb}$ for Whh (W^+hh : $0.329 \pm 0.007 \text{ fb}$, and W^-hh : $0.173 \pm 0.005 \text{ fb}$) and $0.36 \pm 0.01 \text{ fb}$ for Zhh at $\sqrt{s} = 13$ TeV with $m_h = 125 \text{ GeV}$ [9, 10], computed at next-to-next-to-leading-order (NNLO) accuracy in QCD.

Two BSM scenarios are considered for resonant hh production as illustrated in Figures 2(a) and 2(b). The first scenario, labelled as VH , is the ‘Higgstrahlung’ production of a generic neutral CP-even scalar H boson which couples directly to vector bosons and decays into hh , i.e. $VH \rightarrow Vhh$. Examples of such a scalar resonance are the CP-even heavy Higgs boson predicted in the electroweak singlet model [28] or in the type-II 2HDM [29]. In this search, the scalar H is assumed to be a narrow resonance; i.e. its natural width is much smaller than the expected experimental relative mass resolution of approximately 3%. This scenario was explored previously by ATLAS in a VBF hh search [15]. The VH search is complementary because it is sensitive to the HWW and HZZ couplings separately, while the VBF search is sensitive only

¹ Although $W \rightarrow \tau\nu$ and $Z \rightarrow \tau\tau$ are not considered explicitly, final states with electrons or muons from τ decays are included.




Figure 1: Leading-order Feynman diagrams of non-resonant hh production in association with a vector boson V expected in the SM from (a) Higgs boson coupling to vector bosons, (b) Higgs boson self-coupling, and (c) Higgs boson quartic coupling to vector bosons. The coupling modifiers κ_V , κ_λ , and κ_{2V} are discussed in Section 3.




Figure 2: Leading-order Feynman diagrams of resonant hh production in association with a vector boson V predicted in some BSM scenarios from the decay of a heavy scalar H originating from (a) an off-shell vector boson and (b) the decay of a neutral heavy pseudoscalar A .

to their combination. The second scenario, labelled as $A \rightarrow ZH$, is a specific process in the 2HDM which predicts three neutral Higgs bosons: two CP-even scalars h and H (with mass hierarchy $m_H > m_h$), and one CP-odd scalar A . In parts of the 2HDM parameter space where the light Higgs boson h is similar to the SM Higgs boson and has a mass $m_h \sim 125$ GeV favourable for electroweak baryogenesis, the A boson has a mass below about 800 GeV but is heavier than the H boson [30]. If m_H is in the range $2m_h < m_H < 2m_t$, the $H \rightarrow hh$ decay branching ratio could be substantial, leading to a sizeable rate for $gg \rightarrow A \rightarrow ZH \rightarrow Zhh$. Here m_t is the mass of the top quark. For this search, natural widths up to 20% of its mass are considered for A , and a narrow width is assumed for H . Searches for $A \rightarrow ZH$ were performed previously by ATLAS and CMS in the $H \rightarrow bb, \tau\tau$, and WW decay channels [31–34].

2 ATLAS detector

The ATLAS detector [35] at the LHC covers nearly the entire solid angle around the collision point.² It consists of an inner tracking detector surrounded by a thin superconducting solenoid, electromagnetic and hadron calorimeters, and a muon spectrometer incorporating three large superconducting air-core toroidal magnets.

The inner-detector system (ID) is immersed in a 2 T axial magnetic field and provides charged-particle tracking in the range $|\eta| < 2.5$. The high-granularity silicon pixel detector covers the vertex region and typically provides four measurements per track, the first hit normally being in the insertable B-layer (IBL) installed before Run 2 [36, 37]. It is followed by the silicon microstrip tracker (SCT), which usually provides eight measurements per track. These silicon detectors are complemented by the transition radiation tracker (TRT), which enables radially extended track reconstruction up to $|\eta| = 2.0$. The TRT also provides electron identification information based on the fraction of hits (typically 30 in total) above a higher energy-deposit threshold corresponding to transition radiation.

The calorimeter system covers the pseudorapidity range $|\eta| < 4.9$. Within the region $|\eta| < 3.2$, the electromagnetic calorimeter (ECAL) consists of barrel and endcap high-granularity lead/liquid-argon (LAr) calorimeters, with an additional thin LAr presampler covering $|\eta| < 1.8$ to correct for energy loss in material upstream of the calorimeters. Hadron calorimetry is provided by the steel/scintillator-tile calorimeter, segmented into three barrel structures within $|\eta| < 1.7$, and two copper/LAr hadron endcap calorimeters. The solid angle coverage is completed with forward copper/LAr and tungsten/LAr calorimeter modules optimised for electromagnetic and hadronic energy measurements respectively.

The muon spectrometer (MS) comprises separate trigger and high-precision tracking chambers measuring the deflection of muons in a magnetic field generated by the superconducting air-core toroidal magnets. The field integral of the toroids ranges between 2.0 and 6.0 T m across most of the detector. Three layers of precision chambers, each consisting of layers of monitored drift tubes, covers the region $|\eta| < 2.7$, complemented by cathode-strip chambers in the forward region, where the background is highest. The muon trigger system covers the range $|\eta| < 2.4$ with resistive-plate chambers in the barrel, and thin-gap chambers in the endcap regions.

Interesting events are selected by the first-level trigger system implemented in custom hardware, followed by selections made by algorithms implemented in software in the high-level trigger [38]. The first-level trigger accepts events from the 40 MHz bunch crossings at a rate below 100 kHz, which the high-level trigger further reduces in order to record events to disk at about 1 kHz.

An extensive software suite [39] is used in data simulation, in the reconstruction and analysis of real and simulated data, in detector operations, and in the trigger and data acquisition systems of the experiment.

² ATLAS uses a right-handed coordinate system with its origin at the nominal interaction point (IP) in the centre of the detector and the z -axis along the beam pipe. The x -axis points from the IP to the centre of the LHC ring, and the y -axis points upwards. Cylindrical coordinates (r, ϕ) are used in the transverse plane, ϕ being the azimuthal angle around the z -axis. The pseudorapidity is defined in terms of the polar angle θ as $\eta = -\ln \tan(\theta/2)$. Angular distance is measured in units of $\Delta R \equiv \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$.

3 Data and Monte Carlo samples

The data used in this analysis were collected using unrescaled single-lepton, missing transverse momentum (E_T^{miss}), or single-photon triggers. The single-lepton trigger requirements are applied as a logical OR of single-electron or single-muon triggers [40, 41], with transverse momentum (p_T) thresholds that started at 20 GeV or 24 GeV in 2015 for muons or electrons, respectively, and increased to 26 GeV in 2016–2018. The E_T^{miss} trigger [42] threshold was raised from 70 GeV to 110 GeV between the 2015 and 2018 data-taking periods. The single-photon trigger [40] was only used in this analysis for background estimation and had a threshold of 140 GeV for the full data-taking period. Events are selected for analysis only if they are of good quality and if all the relevant detector components are known to have been in good operating condition [43], which corresponds to a total integrated luminosity of 139.0 fb^{-1} . The uncertainty in the combined 2015–2018 integrated luminosity is 1.7% [44], obtained using the LUCID-2 detector [27] for the primary luminosity measurements. The recorded events contain an average of 34 inelastic pp collisions per bunch-crossing.

Monte Carlo (MC) simulations are used to optimise the search sensitivities and to estimate background contributions. MC samples were produced with various event generators, interfaced to different programs for parton showers, hadronisation, and underlying-event simulations. No potential interference between signal and background processes is considered. Table 1 summarises the simulation of the signal and background processes relevant for the searches described in this paper. All MC samples are passed through the ATLAS detector simulation program [45] based on GEANT4 [46]. Simulated processes are normalised to the most accurate theoretical cross-section predictions currently available. The effects of multiple interactions in the same or nearby bunch crossings (pile-up) were modelled by overlaying minimum-bias events, simulated using the soft QCD processes of PYTHIA 8.186 [47] with the A3 [48] set of tuned parameters and the NNPDF2.3LO [49] PDF. For all samples of simulated events, except for those generated using SHERPA [50], the EVTGEN 1.6.0 program [51] was used to describe the decays of bottom and charm hadrons.

3.1 Non-resonant signal samples

As shown in Figure 1, three types of the Higgs boson couplings are responsible for non-resonant hh production in the SM. Although these couplings are predicted in the SM once the Higgs boson mass m_h is known, their values should be determined experimentally. Deviations from their SM values are traditionally parameterised using the coupling modifiers [71] denoted, in this paper, by κ_V , κ_λ , and κ_{2V} for the hVV , hhh , and $hhVV$ vertices, respectively, assuming the same coupling modifiers for the W and Z bosons.³

While it is impractical to produce signal samples for arbitrary values of the coupling modifiers probed by this analysis, such samples can be constructed from six independent samples composed of different combinations of the leading-order (LO) processes shown in Figure 1. These component samples were produced with $\kappa_V = \kappa_\lambda = \kappa_{2V} = 1$ for the following couplings and their combinations:

- (1) κ_V : diagram without the hhh and $hhVV$ vertices, Figure 1(a);
- (2) κ_λ : diagram with the hhh vertex, Figure 1(b);
- (3) κ_{2V} : diagram with the $hhVV$ vertex, Figure 1(c);
- (4) $\kappa_V + \kappa_\lambda$: all diagrams except those with an $hhVV$ vertex, Figures 1(a) and 1(b);

³ In the SM, $\kappa_V = 1$, $\kappa_\lambda = 1$, and $\kappa_{2V} = 1$.

Table 1: List of MC event generators, PDFs, and parton shower, hadronisation and underlying event (UE) models used to simulate signal and background processes. Different versions are the results of matching different calculations and models at the time the samples were produced. Here V_ℓ and V_h , respectively, denote V decaying leptonically or hadronically. The last column shows the calculation orders of cross-sections used. Cross-section orders in the last column can be leading order (LO), next-to-leading order (NLO), or next-to-next-to-leading order (NNLO) in terms of QCD or electroweak (EW) accuracy. The mass of the Higgs boson h is set to 125 GeV in the simulation.

Process	Matrix element		Parton shower	UE model	Cross-section order
	generator	order PDF			
Signal samples					
Non-resonant Vhh	MADGRAPH5_AMC@NLO [52]	LO NNPDF2.3LO [49]	PYTHIA 8.244 [53]	A14 [54]	NNLO [9]
Resonant VH	MADGRAPH5_AMC@NLO	LO NNPDF2.3LO	PYTHIA 8.244	A14	LO
Resonant $A \rightarrow ZH$	MADGRAPH5_AMC@NLO	LO NNPDF2.3LO	PYTHIA 8.244	A14	LO
Top quark processes					
$t\bar{t}$	POWHEG Box v2 [55]	NLO NNPDF3.0NLO [56]	PYTHIA 8.230	A14	NNLO +NNLL [57]
Single- t : s-channel	POWHEG Box v2 [58]	NLO NNPDF3.0NLO	PYTHIA 8.230	A14	NLO [59]
Single- t : t-channel	POWHEG Box v2 [58]	NLO NNPDF3.0NLO	PYTHIA 8.230	A14	NLO [60]
Single- t : Wt	POWHEG Box v2 [61]	NLO NNPDF3.0NLO	PYTHIA 8.230	A14	NNLO [62]
ttH	POWHEG Box v2	NLO NNPDF3.0NLO	PYTHIA 8.230	A14	NLO (QCD+EW) [10]
ttV	MADGRAPH5_AMC@NLO	NLO NNPDF3.0NLO	PYTHIA 8.230	A14	NLO (QCD+EW)
$tt\ell\ell^\S$	MADGRAPH5_AMC@NLO	NLO NNPDF3.0NLO	PYTHIA 8.230	A14	NLO
$tttt$	MADGRAPH5_AMC@NLO	NLO NNPDF3.1LO	PYTHIA 8.230	A14	NLO (QCD+EW)
ttt	MADGRAPH5_AMC@NLO	LO NNPDF2.3LO	PYTHIA 8.230	A14	LO
$ttWW$	MADGRAPH5_AMC@NLO	LO NNPDF2.3LO	PYTHIA 8.230	A14	LO
tWZ	MADGRAPH5_AMC@NLO	NLO NNPDF2.3LO	PYTHIA 8.212	A14	NLO
tZ	MADGRAPH5_AMC@NLO	LO NNPDF2.3LO	PYTHIA 8.212	A14	LO
Single- and multi-boson production					
V + jets	SHERPA 2.2.1 [63–65]	NLO \dagger NNPDF3.0NNLO [56]	SHERPA 2.2.1 [66, 67]	Default	NNLO [68]
γ + jets	SHERPA 2.2.2	NLO \dagger NNPDF3.0NNLO	SHERPA 2.2.2	Default	NNLO
$qq \rightarrow V_\ell V_\ell$ or h	SHERPA 2.2.1	NLO NNPDF3.0NNLO	SHERPA 2.2.1	Default	NLO \ddagger
$gg \rightarrow V_\ell V_\ell$ or h	SHERPA 2.2.2	NLO NNPDF3.0NNLO	SHERPA 2.2.2	Default	NLO \ddagger
$V_\ell V_\ell V$	SHERPA 2.2.2	NLO NNPDF3.0NNLO	SHERPA 2.2.2	Default	LO
$V_\ell V_h V_h$	MADGRAPH5_AMC@NLO	LO NNPDF2.3LO	PYTHIA 8.243	A14	LO
$qq \rightarrow VH$	POWHEG Box v2 [69]	NLO NNPDF3.0NLO	PYTHIA 8.212	AZNLO [70]	NNLO(QCD) +NLO(EW)
$gg \rightarrow ZH$	POWHEG Box v2	NLO NNPDF3.0NLO	PYTHIA 8.212	AZNLO	NLO+NLL

\S Here, the same-flavour $\ell\ell$ pairs are from $Z^*/\gamma^* \rightarrow \ell\ell$.

\dagger Produced with up to two extra jets at NLO and up to four extra jets at LO.

\ddagger For the diboson samples the cross-sections are calculated by the Monte Carlo generator at NLO accuracy in QCD.

(5) $\kappa_V + \kappa_{2V}$: all diagrams except those with an hhh vertex, Figures 1(a) and 1(c);

(6) $\kappa_\lambda + \kappa_{2V}$: all diagrams with either an hhh or an $hhVV$ vertex, Figures 1(b) and 1(c).

Samples 1–3 determine the contributions from their respective diagrams, while samples 4–6 allow the determination of the contributions from interference between the three diagrams. Non-resonant signal samples for any coupling modifier deviations from the SM values can be built from the combinations of these component samples, weighted by the coupling modifiers. A SM Vhh sample with $\kappa_V = \kappa_\lambda = \kappa_{2V} = 1$ was also produced to validate this procedure. Kinematic distributions of the reweighted sample are found

to agree with those of the directly produced sample. Since the hVV coupling has been constrained through the single Higgs boson measurements [4], $\kappa_V = 1$ is assumed in this paper.

In addition to the quark-initiated diagrams shown in Figure 1, Zhh can also be produced via the gluon-initiated $gg \rightarrow Zhh$ diagram. Although $gg \rightarrow Zhh$ is technically a higher-order process for Zhh production, its cross-section is predicted to be approximately 24% of that of the $qq \rightarrow Zhh$ process in the SM [10]. A correction is applied to the non-resonant $qq \rightarrow Zhh$ cross-section, as a function of κ_λ and κ_{2V} , to account for the $gg \rightarrow Zhh$ contribution. The generator-level difference, in both normalisation and shape, between the $qq \rightarrow Zhh$ sample and the sum of the $qq \rightarrow Zhh$ and $gg \rightarrow Zhh$ samples is taken as an uncertainty. For a signal as expected in the SM, the normalisation component of this uncertainty has a magnitude of 24%. It tends to be larger than 24% for large κ_{2V} and smaller than 24% for large κ_λ .

3.2 Resonant signal samples

The production of a generic CP-even scalar resonance H in association with a V boson, where the H decays into a pair of the h Higgs bosons, is modelled using the $qq \rightarrow VH \rightarrow Vhh$ process in the 2HDM as shown in Figure 2(a). The heavy Higgs boson is assumed to have a narrow width, i.e. its total decay width is far smaller than the experimental m_{hh} resolution of approximately 3%, but to decay promptly nevertheless. Ten signal samples were produced for each leptonic channel, corresponding to m_H values of 260, 280, 300, 400, 500, 600, 700, 800, 900, and 1000 GeV. The lower bound is dictated by $2m_h$, while the upper bound is limited by the ability to reconstruct two separate jets from the highly boosted $h \rightarrow bb$ decay.

The $gg \rightarrow A \rightarrow ZH \rightarrow Zhh$ signal samples were produced for mass combinations of $m_A = 360, 400, 500, 600, 700, \text{ and } 800$ GeV and $m_H = 260, 300, \text{ and } 400$ GeV subject to the kinematic bound of $m_A > m_H + m_Z$, leading to a total of 15 (m_A, m_H) grid points for each of the 0L and 2L channels. These mass combinations cover the 2HDM parameter space relevant for this search, but unexplored by previous $gg \rightarrow A \rightarrow ZH$ searches [31–34]. The CP-odd A boson could have a substantial total decay width, depending on the 2HDM parameter values. Two sets of samples were produced, one for a narrow-width (NW) A boson and the other for a large-width (LW) A boson. In both scenarios, the total decay width of the heavy CP-even H boson is assumed to be narrow. For the LW samples, the A boson width is assumed to be 20% of its mass. To perform searches for A bosons with different widths, MC distributions for an A boson with width equal to 5%, 10%, or 15% of its mass are derived through reweighting. The generator-level A boson mass distribution for each of these intermediate widths is compared with that of the LW A boson to derive reweighting factors which are then applied to the distributions of the simulated LW MC samples to obtain the corresponding distributions for the intermediate widths.

4 Object reconstruction and identification

As discussed in Section 1, the signal events are characterised by the products of the targeted decays of the vector and Higgs bosons: electrons or muons, jets (b -jets in particular), and missing transverse momentum. Additionally, events with identified hadronically decaying τ -leptons are vetoed to reduce backgrounds. Events with photons are used for background estimation. The reconstruction and identification of these physics objects are described in this section.

Electrons are reconstructed by matching topological energy clusters in the ECAL with tracks in the ID [72] and are required to have $p_T > 7$ GeV within the pseudorapidity range $|\eta| < 2.47$. They are identified

using likelihood-based identification criteria that combined the requirements of calorimeter shower shape, track-to-cluster matching, and associated track qualities. Electron candidates are required to satisfy the *tight* criterion in the 1L channel and the *loose* criterion for the rest. All candidates must satisfy a loose track- and calorimeter-based isolation criterion to minimise the number of jets misidentified as electrons.

Similarly to electrons, photon reconstruction starts with topological energy clusters in the ECAL [72], but the clusters are required to have either no matching tracks (unconverted photons) or one or two matching tracks consistent with a conversion vertex (converted photons). Photon candidates are required to have $p_T > 10$ GeV within the pseudorapidity range $|\eta| < 1.37$ or $1.52 < |\eta| < 2.37$ and satisfy the *tight* identification criterion as well as a calorimeter-based isolation requirement called ‘*TightCaloOnly*’ [72] to suppress backgrounds.

Muons are reconstructed by matching tracks in the ID to either full tracks or track segments in the MS or, for $|\eta| < 0.1$ only, to energy deposits in the calorimeter [73].⁴ They must have $p_T > 7$ GeV and be within $|\eta| < 2.5$, the combined acceptance of the ID and MS. Muons must satisfy the *medium* quality criterion in the 1L channel and the *loose* criterion in others. The same isolation requirement used for the electron selection is applied to all muon candidates to reduce the rate of muons from heavy-flavour decays.

Electrons (muons) are required to have associated tracks satisfying $|d_0/\sigma_{d_0}| < 5$ (3) and $|z_0 \sin \theta| < 0.5$ mm, where d_0 is the transverse impact parameter relative to the beam line, σ_{d_0} is its uncertainty, and z_0 is the distance between the longitudinal position of the track along the beam line at the point where d_0 is measured and the longitudinal position of the primary pp collision vertex.

Jets are reconstructed from particle-flow objects using the anti- k_t algorithm [74, 75] with a radius parameter of $R = 0.4$ and are calibrated as described in Ref. [76]. They are required to have $p_T > 20$ GeV and $|\eta| < 4.5$. Jets with $p_T < 60$ GeV and $|\eta| < 2.4$ originating from pile-up are suppressed with the jet-vertex-tagger [77], a likelihood discriminant based on matching enough of the jet’s tracks to the primary vertex.

Jets containing b -hadrons (b -jets) are identified with the DL1r algorithm [78]. The algorithm is based on information such as properties of displaced tracks and reconstructed secondary vertices in the jet. A jet is b -tagged if the DL1r value is above a preset threshold. Four thresholds, referred to as working points (WP), are defined with average efficiencies of 60%, 70%, 77%, and 85% for tagging b -jets from simulated $t\bar{t}$ events. A pseudo-continuous b -tagging score is defined for each jet as the number of WPs it satisfies, with zero being failing, and four passing, all WPs. In some cases, instead of directly applying b -tagging in MC simulation, events with c - and light-flavor jets are weighted by the probability that these jets pass the b -tagging requirement [79] and a b -tagging score is chosen for each jet based on its b -tagging probability. This procedure, referred to as ‘truth’-tagging, increases the effective MC sample size for subdominant backgrounds. A momentum correction is applied to b -tagged jets to account for the energy lost to soft radiation and to muons and neutrinos in semileptonic b -hadron decays, following the procedure used in Ref. [79]. Furthermore, correction factors are applied to the simulated samples to compensate for the differences between the b -tagging efficiencies in data and simulation [78].

Hadronically decaying τ -leptons, τ_h , are identified from the reconstructed jets [80–82]. A Recurrent Neural Network multivariate discriminant is used to select jets with energy-deposit profiles consistent with those expected from τ_h decay products and to match tracks in the ID to the τ_h candidates, using the ‘medium’ τ_h criteria. The τ_h candidates must have one or three associated tracks and satisfy the requirements of $p_T > 20$ GeV and $|\eta| < 2.5$, excluding the region $1.37 < |\eta| < 1.52$.

⁴ Designed to compensate for the low muon reconstruction efficiency due to a gap in the MS coverage.

The missing transverse momentum $\mathbf{p}_T^{\text{miss}}$, with magnitude E_T^{miss} , is calculated as the negative vectorial sum of the transverse momenta of reconstructed physics objects, namely electrons, muons, photons, hadronically decaying τ -leptons, and jets [83, 84]. A component, called the ‘soft-term’, from energy deposits due to the underlying event and other soft radiation not included in the physics objects is added in the $\mathbf{p}_T^{\text{miss}}$ calculation. A $\mathbf{p}_T^{\text{miss}}$ significance, $\mathcal{S}(E_T^{\text{miss}})$, is defined in order to test whether the measured $\mathbf{p}_T^{\text{miss}}$ is incompatible with zero real $\mathbf{p}_T^{\text{miss}}$. It is calculated from the measured $\mathbf{p}_T^{\text{miss}}$, its resolution, and the correlation between measurements parallel and perpendicular to the $\mathbf{p}_T^{\text{miss}}$ direction [85].

A sequential overlap-removal procedure is applied to ensure that energy deposits in the calorimeter and tracks in the ID are not included in two or more different reconstructed objects. If two electrons share a track, the electron with lower p_T is removed. If an electron and muon share an inner-detector track, the muon is removed if it is calorimeter-tagged, and the electron is removed otherwise. The closest jet within $\Delta R = 0.2$ of a selected electron is removed. If the nearest jet surviving that selection is within $\Delta R = 0.4$ of the electron, the electron is discarded. Muons are usually removed if they are separated from the nearest jet by $\Delta R < 0.4 + 10 \text{ GeV}/p_T^\mu$, since this reduces the background from heavy-flavour decays inside jets. However, if this jet has fewer than three associated tracks, the muon is kept and the jet is removed instead; this avoids an inefficiency for high-energy muons undergoing significant energy loss in the calorimeter. A τ_h candidate is rejected if it is separated by $\Delta R < 0.2$ from any selected electron, muon, or jet.

5 Analysis

The non-resonant and resonant signal models targeted in this search share the same Vhh final state but have different event kinematics. The search starts with the selection of the Vhh final states which are characterised by a leptonically decaying V boson and four b -jets from the decays of the two Higgs bosons. Three leptonic channels (0L, 1L, 2L), one for each leptonic decay mode of the W and Z bosons ($Z \rightarrow \nu\nu$, $W \rightarrow \ell\nu$, $Z \rightarrow \ell\ell$), define the signal regions (SRs) of the search. Multivariate techniques based on boosted decision trees (BDT), trained to distinguish between signal and background events in each SR and for each signal model, provide the final discriminants to extract potential signal contributions. This is achieved through simultaneous fits to the BDT distributions observed in data with the hypotheses of signal plus background contributions. For the resonant models, mass requirements are applied to the resonance candidates before the fits to the BDT distributions. Background contributions are estimated using both data control regions (CRs) and MC simulations. The SR event selections, background estimations, the designs and trainings of the BDTs, and the mass requirements for resonant models are described in this section.

5.1 Signal region event selection

The 0L channel is intended for the $Zhh \rightarrow \nu\nu bbbb$ final state. Candidate events are required to have $E_T^{\text{miss}} > 150 \text{ GeV}$ and $\mathcal{S}(E_T^{\text{miss}}) > 12$. The high E_T^{miss} criterion is necessitated by the trigger requirement. Events with identified *loose* leptons or τ_h are vetoed. To suppress multi-jet backgrounds with mismeasured E_T^{miss} , the minimum azimuthal opening angle between $\mathbf{p}_T^{\text{miss}}$ and the Higgs boson candidates (see below) must satisfy $\min[\Delta\phi(\mathbf{p}_T^{\text{miss}}, \mathbf{h})] > 1$. The ratio of the E_T^{miss} trigger efficiencies in data and simulation, measured in single-muon events as a function of $\mathbf{p}_T^{\text{miss}} + \mathbf{p}_T^\mu$, is applied as an event-weight scale factor to simulated events [42], with corresponding uncertainties described in Section 6.1.

The 1L channel is designed for the $Whh \rightarrow \ell\nu bbbb$ final state. In this channel, candidate events are selected by requiring exactly one *tight* electron with $p_T > 27$ GeV or one *medium* muon with $p_T > 25$ GeV. In addition, the events must have $E_T^{\text{miss}} > 30$ GeV. Candidates with additional *loose* leptons or identified τ_h are removed. The 1L channel is split into two SRs based on the charge of the lepton, 1L+ and 1L-, motivated by the expected large difference between the signal W^+hh and W^-hh cross-sections⁵ in contrast to the mostly charge-symmetric backgrounds.

The 2L channel targets the $Zhh \rightarrow \ell\ell bbbb$ final state. The $Z \rightarrow \ell\ell$ candidates are selected by requiring exactly two oppositely charged *loose* leptons with the same flavour, e^+e^- or $\mu^+\mu^-$, at least one of which has $p_T > 27$ GeV. The invariant mass of the lepton pair must satisfy $81 < m_{\ell\ell} < 101$ GeV for compatibility with the Z boson mass.

Candidate $hh \rightarrow bbbb$ decays are selected by requiring at least four jets passing the 85% b -tagging WP. Within the four-jet combination with the highest pseudo-continuous b -tagging scores (considering all combinations of four jets when more than one combination satisfies this requirement), the jets are paired to form the two $h \rightarrow bb$ candidates, h_1 and h_2 , by minimising the value of $|m_{h_1} - 120 \text{ GeV}| + |m_{h_2} - 120 \text{ GeV}|$. Here m_{h_1} and m_{h_2} are the invariant masses of the two candidates and 120 GeV is their most probable value in simulation. The pair with the higher p_T is labelled as h_1 and the other as h_2 . The efficiency for correctly identifying and pairing the four b -jets into two $h \rightarrow bb$ decays amongst the selected b -jets depends on the signal model and, if applicable, the resonance mass values. For the non-resonant SM Vhh signal, the efficiency is 73%. For the resonant VH signal, the efficiency varies from 63% at $m_H = 260$ GeV to 85% at $m_H = 1000$ GeV. The efficiencies for the $A \rightarrow ZH$ signals are similar to those for the VH signals.

Table 2 summarises the selections that define the SRs for the three leptonic channels. Also included in the table are selections for control regions discussed below. The products of the acceptances and efficiencies of the SR selections, $\mathcal{A} \times \epsilon$, are shown in Figure 3 as functions of the model parameters for a few selected signal models.

Table 2: Selections for the 0L, 1L, and 2L signal regions, and $t\bar{t}$ and V +jets control regions. The ‘—’ symbol indicates no selection is applied.

	Signal regions			Control regions	
	0L	1L (1L+/1L-)	2L	$t\bar{t}$	V +jets
Trigger	E_T^{miss}	single-lepton	single-lepton	single-lepton	single-photon
Lepton or photon	0 <i>loose</i> leptons, 0 identified τ_h	= 1 <i>tight</i> electron with $p_T > 27$ GeV OR 1 <i>medium</i> muon with $p_T > 25$ GeV, 0 additional <i>loose</i> leptons, 0 identified τ_h	= 2 <i>loose</i> leptons (e^+e^- or $\mu^+\mu^-$), ≥ 1 lepton with $p_T > 27$ GeV, $81 < m_{\ell\ell} < 101$ GeV	= 2 <i>loose</i> leptons ($e^\pm\mu^\mp$), ≥ 1 lepton with $p_T > 27$ GeV	= 1 photon with $p_T > 150$ GeV, 0 <i>loose</i> leptons, 0 identified τ_h
p_T^{miss}	$E_T^{\text{miss}} > 150$ GeV, $\mathcal{S}(E_T^{\text{miss}}) > 12$, $ \Delta\phi(\mathbf{p}_T^{\text{miss}}, h) > 1$	$E_T^{\text{miss}} > 30$ GeV	—	—	—
Jets		≥ 4 jets with $p_T > 20$ GeV and passing the 85% b -tagging WP			

⁵ The cross-section ratio of W^+hh and W^-hh is 1.9 in the SM [10]. For the resonant signal models, the ratio varies between 1.8 and 2.3, depending on the resonance masses.




Figure 3: Example dependences of acceptance times efficiency ($\mathcal{A} \times \epsilon$) on signal model parameters: (a) m_A for the $A \rightarrow ZH$ search in the 0L channel for the case of $m_H = 260$ GeV and a LW A boson, (b) m_H for the WH search in the 1L channel, and (c) κ_{2V} and κ_λ for the non-resonant Zhh search in the 2L channel. Each $\mathcal{A} \times \epsilon$ value is calculated for the respective $\nu\nu bbbb$, $\ell\nu bbbb$, and $\ell\ell bbbb$ final state, with ℓ including the e , μ , and τ leptons here. For the resonant VH and $A \rightarrow ZH$ searches, the $\mathcal{A} \times \epsilon$ value is shown both with and without the mass requirements on the reconstructed resonances. Similarly, for the Zhh search, the $\mathcal{A} \times \epsilon$ value is shown both with and without a BDT requirement. The small decreases in $\mathcal{A} \times \epsilon$ at high m_H values in (b) are due to the merging of b -jets from highly boosted $h \rightarrow bb$ decays. The structures in $\mathcal{A} \times \epsilon$ around small values of κ in (c) reflect large changes in the relative contributions of the three production diagrams shown in Figure 1.

5.2 Background estimations

Major background sources in the Vhh search are the production of top-quark pairs ($t\bar{t}$), single top quarks, vector bosons in association with jets ($V + \text{jets}$), diboson and multi-boson events, and multi-jet processes.

Their relative contributions depend on the channel. In the 0L and 1L channels, the $t\bar{t}$ background dominates, with the subleading contribution being from V + jets (Z + jets for 0L and W + jets for 1L). In the 2L channel, Z + jets and $t\bar{t}$ are the two leading sources. A mixture of data-driven and simulation-based methods are used to estimate these backgrounds. MC simulations are used to model the kinematics of background processes except for the multi-jet backgrounds. Contributions from $t\bar{t}$ and V + jets processes are normalised to the data through the use of CRs, while the rest of the non-multi-jet sources are normalised to their theoretical cross-sections within the estimated uncertainties. For the multi-jet backgrounds, their normalisations and kinematic models are derived from auxiliary data samples. These data-driven procedures are described below.

A $t\bar{t}$ control region ($\text{CR}_{t\bar{t}}$) is used to constrain the normalisations of the leptonic $t\bar{t}$ events⁶ in the SRs. The control region is defined by using a selection similar to that for the 2L SR, but requiring a different-flavour opposite-sign lepton pair ($e^\pm\mu^\mp$) instead of a same-flavour opposite-sign lepton pair (e^+e^- or $\mu^+\mu^-$) and removing the dilepton mass requirement to increase the sample size. The jet requirements are the same as for the SR. These selections are summarised in Table 2.

Top-quark-pair events have two b -jets from top-quark decays and can mimic signal events if there are two additional genuine b -jets from radiation or if there are misidentified c -jets or light-flavour jets (j) from radiation or hadronic W boson decays. The flavour content of this radiation is difficult to model, particularly for heavy flavours (b or c). To allow for flavour-dependent variations of the $t\bar{t}$ background contribution in the signal extraction, simulated $t\bar{t}$ events are labelled according to the ‘truth’ flavours of jets from the radiation: events with one or more b -hadrons ($t\bar{t} + \geq 1b$), events with one or more c -hadrons but no b -hadrons ($t\bar{t} + \geq 1c$), and events with no b -hadrons or c -hadrons ($t\bar{t} + j$). The ‘truth’ flavour of a jet is determined from hadrons with $p_T > 5$ GeV found within a cone of size $\Delta R = 0.3$ around the jet axis [78].

The V + jets process is a major background source in all three channels (Z + jets for 0L and 2L and W + jets for 1L). A V + jets CR ($\text{CR}_{V+\text{jets}}$), defined using γ + jets events, is used to assess their contributions. As summarised in Table 2, events in $\text{CR}_{V+\text{jets}}$ are selected by requiring exactly one photon with $p_T > 150$ GeV and no identified e , μ , or τ_h from the data collected using a single-photon trigger. They must pass the same jet requirements as events in the SRs. At $p_T \gtrsim m_V$, γ + jets events and V + jets events are expected to have similar kinematics because they originate from comparable diagrams and the impact of the finite mass of the V bosons becomes small at high p_T . Residual differences between γ + jets and V + jets, driven mainly by lower- p_T events, are taken into account by extrapolation uncertainties (see Section 6).

The V + jets events can be selected as signal events if they have four jets passing the b -tagging requirement, either from genuine heavy-flavour jets or from misidentified light-flavour jets. Like the $t\bar{t}$ events, modelling the jet flavour composition is challenging. The simulated γ + jets events are categorised according to ‘truth’ flavour and jet matching: events with three or more b -hadrons ($V + \geq 3b$), events with ≥ 1 c -hadron but with ≤ 2 b -hadrons ($V + \geq 1c$), and events with zero c -hadrons and ≤ 2 b -hadrons ($V + j$).

The contribution from multi-jet events, including hadronic $t\bar{t}$ events,⁷ is negligible in the 0L and 2L SRs as well as in $\text{CR}_{t\bar{t}}$, minor in the 1L SRs, but significant in $\text{CR}_{V+\text{jets}}$. Both its rate and kinematics are difficult to simulate. Auxiliary data samples rich in multi-jet events are selected to model the corresponding contributions in the 1L SRs and $\text{CR}_{V+\text{jets}}$. For $\text{CR}_{V+\text{jets}}$, the auxiliary sample is defined by inverting the photon isolation requirement while keeping the rest of the selection the same. The kinematic distributions of multi-jet events in $\text{CR}_{V+\text{jets}}$ are modelled by the selected events in this auxiliary sample after subtracting

⁶ Referring to $t\bar{t}$ events with at least one W boson decaying leptonically.

⁷ Referring to $t\bar{t}$ events with both W bosons decaying hadronically.

approximately 10% non-multi-jet contributions taken from simulation, and an uncertainty in the shapes of these distributions is defined by varying the non-multi-jet contributions in the auxiliary sample by $\pm 100\%$. These distributions are assigned a pre-fit normalisation equal to the difference between the number of data events and the non-multi-jet contributions in $\text{CR}_{V+\text{jets}}$, with an uncertainty of 100%. The multi-jet modelling in the 1L SRs follows the same approach, except that the auxiliary sample is selected by inverting the lepton isolation requirement and the multi-jet contribution in the 1L SRs is normalised through a fit to the distribution of the transverse mass of the lepton and $E_{\text{T}}^{\text{miss}}$ system,⁸ in events selected with the 1L SR requirements but with the $E_{\text{T}}^{\text{miss}}$ criteria reversed to $E_{\text{T}}^{\text{miss}} < 30 \text{ GeV}$, which leads to negligible contamination of potential signals produced with cross-sections comparable to the search sensitivity.

Jet b -tagging scores offer the most sensitive information for separating the different flavour contributions of the $t\bar{t}$ and $V + \text{jets}$ backgrounds. Thus the distributions of the sum of the pseudo-continuous b -tagging scores of the four jets with the highest scores, $\sum s_{b\text{-tag}}^{\text{pc}}$, which ranges from 4 to 16, are used to disentangle contributions from the three flavour components. The $\sum s_{b\text{-tag}}^{\text{pc}}$ distributions observed in $\text{CR}_{t\bar{t}}$ and $\text{CR}_{V+\text{jets}}$, divided into three bins (4–9, 10–12, and 13–16), are included in the fits to determine the potential signal contributions as discussed in Section 7. In the fits, the flavour components of the $t\bar{t}$ and $V + \text{jets}$ contributions are normalised with their separate normalisation factors. The $\sum s_{b\text{-tag}}^{\text{pc}}$ template, i.e. the shape of the $\sum s_{b\text{-tag}}^{\text{pc}}$ distribution, of each component is taken from simulation except for the multi-jet contribution in $\text{CR}_{V+\text{jets}}$ discussed above. Figure 4 compares the $\sum s_{b\text{-tag}}^{\text{pc}}$ distributions of the $\text{CR}_{t\bar{t}}$ and $\text{CR}_{V+\text{jets}}$ observed in the data with their post-fit background expectations. The $t\bar{t}$ and $V + \text{jets}$ contributions comprise 92.4% and 79.7% of the total event yields in their respective CRs.

Validation regions (VRs) are defined for each of the SRs and CRs ($\text{CR}_{t\bar{t}}$ and $\text{CR}_{V+\text{jets}}$) to study the background modelling uncertainties (see Section 6). Events in VRs are selected in the same way as those for SRs and CRs but with different jet requirements. Instead of four b -tagged jets, events are required to have at least four jets, exactly three of which pass the 85% b -tagging WP. The highest- p_{T} non- b -tagged jet in $|\eta| < 2.5$ is taken as the fourth jet to build hh candidates and is assigned a pseudo-continuous b -tagging score of 4, which is found to best reproduce the multivariate discriminants in the signal regions.

5.3 Multivariate discriminant

BDTs are used to exploit the kinematic differences between signal and background events passing the SR event selection. One BDT is constructed for each channel (0L, 1L, 2L) and each signal model (Vhh , VH , $A \rightarrow ZH$), resulting in eight BDTs in total (three each for 0L and 2L, two for 1L). To minimise the complexity of the analysis, these BDTs are built in the same way where possible. They differ in having channel-dependent and signal-model-dependent variables. BDT input variables are chosen through extensive comparisons between their distributions in signal and background events while minimising correlations among the variables. Moreover, for the resonant VH and $A \rightarrow ZH$ searches, only variables with weak correlations with the resonance mass are considered so as to lessen the dependence of the BDTs on any particular hypothesised signal mass value. Requirements on the reconstructed resonance masses are applied separately (Section 5.4). Table 3 summarises the input variables used for the eight BDTs.

A common set of seven input variables are used for all eight BDTs. These variables are

⁸ The transverse mass of a system is defined as $\sqrt{m^2 + p_{\text{T}}^2}$, where m and p_{T} are the mass and transverse momentum of the system, respectively.




Figure 4: The background-only post-fit distributions of the sum of the pseudo-continuous b -tagging scores of the four jets with the highest scores in the (a) $\text{CR}_{t\bar{t}}$ and (b) CR_{V+jets} control regions. The bottom panels show the ratios of the data to the total background expectations. The hashed bands represent the combined statistical and systematic uncertainties in the total background predictions.

- the sum ($m_{h_1} + m_{h_2}$) and the difference ($m_{h_1} - m_{h_2}$) of the masses of the Higgs boson candidates h_1 and h_2 ,
- the sum of the b -tagging scores $\sum s_{b\text{-tag}}^{\text{pc}}$,
- the number of jets (N_{jets}),
- the sum of the transverse energy of jets excluding the four selected b -jets (H_{T}^{ex}),
- $m_{h_1}^{\text{FSR}}$ and $m_{h_2}^{\text{FSR}}$, the masses of the two Higgs boson candidates calculated by including additional jets in a cone of size $\Delta R = 0.8$ around each b -jet's axis. This calculation is intended to recover jet energy lost due to final-state radiation (FSR).

Input variables specific to channels or signal models are the invariant mass m_{hh} and transverse momentum p_{T}^{hh} of the reconstructed Higgs boson pair, $E_{\text{T}}^{\text{miss}}$, the transverse momentum of the vector boson (p_{T}^V), the transverse mass of the lepton– $E_{\text{T}}^{\text{miss}}$ system (m_{T}^W), and four variables exploiting differences between the angular distributions of signal and background events. Because of its strong correlation with the resonance mass, m_{hh} is not used in the BDTs for the resonant searches. Instead, requirements on m_{hh} are applied afterwards (see Section 5.4). The variable p_{T}^{hh} is used in all BDTs except those for the $A \rightarrow ZH$ search, due to its correlation with m_A . For the same reason, $E_{\text{T}}^{\text{miss}}$ (p_{T}^V) is not used in the 0L (2L) channel of the $A \rightarrow ZH$ search. The variables $\cosh(\Delta\eta)_1 - \cos(\Delta\phi)_1$ and $\cosh(\Delta\eta)_2 - \cos(\Delta\phi)_2$ ⁹ of the h_1 and h_2 candidates exploit the angular differences between the signal $h \rightarrow bb$ decay and the

⁹ $\Delta\eta$ and $\Delta\phi$ are, respectively, the differences in pseudorapidity and azimuth of the two b -jets of the $h \rightarrow bb$ candidate.

background gluon-splitting $g^* \rightarrow bb$ process and are motivated by $m_{h_1}^2 \propto \cosh(\Delta\eta)_1 - \cos(\Delta\phi)_1$ and $m_{h_2}^2 \propto \cosh(\Delta\eta)_2 - \cos(\Delta\phi)_2$ in the approximation that jets are massless. Similarly, the Higgs bosons and vector bosons are expected to be produced more centrally for signal events and more forwardly for background events. The two rapidity difference variables, between h_1 and h_2 ($|y_{h_1} - y_{h_2}|$) and between V and hh ($|y_V - y_{hh}|$), are designed to take advantage of these subtle differences.

Figure 5 compares the m_{hh} , $m_{h_1} - m_{h_2}$, and H_T^{ex} distributions in data with those expected from background processes for events passing the SR selections in the three leptonic channels. The background distributions are obtained from the fit to the background-only hypothesis discussed in Section 7. Also shown are the expected distributions from example signal models.

Table 3: Variables used in the BDT discriminant in each of the channels and for each signal model; see text for the variable definitions. The \checkmark symbol indicates the inclusion of the variable. The BDTs for the VH and $A \rightarrow ZH$ searches exclude variables strongly correlated with the resonance mass.

Variable	Channel and signal model								
	0L			1L		2L			
	Vhh	VH	$A \rightarrow ZH$	Vhh	VH	Vhh	VH	$A \rightarrow ZH$	
$m_{h_1} + m_{h_2}$	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark
$m_{h_1} - m_{h_2}$	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark
N_{jets}	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark
H_T^{ex}	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark
$\sum s_{b\text{-tag}}^{\text{pc}}$	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark
$m_{h_1}^{\text{FSR}}$	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark
$m_{h_2}^{\text{FSR}}$	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark
m_{hh}	\checkmark			\checkmark		\checkmark			
p_T^{hh}	\checkmark	\checkmark		\checkmark	\checkmark	\checkmark	\checkmark		
E_T^{miss}	\checkmark	\checkmark		\checkmark	\checkmark	\checkmark	\checkmark		\checkmark
p_T^V				\checkmark	\checkmark	\checkmark	\checkmark		
m_T^W				\checkmark					
$\cosh(\Delta\eta)_1 - \cos(\Delta\phi)_1$	\checkmark	\checkmark		\checkmark	\checkmark	\checkmark	\checkmark		
$\cosh(\Delta\eta)_2 - \cos(\Delta\phi)_2$	\checkmark	\checkmark		\checkmark	\checkmark	\checkmark	\checkmark		
$ y_{h_1} - y_{h_2} $	\checkmark	\checkmark		\checkmark	\checkmark	\checkmark	\checkmark		
$ y_V - y_{hh} $						\checkmark	\checkmark		

The BDTs are trained in the TMVA [86] framework. All background processes modelled using simulation are included in the training. In order to make use of the complete set of simulated MC events for the BDT training and evaluation in an unbiased way, the MC events are split into two samples of equal size. A BDT is trained on each of the two samples and applied to the other, such that the same events are never used for both the BDT training and evaluation. Similarly, half of the events in data are evaluated with one of the two BDTs, and half are evaluated with the other. As a result, while significant overtraining is not observed, the effect of any potential overtraining is minimised by following this method. For each BDT, event weights are included in the training so that the relative importance of each background process is taken into account.




Figure 5: Example distributions of kinematic variables used in the BDT trainings: the invariant mass m_{hh} (a, b, c) and the mass difference $m_{h_1} - m_{h_2}$ (d, e, f) of the two Higgs boson candidates, and H_T^{ex} (g, h, i) of events passing the 0L (a, d, g), 1L (b, e, h), and 2L (c, f, i) SR selections. The 1L SR combines 1L+ and 1L- categories. The expected distributions from example non-resonant and resonant signal models, normalised to the total background expectations, are overlaid. No resonant signal distributions are shown for m_{hh} , which is used in the BDTs for non-resonant production only. The final bins include overflows. The bottom panels show the ratios of the data to the total background expectations. The hatched bands represent the combined statistical and systematic uncertainties in the total background predictions.

For the search for non-resonant Vhh production, the SM, κ_λ -only, and κ_{2V} -only signal samples are added together, with equal weights, to form a combined signal sample for the BDT training. The inclusion of the κ_λ and κ_{2V} samples in the training improves the sensitivity to their respective couplings. Three different BDTs are trained, one for each leptonic channel. Little degradation in sensitivity is observed with this strategy relative to training BDTs for each signal sample separately.

For the search for resonant Vhh signals, MC samples produced with different values of the resonance mass are combined with equal weights to form the signal sample for the BDT training in each channel. It results in a BDT that is not strongly correlated with the resonance mass and therefore has good sensitivity to signals with different mass values. This strategy is applied to both the VH and $A \rightarrow ZH$ searches; the combination for the latter includes samples with different m_A and m_H values as well as A bosons with narrow and large widths.

5.4 Mass requirements for resonance searches

For the resonant VH and $A \rightarrow ZH$ searches, the Higgs boson pair hh is produced from the decay of a new heavy scalar H . Consequently, the signal m_{hh} distributions are expected to peak around m_H with a width equal to the natural width of the new scalar convolved with the detector resolution. Therefore, m_{hh} is a powerful discriminant against the expected continuum SM backgrounds.

Since the lighter Higgs boson h is a narrow resonance with known mass, the m_{hh} resolution can be improved by constraining the measured masses of the two Higgs boson candidates, m_{h_1} and m_{h_2} , to their expected value of 125 GeV. This is achieved by scaling the momenta of the b -jets from each Higgs boson candidate by the ratio of 125 GeV to the measured di- b -jet mass. Figure 6(a) compares the m_{hh} distributions before and after the rescaling for a few selected m_H values in the 2L channel of the ZH search. The rescaling improves the relative m_{hh} resolution from 12.1% (6.1%) to 3.5% (2.6%) at $m_H = 300$ (1000) GeV. Similar improvements are obtained in the other leptonic channels of both the VH and $A \rightarrow ZH$ searches. Moreover, in the 2L channel of the $A \rightarrow ZH$ search the rescaling leads to improvement in the mass resolution of Zhh from the decay of a narrow A resonance, as shown in Figure 6(b). At $m_H = 300$ GeV, the relative Zhh mass resolution improves from 9.4% (5.4%) to 2.8% (2.5%) at $m_A = 400$ (800) GeV. For LW A bosons, the rescaling has negligible impact on the width of the Zhh mass distribution. In the following analysis, the rescaled b -jet momenta are used solely to calculate the masses of resonance candidates.

As discussed in Section 5.3, the m_{hh} variable is not used to construct BDTs for the VH and $A \rightarrow ZH$ searches. Instead, m_{hh} is required to be in a window around the target m_H value afterwards. The window sizes are optimised to improve the search sensitivities. They vary from 30 GeV at $m_H = 260$ GeV to 220 GeV at $m_H = 1000$ GeV, corresponding to approximately 3–8 times of the expected m_{hh} resolution after the rescaling. At high resonance masses the backgrounds are small, so the mass windows are widened (relative to the resolution) to increase signal efficiencies. The same m_H -dependent m_{hh} windows are used for all channels and for both the VH and $A \rightarrow ZH$ searches.

The $A \rightarrow ZH$ model has an additional resonance, the A boson. In the 2L channel, the $A \rightarrow ZH \rightarrow \ell\ell hh$ decay can be fully reconstructed. For A bosons with a width significantly smaller than the detector mass resolution, i.e. the NW case, the width of the Zhh mass distribution is dominated by the m_{hh} resolution as a result of the good lepton momentum resolution and the relatively narrow width of the Z boson. Therefore, the invariant mass of the Zhh candidate, m_{Zhh} , is required to fall in a window of the same size as the m_{hh} window for a given m_H , but shifted from the m_H value to the targeted m_A value. For A bosons with a width equal to 20% of the boson mass, i.e. the LW case, such requirements reduce the signal efficiencies




Figure 6: Reconstructed invariant mass distributions of (a) the hh system in the ZH search and (b) the Zhh system in the $A \rightarrow ZH$ search with a NW A boson for a few representative signal mass points in the 2L channel. Distributions before the m_h rescaling are shown as open circles and dashed lines while those after the rescaling are shown as solid circles and solid lines. All distributions are normalised to unity. A NW A boson has a negligible total decay width compared with the experimental mass resolution.

substantially as the m_{Zhh} distributions are broadened by the A boson width. In this case, $m_{Zhh} > 475$ GeV is required only for $m_A \geq 500$ GeV to reduce backgrounds which are present mostly at low m_{Zhh} values. In the 0L channel, the $A \rightarrow ZH \rightarrow \nu\nu hh$ decay cannot be reconstructed fully, due to the escaping neutrinos. Instead, the transverse mass of the hh and E_T^{miss} system, m_T^{Zhh} , is used. However, the m_T^{Zhh} distributions are either too broad in the case of LW A bosons or too severely sculpted by the phase-space limitation for NW A bosons with small mass-splittings between the A and H bosons to be effective in discriminating between signal and backgrounds. Thus a requirement on m_T^{Zhh} is applied only for NW A bosons with $m_A - m_H \geq 200$ GeV. In this case, m_T^{Zhh} must be in the window of $[m_A - 150, m_A + 50]$ GeV for simplicity. Figure 7 compares the data and expected background distributions of m_{hh} and m_T^{Zhh} in the 0L SR, and those of m_{hh} and m_{Zhh} in the 2L SR. Expected distributions from an example signal model are overlaid.

6 Systematic uncertainties

Systematic uncertainties are divided into four categories: experimental uncertainties, theoretical uncertainties of the overall background normalisations, theoretical uncertainties of acceptances and BDT shapes, and data-driven background modelling uncertainties.

6.1 Experimental systematic uncertainties

Jet energy scale (JES) and jet energy resolution (JER) uncertainties comprise the largest group of experimental uncertainties. The JES uncertainties are primarily determined using data-based Z -boson-jet, photon-jet, and multi-jet p_T -balance techniques [87]. Additional uncertainties are applied for the energy




Figure 7: Comparisons between mass distributions of the data and the expected backgrounds: (a) m_{hh} and (b) m_{T}^{Zhh} of the 0L SR, and (c) m_{hh} and (d) m_{Zhh}^{Zhh} of the 2L SR. All masses are calculated after rescaling the measured Higgs boson candidate mass to 125 GeV. The backgrounds are obtained from the background-only fits to the control and signal regions discussed in Section 7. Expected distributions from the $A \rightarrow ZH$ signal at $(m_A, m_H) = (800, 300)$ GeV for a NW A boson, normalised to the total background expectations, are overlaid. The final bins include overflows. The bottom panels show the ratios of the data to the total background expectations. The hatched bands represent the combined statistical and systematic uncertainties in the total background predictions.

scale of jets containing b -quarks. The impact of the JES uncertainties is estimated by scaling the jet energies within their uncertainties. JER uncertainties are also determined from in situ measurements of Z -boson–jet, photon–jet, and dijet p_T balance [87]. The effect of the JER uncertainties is calculated by increasing the resolution within its uncertainties, smearing the jet energy by the resulting change in resolution, and comparing the result with the nominal shape and normalisation in simulation.

Subdominant experimental uncertainties originate from the b -tagging correction factors. The b -tagging correction factors, determined from the difference between the efficiencies measured in data and simulation, are evaluated in five DL1r discriminant bins and are derived separately for b -jets, c -jets, and light-flavour jets [78, 88, 89]. All of the correction factors for the three jet flavours have uncertainties estimated from multiple measurements, which are decomposed into uncorrelated components that are then treated independently. Extra uncertainties are applied when ‘truth’-tagging is used (in the highest BDT-score bins for subleading backgrounds), and these are decorrelated between channels and defined as an overall 30% uncertainty in the yield for these backgrounds. The ‘truth’-tagging uncertainties are designed to cover any differences in BDT shapes between background samples when ‘truth’-tagging is or is not applied.

Uncertainties in the reconstruction, identification, isolation, and trigger efficiencies of electrons [90] and muons [91] are considered, along with the uncertainty in their energy scale and resolution. These are found to have only a small impact on the results. The uncertainties in the energy scale and resolution of the jets and leptons are propagated to the calculation of E_T^{miss} , which also has additional uncertainties from the modelling of the underlying event and the momentum scale, momentum resolution, and reconstruction efficiency of the tracks used to compute the soft-term (see Section 4) [83, 84]. An uncertainty is assigned to the E_T^{miss} trigger correction factors in the OL channel, defined as the entire difference between the trigger efficiencies for data and simulated events. The uncertainty in the combined 2015–2018 integrated luminosity is 1.7%, as described in Section 3. The average number of interactions per bunch crossing in the simulation is rescaled by 1.03 to improve agreement between the data and the simulation, based on a measurement of the visible cross-section in minimum-bias events [92], and an uncertainty, as large as the correction, is included.

6.2 Background normalisation systematic uncertainties

For all background processes, uncertainties are included in the overall normalisations following a similar approach taken for the ATLAS $t\bar{t}h(\rightarrow bb)$ measurement [93]. These uncertainties are correlated across the three leptonic channels. The $t\bar{t} + \geq 1b$ and $V + \geq 3b$ flavour components are free to float in the fit. The $t\bar{t} + \geq 1c$ and $V + \geq 1c$ flavour components are assigned a large uncertainty, 100%, intended to be conservative in case there is significant mismodelling of these processes. These uncertainties are always constrained by the fits. The light-flavour $t\bar{t} + j$ and $V + j$ components are assigned a smaller uncertainty, 10%, as these are better-measured processes. In practice, these overall $t\bar{t} + j$ and $V + j$ uncertainties have little impact on the analysis, since the yields from these backgrounds, particularly at high BDT score, are small. These uncertainties are all constrained by the $\sum s_{b\text{-tag}}^{\text{pc}}$ distributions in $\text{CR}_{t\bar{t}}$ and CR_{V+jets} , as well as by the low BDT-score regions of the SRs.

The above $V + jets$ uncertainties are correlated between $\gamma + jets$, $Z + jets$, and $W + jets$ backgrounds. Extra uncertainties are defined for extrapolations between these backgrounds to account for their kinematic differences. Each flavour component of $Z + jets$ is assigned an extra 20% uncertainty, and each component of $W + jets$ is assigned an extra 30% uncertainty. The sizes of the uncertainties are defined so as to cover flavour-composition differences between the $\gamma + jets$, $Z + jets$, and $W + jets$ backgrounds.

The remaining normalisation uncertainties are subdominant in the analysis. A 100% uncertainty is applied to the multi-jet backgrounds in both the 1L SRs and CR_{V+jets} . The 1L SR multi-jet uncertainty has a small impact due to the size of the background, and the CR_{V+jets} multi-jet background is constrained by the fits to have an uncertainty smaller than 100%. Uncertainties of 25% are applied to each of the remaining small SM backgrounds. Given the small relative yields from these backgrounds, their uncertainties have little impact on the analysis and are thus chosen to be conservative.

6.3 Data-driven background modelling uncertainties

A set of extra data-driven background uncertainties is included. The relative differences of the BDT distributions between data and simulation in the validation region define a set of VR non-closure uncertainties, which are determined prior to any fit. In each of the 0L, 1L (1L+ and 1L-), and 2L regions, two uncertainties are defined: one for normalisation and one for shape (based on the bin-by-bin differences), each of which is correlated between all backgrounds in the channel. In practice, each signal region tends to be dominated by a single background (either $t\bar{t}$ or $V+jets$), such that correlating these uncertainties between backgrounds makes little difference compared to decorrelating the uncertainties between backgrounds. In $CR_{t\bar{t}}$, only the normalisation component of the non-closure is considered, since the individual normalisation factors for each $t\bar{t}$ flavour component can cover any shape difference between data and simulation. In CR_{V+jets} , no non-closure uncertainty is included, as the normalisation component is redundant because of the multi-jet normalisation uncertainty, which by design covers any non-closure between data and simulation. In total, this defines nine components of the validation region non-closure uncertainty, each of which is taken as 100% of the relative difference in normalisation or shape between data and simulation in the validation regions. These uncertainties are designed to be conservative, and in practice they tend to be constrained by the fits.

For the multi-jet backgrounds in CR_{V+jets} and the 1L SRs, shape uncertainties are included in addition to the normalisation uncertainties defined in Section 6.2. The shape uncertainty included in the CR_{V+jets} multi-jet template is obtained by increasing or decreasing the $\gamma+jets$ contamination by 100% in this region. Two shape uncertainties are included in the multi-jet template in the 1L SRs: one by increasing or decreasing the prompt-lepton contamination by 30% when evaluating the template, to account for potential mismodelling of the lepton selection inefficiency, and the other by raising or lowering the m_T^W threshold used to determine this template, to account for the potential impact of the selection on the extracted BDT shapes.

6.4 Theoretical uncertainties in BDT shapes and acceptances

In addition to the normalisation uncertainties defined in Section 6.2, a number of theoretical uncertainties in background and signal modelling are included. These uncertainties tend to be subdominant relative to the background normalisation and data-driven uncertainties, which are in turn subdominant relative to the statistical uncertainty of the data. Each uncertainty is correlated across all analysis regions, and the effects of each uncertainty on the acceptance and BDT shape are correlated.

For each component of the $t\bar{t}$ and $V+jets$ backgrounds and the single-top-quark background, scale uncertainties are included, defined [94] using seven variations of the QCD factorisation and renormalisation scales in the matrix elements by factors of 0.5 and 2, avoiding variations in opposite directions. For each component of the $t\bar{t}$ background and the single-top-quark background, an extra parton-shower uncertainty

is defined by comparing the nominal sample with an alternative sample showered with HERWIG 7.04 [95, 96], and an extra matching uncertainty is defined by comparison with an alternative sample produced with MADGRAPH5_AMC@NLO. For each $t\bar{t}$ component, initial-state radiation (ISR) and final-state radiation (FSR) variations are included, following the procedure described in Ref. [97]. For the Wt component of the single-top-quark background, a comparison between the nominal sample, which uses the diagram removal scheme [98], and an alternative sample, which uses the diagram subtraction scheme [99], defines an extra uncertainty. For $V + \text{jets}$, variations of the CKKW parameter for merging/matching the matrix element with the parton shower are included, as are variations of the resummation scale.

Modelling of the $t\bar{t}$ background is improved by correcting the top-quark p_T distribution to that predicted by calculations of top-quark-pair differential distributions at NNLO QCD and NLO EW accuracy [100]. Previous studies have seen improved agreement between data and prediction in $t\bar{t}$ events, particularly for the top-quark p_T distribution, when comparing the data with the NNLO calculations [101]. For each $t\bar{t}$ component, the change in the BDT shape from the NNLO correction is taken as an uncertainty.

For the signal samples, acceptance uncertainties evaluated at the generator level are included to account for scale variations, PDF variations, and variations of the ISR, FSR, multi-parton interaction (MPI), and colour reconnection parameters [54]. For the non-resonant Zhh signal only, an extra uncertainty is included to cover potential mismodelling of $gg \rightarrow Zhh$. An uncertainty is included in the Zhh normalisation and BDT shape, covering the difference between the $qq \rightarrow Zhh$ process alone and the sum of the $gg \rightarrow Zhh$ and $qq \rightarrow Zhh$ processes (the effects on the normalisation and shape are correlated). No other uncertainties are included in the overall cross-section for any signal sample.

6.5 Impact of systematic uncertainties

The effects of the statistical and systematic uncertainties on the search sensitivities are studied for hypothesised signals following the procedure discussed in Section 7. Table 4 lists the leading sources of uncertainty and shows, for a few selected signal models, the expected relative uncertainties in the fitted signal-strength parameter μ , a factor multiplying the predicted cross-section for the hypothesised signal. In the resonant case, the mass values are chosen to be those with the largest excesses observed in the data as discussed in Section 7. The cross-sections used for signal normalisation correspond approximately to their expected upper limits. In each case, the leading source of uncertainty is the statistical uncertainty of the data.

Table 4: Dominant uncertainties in the best-fit signal-strength parameter μ for hypothesised signals. For this study, the resonance masses are chosen to be those with the largest excesses in the data for illustration: $m_H = 315$ GeV for WH , $m_H = 550$ GeV for ZH , $(m_A, m_H) = (790, 300)$ GeV for $NW A \rightarrow ZH$, and $(m_A, m_H) = (420, 320)$ GeV for $LW A \rightarrow ZH$. The hypothesised signals are normalised to reference cross-sections approximately equal to their expected upper limits.

Model	Vhh like in SM	WH	ZH	$NW A \rightarrow ZH$	$LW A \rightarrow ZH$
Systematic uncertainty source			$\Delta\mu/\mu$ [%]		
Background modelling	+20, -15	+14, -11	+4.7, -3.0	+17, -13	+20, -18
MC statistics	+12, -9.1	+13, -7.8	+4.8, -2.2	+7.2, -4.1	+10, -8.3
Objects	+12, -8.6	+8.0, -5.2	+4.5, -2.2	+19, -11	+16, -12
Signal modelling	+10, -4.7	+12, -4.9	+8.6, -3.0	+14, -5.1	+17, -7.6
VR non-closure	+14, -11	+11, -9.4	+4.4, -3.0	+4.9, -3.7	+12, -10
Total systematic uncertainty	+30, -22	+27, -18	+12, -5.8	+30, -18	+33, -24
Statistical uncertainty	+44, -39	+52, -43	+68, -49	+59, -47	+42, -37
Total	+52, -44	+59, -47	+69, -49	+66, -50	+53, -45

7 Results and interpretations

Potential signal contributions in the data are determined through maximum-likelihood fits to the BDT distributions in the SRs and the $\sum s_{b\text{-tag}}^{\text{pc}}$ distributions in $\text{CR}_{t\bar{t}}$ and $\text{CR}_{V+\text{jets}}$. The procedure is based on the framework described in Refs. [102–104]. A profile-likelihood-ratio test statistic is used to test the signal-plus-background hypothesis, with the signal production rate as the parameter-of-interest. All SRs (0L, 1L, and 2L) are included in the fits for the non-resonant Vhh search and the resonant VH search, while only the 0L and 2L SRs are included for the resonant $A \rightarrow ZH$ search. The BDT distributions are divided into four bins, determined through optimisations of signal sensitivities while maintaining a reasonable number of background MC events in each bin. The binning boundaries are optimised for each channel separately, but are kept the same for different signal models for simplicity.

The $t\bar{t}$ and $V+\text{jets}$ backgrounds in the SRs from MC simulation are decomposed into three jet-flavour categories in the same way as those in the $t\bar{t}$ and $V+\text{jets}$ CRs discussed in Section 5.2. These flavour-dependent contributions share the same normalisation factors (NFs) as their corresponding components in the CRs. In the fits, the NFs are unconstrained for the $t\bar{t} + \geq 1b$ and $V+ \geq 3b$ components and are constrained to unity within their estimated uncertainties for other components. Systematic uncertainties, described in Section 6, are incorporated as additional multiplicative terms, parameterised with nuisance parameters, in the likelihood calculations, where each nuisance parameter is given a prior distribution based on individual studies.

To test the overall compatibility of the data with the background expectations, fits are first performed for the background-only hypothesis. Since BDTs are model dependent, a fit to the BDT and $\sum s_{b\text{-tag}}^{\text{pc}}$ distributions for the non-resonant search is chosen to illustrate the background modelling. Figure 5 compares the data with the background expectations from the fit for a few selected kinematic variables used in training the BDTs. Overall, the post-fit backgrounds are found to reproduce the data well. The fitted NF values of the heavy-flavour components of the $t\bar{t}$ and $V+\text{jets}$ backgrounds are listed in Table 5, along with those from fits to the different signal-plus-background hypotheses discussed below.

The fits are repeated for the signal-plus-background hypotheses for each signal model and resonance mass assumption. The NF values shown in Table 5 from different fits are stable and consistent. Upper limits on

signal production cross-sections are calculated with the CL_s method [105], using the \tilde{q}_μ test statistic in the asymptotic approximation [106].

Global significances are calculated following the procedure given in Ref. [107]. Pseudo-experiments are generated for the background-only hypothesis. For each pseudo-experiment, a scan over all considered mass points is performed to determine the largest local significance for that pseudo-experiment. The fraction of pseudo-experiments with a local significance greater than the maximum local significance found in the scan over the real data defines the global p -value, which in turn defines the global significance.

Table 5: Values of the normalisation factors of the heavy-flavour background components from the simultaneous fits of the signal and control regions to the background-only and different signal-plus-background hypotheses. For this study, the resonance masses with the largest excesses in the data are chosen: $m_H = 315$ GeV for WH , $m_H = 550$ GeV for ZH , $(m_A, m_H) = (790, 300)$ GeV for $NW A \rightarrow ZH$, and $(m_A, m_H) = (420, 320)$ GeV for $LW A \rightarrow ZH$.

Background source	Normalisation constraint	Bkg-only fit	Vhh like in SM	VH		$A \rightarrow ZH$	
				WH	ZH	NW	LW
$t\bar{t} + \geq 1b$	free-float	$1.21^{+0.19}_{-0.16}$	$1.21^{+0.19}_{-0.17}$	$1.20^{+0.21}_{-0.18}$	$1.18^{+0.22}_{-0.18}$	$1.29^{+0.30}_{-0.23}$	$1.27^{+0.26}_{-0.22}$
$t\bar{t} + \geq 1c$	$\pm 100\%$	$1.54^{+0.62}_{-0.54}$	$1.51^{+0.64}_{-0.53}$	$1.56^{+0.68}_{-0.57}$	$1.44^{+0.61}_{-0.49}$	$1.65^{+0.70}_{-0.63}$	$1.85^{+0.57}_{-0.74}$
$V + \geq 3b$	free-float	$0.98^{+0.28}_{-0.30}$	$0.86^{+0.33}_{-0.30}$	$0.83^{+0.35}_{-0.29}$	$0.90^{+0.31}_{-0.30}$	$0.76^{+0.38}_{-0.30}$	$0.72^{+0.38}_{-0.28}$
$V + \geq 1c$	$\pm 100\%$	$1.29^{+0.55}_{-0.40}$	$1.17^{+0.60}_{-0.35}$	$1.09^{+0.61}_{-0.33}$	$1.24^{+0.59}_{-0.39}$	$1.01^{+0.63}_{-0.31}$	$0.97^{+0.60}_{-0.30}$

7.1 Search for non-resonant Vhh production

The BDTs for non-resonant production are used to search for Vhh signals from three production scenarios: ‘SM’, κ_λ , and κ_{2V} . The ‘SM’ scenario assumes SM kinematics but with its cross-section scaled by a signal-strength parameter μ . The κ_λ scenario tests for an anomalous tri-linear hhh coupling, assuming SM couplings for the rest. Similarly, the κ_{2V} scenario tests for an anomalous quartic $hhVV$ coupling. For the κ_λ and κ_{2V} scenarios, both the event kinematics and production cross-section depend on their respective coupling modifier.

Constraints on non-resonant Vhh production are obtained through fits to the signal-plus-background hypotheses described above, assuming the SM value for the $h \rightarrow bb$ decay branching ratio [10]. The three non-resonant scenarios have the same data BDT distribution, but differ in the signal BDT distributions. The background BDT distributions are largely the same, barring small variations in the post-fit background contributions. For ‘SM’ Vhh production, a 95% confidence-level (CL) upper limit of 183 on μ is observed compared with 87^{+41}_{-24} expected. The corresponding post-fit BDT distribution is shown in Figure 8(a). For the κ_λ and κ_{2V} scenarios, Vhh cross-section upper limits are derived from the fits for different values of the coupling modifiers. These limits lead to the observed (expected) 95% CL intervals of $-34.4 < \kappa_\lambda < 33.3$ ($-24.1 < \kappa_\lambda < 22.9$) and $-8.6 < \kappa_{2V} < 10.0$ ($-5.7 < \kappa_{2V} < 7.1$) for the two coupling modifiers. The observed bounds are weaker than the expectations largely because of small excesses of data in the highest BDT bins. These are the first limits derived from the Vhh process, and are considerably weaker than those obtained from the hh searches focused on the ggF and VBF processes [11–16]. In addition, this analysis can search for deviations of the $WWhh$ and $ZZhh$ couplings from their SM values, parameterised by the respective coupling modifiers κ_{2W} and κ_{2Z} (in the SM, $\kappa_{2W} = 1$, and $\kappa_{2Z} = 1$). Cross-section upper limits

are derived from the fits for different values of these coupling modifiers, leading to limits for the observed (expected) 95% CL intervals of $-12.3 < \kappa_{2W} < 13.5$ ($-8.6 < \kappa_{2W} < 9.8$) and $-9.9 < \kappa_{2Z} < 11.3$ ($-7.1 < \kappa_{2Z} < 8.5$) for the two coupling modifiers. Higgs boson couplings other than the one being tested are set to their SM values.

7.2 Searches for $VH \rightarrow Vhh$ production

Constraints on the production of a heavy narrow scalar resonance H in association with a V boson are determined through the fits of the BDT distributions for resonant VH production to the signal-plus-background hypothesis in 5 GeV m_H steps. The step size is chosen to be comparable to, or smaller than, the experimental m_{hh} mass resolution. For each tested m_H value, the BDT distributions are obtained after imposing the m_{hh} mass window requirement discussed in Section 5.4. For m_H values with no corresponding MC signal sample, the BDT distributions are linearly interpolated from those of the two closest neighbouring mass points with MC signal samples. To validate this interpolation, the results of fits performed at m_H points with a MC signal sample are compared with the results obtained when the BDT distribution is an interpolation between those of neighbouring points with MC signal samples.

Fits are performed separately for the $WH \rightarrow \ell\nu hh$ and $ZH \rightarrow (\ell\ell/\nu\nu)hh$ searches. All three channels are included in the fits for both searches. The WH signal contributes mostly to the 1L channel, but with a sizeable contribution in the 0L channel due to an inefficiency in lepton identification. Its contribution to the 2L channel is negligible. Inclusion of the 2L channel in the fit effectively makes the channel an additional CR for the $WH \rightarrow \ell\nu hh$ search, further constraining the Z + jets background. Similarly, the ZH signal contributes mostly to the 0L and 2L channels. Inclusion of the 1L channel helps to constrain the W + jets background.

The data are found to be consistent with the estimated background contributions. The largest upward deviations from the background expectations are at $m_H = 315$ GeV with a local (global) significance of 2.5 (1.3) standard deviations (σ) in the WH search and at $m_H = 550$ GeV with a local (global) significance of 2.7σ (1.3σ) in the ZH search. These small excesses are largely correlated with those observed in the non-resonant Vhh search. The post-fit BDT distribution in the WH search at $m_H = 315$ GeV is illustrated in Figure 8(b). The heavy-flavour background NFs from the fits are compared with those from other fits in Table 5.

The observed and expected 95% CL upper limits on the cross-section $\sigma(VH) \times B(H \rightarrow hh \rightarrow bbbb)$ as a function of m_H are shown in Figure 9. The resonant VH search is sensitive to the HWW and HZZ couplings separately. Compared with the search in the VBF channel [15], which is sensitive to the combination of the two couplings, the VH search has better sensitivity for m_H up to ~ 450 GeV, assuming that SU(2) custodial symmetry (like that in the SM) applies to the HWW and HZZ couplings [109].

7.3 Search for $A \rightarrow ZH$ production

The $A \rightarrow ZH$ search follows a strategy similar to that used in the VH search. The $A \rightarrow ZH$ BDT distributions in the 0L and 2L channels, after applying the mass requirements discussed in Section 5.4, are used to constrain $gg \rightarrow A \rightarrow ZH \rightarrow Zhh \rightarrow Zbbbb$ production for each (m_A, m_H) hypothesis with 10 GeV steps in both masses, separately for NW and LW A bosons. The BDT distributions for (m_A, m_H) hypotheses without MC samples are linearly interpolated from those of the four closest neighbouring mass points with MC samples. For a NW A boson, the expected and observed upper limits in the (m_A, m_H)


plane are shown in Figure 10(a) and Figure 10(b) respectively. The upper limits for a LW A boson, with a width of 20%, are shown in Figure 10(c) (expected) and Figure 10(d) (observed). The ratios of the observed and the expected upper bounds in the (m_A, m_H) plane are shown in Figure 11 for the two width scenarios.

The most significant excesses are observed at $(m_A, m_H) = (790, 300)$ GeV with a local (global) significance of 3.9σ (2.1σ) in the NW scenario and at $(m_A, m_H) = (420, 320)$ GeV with a local (global) significance of 3.8σ (2.8σ) in the LW scenario. Figure 8(c) shows the post-fit BDT distributions in the LW scenario from the search at $(m_A, m_H) = (420, 320)$ GeV. Due to the selection for the LW scenario, there is little sensitivity to either the A boson mass or width, and a real LW signal would likely produce a higher-than-otherwise-expected cross-section upper limit over a wide range of probed A boson mass values. Thus the broad excess in the A boson mass distribution in the LW scenario is consistent with the expectation for a signal.


The 2HDM benchmark used for interpretations has four free parameters: m_A , m_H , $\tan\beta$, and $\cos(\beta - \alpha)$, where $\tan\beta$ is the ratio of the vacuum expectation values of the two doublets and α is the mixing angle of the CP-even Higgs bosons. The limiting case of $\cos(\beta - \alpha) \rightarrow 0$ corresponds to the 2HDM weak decoupling limit [110] in which the lightest CP-even Higgs boson has the same couplings as the SM Higgs boson at the lowest order. The $A \rightarrow ZH$ search results are interpreted as constraints in the plane defined by $\cos(\beta - \alpha)$ and m_A for given m_H and $\tan\beta$ values. For the remaining 2HDM parameters, the mass of the charged Higgs boson is set to be equal to m_A and the potential parameter m_{12}^2 is set to $m_A^2 \tan\beta / (1 + \tan^2\beta)$.

For regions relevant to the sensitivity of this search, the natural width of the H boson remains narrow, especially where $\cos(\beta - \alpha)$ is close to zero. It is estimated that the cross-section upper limits should be valid as long as the natural width of the H boson is less than 1% and, therefore, the search constrains only parts of the 2HDM parameter space that conform with this requirement. On the other hand, the natural width of the A boson varies from narrow to about 20%. This means that cross-section limits have to be calculated for a range of A boson natural widths in the (m_A, m_H) plane, which are subsequently interpreted in the 2HDM planes discussed previously. The signal hypothesis is tested for several values of the A boson natural width, and cross-section upper limits for those widths are derived as a function of m_A . Linear interpolation is used to derive the limit for any natural width between a pair of tested width values.


To interpret these results in the 2HDM, the upper limits on the cross-section are compared with the theoretical predictions of the model. In the type-I and lepton-specific 2HDMs, only gluon–gluon fusion production of the A boson, $gg \rightarrow A$, is relevant, and it is calculated with corrections at up to NNLO in QCD as implemented in SusHi [111–114]. The widths and branching ratios of the Higgs bosons (A , H , and h) are calculated using the 2HDMC code [115]. The procedure used to calculate the cross-sections and branching ratios, as well as to choose the 2HDM parameter values, follows Ref. [10]. The upper limits are shown for some representative values in Figure 12 for the type-I 2HDM and in Figure 13 for the lepton-specific 2HDM. The Hhh coupling vanishes at $\cos(\beta - \alpha) = 0$, a feature which is reflected by the inability of this analysis to exclude this region of the $(\cos(\beta - \alpha), m_A)$ plane. For $\tan\beta = 1$ the sensitivity is the same for both 2HDM scenarios and therefore the corresponding results are omitted from Figure 13. The sensitivity of this search is complementary to that of the $A \rightarrow ZH \rightarrow \ell\ell bb / \ell\ell WW$ search [34] and to the constraints from the Higgs boson coupling measurements [4].



(a)



(b)



(c)

Figure 8: The post-fit BDT distributions of the signal-plus-background hypotheses from the searches for (a) non-resonant Vhh production like in the SM, (b) the resonant WH process at $m_H = 315$ GeV, and (c) the resonant $A \rightarrow ZH$ process at $(m_A, m_H) = (420, 320)$ GeV for a LW A boson. The chosen mass points in (b) and (c) correspond to the most significant excesses of data over the background expectations in their respective search. The fitted signal and background contributions are shown as stacked histograms. Bins without data points have zero observed events. The hatched bands represent the combined statistical and systematic uncertainties in the total background predictions. The bottom panels show the observed bin-by-bin significances for the background-only hypotheses, calculated following the prescription of Ref. [108]. The LW A boson has a total decay width equal to 20% of its mass.




Figure 9: Observed (black solid curve) and expected (black dashed curve) 95% CL upper limits on the production cross-section at $\sqrt{s} = 13$ TeV of a heavy narrow scalar resonance H in the decay mode $H \rightarrow hh \rightarrow bbbb$ in association with (a) a W boson and (b) a Z boson as a function of the resonance mass. The green (inner) and yellow (outer) bands represent $\pm 1\sigma$ and $\pm 2\sigma$ uncertainty in the expected limits.




Figure 10: Upper bounds at 95% CL on $\sigma(A) \times B(A \rightarrow ZH \rightarrow Zhh \rightarrow Zbbbb)$ in the (m_A, m_H) plane for (a, b) a NW A boson and (c, d) a LW A boson. The expected upper limits are shown in (a) and (c) while the observed limits are shown in (b) and (d). The A boson has a total decay width that is negligible compared to the experimental mass resolution in the NW case and is 20% of its mass in the LW case.




Figure 11: Ratios of the observed and the expected 95% CL upper bounds on $\sigma(A) \times B(A \rightarrow ZH \rightarrow Zhh \rightarrow Zbbbb)$ in the (m_A, m_H) plane of the $A \rightarrow ZH$ search with a (a) NW and (b) LW A boson. The A boson has a total decay width that is negligible compared to the experimental mass resolution in the NW case and is 20% of its mass in the LW case.




Figure 12: Interpretation of the upper limits on $\sigma(A) \times B(A \rightarrow ZH \rightarrow Zhh \rightarrow Zbbbb)$ in the parameter space of the type-I 2HDM. The shaded areas correspond to 95% CL exclusion regions in the $(\cos(\beta - \alpha), m_A)$ plane for a given m_H and $\tan\beta$. The hatched area corresponds to natural widths of the H boson for which the upper limits are not valid. Panels (a) and (b) refer to $m_H = 260$ GeV and to $\tan\beta = 1$ and 10, respectively; (c) and (d) refer to $m_H = 350$ GeV and to $\tan\beta = 1$ and 10, respectively.




Figure 13: Interpretation of the upper limits on $\sigma(A) \times B(A \rightarrow ZH \rightarrow Zhh \rightarrow Zbbbb)$ in the parameter space of the lepton-specific 2HDM. The same notation as in Figure 12 is used. Panel (a) refers to $m_H = 260$ GeV, $\tan\beta = 10$ and (b) refers to $m_H = 350$ GeV, $\tan\beta = 5$.

8 Summary

Searches for Higgs boson pair production in association with a vector boson in pp collisions at $\sqrt{s} = 13$ TeV are performed using a data sample corresponding to an integrated luminosity of 139 fb^{-1} , recorded by the ATLAS experiment between 2015 and 2018 at the LHC. The Higgs bosons are identified via their decays into a pair of b -quarks and the vector bosons are required to decay into leptons, leading to final states with zero, one or two charged leptons along with four b -jets. The searches target both SM-inspired non-resonant hh production and BSM-motivated resonant hh production. The non-resonant Vhh search is carried out for scenarios with either SM kinematics but an enhanced production cross-section or modified Higgs boson couplings to vector bosons or itself. The resonant Vhh searches are designed for the production of a vector boson along with a heavy neutral scalar Higgs boson H decaying into hh , either directly or indirectly from the decay of another heavier neutral pseudoscalar Higgs boson A .

In general, the data are found to be in good agreement with the estimated background contributions, except for a few notable excesses. The most significant global excess is observed in the $gg \rightarrow A \rightarrow ZH \rightarrow Zhh$ search for a large-width A boson at $(m_A, m_H) = (420, 320)$ GeV, where the local (global) significance is 3.8 (2.8) standard deviations. More data are needed to ascertain the nature of this excess. Upper bounds on the Vhh production cross-sections are derived. For non-resonant production with SM kinematics, a 95% CL upper limit of 183 (87) is observed (expected) for the Vhh cross-section relative to its SM prediction. For resonant production, the observed (expected) upper limits are presented as a function of m_H in the range 260–1000 GeV for WH and ZH separately, and in the (m_A, m_H) plane for $A \rightarrow ZH$, covering the m_A range 360–800 GeV and m_H range 260–400 GeV. The constraints on $A \rightarrow ZH$ production are also interpreted in the $(\cos(\beta - \alpha), m_A)$ parameter space of type-I and lepton-specific two-Higgs-doublet models.

Acknowledgements

We thank CERN for the very successful operation of the LHC, as well as the support staff from our institutions without whom ATLAS could not be operated efficiently.

We acknowledge the support of ANPCyT, Argentina; YerPhI, Armenia; ARC, Australia; BMWFW and FWF, Austria; ANAS, Azerbaijan; CNPq and FAPESP, Brazil; NSERC, NRC and CFI, Canada; CERN; ANID, Chile; CAS, MOST and NSFC, China; Minciencias, Colombia; MEYS CR, Czech Republic; DNRf and DNSRC, Denmark; IN2P3-CNRS and CEA-DRF/IRFU, France; SRNSFG, Georgia; BMBF, HGF and MPG, Germany; GSRI, Greece; RGC and Hong Kong SAR, China; ISF and Benozziyo Center, Israel; INFN, Italy; MEXT and JSPS, Japan; CNRST, Morocco; NWO, Netherlands; RCN, Norway; MEiN, Poland; FCT, Portugal; MNE/IFA, Romania; MESTD, Serbia; MSSR, Slovakia; ARRS and MIZŠ, Slovenia; DSI/NRF, South Africa; MICINN, Spain; SRC and Wallenberg Foundation, Sweden; SERI, SNSF and Cantons of Bern and Geneva, Switzerland; MOST, Taiwan; TENMAK, Türkiye; STFC, United Kingdom; DOE and NSF, United States of America. In addition, individual groups and members have received support from BCKDF, CANARIE, Compute Canada and CRC, Canada; PRIMUS 21/SCI/017 and UNCE SCI/013, Czech Republic; COST, ERC, ERDF, Horizon 2020 and Marie Skłodowska-Curie Actions, European Union; Investissements d’Avenir Labex, Investissements d’Avenir Idex and ANR, France; DFG and AvH Foundation, Germany; Herakleitos, Thales and Aristeia programmes co-financed by EU-ESF and the Greek NSRF, Greece; BSF-NSF and MINERVA, Israel; Norwegian Financial Mechanism 2014-2021, Norway; NCN and NAWA, Poland; La Caixa Banking Foundation, CERCA Programme Generalitat de

Catalunya and PROMETEO and GenT Programmes Generalitat Valenciana, Spain; Göran Gustafssons Stiftelse, Sweden; The Royal Society and Leverhulme Trust, United Kingdom.

The crucial computing support from all WLCG partners is acknowledged gratefully, in particular from CERN, the ATLAS Tier-1 facilities at TRIUMF (Canada), NDGF (Denmark, Norway, Sweden), CC-IN2P3 (France), KIT/GridKA (Germany), INFN-CNAF (Italy), NL-T1 (Netherlands), PIC (Spain), ASGC (Taiwan), RAL (UK) and BNL (USA), the Tier-2 facilities worldwide and large non-WLCG resource providers. Major contributors of computing resources are listed in Ref. [116].

References

- [1] ATLAS Collaboration, *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, *Phys. Lett. B* **716** (2012) 1, arXiv: [1207.7214 \[hep-ex\]](#).
- [2] CMS Collaboration, *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, *Phys. Lett. B* **716** (2012) 30, arXiv: [1207.7235 \[hep-ex\]](#).
- [3] ATLAS and CMS Collaborations, *Measurements of the Higgs boson production and decay rates and constraints on its couplings from a combined ATLAS and CMS analysis of the LHC pp collision data at $\sqrt{s} = 7$ and 8 TeV*, *JHEP* **08** (2016) 045, arXiv: [1606.02266 \[hep-ex\]](#).
- [4] ATLAS Collaboration, *A detailed map of Higgs boson interactions by the ATLAS experiment ten years after the discovery*, *Nature* **607** (2022) 52, arXiv: [2207.00092 \[hep-ex\]](#).
- [5] CMS Collaboration, *A portrait of the Higgs boson by the CMS experiment ten years after the discovery*, *Nature* **607** (2022) 60, arXiv: [2207.00043 \[hep-ex\]](#).
- [6] F. Englert and R. Brout, *Broken Symmetry and the Mass of Gauge Vector Mesons*, *Phys. Rev. Lett.* **13** (1964) 321.
- [7] P. W. Higgs, *Broken Symmetries and the Masses of Gauge Bosons*, *Phys. Rev. Lett.* **13** (1964) 508.
- [8] G. S. Guralnik, C. R. Hagen and T. W. B. Kibble, *Global Conservation Laws and Massless Particles*, *Phys. Rev. Lett.* **13** (1964) 585.
- [9] J. Baglio et al., *The measurement of the Higgs self-coupling at the LHC: theoretical status*, *JHEP* **04** (2013) 151, arXiv: [1212.5581 \[hep-ph\]](#).
- [10] D. de Florian et al., *Handbook of LHC Higgs Cross Sections: 4. Deciphering the Nature of the Higgs Sector*, (2016), arXiv: [1610.07922 \[hep-ph\]](#).
- [11] ATLAS Collaboration, *Combination of searches for Higgs boson pairs in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Lett. B* **800** (2020) 135103, arXiv: [1906.02025 \[hep-ex\]](#).
- [12] CMS Collaboration, *Combination of Searches for Higgs Boson Pair Production in Proton–Proton Collisions at $\sqrt{s} = 13$ TeV*, *Phys. Rev. Lett.* **122** (2019) 121803, arXiv: [1811.09689 \[hep-ex\]](#).
- [13] ATLAS Collaboration, *Search for Higgs boson pair production in the two bottom quarks plus two photons final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Rev. D* **106** (2021) 052001, arXiv: [2112.11876 \[hep-ex\]](#).
- [14] CMS Collaboration, *Search for nonresonant Higgs boson pair production in final states with two bottom quarks and two photons in proton–proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **03** (2021) 257, arXiv: [2011.12373 \[hep-ex\]](#).
- [15] ATLAS Collaboration, *Search for the $HH \rightarrow b\bar{b}b\bar{b}$ process via vector-boson fusion production using proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *JHEP* **07** (2020) 108, arXiv: [2001.05178 \[hep-ex\]](#), Erratum: *JHEP* **01** (2021) 145.

- [16] CMS Collaboration, *Search for Higgs Boson Pair Production in the Four b Quark Final State in Proton–Proton Collisions at $\sqrt{s} = 13$ TeV*, *Phys. Rev. Lett.* **129** (2022) 081802, arXiv: [2202.09617 \[hep-ex\]](#).
- [17] CMS Collaboration, *Search for nonresonant pair production of highly energetic Higgs bosons decaying to bottom quarks*, (2022), arXiv: [2205.06667 \[hep-ex\]](#).
- [18] T. D. Lee, *A Theory of Spontaneous T Violation*, *Phys. Rev. D* **8** (1973) 1226.
- [19] G. C. Branco et al., *Theory and phenomenology of two-Higgs-doublet models*, *Phys. Rept.* **516** (2012) 1, arXiv: [1106.0034 \[hep-ph\]](#).
- [20] F. Kling, J. M. No and S. Su, *Anatomy of exotic Higgs decays in 2HDM*, *JHEP* **09** (2016) 093, arXiv: [1604.01406 \[hep-ph\]](#).
- [21] L. Randall and R. Sundrum, *Large Mass Hierarchy from a Small Extra Dimension*, *Phys. Rev. Lett.* **83** (1999) 3370, arXiv: [hep-ph/9905221](#).
- [22] A. L. Fitzpatrick, J. Kaplan, L. Randall and L.-T. Wang, *Searching for the Kaluza-Klein graviton in bulk RS models*, *JHEP* **09** (2007) 013, arXiv: [hep-ph/0701150](#).
- [23] K. Agashe, H. Davoudiasl, G. Perez and A. Soni, *Warped gravitons at the CERN LHC and beyond*, *Phys. Rev. D* **76** (2007) 036006, arXiv: [hep-ph/0701186](#).
- [24] ATLAS Collaboration, *Search for resonant pair production of Higgs bosons in the $b\bar{b}b\bar{b}$ final state using pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Rev. D* **105** (2022) 092002, arXiv: [2202.07288 \[hep-ex\]](#).
- [25] CMS Collaboration, *Search for resonant pair production of Higgs bosons decaying to bottom quark-antiquark pairs in proton–proton collisions at 13 TeV*, *JHEP* **08** (2018) 152, arXiv: [1806.03548 \[hep-ex\]](#).
- [26] CMS Collaboration, *Search for a heavy Higgs boson decaying into two lighter Higgs bosons in the $\tau\tau bb$ final state at 13 TeV*, *JHEP* **11** (2021) 057, arXiv: [2106.10361 \[hep-ex\]](#).
- [27] G. Avoni et al., *The new LUCID-2 detector for luminosity measurement and monitoring in ATLAS*, *JINST* **13** (2018) P07017.
- [28] M. Mühlleitner, M. O. P. Sampaio, R. Santos and J. Wittbrodt, *Phenomenological comparison of models with extended Higgs sectors*, *JHEP* **08** (2017) 132, arXiv: [1703.07750 \[hep-ph\]](#).
- [29] B. Coleppa, F. Kling and S. Su, *Constraining type II 2HDM in light of LHC Higgs searches*, *JHEP* **01** (2014) 161, arXiv: [1305.0002 \[hep-ph\]](#).
- [30] G. C. Dorsch, S. J. Huber, K. Mimasu and J. M. No, *Echoes of the Electroweak Phase Transition: Discovering a Second Higgs Doublet through $A_0 \rightarrow ZH_0$* , *Phys. Rev. Lett.* **113** (2014) 211802, arXiv: [1405.5537 \[hep-ph\]](#).
- [31] CMS Collaboration, *Search for neutral resonances decaying into a Z boson and a pair of b jets or τ leptons*, *Phys. Lett. B* **759** (2016) 369, arXiv: [1603.02991 \[hep-ex\]](#).
- [32] ATLAS Collaboration, *Search for a heavy Higgs boson decaying into a Z boson and another heavy Higgs boson in the $\ell\ell bb$ final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Lett. B* **783** (2018) 392, arXiv: [1804.01126 \[hep-ex\]](#).

- [33] CMS Collaboration, *Search for new neutral Higgs bosons through the $H \rightarrow ZA \rightarrow \ell^+ \ell^- b \bar{b}$ process in pp collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **03** (2020) 055, arXiv: 1911.03781 [hep-ex].
- [34] ATLAS Collaboration, *Search for a heavy Higgs boson decaying into a Z boson and another heavy Higgs boson in the $\ell\ell b\bar{b}$ and $\ell\ell WW$ final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Eur. Phys. J. C* **81** (2020) 396, arXiv: 2011.05639 [hep-ex].
- [35] ATLAS Collaboration, *The ATLAS Experiment at the CERN Large Hadron Collider*, *JINST* **3** (2008) S08003.
- [36] ATLAS Collaboration, *ATLAS Insertable B-Layer: Technical Design Report*, ATLAS-TDR-19; CERN-LHCC-2010-013, 2010, URL: <https://cds.cern.ch/record/1291633>, Addendum: ATLAS-TDR-19-ADD-1; CERN-LHCC-2012-009, 2012, URL: <https://cds.cern.ch/record/1451888>.
- [37] B. Abbott et al., *Production and integration of the ATLAS Insertable B-Layer*, *JINST* **13** (2018) T05008, arXiv: 1803.00844 [physics.ins-det].
- [38] ATLAS Collaboration, *Performance of the ATLAS trigger system in 2015*, *Eur. Phys. J. C* **77** (2017) 317, arXiv: 1611.09661 [hep-ex].
- [39] ATLAS Collaboration, *The ATLAS Collaboration Software and Firmware*, ATL-SOFT-PUB-2021-001, 2021, URL: <https://cds.cern.ch/record/2767187>.
- [40] ATLAS Collaboration, *Performance of electron and photon triggers in ATLAS during LHC Run 2*, *Eur. Phys. J. C* **80** (2020) 47, arXiv: 1909.00761 [hep-ex].
- [41] ATLAS Collaboration, *Performance of the ATLAS muon triggers in Run 2*, *JINST* **15** (2020) P09015, arXiv: 2004.13447 [hep-ex].
- [42] ATLAS Collaboration, *Performance of the missing transverse momentum triggers for the ATLAS detector during Run-2 data taking*, *JHEP* **08** (2020) 080, arXiv: 2005.09554 [hep-ex].
- [43] ATLAS Collaboration, *ATLAS data quality operations and performance for 2015–2018 data-taking*, *JINST* **15** (2020) P04003, arXiv: 1911.04632 [physics.ins-det].
- [44] ATLAS Collaboration, *Luminosity determination in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector at the LHC*, ATLAS-CONF-2019-021, 2019, URL: <https://cds.cern.ch/record/2677054>.
- [45] ATLAS Collaboration, *The ATLAS Simulation Infrastructure*, *Eur. Phys. J. C* **70** (2010) 823, arXiv: 1005.4568 [physics.ins-det].
- [46] S. Agostinelli et al., *GEANT4 – a simulation toolkit*, *Nucl. Instrum. Meth. A* **506** (2003) 250.
- [47] T. Sjöstrand, S. Mrenna and P. Z. Skands, *A brief introduction to PYTHIA 8.1*, *Comput. Phys. Commun.* **178** (2008) 852, arXiv: 0710.3820 [hep-ph].
- [48] ATLAS Collaboration, *The Pythia 8 A3 tune description of ATLAS minimum bias and inelastic measurements incorporating the Donnachie–Landshoff diffractive model*, ATL-PHYS-PUB-2016-017, 2016, URL: <https://cds.cern.ch/record/2206965>.
- [49] NNPDF Collaboration, R. D. Ball et al., *Parton distributions with LHC data*, *Nucl. Phys. B* **867** (2013) 244, arXiv: 1207.1303 [hep-ph].
- [50] E. Bothmann et al., *Event Generation with Sherpa 2.2*, *SciPost Phys.* **7** (2019) 034, arXiv: 1905.09127 [hep-ph].

- [51] D. J. Lange, *The EvtGen particle decay simulation package*, *Nucl. Instrum. Meth. A* **462** (2001) 152.
- [52] J. Alwall et al., *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, *JHEP* **07** (2014) 079, arXiv: [1405.0301 \[hep-ph\]](#).
- [53] T. Sjöstrand et al., *An introduction to PYTHIA 8.2*, *Comput. Phys. Commun.* **191** (2015) 159, arXiv: [1410.3012 \[hep-ph\]](#).
- [54] ATLAS Collaboration, *ATLAS Pythia 8 tunes to 7 TeV data*, ATL-PHYS-PUB-2014-021, 2014, URL: <https://cds.cern.ch/record/1966419>.
- [55] S. Frixione, G. Ridolfi and P. Nason, *A positive-weight next-to-leading-order Monte Carlo for heavy flavour hadroproduction*, *JHEP* **09** (2007) 126, arXiv: [0707.3088 \[hep-ph\]](#).
- [56] The NNPDF Collaboration, R. D. Ball et al., *Parton distributions for the LHC run II*, *JHEP* **04** (2015) 040, arXiv: [1410.8849 \[hep-ph\]](#).
- [57] M. Czakon and A. Mitov, *Top++: A program for the calculation of the top-pair cross-section at hadron colliders*, *Comput. Phys. Commun.* **185** (2014) 2930, arXiv: [1112.5675 \[hep-ph\]](#).
- [58] S. Alioli, P. Nason, C. Oleari and E. Re, *NLO single-top production matched with shower in POWHEG: s- and t-channel contributions*, *JHEP* **09** (2009) 111, arXiv: [0907.4076 \[hep-ph\]](#), Erratum: *JHEP* **02** (2010) 011.
- [59] N. Kidonakis, *NNLL resummation for s-channel single top quark production*, *Phys. Rev. D* **81** (2010) 054028, arXiv: [1001.5034 \[hep-ph\]](#).
- [60] N. Kidonakis, *Next-to-next-to-leading-order collinear and soft gluon corrections for t-channel single top quark production*, *Phys. Rev. D* **83** (2011) 091503, arXiv: [1103.2792 \[hep-ph\]](#).
- [61] E. Re, *Single-top Wt-channel production matched with parton showers using the POWHEG method*, *Eur. Phys. J. C* **71** (2011) 1547, arXiv: [1009.2450 \[hep-ph\]](#).
- [62] N. Kidonakis, *Two-loop soft anomalous dimensions for single top quark associated production with a W^- or H^-* , *Phys. Rev. D* **82** (2010) 054018, arXiv: [1005.4451 \[hep-ph\]](#).
- [63] E. Bothmann et al., *Event generation with Sherpa 2.2*, *SciPost Phys.* **7** (2019) 034, arXiv: [1905.09127 \[hep-ph\]](#).
- [64] T. Gleisberg and S. Höche, *Comix, a new matrix element generator*, *JHEP* **12** (2008) 039, arXiv: [0808.3674 \[hep-ph\]](#).
- [65] F. Cascioli, P. Maierhöfer and S. Pozzorini, *Scattering Amplitudes with Open Loops*, *Phys. Rev. Lett.* **108** (2012) 111601, arXiv: [1111.5206 \[hep-ph\]](#).
- [66] S. Schumann and F. Krauss, *A parton shower algorithm based on Catani–Seymour dipole factorisation*, *JHEP* **03** (2008) 038, arXiv: [0709.1027 \[hep-ph\]](#).
- [67] S. Höche, F. Krauss, M. Schönherr and F. Siegert, *QCD matrix elements + parton showers. The NLO case*, *JHEP* **04** (2013) 027, arXiv: [1207.5030 \[hep-ph\]](#).

- [68] S. Catani, L. Cieri, G. Ferrera, D. de Florian and M. Grazzini, *Vector Boson Production at Hadron Colliders: A Fully Exclusive QCD Calculation at Next-to-Next-to-Leading Order*, [Phys. Rev. Lett. **103** \(2009\) 082001](#), arXiv: [0903.2120 \[hep-ph\]](#).
- [69] S. Alioli, P. Nason, C. Oleari and E. Re, *A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX*, [JHEP **06** \(2010\) 043](#), arXiv: [1002.2581 \[hep-ph\]](#).
- [70] ATLAS Collaboration, *Measurement of the Z/γ^* boson transverse momentum distribution in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector*, [JHEP **09** \(2014\) 145](#), arXiv: [1406.3660 \[hep-ex\]](#).
- [71] J. R. Andersen et al., *Handbook of LHC Higgs Cross Sections: 3. Higgs Properties*, (2013), ed. by S. Heinemeyer, C. Mariotti, G. Passarino and R. Tanaka, arXiv: [1307.1347 \[hep-ph\]](#).
- [72] ATLAS Collaboration, *Electron and photon performance measurements with the ATLAS detector using the 2015–2017 LHC proton–proton collision data*, [JINST **14** \(2019\) P12006](#), arXiv: [1908.00005 \[hep-ex\]](#).
- [73] ATLAS Collaboration, *Muon reconstruction and identification efficiency in ATLAS using the full Run 2 pp collision data set at $\sqrt{s} = 13$ TeV*, [Eur. Phys. J. C **81** \(2021\) 578](#), arXiv: [2012.00578 \[hep-ex\]](#).
- [74] M. Cacciari, G. P. Salam and G. Soyez, *The anti- k_t jet clustering algorithm*, [JHEP **04** \(2008\) 063](#), arXiv: [0802.1189 \[hep-ph\]](#).
- [75] M. Cacciari, G. P. Salam and G. Soyez, *FastJet user manual*, [Eur. Phys. J. C **72** \(2012\) 1896](#), arXiv: [1111.6097 \[hep-ph\]](#).
- [76] ATLAS Collaboration, *Jet energy scale measurements and their systematic uncertainties in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, [Phys. Rev. D **96** \(2017\) 072002](#), arXiv: [1703.09665 \[hep-ex\]](#).
- [77] ATLAS Collaboration, *Performance of pile-up mitigation techniques for jets in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector*, [Eur. Phys. J. C **76** \(2016\) 581](#), arXiv: [1510.03823 \[hep-ex\]](#).
- [78] ATLAS Collaboration, *ATLAS b -jet identification performance and efficiency measurement with $t\bar{t}$ events in pp collisions at $\sqrt{s} = 13$ TeV*, [Eur. Phys. J. C **79** \(2019\) 970](#), arXiv: [1907.05120 \[hep-ex\]](#).
- [79] ATLAS Collaboration, *Evidence for the $H \rightarrow b\bar{b}$ decay with the ATLAS detector*, [JHEP **12** \(2017\) 024](#), arXiv: [1708.03299 \[hep-ex\]](#).
- [80] ATLAS Collaboration, *Identification and energy calibration of hadronically decaying tau leptons with the ATLAS experiment in pp collisions at $\sqrt{s} = 8$ TeV*, [Eur. Phys. J. C **75** \(2015\) 303](#), arXiv: [1412.7086 \[hep-ex\]](#).
- [81] ATLAS Collaboration, *Identification of hadronic tau lepton decays using neural networks in the ATLAS experiment*, ATL-PHYS-PUB-2019-033, 2019, URL: <https://cds.cern.ch/record/2688062>.
- [82] ATLAS Collaboration, *Measurement of the tau lepton reconstruction and identification performance in the ATLAS experiment using pp collisions at $\sqrt{s} = 13$ TeV*, ATLAS-CONF-2017-029, 2017, URL: <https://cds.cern.ch/record/2261772>.

- [83] ATLAS Collaboration, *Performance of missing transverse momentum reconstruction with the ATLAS detector using proton–proton collisions at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **78** (2018) 903, arXiv: [1802.08168](https://arxiv.org/abs/1802.08168) [[hep-ex](#)].
- [84] ATLAS Collaboration, *E_T^{miss} performance in the ATLAS detector using 2015–2016 LHC pp collisions*, ATLAS-CONF-2018-023, 2018, URL: <https://cds.cern.ch/record/2625233>.
- [85] ATLAS Collaboration, *Object-based missing transverse momentum significance in the ATLAS Detector*, ATLAS-CONF-2018-038, 2018, URL: <https://cds.cern.ch/record/2630948>.
- [86] A. Hoecker et al., *TMVA - Toolkit for Multivariate Data Analysis*, 2007, arXiv: [physics/0703039](https://arxiv.org/abs/physics/0703039) [[physics.data-an](#)].
- [87] ATLAS Collaboration, *Jet energy scale and resolution measured in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Eur. Phys. J. C* **81** (2020) 689, arXiv: [2007.02645](https://arxiv.org/abs/2007.02645) [[hep-ex](#)].
- [88] ATLAS Collaboration, *Measurement of b -tagging efficiency of c -jets in $t\bar{t}$ events using a likelihood approach with the ATLAS detector*, ATLAS-CONF-2018-001, 2018, URL: <https://cds.cern.ch/record/2306649>.
- [89] ATLAS Collaboration, *Calibration of light-flavour b -jet mistagging rates using ATLAS proton–proton collision data at $\sqrt{s} = 13$ TeV*, ATLAS-CONF-2018-006, 2018, URL: <https://cds.cern.ch/record/2314418>.
- [90] ATLAS Collaboration, *Electron reconstruction and identification in the ATLAS experiment using the 2015 and 2016 LHC proton–proton collision data at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **79** (2019) 639, arXiv: [1902.04655](https://arxiv.org/abs/1902.04655) [[hep-ex](#)].
- [91] ATLAS Collaboration, *Muon reconstruction performance of the ATLAS detector in proton–proton collision data at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **76** (2016) 292, arXiv: [1603.05598](https://arxiv.org/abs/1603.05598) [[hep-ex](#)].
- [92] ATLAS Collaboration, *Measurement of the Inelastic Proton–Proton Cross Section at $\sqrt{s} = 13$ TeV with the ATLAS Detector at the LHC*, *Phys. Rev. Lett.* **117** (2016) 182002, arXiv: [1606.02625](https://arxiv.org/abs/1606.02625) [[hep-ex](#)].
- [93] ATLAS Collaboration, *Measurement of Higgs boson decay into b -quarks in associated production with a top-quark pair in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *JHEP* **06** (2021) 097, arXiv: [2111.06712](https://arxiv.org/abs/2111.06712) [[hep-ex](#)].
- [94] E. Bothmann, M. Schönherr and S. Schumann, *Reweighting QCD matrix-element and parton-shower calculations*, *Eur. Phys. J. C* **76** (2016) 590, arXiv: [1606.08753](https://arxiv.org/abs/1606.08753) [[hep-ph](#)].
- [95] M. Bähr et al., *Herwig++ physics and manual*, *Eur. Phys. J. C* **58** (2008) 639, arXiv: [0803.0883](https://arxiv.org/abs/0803.0883) [[hep-ph](#)].
- [96] J. Bellm et al., *Herwig 7.0/Herwig++ 3.0 release note*, *Eur. Phys. J. C* **76** (2016) 196, arXiv: [1512.01178](https://arxiv.org/abs/1512.01178) [[hep-ph](#)].
- [97] ATLAS Collaboration, *Study of top-quark pair modelling and uncertainties using ATLAS measurements at $\sqrt{s} = 13$ TeV*, ATL-PHYS-PUB-2020-023, 2020, URL: <https://cds.cern.ch/record/2730443>.

- [98] S. Frixione, E. Laenen, P. Motylinski, C. White and B. R. Webber, *Single-top hadroproduction in association with a W boson*, [JHEP **07** \(2008\) 029](#), arXiv: [0805.3067 \[hep-ph\]](#).
- [99] S. Frixione, P. Nason and C. Oleari, *Matching NLO QCD computations with parton shower simulations: the POWHEG method*, [JHEP **11** \(2007\) 070](#), arXiv: [0709.2092 \[hep-ph\]](#).
- [100] M. Czakon et al., *Top-pair production at the LHC through NNLO QCD and NLO EW*, [JHEP **10** \(2017\) 186](#), arXiv: [1705.04105 \[hep-ph\]](#).
- [101] ATLAS Collaboration, *Measurements of top-quark pair differential cross-sections in the lepton+jets channel in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector*, [Eur. Phys. J. C **76** \(2016\) 538](#), arXiv: [1511.04716 \[hep-ex\]](#).
- [102] ATLAS Collaboration, *Combined search for the Standard Model Higgs boson in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector*, [Phys. Rev. D **86** \(2012\) 032003](#), arXiv: [1207.0319 \[hep-ex\]](#).
- [103] L. Moneta et al., *The RooStats Project*, [PoS **ACAT2010** \(2010\) 057](#), ed. by T. Speer et al., arXiv: [1009.1003 \[physics.data-an\]](#).
- [104] W. Verkerke and D. P. Kirkby, *The RooFit toolkit for data modeling*, eConf [C0303241](#) (2003) MOLT007, ed. by L. Lyons and M. Karagoz, arXiv: [physics/0306116](#).
- [105] A. L. Read, *Presentation of search results: the CL_s technique*, [J. Phys. G **28** \(2002\) 2693](#).
- [106] G. Cowan, K. Cranmer, E. Gross and O. Vitells, *Asymptotic formulae for likelihood-based tests of new physics*, [Eur. Phys. J. C **71** \(2011\) 1554](#), arXiv: [1007.1727 \[physics.data-an\]](#), Erratum: [Eur. Phys. J. C **73** \(2013\) 2501](#).
- [107] E. Gross and O. Vitells, *Trial factors for the look elsewhere effect in high energy physics*, [Eur. Phys. J. C **70** \(2010\) 525](#), arXiv: [1005.1891 \[physics.data-an\]](#).
- [108] ATLAS Collaboration, *Formulae for Estimating Significance*, ATL-PHYS-PUB-2020-025, 2020, URL: <https://cds.cern.ch/record/2736148>.
- [109] P. Sikivie, L. Susskind, M. Voloshin and V. Zakharov, *Isospin breaking in technicolor models*, [Nucl. Phys. B **173** \(1980\) 189](#).
- [110] J. F. Gunion and H. E. Haber, *CP-conserving two-Higgs-doublet model: the approach to the decoupling limit*, [Phys. Rev. D **67** \(2003\) 075019](#), arXiv: [hep-ph/0207010](#).
- [111] R. V. Harlander, S. Liebler and H. Mantler, *SusHi: A program for the calculation of Higgs production in gluon fusion and bottom-quark annihilation in the Standard Model and the MSSM*, [Comput. Phys. Commun. **184** \(2013\) 1605](#), arXiv: [1212.3249 \[hep-ph\]](#).
- [112] R. V. Harlander and P. Kant, *Higgs production and decay: analytic results at next-to-leading order QCD*, [JHEP **12** \(2005\) 015](#), arXiv: [hep-ph/0509189](#).
- [113] R. V. Harlander and W. B. Kilgore, *Higgs boson production in bottom quark fusion at next-to-next-to-leading order*, [Phys. Rev. D **68** \(2003\) 013001](#), arXiv: [hep-ph/0304035](#).

- [114] R. V. Harlander and W. B. Kilgore,
Next-to-Next-to-Leading Order Higgs Production at Hadron Colliders,
Phys. Rev. Lett. **88** (2002) 201801, arXiv: [hep-ph/0201206](https://arxiv.org/abs/hep-ph/0201206).
- [115] D. Eriksson, J. Rathsman and O. Stål,
2HDMC - two-Higgs-doublet model calculator physics and manual,
Comput. Phys. Commun. **181** (2010) 189, arXiv: [0902.0851](https://arxiv.org/abs/0902.0851) [[hep-ph](#)].
- [116] ATLAS Collaboration, *ATLAS Computing Acknowledgements*, ATL-SOFT-PUB-2021-003, 2021,
URL: <https://cds.cern.ch/record/2776662>.

The ATLAS Collaboration

G. Aad ¹⁰¹, B. Abbott ¹¹⁹, D.C. Abbott ¹⁰², K. Abeling ⁵⁵, S.H. Abidi ²⁹, A. Aboulhorma ^{35e}, H. Abramowicz ¹⁵⁰, H. Abreu ¹⁴⁹, Y. Abulaiti ¹¹⁶, A.C. Abusleme Hoffman ^{136a}, B.S. Acharya ^{68a,68b,o}, B. Achkar ⁵⁵, C. Adam Bourdarios ⁴, L. Adamczyk ^{84a}, L. Adamek ¹⁵⁴, S.V. Addepalli ²⁶, J. Adelman ¹¹⁴, A. Adiguzel ^{21c}, S. Adorni ⁵⁶, T. Adye ¹³³, A.A. Affolder ¹³⁵, Y. Afik ³⁶, M.N. Agaras ¹³, J. Agarwala ^{72a,72b}, A. Aggarwal ⁹⁹, C. Agheorghiesei ^{27c}, J.A. Aguilar-Saavedra ^{129f}, A. Ahmad ³⁶, F. Ahmadov ^{38,x}, W.S. Ahmed ¹⁰³, S. Ahuja ⁹⁴, X. Ai ⁴⁸, G. Aielli ^{75a,75b}, I. Aizenberg ¹⁶⁷, M. Akbiyik ⁹⁹, T.P.A. Åkesson ⁹⁷, A.V. Akimov ³⁷, K. Al Khoury ⁴¹, G.L. Alberghi ^{23b}, J. Albert ¹⁶³, P. Albicocco ⁵³, S. Alderweireldt ⁵², M. Aleksa ³⁶, I.N. Aleksandrov ³⁸, C. Alexa ^{27b}, T. Alexopoulos ¹⁰, A. Alfonsi ¹¹³, F. Alfonsi ^{23b}, M. Alhroob ¹¹⁹, B. Ali ¹³¹, S. Ali ¹⁴⁷, M. Aliev ³⁷, G. Alimonti ^{70a}, W. Alkahi ⁵⁵, C. Allaire ⁶⁶, B.M.M. Allbrooke ¹⁴⁵, P.P. Allport ²⁰, A. Aloisio ^{71a,71b}, F. Alonso ⁸⁹, C. Alpigiani ¹³⁷, E. Alunno Camelia ^{75a,75b}, M. Alvarez Estevez ⁹⁸, M.G. Alvigi ^{71a,71b}, M. Aly ¹⁰⁰, Y. Amaral Coutinho ^{81b}, A. Ambler ¹⁰³, C. Amelung ³⁶, M. Amerl ¹, C.G. Ames ¹⁰⁸, D. Amidei ¹⁰⁵, S.P. Amor Dos Santos ^{129a}, S. Amoroso ⁴⁸, K.R. Amos ¹⁶¹, V. Ananiev ¹²⁴, C. Anastopoulos ¹³⁸, T. Andeen ¹¹, J.K. Anders ³⁶, S.Y. Andrean ^{47a,47b}, A. Andreatza ^{70a,70b}, S. Angelidakis ⁹, A. Angerami ^{41,z}, A.V. Anisenkov ³⁷, A. Annovi ^{73a}, C. Antel ⁵⁶, M.T. Anthony ¹³⁸, E. Antipov ¹²⁰, M. Antonelli ⁵³, D.J.A. Antrim ^{17a}, F. Anulli ^{74a}, M. Aoki ⁸², T. Aoki ¹⁵², J.A. Aparisi Pozo ¹⁶¹, M.A. Aparo ¹⁴⁵, L. Aperio Bella ⁴⁸, C. Appelt ¹⁸, N. Aranzabal ³⁶, V. Araujo Ferraz ^{81a}, C. Arcangeletti ⁵³, A.T.H. Arce ⁵¹, E. Arena ⁹¹, J-F. Arguin ¹⁰⁷, S. Argyropoulos ⁵⁴, J.-H. Arling ⁴⁸, A.J. Armbruster ³⁶, O. Arnaez ¹⁵⁴, H. Arnold ¹¹³, Z.P. Arrubarrena Tame ¹⁰⁸, G. Artoni ^{74a,74b}, H. Asada ¹¹⁰, K. Asai ¹¹⁷, S. Asai ¹⁵², N.A. Asbah ⁶¹, J. Assahsah ^{35d}, K. Assamagan ²⁹, R. Astalos ^{28a}, R.J. Atkin ^{33a}, M. Atkinson ¹⁶⁰, N.B. Atlay ¹⁸, H. Atmani ^{62b}, P.A. Atmasiddha ¹⁰⁵, K. Augsten ¹³¹, S. Auricchio ^{71a,71b}, A.D. Auriol ²⁰, V.A. Austrup ¹⁶⁹, G. Avner ¹⁴⁹, G. Avolio ³⁶, K. Axiotis ⁵⁶, M.K. Ayoub ^{14c}, G. Azuelos ^{107,ac}, D. Babal ^{28a}, H. Bachacou ¹³⁴, K. Bachas ^{151,q}, A. Bachiu ³⁴, F. Backman ^{47a,47b}, A. Badea ⁶¹, P. Bagnaia ^{74a,74b}, M. Bahmani ¹⁸, A.J. Bailey ¹⁶¹, V.R. Bailey ¹⁶⁰, J.T. Baines ¹³³, C. Bakalis ¹⁰, O.K. Baker ¹⁷⁰, P.J. Bakker ¹¹³, E. Bakos ¹⁵, D. Bakshi Gupta ⁸, S. Balaji ¹⁴⁶, R. Balasubramanian ¹¹³, E.M. Baldin ³⁷, P. Balek ¹³², E. Ballabene ^{70a,70b}, F. Balli ¹³⁴, L.M. Baltes ^{63a}, W.K. Balunas ³², J. Balz ⁹⁹, E. Banas ⁸⁵, M. Bandieramonte ¹²⁸, A. Bandyopadhyay ²⁴, S. Bansal ²⁴, L. Barak ¹⁵⁰, E.L. Barberio ¹⁰⁴, D. Barberis ^{57b,57a}, M. Barbero ¹⁰¹, G. Barbour ⁹⁵, K.N. Barends ^{33a}, T. Barillari ¹⁰⁹, M-S. Barisits ³⁶, T. Barklow ¹⁴², R.M. Barnett ^{17a}, P. Baron ¹²¹, D.A. Baron Moreno ¹⁰⁰, A. Baroncelli ^{62a}, G. Barone ²⁹, A.J. Barr ¹²⁵, L. Barranco Navarro ^{47a,47b}, F. Barreiro ⁹⁸, J. Barreiro Guimarães da Costa ^{14a}, U. Barron ¹⁵⁰, M.G. Barros Teixeira ^{129a}, S. Barsov ³⁷, F. Bartels ^{63a}, R. Bartoldus ¹⁴², A.E. Barton ⁹⁰, P. Bartos ^{28a}, A. Basalae ⁴⁸, A. Basan ⁹⁹, M. Baselga ⁴⁹, I. Bashta ^{76a,76b}, A. Bassalat ⁶⁶, M.J. Basso ¹⁵⁴, C.R. Basson ¹⁰⁰, R.L. Bates ⁵⁹, S. Batlamous ^{35e}, J.R. Batley ³², B. Batool ¹⁴⁰, M. Battaglia ¹³⁵, D. Battulga ¹⁸, M. Baucé ^{74a,74b}, P. Bauer ²⁴, A. Bayirli ^{21a}, J.B. Beacham ⁵¹, T. Beau ¹²⁶, P.H. Beauchemin ¹⁵⁷, F. Becherer ⁵⁴, P. Bechtel ²⁴, H.P. Beck ^{19,p}, K. Becker ¹⁶⁵, A.J. Beddall ^{21d}, V.A. Bednyakov ³⁸, C.P. Bee ¹⁴⁴, L.J. Beemster ¹⁵, T.A. Beermann ³⁶, M. Begalli ^{81d,81d}, M. Begel ²⁹, A. Behera ¹⁴⁴, J.K. Behr ⁴⁸, C. Beirao Da Cruz E Silva ³⁶, J.F. Beirer ^{55,36}, F. Beisiegel ²⁴, M. Belfkir ^{115b}, G. Bella ¹⁵⁰, L. Bellagamba ^{23b}, A. Bellerive ³⁴, P. Bellos ²⁰, K. Beloborodov ³⁷, K. Belotskiy ³⁷, N.L. Belyaev ³⁷, D. Benckekroun ^{35a}, F. Bendebba ^{35a}, Y. Benhammou ¹⁵⁰, D.P. Benjamin ²⁹,

M. Benoit ²⁹, J.R. Bensinger ²⁶, S. Bentvelsen ¹¹³, L. Beresford ³⁶, M. Beretta ⁵³, D. Berge ¹⁸,
E. Bergeaas Kuutmann ¹⁵⁹, N. Berger ⁴, B. Bergmann ¹³¹, J. Beringer ^{17a}, S. Berlendis ⁷,
G. Bernardi ⁵, C. Bernius ¹⁴², F.U. Bernlochner ²⁴, T. Berry ⁹⁴, P. Berta ¹³², A. Berthold ⁵⁰,
I.A. Bertram ⁹⁰, S. Bethke ¹⁰⁹, A. Betti ^{74a,74b}, A.J. Bevan ⁹³, M. Bhamjee ^{33c}, S. Bhatta ¹⁴⁴,
D.S. Bhattacharya ¹⁶⁴, P. Bhattarai ²⁶, V.S. Bhopatkar ¹²⁰, R. Bi ^{29,af}, R.M. Bianchi ¹²⁸,
O. Biebel ¹⁰⁸, R. Bielski ¹²², M. Biglietti ^{76a}, T.R.V. Billoud ¹³¹, M. Bindi ⁵⁵, A. Bingul ^{21b},
C. Bini ^{74a,74b}, S. Biondi ^{23b,23a}, A. Biondini ⁹¹, C.J. Birch-sykes ¹⁰⁰, G.A. Bird ^{20,133},
M. Birman ¹⁶⁷, T. Bisanz ³⁶, E. Bisceglie ^{43b,43a}, D. Biswas ^{168,k}, A. Bitadze ¹⁰⁰, K. Bjørke ¹²⁴,
I. Bloch ⁴⁸, C. Blocker ²⁶, A. Blue ⁵⁹, U. Blumenschein ⁹³, J. Blumenthal ⁹⁹, G.J. Bobbink ¹¹³,
V.S. Bobrovnikov ³⁷, M. Boehler ⁵⁴, D. Bogavac ³⁶, A.G. Bogdanchikov ³⁷, C. Bohm ^{47a},
V. Boisvert ⁹⁴, P. Bokan ⁴⁸, T. Bold ^{84a}, M. Bomben ⁵, M. Bona ⁹³, M. Boonekamp ¹³⁴,
C.D. Booth ⁹⁴, A.G. Borbély ⁵⁹, H.M. Borecka-Bielska ¹⁰⁷, L.S. Borgna ⁹⁵, G. Borissov ⁹⁰,
D. Bortoletto ¹²⁵, D. Boscherini ^{23b}, M. Bosman ¹³, J.D. Bossio Sola ³⁶, K. Bouaouda ^{35a},
N. Bouchhar ¹⁶¹, J. Boudreau ¹²⁸, E.V. Bouhova-Thacker ⁹⁰, D. Boumediene ⁴⁰, R. Bouquet ⁵,
A. Boveia ¹¹⁸, J. Boyd ³⁶, D. Boye ²⁹, I.R. Boyko ³⁸, J. Bracinik ²⁰, N. Brahimi ^{62d},
G. Brandt ¹⁶⁹, O. Brandt ³², F. Braren ⁴⁸, B. Brau ¹⁰², J.E. Brau ¹²², K. Brendlinger ⁴⁸,
R. Brenner ¹⁶⁷, L. Brenner ¹¹³, R. Brenner ¹⁵⁹, S. Bressler ¹⁶⁷, B. Brickwedde ⁹⁹, D. Britton ⁵⁹,
D. Britzger ¹⁰⁹, I. Brock ²⁴, G. Brooijmans ⁴¹, W.K. Brooks ^{136f}, E. Brost ²⁹, T.L. Bruckler ¹²⁵,
P.A. Bruckman de Renstrom ⁸⁵, B. Brüers ⁴⁸, D. Bruncko ^{28b,*}, A. Bruni ^{23b}, G. Bruni ^{23b},
M. Bruschi ^{23b}, N. Bruscinò ^{74a,74b}, L. Bryngemark ¹⁴², T. Buanes ¹⁶, Q. Buat ¹³⁷,
P. Buchholz ¹⁴⁰, A.G. Buckley ⁵⁹, I.A. Budagov ^{38,*}, M.K. Bugge ¹²⁴, O. Bulekov ³⁷,
B.A. Bullard ⁶¹, S. Burdin ⁹¹, C.D. Burgard ⁴⁸, A.M. Burger ⁴⁰, B. Burghgrave ⁸, J.T.P. Burr ³²,
C.D. Burton ¹¹, J.C. Burzynski ¹⁴¹, E.L. Busch ⁴¹, V. Büscher ⁹⁹, P.J. Bussey ⁵⁹, J.M. Butler ²⁵,
C.M. Buttar ⁵⁹, J.M. Butterworth ⁹⁵, W. Buttinger ¹³³, C.J. Buxo Vazquez ¹⁰⁶, A.R. Buzykaev ³⁷,
G. Cabras ^{23b}, S. Cabrera Urbán ¹⁶¹, D. Caforio ⁵⁸, H. Cai ¹²⁸, Y. Cai ^{14a,14d}, V.M.M. Cairo ³⁶,
O. Cakir ^{3a}, N. Calace ³⁶, P. Calafiura ^{17a}, G. Calderini ¹²⁶, P. Calfayan ⁶⁷, G. Callea ⁵⁹,
L.P. Caloba ^{81b}, D. Calvet ⁴⁰, S. Calvet ⁴⁰, T.P. Calvet ¹⁰¹, M. Calvetti ^{73a,73b},
R. Camacho Toro ¹²⁶, S. Camarda ³⁶, D. Camarero Munoz ²⁶, P. Camarri ^{75a,75b},
M.T. Camerlingo ^{76a,76b}, D. Cameron ¹²⁴, C. Camincher ¹⁶³, M. Campanelli ⁹⁵, A. Camplani ⁴²,
V. Canale ^{71a,71b}, A. Canesse ¹⁰³, M. Cano Bret ⁷⁹, J. Cantero ¹⁶¹, Y. Cao ¹⁶⁰, F. Capocasa ²⁶,
M. Capua ^{43b,43a}, A. Carbone ^{70a,70b}, R. Cardarelli ^{75a}, J.C.J. Cardenas ⁸, F. Cardillo ¹⁶¹,
T. Carli ³⁶, G. Carlino ^{71a}, J.I. Carlotto ¹³, B.T. Carlson ^{128,r}, E.M. Carlson ^{163,155a},
L. Carminati ^{70a,70b}, M. Carnesale ^{74a,74b}, S. Caron ¹¹², E. Carquin ^{136f}, S. Carrá ^{70a,70b},
G. Carratta ^{23b,23a}, F. Carri Argos ^{33g}, J.W.S. Carter ¹⁵⁴, T.M. Carter ⁵², M.P. Casado ^{13,h},
A.F. Casha ¹⁵⁴, E.G. Castiglia ¹⁷⁰, F.L. Castillo ^{63a}, L. Castillo Garcia ¹³, V. Castillo Gimenez ¹⁶¹,
N.F. Castro ^{129a,129e}, A. Catinaccio ³⁶, J.R. Catmore ¹²⁴, V. Cavaliere ²⁹, N. Cavalli ^{23b,23a},
V. Cavasinni ^{73a,73b}, E. Celebi ^{21a}, F. Celli ¹²⁵, M.S. Centonze ^{69a,69b}, K. Cerny ¹²¹,
A.S. Cerqueira ^{81a}, A. Cerri ¹⁴⁵, L. Cerrito ^{75a,75b}, F. Cerutti ^{17a}, A. Cervelli ^{23b}, S.A. Cetin ^{21d},
Z. Chadi ^{35a}, D. Chakraborty ¹¹⁴, M. Chala ^{129f}, J. Chan ¹⁶⁸, W.Y. Chan ¹⁵², J.D. Chapman ³²,
B. Chargeishvili ^{148b}, D.G. Charlton ²⁰, T.P. Charman ⁹³, M. Chatterjee ¹⁹, S. Chekanov ⁶,
S.V. Chekulaev ^{155a}, G.A. Chelkov ^{38,a}, A. Chen ¹⁰⁵, B. Chen ¹⁵⁰, B. Chen ¹⁶³, C. Chen ^{62a},
H. Chen ^{14c}, H. Chen ²⁹, J. Chen ^{62c}, J. Chen ²⁶, S. Chen ¹⁵², S.J. Chen ^{14c}, X. Chen ^{62c},
X. Chen ^{14b,ab}, Y. Chen ^{62a}, C.L. Cheng ¹⁶⁸, H.C. Cheng ^{64a}, S. Cheong ¹⁴², A. Cheplakov ³⁸,
E. Cheremushkina ⁴⁸, E. Cherepanova ¹¹³, R. Cherkaoui El Moursli ^{35e}, E. Cheu ⁷, K. Cheung ⁶⁵,
L. Chevalier ¹³⁴, V. Chiarella ⁵³, G. Chiarelli ^{73a}, N. Chiedde ¹⁰¹, G. Chiodini ^{69a},
A.S. Chisholm ²⁰, A. Chitan ^{27b}, M. Chitishvili ¹⁶¹, Y.H. Chiu ¹⁶³, M.V. Chizhov ³⁸, K. Choi ¹¹,
A.R. Chomont ^{74a,74b}, Y. Chou ¹⁰², E.Y.S. Chow ¹¹³, T. Chowdhury ^{33g}, L.D. Christopher ^{33g},

K.L. Chu^{64a}, M.C. Chu^{64a}, X. Chu^{14a,14d}, J. Chudoba¹³⁰, J.J. Chwastowski⁸⁵, D. Cieri¹⁰⁹,
 K.M. Ciesla^{84a}, V. Cindro⁹², A. Ciocio^{17a}, F. Ciroto^{71a,71b}, Z.H. Citron^{167,1}, M. Citterio^{70a},
 D.A. Ciubotaru^{27b}, B.M. Ciungu¹⁵⁴, A. Clark⁵⁶, P.J. Clark⁵², J.M. Clavijo Columbie⁴⁸,
 S.E. Clawson¹⁰⁰, C. Clement^{47a,47b}, J. Clercx⁴⁸, L. Clissa^{23b,23a}, Y. Coadou¹⁰¹,
 M. Cobal^{68a,68c}, A. Coccaro^{57b}, R.F. Coelho Barrue^{129a}, R. Coelho Lopes De Sa¹⁰²,
 S. Coelli^{70a}, H. Cohen¹⁵⁰, A.E.C. Coimbra^{70a,70b}, B. Cole⁴¹, J. Collot⁶⁰,
 P. Conde Muiño^{129a,129g}, M.P. Connell^{33c}, S.H. Connell^{33c}, I.A. Connelly⁵⁹, E.I. Conroy¹²⁵,
 F. Conventi^{71a,ad}, H.G. Cooke²⁰, A.M. Cooper-Sarkar¹²⁵, F. Cormier¹⁶², L.D. Corpe³⁶,
 M. Corradi^{74a,74b}, E.E. Corrigan⁹⁷, F. Corriveau^{103,v}, A. Cortes-Gonzalez¹⁸, M.J. Costa¹⁶¹,
 F. Costanza⁴, D. Costanzo¹³⁸, B.M. Cote¹¹⁸, G. Cowan⁹⁴, J.W. Cowley³², K. Cranmer¹¹⁶,
 S. Crépe-Renaudin⁶⁰, F. Crescioli¹²⁶, M. Cristinziani¹⁴⁰, M. Cristoforetti^{77a,77b,c}, V. Croft¹⁵⁷,
 G. Crosetti^{43b,43a}, A. Cueto³⁶, T. Cuhadar Donszelmann¹⁵⁸, H. Cui^{14a,14d}, Z. Cui⁷,
 A.R. Cukierman¹⁴², W.R. Cunningham⁵⁹, F. Curcio^{43b,43a}, P. Czodrowski³⁶, M.M. Czurylo^{63b},
 M.J. Da Cunha Sargedas De Sousa^{62a}, J.V. Da Fonseca Pinto^{81b}, C. Da Via¹⁰⁰, W. Dabrowski^{84a},
 T. Dado⁴⁹, S. Dahbi^{33g}, T. Dai¹⁰⁵, C. Dallapiccola¹⁰², M. Dam⁴², G. D'amen²⁹,
 V. D'Amico¹⁰⁸, J. Damp⁹⁹, J.R. Dandoy¹²⁷, M.F. Daneri³⁰, M. Danninger¹⁴¹, V. Dao³⁶,
 G. Darbo^{57b}, S. Darmora⁶, S.J. Das²⁹, S. D'Auria^{70a,70b}, C. David^{155b}, T. Davidek¹³²,
 D.R. Davis⁵¹, B. Davis-Purcell³⁴, I. Dawson⁹³, K. De⁸, R. De Asmundis^{71a},
 M. De Beurs¹¹³, N. De Biase⁴⁸, S. De Castro^{23b,23a}, N. De Groot¹¹², P. de Jong¹¹³,
 H. De la Torre¹⁰⁶, A. De Maria^{14c}, A. De Salvo^{74a}, U. De Sanctis^{75a,75b}, A. De Santo¹⁴⁵,
 J.B. De Vivie De Regie⁶⁰, D.V. Dedovich³⁸, J. Degens¹¹³, A.M. Deiana⁴⁴, F. Del Corso^{23b,23a},
 J. Del Peso⁹⁸, F. Del Rio^{63a}, F. Deliot¹³⁴, C.M. Delitzsch⁴⁹, M. Della Pietra^{71a,71b},
 D. Della Volpe⁵⁶, A. Dell'Acqua³⁶, L. Dell'Asta^{70a,70b}, M. Delmastro⁴, P.A. Delsart⁶⁰,
 S. Demers¹⁷⁰, M. Demichev³⁸, S.P. Denisov³⁷, L. D'Eramo¹¹⁴, D. Derendarz⁸⁵,
 F. Derue¹²⁶, P. Dervan⁹¹, K. Desch²⁴, K. Dette¹⁵⁴, C. Deutsch²⁴, P.O. Deviveiros³⁶,
 F.A. Di Bello^{57b,57a}, A. Di Ciaccio^{75a,75b}, L. Di Ciaccio⁴, A. Di Domenico^{74a,74b},
 C. Di Donato^{71a,71b}, A. Di Girolamo³⁶, G. Di Gregorio⁵, A. Di Luca^{77a,77b},
 B. Di Micco^{76a,76b}, R. Di Nardo^{76a,76b}, C. Diaconu¹⁰¹, F.A. Dias¹¹³, T. Dias Do Vale¹⁴¹,
 M.A. Diaz^{136a,136b}, F.G. Diaz Capriles²⁴, M. Didenko¹⁶¹, E.B. Diehl¹⁰⁵, L. Diehl⁵⁴,
 S. Díez Cornell⁴⁸, C. Diez Pardos¹⁴⁰, C. Dimitriadi^{24,159}, A. Dimitrievska^{17a}, W. Ding^{14b},
 J. Dingfelder²⁴, I-M. Dinu^{27b}, S.J. Dittmeier^{63b}, F. Dittus³⁶, F. Djama¹⁰¹, T. Djobava^{148b},
 J.I. Djuvsland¹⁶, C. Doglioni^{100,97}, J. Dolejsi¹³², Z. Dolezal¹³², M. Donadelli^{81c},
 B. Dong^{62c}, J. Donini⁴⁰, A. D'Onofrio^{14c}, M. D'Onofrio⁹¹, J. Dopke¹³³, A. Doria^{71a},
 M.T. Dova⁸⁹, A.T. Doyle⁵⁹, M.A. Draguet¹²⁵, E. Drechsler¹⁴¹, E. Dreyer¹⁶⁷,
 I. Drivas-koulouris¹⁰, A.S. Drobac¹⁵⁷, M. Drozdova⁵⁶, D. Du^{62a}, T.A. du Pree¹¹³,
 F. Dubinin³⁷, M. Dubovsky^{28a}, E. Duchovni¹⁶⁷, G. Duckeck¹⁰⁸, O.A. Ducu^{27b}, D. Duda¹⁰⁹,
 A. Dudarev³⁶, M. D'uffizi¹⁰⁰, L. Dufлот⁶⁶, M. Dührssen³⁶, C. Dülsen¹⁶⁹, A.E. Dumitriu^{27b},
 M. Dunford^{63a}, S. Dungs⁴⁹, K. Dunne^{47a,47b}, A. Duperrin¹⁰¹, H. Duran Yildiz^{3a},
 M. Düren⁵⁸, A. Durglishvili^{148b}, B.L. Dwyer¹¹⁴, G.I. Dyckes^{17a}, M. Dyndal^{84a},
 S. Dysch¹⁰⁰, B.S. Dziedzic⁸⁵, Z.O. Earnshaw¹⁴⁵, B. Eckerova^{28a}, M.G. Eggleston⁵¹,
 E. Egidio Purcino De Souza^{81b}, L.F. Ehrke⁵⁶, G. Eigen¹⁶, K. Einsweiler^{17a}, T. Ekelof¹⁵⁹,
 P.A. Ekman⁹⁷, Y. El Ghazali^{35b}, H. El Jarrari^{35c,147}, A. El Moussaouy^{35a}, V. Ellajosyula¹⁵⁹,
 M. Ellert¹⁵⁹, F. Ellinghaus¹⁶⁹, A.A. Elliot⁹³, N. Ellis³⁶, J. Elmsheuser²⁹, M. Elsing³⁶,
 D. Emelianov¹³³, A. Emerman⁴¹, Y. Enari¹⁵², I. Ene^{17a}, S. Epari¹³, J. Erdmann^{49,aa},
 A. Ereditato¹⁹, P.A. Erland⁸⁵, M. Errenst¹⁶⁹, M. Escalier⁶⁶, C. Escobar¹⁶¹, E. Etzion¹⁵⁰,
 G. Evans^{129a}, H. Evans⁶⁷, M.O. Evans¹⁴⁵, A. Ezhilov³⁷, S. Ezzarqtouni^{35a}, F. Fabbri⁵⁹,
 L. Fabbri^{23b,23a}, G. Facini⁹⁵, V. Fadeyev¹³⁵, R.M. Fakhruddinov³⁷, S. Falciano^{74a},

P.J. Falke [ID24](#), S. Falke [ID36](#), J. Faltova [ID132](#), Y. Fan [ID14a](#), Y. Fang [ID14a,14d](#), G. Fanourakis [ID46](#),
 M. Fanti [ID70a,70b](#), M. Faraj [ID68a,68b](#), Z. Farazpay [ID96](#), A. Farbin [ID8](#), A. Farilla [ID76a](#), T. Farooque [ID106](#),
 S.M. Farrington [ID52](#), F. Fassi [ID35e](#), D. Fassouliotis [ID9](#), M. Faucci Giannelli [ID75a,75b](#), W.J. Fawcett [ID32](#),
 L. Fayard [ID66](#), P. Federicova [ID130](#), O.L. Fedin [ID37,a](#), G. Fedotov [ID37](#), M. Feickert [ID168](#), L. Feligioni [ID101](#),
 A. Fell [ID138](#), D.E. Fellers [ID122](#), C. Feng [ID62b](#), M. Feng [ID14b](#), Z. Feng [ID113](#), M.J. Fenton [ID158](#),
 A.B. Fenyuk [ID37](#), L. Ferencz [ID48](#), S.W. Ferguson [ID45](#), J.A. Fernandez Pretel [ID54](#), J. Ferrando [ID48](#),
 A. Ferrari [ID159](#), P. Ferrari [ID113,112](#), R. Ferrari [ID72a](#), D. Ferrere [ID56](#), C. Ferretti [ID105](#), F. Fiedler [ID99](#),
 A. Filipčič [ID92](#), E.K. Filmer [ID1](#), F. Filthaut [ID112](#), M.C.N. Fiolhais [ID129a,129c,b](#), L. Fiorini [ID161](#),
 F. Fischer [ID140](#), W.C. Fisher [ID106](#), T. Fitschen [ID100](#), I. Fleck [ID140](#), P. Fleischmann [ID105](#), T. Flick [ID169](#),
 L. Flores [ID127](#), M. Flores [ID33d](#), L.R. Flores Castillo [ID64a](#), F.M. Follega [ID77a,77b](#), N. Fomin [ID16](#),
 J.H. Foo [ID154](#), B.C. Forland [ID67](#), A. Formica [ID134](#), A.C. Forti [ID100](#), E. Fortin [ID101](#), A.W. Fortman [ID61](#),
 M.G. Foti [ID17a](#), L. Fountas [ID9](#), D. Fournier [ID66](#), H. Fox [ID90](#), P. Francavilla [ID73a,73b](#), S. Francescato [ID61](#),
 S. Franchellucci [ID56](#), M. Franchini [ID23b,23a](#), S. Franchino [ID63a](#), D. Francis [ID36](#), L. Franco [ID112](#),
 L. Franconi [ID19](#), M. Franklin [ID61](#), G. Frattari [ID26](#), A.C. Freegard [ID93](#), P.M. Freeman [ID20](#), W.S. Freund [ID81b](#),
 N. Fritzsche [ID50](#), A. Froch [ID54](#), D. Froidevaux [ID36](#), J.A. Frost [ID125](#), Y. Fu [ID62a](#), M. Fujimoto [ID117](#),
 E. Fullana Torregrosa [ID161,*](#), J. Fuster [ID161](#), A. Gabrielli [ID23b,23a](#), A. Gabrielli [ID154](#), P. Gadow [ID48](#),
 G. Gagliardi [ID57b,57a](#), L.G. Gagnon [ID17a](#), G.E. Gallardo [ID125](#), E.J. Gallas [ID125](#), B.J. Gallop [ID133](#),
 R. Gamboa Goni [ID93](#), K.K. Gan [ID118](#), S. Ganguly [ID152](#), J. Gao [ID62a](#), Y. Gao [ID52](#),
 F.M. Garay Walls [ID136a,136b](#), B. Garcia [ID29,af](#), C. García [ID161](#), J.E. García Navarro [ID161](#),
 J.A. García Pascual [ID14a](#), M. Garcia-Sciveres [ID17a](#), R.W. Gardner [ID39](#), D. Garg [ID79](#), R.B. Garg [ID142](#),
 S. Gargiulo [ID54](#), C.A. Garner [ID154](#), V. Garonne [ID29](#), S.J. Gasiorowski [ID137](#), P. Gaspar [ID81b](#), G. Gaudio [ID72a](#),
 V. Gautam [ID13](#), P. Gauzzi [ID74a,74b](#), I.L. Gavrilenko [ID37](#), A. Gavrilyuk [ID37](#), C. Gay [ID162](#), G. Gaycken [ID48](#),
 E.N. Gazis [ID10](#), A.A. Geanta [ID27b,27e](#), C.M. Gee [ID135](#), J. Geisen [ID97](#), M. Geisen [ID99](#), C. Gemme [ID57b](#),
 M.H. Genest [ID60](#), S. Gentile [ID74a,74b](#), S. George [ID94](#), W.F. George [ID20](#), T. Geralis [ID46](#), L.O. Gerlach [ID55](#),
 P. Gessinger-Befurt [ID36](#), M. Ghasemi Bostanabad [ID163](#), M. Ghneimat [ID140](#), K. Ghorbanian [ID93](#),
 A. Ghosal [ID140](#), A. Ghosh [ID158](#), A. Ghosh [ID7](#), B. Giacobbe [ID23b](#), S. Giagu [ID74a,74b](#),
 N. Giangiacomi [ID154](#), P. Giannetti [ID73a](#), A. Giannini [ID62a](#), S.M. Gibson [ID94](#), M. Gignac [ID135](#),
 D.T. Gil [ID84b](#), A.K. Gilbert [ID84a](#), B.J. Gilbert [ID41](#), D. Gillberg [ID34](#), G. Gilles [ID113](#), N.E.K. Gillwald [ID48](#),
 L. Ginabat [ID126](#), D.M. Gingrich [ID2,ac](#), M.P. Giordani [ID68a,68c](#), P.F. Giraud [ID134](#), G. Giugliarelli [ID68a,68c](#),
 D. Giugni [ID70a](#), F. Giuli [ID36](#), I. Gkialas [ID9,i](#), L.K. Gladilin [ID37](#), C. Glasman [ID98](#), G.R. Gledhill [ID122](#),
 M. Glisic [ID122](#), I. Gnesi [ID43b,e](#), Y. Go [ID29,af](#), M. Goblirsch-Kolb [ID26](#), B. Gocke [ID49](#), D. Godin [ID107](#),
 S. Goldfarb [ID104](#), T. Golling [ID56](#), M.G.D. Gololo [ID33g](#), D. Golubkov [ID37](#), J.P. Gombas [ID106](#),
 A. Gomes [ID129a,129b](#), G. Gomes Da Silva [ID140](#), A.J. Gomez Delegido [ID161](#), R. Goncalves Gama [ID55](#),
 R. Gonçalves [ID129a,129c](#), G. Gonella [ID122](#), L. Gonella [ID20](#), A. Gongadze [ID38](#), F. Gonnella [ID20](#),
 J.L. Gonski [ID41](#), S. González de la Hoz [ID161](#), S. Gonzalez Fernandez [ID13](#), R. Gonzalez Lopez [ID91](#),
 C. Gonzalez Renteria [ID17a](#), R. Gonzalez Suarez [ID159](#), S. Gonzalez-Sevilla [ID56](#),
 G.R. Gonzalvo Rodriguez [ID161](#), R.Y. González Andana [ID52](#), L. Goossens [ID36](#), N.A. Gorasia [ID20](#),
 P.A. Gorbounov [ID37](#), B. Gorini [ID36](#), E. Gorini [ID69a,69b](#), A. Gorišek [ID92](#), A.T. Goshaw [ID51](#),
 M.I. Gostkin [ID38](#), C.A. Gottardo [ID36](#), M. Goughri [ID35b](#), V. Goumarre [ID48](#), A.G. Goussiou [ID137](#),
 N. Govender [ID33c](#), C. Goy [ID4](#), I. Grabowska-Bold [ID84a](#), K. Graham [ID34](#), E. Gramstad [ID124](#),
 S. Grancagnolo [ID18](#), M. Grandi [ID145](#), V. Gratchev [ID37,*](#), P.M. Gravila [ID27f](#), F.G. Gravili [ID69a,69b](#),
 H.M. Gray [ID17a](#), M. Greco [ID69a,69b](#), C. Grefe [ID24](#), I.M. Gregor [ID48](#), P. Grenier [ID142](#), C. Grieco [ID13](#),
 A.A. Grillo [ID135](#), K. Grimm [ID31,m](#), S. Grinstein [ID13,t](#), J.-F. Grivaz [ID66](#), E. Gross [ID167](#),
 J. Grosse-Knetter [ID55](#), C. Grud [ID105](#), A. Grummer [ID111](#), J.C. Grundy [ID125](#), L. Guan [ID105](#), W. Guan [ID168](#),
 C. Gubbels [ID162](#), J.G.R. Guerrero Rojas [ID161](#), G. Guerrieri [ID68a,68b](#), F. Guescini [ID109](#), R. Gugel [ID99](#),
 J.A.M. Guhit [ID105](#), A. Guida [ID48](#), T. Guillemin [ID4](#), E. Guilloton [ID165,133](#), S. Guindon [ID36](#),
 F. Guo [ID14a,14d](#), J. Guo [ID62c](#), L. Guo [ID66](#), Y. Guo [ID105](#), R. Gupta [ID48](#), S. Gurbuz [ID24](#), S.S. Gurdasani [ID54](#),

G. Gustavino ^{id36}, M. Guth ^{id56}, P. Gutierrez ^{id119}, L.F. Gutierrez Zagazeta ^{id127}, C. Gutschow ^{id95}, C. Guyot ^{id134}, C. Gwenlan ^{id125}, C.B. Gwilliam ^{id91}, E.S. Haaland ^{id124}, A. Haas ^{id116}, M. Habedank ^{id48}, C. Haber ^{id17a}, H.K. Hadavand ^{id8}, A. Hadeef ^{id99}, S. Hadzic ^{id109}, E.H. Haines ^{id95}, M. Haleem ^{id164}, J. Haley ^{id120}, J.J. Hall ^{id138}, G.D. Hallewell ^{id101}, L. Halser ^{id19}, K. Hamano ^{id163}, H. Hamdaoui ^{id35e}, M. Hamer ^{id24}, G.N. Hamity ^{id52}, J. Han ^{id62b}, K. Han ^{id62a}, L. Han ^{id14c}, L. Han ^{id62a}, S. Han ^{id17a}, Y.F. Han ^{id154}, K. Hanagaki ^{id82}, M. Hance ^{id135}, D.A. Hangal ^{id41,z}, H. Hanif ^{id141}, M.D. Hank ^{id39}, R. Hankache ^{id100}, J.B. Hansen ^{id42}, J.D. Hansen ^{id42}, P.H. Hansen ^{id42}, K. Hara ^{id156}, D. Harada ^{id56}, T. Harenberg ^{id169}, S. Harkusha ^{id37}, Y.T. Harris ^{id125}, N.M. Harrison ^{id118}, P.F. Harrison ^{id165}, N.M. Hartman ^{id142}, N.M. Hartmann ^{id108}, Y. Hasegawa ^{id139}, A. Hasib ^{id52}, S. Haug ^{id19}, R. Hauser ^{id106}, M. Havranek ^{id131}, C.M. Hawkes ^{id20}, R.J. Hawkins ^{id36}, S. Hayashida ^{id110}, D. Hayden ^{id106}, C. Hayes ^{id105}, R.L. Hayes ^{id162}, C.P. Hays ^{id125}, J.M. Hays ^{id93}, H.S. Hayward ^{id91}, F. He ^{id62a}, Y. He ^{id153}, Y. He ^{id126}, M.P. Heath ^{id52}, V. Hedberg ^{id97}, A.L. Heggelund ^{id124}, N.D. Hehir ^{id93}, C. Heidegger ^{id54}, K.K. Heidegger ^{id54}, W.D. Heidorn ^{id80}, J. Heilman ^{id34}, S. Heim ^{id48}, T. Heim ^{id17a}, J.G. Heinlein ^{id127}, J.J. Heinrich ^{id122}, L. Heinrich ^{id109}, J. Hejbal ^{id130}, L. Helary ^{id48}, A. Held ^{id168}, S. Hellesund ^{id124}, C.M. Helling ^{id162}, S. Hellman ^{id47a,47b}, C. Hensens ^{id36}, R.C.W. Henderson ^{id90}, L. Henkelmann ^{id32}, A.M. Henriques Correia ^{id36}, H. Herde ^{id97}, Y. Hernández Jiménez ^{id144}, L.M. Herrmann ^{id24}, M.G. Herrmann ^{id108}, T. Herrmann ^{id50}, G. Herten ^{id54}, R. Hertenberger ^{id108}, L. Hervas ^{id36}, N.P. Hessey ^{id155a}, H. Hibi ^{id83}, E. Higón-Rodríguez ^{id161}, S.J. Hillier ^{id20}, I. Hinchliffe ^{id17a}, F. Hinterkeuser ^{id24}, M. Hirose ^{id123}, S. Hirose ^{id156}, D. Hirschbuehl ^{id169}, T.G. Hitchings ^{id100}, B. Hiti ^{id92}, J. Hobbs ^{id144}, R. Hobincu ^{id27e}, N. Hod ^{id167}, M.C. Hodgkinson ^{id138}, B.H. Hodgkinson ^{id32}, A. Hoecker ^{id36}, J. Hofer ^{id48}, D. Hohn ^{id54}, T. Holm ^{id24}, M. Holzbock ^{id109}, L.B.A.H. Hommels ^{id32}, B.P. Honan ^{id100}, J. Hong ^{id62c}, T.M. Hong ^{id128}, J.C. Honig ^{id54}, A. Hönle ^{id109}, B.H. Hooberman ^{id160}, W.H. Hopkins ^{id6}, Y. Horii ^{id110}, S. Hou ^{id147}, A.S. Howard ^{id92}, J. Howarth ^{id59}, J. Hoya ^{id6}, M. Hrabovsky ^{id121}, A. Hrynevich ^{id48}, T. Hryn'ova ^{id4}, P.J. Hsu ^{id65}, S.-C. Hsu ^{id137}, Q. Hu ^{id41,z}, Y.F. Hu ^{id14a,14d,ac}, D.P. Huang ^{id95}, S. Huang ^{id64b}, X. Huang ^{id14c}, Y. Huang ^{id62a}, Y. Huang ^{id14a}, Z. Huang ^{id100}, Z. Hubacek ^{id131}, M. Huebner ^{id24}, F. Huegging ^{id24}, T.B. Huffman ^{id125}, M. Huhtinen ^{id36}, S.K. Huiberts ^{id16}, R. Hulsken ^{id103}, N. Huseynov ^{id12,a}, J. Huston ^{id106}, J. Huth ^{id61}, R. Hyneman ^{id142}, S. Hyrych ^{id28a}, G. Iacobucci ^{id56}, G. Iakovidis ^{id29}, I. Ibragimov ^{id140}, L. Iconomidou-Fayard ^{id66}, P. Iengo ^{id71a,71b}, R. Iguchi ^{id152}, T. Iizawa ^{id56}, Y. Ikegami ^{id82}, A. Ilg ^{id19}, N. Ilic ^{id154}, H. Imam ^{id35a}, T. Ingebretsen Carlson ^{id47a,47b}, G. Introzzi ^{id72a,72b}, M. Iodice ^{id76a}, V. Ippolito ^{id74a,74b}, M. Ishino ^{id152}, W. Islam ^{id168}, C. Issever ^{id18,48}, S. Istin ^{id21a,ah}, H. Ito ^{id166}, J.M. Iturbe Ponce ^{id64a}, R. Iuppa ^{id77a,77b}, A. Ivina ^{id167}, J.M. Izen ^{id45}, V. Izzo ^{id71a}, P. Jacka ^{id130,131}, P. Jackson ^{id1}, R.M. Jacobs ^{id48}, B.P. Jaeger ^{id141}, C.S. Jagfeld ^{id108}, G. Jäkel ^{id169}, K. Jakobs ^{id54}, T. Jakoubek ^{id167}, J. Jamieson ^{id59}, K.W. Janas ^{id84a}, G. Jarlskog ^{id97}, A.E. Jaspán ^{id91}, M. Javurkova ^{id102}, F. Jeanneau ^{id134}, L. Jeanty ^{id122}, J. Jejelava ^{id148a,y}, P. Jenni ^{id54,f}, C.E. Jessiman ^{id34}, S. Jézéquel ^{id4}, J. Jia ^{id144}, X. Jia ^{id61}, X. Jia ^{id14a,14d}, Z. Jia ^{id14c}, Y. Jiang ^{id62a}, S. Jiggins ^{id52}, J. Jimenez Pena ^{id109}, S. Jin ^{id14c}, A. Jinaru ^{id27b}, O. Jinnouchi ^{id153}, P. Johansson ^{id138}, K.A. Johns ^{id7}, D.M. Jones ^{id32}, E. Jones ^{id165}, P. Jones ^{id32}, R.W.L. Jones ^{id90}, T.J. Jones ^{id91}, R. Joshi ^{id118}, J. Jovicevic ^{id15}, X. Ju ^{id17a}, J.J. Junggeburth ^{id36}, A. Juste Rozas ^{id13,t}, S. Kabana ^{id136e}, A. Kaczmarska ^{id85}, M. Kado ^{id74a,74b}, H. Kagan ^{id118}, M. Kagan ^{id142}, A. Kahn ^{id41}, A. Kahn ^{id127}, C. Kahra ^{id99}, T. Kaji ^{id166}, E. Kajomovitz ^{id149}, N. Kakati ^{id167}, C.W. Kalderon ^{id29}, A. Kamenshchikov ^{id154}, S. Kanayama ^{id153}, N.J. Kang ^{id135}, Y. Kano ^{id110}, D. Kar ^{id33g}, K. Karava ^{id125}, M.J. Kareem ^{id155b}, E. Karentzos ^{id54}, I. Karkanas ^{id151}, S.N. Karpov ^{id38}, Z.M. Karpova ^{id38}, V. Kartvelishvili ^{id90}, A.N. Karyukhin ^{id37}, E. Kasimi ^{id151}, C. Kato ^{id62d}, J. Katzy ^{id48}, S. Kaur ^{id34}, K. Kawade ^{id139}, K. Kawagoe ^{id88}, T. Kawamoto ^{id134}, G. Kawamura ^{id55}, E.F. Kay ^{id163}, F.I. Kaya ^{id157}, S. Kazakos ^{id13}, V.F. Kazanin ^{id37}, Y. Ke ^{id144}, J.M. Keaveney ^{id33a}, R. Keeler ^{id163}, G.V. Kehris ^{id61}, J.S. Keller ^{id34}, A.S. Kelly ^{id95}, D. Kelsey ^{id145}, J.J. Kempster ^{id20}, K.E. Kennedy ^{id41}, P.D. Kennedy ^{id99},

O. Kepka ¹³⁰, B.P. Kerridge ¹⁶⁵, S. Kersten ¹⁶⁹, B.P. Kerševan ⁹², S. Keshri ⁶⁶,
L. Keszeghova ^{28a}, S. Ketabchi Haghghat ¹⁵⁴, M. Khandoga ¹²⁶, A. Khanov ¹²⁰,
A.G. Kharlamov ³⁷, T. Kharlamova ³⁷, E.E. Khoda ¹³⁷, T.J. Khoo ¹⁸, G. Khoraiuli ¹⁶⁴,
J. Khubua ^{148b}, Y.A.R. Khwaira ⁶⁶, M. Kiehn ³⁶, A. Kilgallon ¹²², D.W. Kim ^{47a,47b},
E. Kim ¹⁵³, Y.K. Kim ³⁹, N. Kimura ⁹⁵, A. Kirchhoff ⁵⁵, D. Kirchmeier ⁵⁰, C. Kirfel ²⁴,
J. Kirk ¹³³, A.E. Kiryunin ¹⁰⁹, T. Kishimoto ¹⁵², D.P. Kisliuk ¹⁵⁴, C. Kitsaki ¹⁰, O. Kivernyk ²⁴,
M. Klassen ^{63a}, C. Klein ³⁴, L. Klein ¹⁶⁴, M.H. Klein ¹⁰⁵, M. Klein ⁹¹, S.B. Klein ⁵⁶,
U. Klein ⁹¹, P. Klimek ³⁶, A. Klimentov ²⁹, F. Klimpel ¹⁰⁹, T. Klingl ²⁴, T. Klioutchnikova ³⁶,
F.F. Klitzner ¹⁰⁸, P. Kluit ¹¹³, S. Kluth ¹⁰⁹, E. Kneringer ⁷⁸, T.M. Knight ¹⁵⁴, A. Knue ⁵⁴,
D. Kobayashi⁸⁸, R. Kobayashi ⁸⁶, M. Kocian ¹⁴², P. Kodyš ¹³², D.M. Koeck ¹⁴⁵, P.T. Koenig ²⁴,
T. Koffas ³⁴, M. Kolb ¹³⁴, I. Koletsou ⁴, T. Komarek ¹²¹, K. Köneke ⁵⁴, A.X.Y. Kong ¹,
T. Kono ¹¹⁷, N. Konstantinidis ⁹⁵, B. Konya ⁹⁷, R. Kopeliansky ⁶⁷, S. Koperny ^{84a}, K. Korcyl ⁸⁵,
K. Kordas ¹⁵¹, G. Koren ¹⁵⁰, A. Korn ⁹⁵, S. Korn ⁵⁵, I. Korolkov ¹³, N. Korotkova ³⁷,
B. Kortman ¹¹³, O. Kortner ¹⁰⁹, S. Kortner ¹⁰⁹, W.H. Kostecka ¹¹⁴, V.V. Kostyukhin ¹⁴⁰,
A. Kotsokechagia ¹³⁴, A. Kotwal ⁵¹, A. Koulouris ³⁶, A. Kourkoumeli-Charalampidi ^{72a,72b},
C. Kourkoumelis ⁹, E. Kourlitis ⁶, O. Kovanda ¹⁴⁵, R. Kowalewski ¹⁶³, W. Kozanecki ¹³⁴,
A.S. Kozhin ³⁷, V.A. Kramarenko ³⁷, G. Kramberger ⁹², P. Kramer ⁹⁹, M.W. Krasny ¹²⁶,
A. Krasznahorkay ³⁶, J.A. Kremer ⁹⁹, T. Kresse ⁵⁰, J. Kretzschmar ⁹¹, K. Kreul ¹⁸,
P. Krieger ¹⁵⁴, F. Krieter ¹⁰⁸, S. Krishnamurthy ¹⁰², A. Krishnan ^{63b}, M. Krivos ¹³²,
K. Krizka ^{17a}, K. Kroeninger ⁴⁹, H. Kroha ¹⁰⁹, J. Kroll ¹³⁰, J. Kroll ¹²⁷, K.S. Krowpman ¹⁰⁶,
U. Kruchonak ³⁸, H. Krüger ²⁴, N. Krumnack⁸⁰, M.C. Kruse ⁵¹, J.A. Krzysiak ⁸⁵, A. Kubota ¹⁵³,
O. Kuchinskaia ³⁷, S. Kuday ^{3a}, D. Kuechler ⁴⁸, J.T. Kuechler ⁴⁸, S. Kuehn ³⁶, T. Kuhl ⁴⁸,
V. Kukhtin ³⁸, Y. Kulchitsky ^{37,a}, S. Kuleshov ^{136d,136b}, M. Kumar ^{33g}, N. Kumari ¹⁰¹,
A. Kupco ¹³⁰, T. Kupfer⁴⁹, A. Kupich ³⁷, O. Kuprash ⁵⁴, H. Kurashige ⁸³, L.L. Kurchaninov ^{155a},
Y.A. Kurochkin ³⁷, A. Kurova ³⁷, M. Kuze ¹⁵³, A.K. Kvam ¹⁰², J. Kvita ¹²¹, T. Kwan ¹⁰³,
K.W. Kwok ^{64a}, N.G. Kyriacou ¹⁰⁵, L.A.O. Laatu ¹⁰¹, C. Lacasta ¹⁶¹, F. Lacava ^{74a,74b},
H. Lacker ¹⁸, D. Lacour ¹²⁶, N.N. Lad ⁹⁵, E. Ladygin ³⁸, B. Laforge ¹²⁶, T. Lagouri ^{136e},
S. Lai ⁵⁵, I.K. Lakomic ^{84a}, N. Lalloue ⁶⁰, J.E. Lambert ¹¹⁹, S. Lammers ⁶⁷, W. Lampl ⁷,
C. Lampoudis ¹⁵¹, A.N. Lancaster ¹¹⁴, E. Lançon ²⁹, U. Landgraf ⁵⁴, M.P.J. Landon ⁹³,
V.S. Lang ⁵⁴, R.J. Langenberg ¹⁰², A.J. Lankford ¹⁵⁸, F. Lanni ³⁶, K. Lantzsch ²⁴, A. Lanza ^{72a},
A. Lapertosa ^{57b,57a}, J.F. Laporte ¹³⁴, T. Lari ^{70a}, F. Lasagni Manghi ^{23b}, M. Lassnig ³⁶,
V. Latonova ¹³⁰, T.S. Lau ^{64a}, A. Laudrain ⁹⁹, A. Laurier ³⁴, S.D. Lawlor ⁹⁴, Z. Lawrence ¹⁰⁰,
M. Lazzaroni ^{70a,70b}, B. Le¹⁰⁰, B. Leban ⁹², A. Lebedev ⁸⁰, M. LeBlanc ³⁶, T. LeCompte ⁶,
F. Ledroit-Guillon ⁶⁰, A.C.A. Lee⁹⁵, G.R. Lee ¹⁶, L. Lee ⁶¹, S.C. Lee ¹⁴⁷, S. Lee ^{47a,47b},
T.F. Lee ⁹¹, L.L. Leeuw ^{33c}, H.P. Lefebvre ⁹⁴, M. Lefebvre ¹⁶³, C. Leggett ^{17a}, K. Lehmann ¹⁴¹,
G. Lehmann Miotto ³⁶, M. Leigh ⁵⁶, W.A. Leight ¹⁰², A. Leisos ^{151,s}, M.A.L. Leite ^{81c},
C.E. Leitgeb ⁴⁸, R. Leitner ¹³², K.J.C. Leney ⁴⁴, T. Lenz ²⁴, S. Leone ^{73a}, C. Leonidopoulos ⁵²,
A. Leopold ¹⁴³, C. Leroy ¹⁰⁷, R. Les ¹⁰⁶, C.G. Lester ³², M. Levchenko ³⁷, J. Levêque ⁴,
D. Levin ¹⁰⁵, L.J. Levinson ¹⁶⁷, M.P. Lewicki ⁸⁵, D.J. Lewis ⁴, A. Li ⁵, B. Li ^{14b}, B. Li ^{62b},
C. Li^{62a}, C-Q. Li ^{62c}, H. Li ^{62a}, H. Li ^{62b}, H. Li ^{14c}, H. Li ^{62b}, J. Li ^{62c}, K. Li ¹³⁷, L. Li ^{62c},
M. Li ^{14a,14d}, Q.Y. Li ^{62a}, S. Li ^{14a,14d}, S. Li ^{62d,62c,d}, T. Li ^{62b}, X. Li ¹⁰³, Z. Li ^{62b}, Z. Li ¹²⁵,
Z. Li ¹⁰³, Z. Li ⁹¹, Z. Li ^{14a,14d}, Z. Liang ^{14a}, M. Liberatore ⁴⁸, B. Liberti ^{75a}, K. Lie ^{64c},
J. Lieber Marin ^{81b}, K. Lin ¹⁰⁶, R.A. Linck ⁶⁷, R.E. Lindley ⁷, J.H. Lindon ², A. Linss ⁴⁸,
E. Lipeles ¹²⁷, A. Lipniacka ¹⁶, A. Lister ¹⁶², J.D. Little ⁴, B. Liu ^{14a}, B.X. Liu ¹⁴¹,
D. Liu ^{62d,62c}, J.B. Liu ^{62a}, J.K.K. Liu ³², K. Liu ^{62d,62c}, M. Liu ^{62a}, M.Y. Liu ^{62a}, P. Liu ^{14a},
Q. Liu ^{62d,137,62c}, X. Liu ^{62a}, Y. Liu ⁴⁸, Y. Liu ^{14c,14d}, Y.L. Liu ¹⁰⁵, Y.W. Liu ^{62a},
M. Livan ^{72a,72b}, J. Llorente Merino ¹⁴¹, S.L. Lloyd ⁹³, E.M. Lobodzinska ⁴⁸, P. Loch ⁷,




































































































































S. Loffredo ^{75a,75b}, T. Lohse ¹⁸, K. Lohwasser ¹³⁸, M. Lokajicek ¹³⁰, J.D. Long ¹⁶⁰,
 I. Longarini ^{74a,74b}, L. Longo ^{69a,69b}, R. Longo ¹⁶⁰, I. Lopez Paz ³⁶, A. Lopez Solis ⁴⁸,
 J. Lorenz ¹⁰⁸, N. Lorenzo Martinez ⁴, A.M. Lory ¹⁰⁸, A. Lösle ⁵⁴, X. Lou ^{47a,47b}, X. Lou ^{14a,14d},
 A. Lounis ⁶⁶, J. Love ⁶, P.A. Love ⁹⁰, J.J. Lozano Bahilo ¹⁶¹, G. Lu ^{14a,14d}, M. Lu ⁷⁹, S. Lu ¹²⁷,
 Y.J. Lu ⁶⁵, H.J. Lubatti ¹³⁷, C. Luci ^{74a,74b}, F.L. Lucio Alves ^{14c}, A. Lucotte ⁶⁰, F. Luehring ⁶⁷,
 I. Luise ¹⁴⁴, O. Lukianchuk ⁶⁶, O. Lundberg ¹⁴³, B. Lund-Jensen ¹⁴³, N.A. Luongo ¹²²,
 M.S. Lutz ¹⁵⁰, D. Lynn ²⁹, H. Lyons⁹¹, R. Lysak ¹³⁰, E. Lytken ⁹⁷, F. Lyu ^{14a}, V. Lyubushkin ³⁸,
 T. Lyubushkina ³⁸, H. Ma ²⁹, L.L. Ma ^{62b}, Y. Ma ⁹⁵, D.M. Mac Donell ¹⁶³, G. Maccarrone ⁵³,
 J.C. MacDonald ¹³⁸, R. Madar ⁴⁰, W.F. Mader ⁵⁰, J. Maeda ⁸³, T. Maeno ²⁹, M. Maerker ⁵⁰,
 V. Magerl ⁵⁴, H. Maguire ¹³⁸, D.J. Mahon ⁴¹, C. Maidantchik ^{81b}, A. Maio ^{129a,129b,129d},
 K. Maj ^{84a}, O. Majersky ^{28a}, S. Majewski ¹²², N. Makovec ⁶⁶, V. Maksimovic ¹⁵,
 B. Malaescu ¹²⁶, Pa. Malecki ⁸⁵, V.P. Maleev ³⁷, F. Malek ⁶⁰, D. Malito ^{43b,43a}, U. Mallik ⁷⁹,
 C. Malone ³², S. Maltezos¹⁰, S. Malyukov³⁸, J. Mamuzic ¹³, G. Mancini ⁵³, G. Manco ^{72a,72b},
 J.P. Mandalia ⁹³, I. Mandić ⁹², L. Manhaes de Andrade Filho ^{81a}, I.M. Maniatis ¹⁵¹,
 M. Manisha ¹³⁴, J. Manjarres Ramos ⁵⁰, D.C. Mankad ¹⁶⁷, A. Mann ¹⁰⁸, B. Mansoulie ¹³⁴,
 S. Manzoni ³⁶, A. Marantis ¹⁵¹, G. Marchiori ⁵, M. Marcisovsky ¹³⁰, L. Marcoccia ^{75a,75b},
 C. Marcon ^{70a,70b}, M. Marinescu ²⁰, M. Marjanovic ¹¹⁹, Z. Marshall ^{17a}, S. Marti-Garcia ¹⁶¹,
 T.A. Martin ¹⁶⁵, V.J. Martin ⁵², B. Martin dit Latour ¹⁶, L. Martinelli ^{74a,74b}, M. Martinez ^{13,t},
 P. Martinez Agullo ¹⁶¹, V.I. Martinez Outschoorn ¹⁰², P. Martinez Suarez ¹³, S. Martin-Haugh ¹³³,
 V.S. Martoiu ^{27b}, A.C. Martyniuk ⁹⁵, A. Marzin ³⁶, S.R. Maschek ¹⁰⁹, L. Masetti ⁹⁹,
 T. Mashimo ¹⁵², J. Masik ¹⁰⁰, A.L. Maslennikov ³⁷, L. Massa ^{23b}, P. Massarotti ^{71a,71b},
 P. Mastrandrea ^{73a,73b}, A. Mastroberardino ^{43b,43a}, T. Masubuchi ¹⁵², T. Mathisen ¹⁵⁹,
 N. Matsuzawa¹⁵², J. Maurer ^{27b}, B. Maček ⁹², D.A. Maximov ³⁷, R. Mazini ¹⁴⁷, I. Maznas ¹⁵¹,
 M. Mazza ¹⁰⁶, S.M. Mazza ¹³⁵, C. Mc Ginn ^{29,af}, J.P. Mc Gowan ¹⁰³, S.P. Mc Kee ¹⁰⁵,
 W.P. McCormack ^{17a}, E.F. McDonald ¹⁰⁴, A.E. McDougall ¹¹³, J.A. Mcfayden ¹⁴⁵,
 G. Mchedlidze ^{148b}, R.P. Mckenzie ^{33g}, T.C. Mclachlan ⁴⁸, D.J. Mclaughlin ⁹⁵, K.D. McLean ¹⁶³,
 S.J. McMahon ¹³³, P.C. McNamara ¹⁰⁴, C.M. Mcpartland ⁹¹, R.A. McPherson ^{163,v}, T. Megy ⁴⁰,
 S. Mehlhase ¹⁰⁸, A. Mehta ⁹¹, B. Meirose ⁴⁵, D. Melini ¹⁴⁹, B.R. Mellado Garcia ^{33g},
 A.H. Melo ⁵⁵, F. Meloni ⁴⁸, E.D. Mendes Gouveia ^{129a}, A.M. Mendes Jacques Da Costa ²⁰,
 H.Y. Meng ¹⁵⁴, L. Meng ⁹⁰, S. Menke ¹⁰⁹, M. Mentink ³⁶, E. Meoni ^{43b,43a}, C. Merlassino ¹²⁵,
 L. Merola ^{71a,71b}, C. Meroni ^{70a}, G. Merz¹⁰⁵, O. Meshkov ³⁷, J.K.R. Meshreki ¹⁴⁰, J. Metcalfe ⁶,
 A.S. Mete ⁶, C. Meyer ⁶⁷, J-P. Meyer ¹³⁴, M. Michetti ¹⁸, R.P. Middleton ¹³³, L. Mijović ⁵²,
 G. Mikenberg ¹⁶⁷, M. Mikestikova ¹³⁰, M. Mikuž ⁹², H. Mildner ¹³⁸, A. Milic ³⁶,
 C.D. Milke ⁴⁴, D.W. Miller ³⁹, L.S. Miller ³⁴, A. Milov ¹⁶⁷, D.A. Milstead^{47a,47b}, T. Min^{14c},
 A.A. Minaenko ³⁷, I.A. Minashvili ^{148b}, L. Mince ⁵⁹, A.I. Mincer ¹¹⁶, B. Mindur ^{84a},
 M. Mineev ³⁸, Y. Mino ⁸⁶, L.M. Mir ¹³, M. Miralles Lopez ¹⁶¹, M. Mironova ¹²⁵,
 M.C. Missio ¹¹², T. Mitani ¹⁶⁶, A. Mitra ¹⁶⁵, V.A. Mitsou ¹⁶¹, O. Miu ¹⁵⁴, P.S. Miyagawa ⁹³,
 Y. Miyazaki⁸⁸, A. Mizukami ⁸², J.U. Mjörnmark ⁹⁷, T. Mkrtchyan ^{63a}, T. Mlinarevic ⁹⁵,
 M. Mlynarikova ³⁶, T. Moa ^{47a,47b}, S. Mobius ⁵⁵, K. Mochizuki ¹⁰⁷, P. Moder ⁴⁸, P. Mogg ¹⁰⁸,
 A.F. Mohammed ^{14a,14d}, S. Mohapatra ⁴¹, G. Mokgatitswane ^{33g}, B. Mondal ¹⁴⁰, S. Mondal ¹³¹,
 K. Mönig ⁴⁸, E. Monnier ¹⁰¹, L. Monsonis Romero¹⁶¹, J. Montejo Berlingen ³⁶, M. Montella ¹¹⁸,
 F. Monticelli ⁸⁹, N. Morange ⁶⁶, A.L. Moreira De Carvalho ^{129a}, M. Moreno Llácer ¹⁶¹,
 C. Moreno Martinez ⁵⁶, P. Morettini ^{57b}, S. Morgenstern ¹⁶⁵, M. Morii ⁶¹, M. Morinaga ¹⁵²,
 V. Morisbak ¹²⁴, A.K. Morley ³⁶, F. Morodei ^{74a,74b}, L. Morvaj ³⁶, P. Moschovakos ³⁶,
 B. Moser ³⁶, M. Mosidze^{148b}, T. Moskalets ⁵⁴, P. Moskvitina ¹¹², J. Moss ^{31,n}, E.J.W. Moyses ¹⁰²,
 S. Muanza ¹⁰¹, J. Mueller ¹²⁸, D. Muenstermann ⁹⁰, R. Müller ¹⁹, G.A. Mullier ⁹⁷, J.J. Mullin¹²⁷,
 D.P. Mungo ¹⁵⁴, J.L. Munoz Martinez ¹³, D. Munoz Perez ¹⁶¹, F.J. Munoz Sanchez ¹⁰⁰,

M. Murin ¹⁰⁰, W.J. Murray ^{165,133}, A. Murrone ^{70a,70b}, J.M. Muse ¹¹⁹, M. Muškinja ^{17a},
C. Mwewa ²⁹, A.G. Myagkov ^{37,a}, A.J. Myers ⁸, A.A. Myers ¹²⁸, G. Myers ⁶⁷, M. Myska ¹³¹,
B.P. Nachman ^{17a}, O. Nackenhorst ⁴⁹, A.Nag Nag ⁵⁰, K. Nagai ¹²⁵, K. Nagano ⁸²,
J.L. Nagle ^{29,af}, E. Nagy ¹⁰¹, A.M. Nairz ³⁶, Y. Nakahama ⁸², K. Nakamura ⁸², H. Nanjo ¹²³,
R. Narayan ⁴⁴, E.A. Narayanan ¹¹¹, I. Naryshkin ³⁷, M. Naseri ³⁴, C. Nass ²⁴, G. Navarro ^{22a},
J. Navarro-Gonzalez ¹⁶¹, R. Nayak ¹⁵⁰, A. Nayaz ¹⁸, P.Y. Nechaeva ³⁷, F. Nechansky ⁴⁸,
L. Nedic ¹²⁵, T.J. Neep ²⁰, A. Negri ^{72a,72b}, M. Negrini ^{23b}, C. Nellist ¹¹², C. Nelson ¹⁰³,
K. Nelson ¹⁰⁵, S. Nemecek ¹³⁰, M. Nessi ^{36,g}, M.S. Neubauer ¹⁶⁰, F. Neuhaus ⁹⁹,
J. Neundorf ⁴⁸, R. Newhouse ¹⁶², P.R. Newman ²⁰, C.W. Ng ¹²⁸, Y.S. Ng ¹⁸, Y.W.Y. Ng ⁴⁸,
B. Ngair ^{35e}, H.D.N. Nguyen ¹⁰⁷, R.B. Nickerson ¹²⁵, R. Nicolaidou ¹³⁴, J. Nielsen ¹³⁵,
M. Niemeyer ⁵⁵, N. Nikipforou ³⁶, V. Nikolaenko ^{37,a}, I. Nikolic-Audit ¹²⁶, K. Nikolopoulos ²⁰,
P. Nilsson ²⁹, H.R. Nindhito ⁵⁶, A. Nisati ^{74a}, N. Nishu ², R. Nisius ¹⁰⁹, J.-E. Nitschke ⁵⁰,
E.K. Nkadimeng ^{33g}, S.J. Noacco Rosende ⁸⁹, T. Nobe ¹⁵², D.L. Noel ³², Y. Noguchi ⁸⁶,
T. Nommensen ¹⁴⁶, M.A. Nomura ²⁹, M.B. Norfolk ¹³⁸, R.R.B. Norisam ⁹⁵, B.J. Norman ³⁴,
J. Novak ⁹², T. Novak ⁴⁸, O. Novgorodova ⁵⁰, L. Novotny ¹³¹, R. Novotny ¹¹¹, L. Nozka ¹²¹,
K. Ntekas ¹⁵⁸, N.M.J. Nunes De Moura Junior ^{81b}, E. Nurse ⁹⁵, F.G. Oakham ^{34,ac}, J. Ocariz ¹²⁶,
A. Ochi ⁸³, I. Ochoa ^{129a}, S. Oerdek ¹⁵⁹, A. Ogrodnik ^{84a}, A. Oh ¹⁰⁰, C.C. Ohm ¹⁴³,
H. Oide ⁸², R. Oishi ¹⁵², M.L. Ojeda ⁴⁸, Y. Okazaki ⁸⁶, M.W. O'Keefe ⁹¹, Y. Okumura ¹⁵²,
A. Olariu ^{27b}, L.F. Oleiro Seabra ^{129a}, S.A. Olivares Pino ^{136e}, D. Oliveira Damazio ²⁹,
D. Oliveira Goncalves ^{81a}, J.L. Oliver ¹⁵⁸, M.J.R. Olsson ¹⁵⁸, A. Olszewski ⁸⁵, J. Olszowska ^{85,*},
Ö.O. Öncel ⁵⁴, D.C. O'Neil ¹⁴¹, A.P. O'Neill ¹⁹, A. Onofre ^{129a,129e}, P.U.E. Onyisi ¹¹,
M.J. Oreglia ³⁹, G.E. Orellana ⁸⁹, D. Orestano ^{76a,76b}, N. Orlando ¹³, R.S. Orr ¹⁵⁴,
V. O'Shea ⁵⁹, R. Ospanov ^{62a}, G. Otero y Garzon ³⁰, H. Otono ⁸⁸, P.S. Ott ^{63a}, G.J. Ottino ^{17a},
M. Ouchrif ^{35d}, J. Ouellette ^{29,af}, F. Ould-Saada ¹²⁴, M. Owen ⁵⁹, R.E. Owen ¹³³,
K.Y. Oyulmaz ^{21a}, V.E. Ozcan ^{21a}, N. Ozturk ⁸, S. Ozturk ^{21d}, J. Pacalt ¹²¹, H.A. Pacey ³²,
K. Pachal ⁵¹, A. Pacheco Pages ¹³, C. Padilla Aranda ¹³, G. Padovano ^{74a,74b}, S. Pagan Griso ^{17a},
G. Palacino ⁶⁷, A. Palazzo ^{69a,69b}, S. Palazzo ⁵², S. Palestini ³⁶, M. Palka ^{84b}, J. Pan ¹⁷⁰,
T. Pan ^{64a}, D.K. Panchal ¹¹, C.E. Pandini ¹¹³, J.G. Panduro Vazquez ⁹⁴, H. Pang ^{14b}, P. Pani ⁴⁸,
G. Panizzo ^{68a,68c}, L. Paolozzi ⁵⁶, C. Papadatos ¹⁰⁷, S. Parajuli ⁴⁴, A. Paramonov ⁶,
C. Paraskevopoulos ¹⁰, D. Paredes Hernandez ^{64b}, T.H. Park ¹⁵⁴, M.A. Parker ³², F. Parodi ^{57b,57a},
E.W. Parrish ¹¹⁴, V.A. Parrish ⁵², J.A. Parsons ⁴¹, U. Parzefall ⁵⁴, B. Pascual Dias ¹⁰⁷,
L. Pascual Dominguez ¹⁵⁰, V.R. Pascuzzi ^{17a}, F. Pasquali ¹¹³, E. Pasqualucci ^{74a}, S. Passaggio ^{57b},
F. Pastore ⁹⁴, P. Pasuwan ^{47a,47b}, P. Patel ⁸⁵, J.R. Pater ¹⁰⁰, T. Pauly ³⁶, J. Pearkes ¹⁴²,
M. Pedersen ¹²⁴, R. Pedro ^{129a}, S.V. Peleganchuk ³⁷, O. Penc ³⁶, E.A. Pender ⁵², C. Peng ^{64b},
H. Peng ^{62a}, K.E. Penski ¹⁰⁸, M. Penzin ³⁷, B.S. Peralva ^{81d,81d}, A.P. Pereira Peixoto ⁶⁰,
L. Pereira Sanchez ^{47a,47b}, D.V. Perepelitsa ^{29,af}, E. Perez Codina ^{155a}, M. Perganti ¹⁰,
L. Perini ^{70a,70b,*}, H. Pernegger ³⁶, S. Perrella ³⁶, A. Perrevoort ¹¹², O. Perrin ⁴⁰, K. Peters ⁴⁸,
R.F.Y. Peters ¹⁰⁰, B.A. Petersen ³⁶, T.C. Petersen ⁴², E. Petit ¹⁰¹, V. Petousis ¹³¹,
C. Petridou ¹⁵¹, A. Petrukhin ¹⁴⁰, M. Pettee ^{17a}, N.E. Pettersson ³⁶, A. Petukhov ³⁷,
K. Petukhova ¹³², A. Peyaud ¹³⁴, R. Pezoa ^{136f}, L. Pezzotti ³⁶, G. Pezzullo ¹⁷⁰, T.M. Pham ¹⁶⁸,
T. Pham ¹⁰⁴, P.W. Phillips ¹³³, M.W. Phipps ¹⁶⁰, G. Piacquadio ¹⁴⁴, E. Pianori ^{17a},
F. Piazza ^{70a,70b}, R. Piegaia ³⁰, D. Pietreanu ^{27b}, A.D. Pilkington ¹⁰⁰, M. Pinamonti ^{68a,68c},
J.L. Pinfeld ², B.C. Pinheiro Pereira ^{129a}, C. Pitman Donaldson ⁹⁵, D.A. Pizzi ³⁴,
L. Pizzimento ^{75a,75b}, A. Pizzini ¹¹³, M.-A. Pleier ²⁹, V. Plesanovs ⁵⁴, V. Pleskot ¹³²,
E. Plotnikova ³⁸, G. Poddar ⁴, R. Poettgen ⁹⁷, L. Poggioli ¹²⁶, I. Pogrebnyak ¹⁰⁶, D. Pohl ²⁴,
I. Pokharel ⁵⁵, S. Polacek ¹³², G. Polesello ^{72a}, A. Poley ^{141,155a}, R. Polifka ¹³¹, A. Polini ^{23b},
C.S. Pollard ¹²⁵, Z.B. Pollock ¹¹⁸, V. Polychronakos ²⁹, E. Pompa Pacchi ^{74a,74b}, D. Ponomarenko ³⁷,

L. Pontecorvo ³⁶, S. Popa ^{27a}, G.A. Popeneciu ^{27d}, D.M. Portillo Quintero ^{155a}, S. Pospisil ¹³¹,
 P. Postolache ^{27c}, K. Potamianos ¹²⁵, I.N. Potrap ³⁸, C.J. Potter ³², H. Potti ¹, T. Poulsen ⁴⁸,
 J. Poveda ¹⁶¹, M.E. Pozo Astigarraga ³⁶, A. Prades Ibanez ¹⁶¹, M.M. Prapa ⁴⁶, D. Price ¹⁰⁰,
 M. Primavera ^{69a}, M.A. Principe Martin ⁹⁸, R. Privara ¹²¹, M.L. Proffitt ¹³⁷, N. Proklova ¹²⁷,
 K. Prokofiev ^{64c}, G. Proto ^{75a,75b}, S. Protopopescu ²⁹, J. Proudfoot ⁶, M. Przybycien ^{84a},
 J.E. Puddefoot ¹³⁸, D. Pudzha ³⁷, P. Puzo ⁶⁶, D. Pyatiizbyantseva ³⁷, J. Qian ¹⁰⁵, D. Qichen ¹⁰⁰,
 Y. Qin ¹⁰⁰, T. Qiu ⁹³, A. Quadt ⁵⁵, M. Queitsch-Maitland ¹⁰⁰, G. Quetant ⁵⁶,
 G. Rabanal Bolanos ⁶¹, D. Rafanoharana ⁵⁴, F. Ragusa ^{70a,70b}, J.L. Rainbolt ³⁹, J.A. Raine ⁵⁶,
 S. Rajagopalan ²⁹, E. Ramakoti ³⁷, K. Ran ^{48,14d}, N.P. Rapheeha ^{33g}, V. Raskina ¹²⁶,
 D.F. Rassloff ^{63a}, S. Rave ⁹⁹, B. Ravina ⁵⁵, I. Ravinovich ¹⁶⁷, M. Raymond ³⁶, A.L. Read ¹²⁴,
 N.P. Readioff ¹³⁸, D.M. Rebuzzi ^{72a,72b}, G. Redlinger ²⁹, K. Reeves ⁴⁵, J.A. Reidelsturz ¹⁶⁹,
 D. Reikher ¹⁵⁰, A. Reiss ⁹⁹, A. Rej ¹⁴⁰, C. Rembser ³⁶, A. Renardi ⁴⁸, M. Renda ^{27b},
 M.B. Rendel ¹⁰⁹, F. Renner ⁴⁸, A.G. Rennie ⁵⁹, S. Resconi ^{70a}, M. Ressegotti ^{57b,57a},
 E.D. Resseguie ^{17a}, S. Rettie ³⁶, B. Reynolds ¹¹⁸, E. Reynolds ^{17a}, M. Rezaei Estabragh ¹⁶⁹,
 O.L. Rezanova ³⁷, P. Reznicek ¹³², E. Ricci ^{77a,77b}, R. Richter ¹⁰⁹, S. Richter ^{47a,47b},
 E. Richter-Was ^{84b}, M. Ridel ¹²⁶, P. Rieck ¹¹⁶, P. Riedler ³⁶, M. Rijssenbeek ¹⁴⁴,
 A. Rimoldi ^{72a,72b}, M. Rimoldi ⁴⁸, L. Rinaldi ^{23b,23a}, T.T. Rinn ²⁹, M.P. Rinnagel ¹⁰⁸,
 G. Ripellino ¹⁴³, I. Riu ¹³, P. Rivadeneira ⁴⁸, J.C. Rivera Vergara ¹⁶³, F. Rizatdinova ¹²⁰,
 E. Rizvi ⁹³, C. Rizzi ⁵⁶, B.A. Roberts ¹⁶⁵, B.R. Roberts ^{17a}, S.H. Robertson ^{103,v}, M. Robin ⁴⁸,
 D. Robinson ³², C.M. Robles Gajardo ^{136f}, M. Robles Manzano ⁹⁹, A. Robson ⁵⁹, A. Rocchi ^{75a,75b},
 C. Roda ^{73a,73b}, S. Rodriguez Bosca ^{63a}, Y. Rodriguez Garcia ^{22a}, A. Rodriguez Rodriguez ⁵⁴,
 A.M. Rodríguez Vera ^{155b}, S. Roe ³⁶, J.T. Roemer ¹⁵⁸, A.R. Roepe-Gier ¹¹⁹, J. Roggel ¹⁶⁹,
 O. Røhne ¹²⁴, R.A. Rojas ¹⁶³, B. Roland ⁵⁴, C.P.A. Roland ⁶⁷, J. Roloff ²⁹, A. Romaniouk ³⁷,
 E. Romano ^{72a,72b}, M. Romano ^{23b}, A.C. Romero Hernandez ¹⁶⁰, N. Rompotis ⁹¹, L. Roos ¹²⁶,
 S. Rosati ^{74a}, B.J. Rosser ³⁹, E. Rossi ⁴, E. Rossi ^{71a,71b}, L.P. Rossi ^{57b}, L. Rossini ⁴⁸,
 R. Rosten ¹¹⁸, M. Rotaru ^{27b}, B. Rottler ⁵⁴, D. Rousseau ⁵⁶, D. Rousso ³², G. Rovelli ^{72a,72b},
 A. Roy ¹⁶⁰, A. Rozanov ¹⁰¹, Y. Rozen ¹⁴⁹, X. Ruan ^{33g}, A. Rubio Jimenez ¹⁶¹, A.J. Ruby ⁹¹,
 V.H. Ruelas Rivera ¹⁸, T.A. Ruggeri ¹, F. Rühr ⁵⁴, A. Ruiz-Martinez ¹⁶¹, A. Rummler ³⁶,
 Z. Rurikova ⁵⁴, N.A. Rusakovich ³⁸, H.L. Russell ¹⁶³, J.P. Rutherford ⁷, K. Rybacki ⁹⁰,
 M. Rybar ¹³², E.B. Rye ¹²⁴, A. Ryzhov ³⁷, J.A. Sabater Iglesias ⁵⁶, P. Sabatini ¹⁶¹,
 L. Sabetta ^{74a,74b}, H.F-W. Sadrozinski ¹³⁵, F. Safai Tehrani ^{74a}, B. Safarzadeh Samani ¹⁴⁵,
 M. Safdari ¹⁴², S. Saha ¹⁰³, M. Sahinsoy ¹⁰⁹, M. Saimpert ¹³⁴, M. Saito ¹⁵², T. Saito ¹⁵²,
 D. Salamani ³⁶, G. Salamanna ^{76a,76b}, A. Salnikov ¹⁴², J. Salt ¹⁶¹, A. Salvador Salas ¹³,
 D. Salvatore ^{43b,43a}, F. Salvatore ¹⁴⁵, A. Salzburger ³⁶, D. Sammel ⁵⁴, D. Sampsonidis ¹⁵¹,
 D. Sampsonidou ^{62d,62c}, J. Sánchez ¹⁶¹, A. Sanchez Pineda ⁴, V. Sanchez Sebastian ¹⁶¹,
 H. Sandaker ¹²⁴, C.O. Sander ⁴⁸, J.A. Sandesara ¹⁰², M. Sandhoff ¹⁶⁹, C. Sandoval ^{22b},
 D.P.C. Sankey ¹³³, A. Sansoni ⁵³, L. Santi ^{74a,74b}, C. Santoni ⁴⁰, H. Santos ^{129a,129b},
 S.N. Santpur ^{17a}, A. Santra ¹⁶⁷, K.A. Saoucha ¹³⁸, J.G. Saraiva ^{129a,129d}, J. Sardain ⁷,
 O. Sasaki ⁸², K. Sato ¹⁵⁶, C. Sauer ^{63b}, F. Sauerburger ⁵⁴, E. Sauvan ⁴, P. Savard ^{154,ac},
 R. Sawada ¹⁵², C. Sawyer ¹³³, L. Sawyer ⁹⁶, I. Sayago Galvan ¹⁶¹, C. Sbarra ^{23b}, A. Sbrizzi ^{23b,23a},
 T. Scanlon ⁹⁵, J. Schaarschmidt ¹³⁷, P. Schacht ¹⁰⁹, D. Schaefer ³⁹, U. Schäfer ⁹⁹,
 A.C. Schaffer ⁶⁶, D. Schaile ¹⁰⁸, R.D. Schamberger ¹⁴⁴, E. Schanet ¹⁰⁸, C. Scharf ¹⁸,
 M.M. Schefer ¹⁹, V.A. Schegelsky ³⁷, D. Scheirich ¹³², F. Schenck ¹⁸, M. Schernau ¹⁵⁸,
 C. Scheulen ⁵⁵, C. Schiavi ^{57b,57a}, Z.M. Schillaci ²⁶, E.J. Schioppa ^{69a,69b}, M. Schioppa ^{43b,43a},
 B. Schlag ⁹⁹, K.E. Schleicher ⁵⁴, S. Schlenker ³⁶, J. Schmeing ¹⁶⁹, M.A. Schmidt ¹⁶⁹,
 K. Schmieden ⁹⁹, C. Schmitt ⁹⁹, S. Schmitt ⁴⁸, L. Schoeffel ¹³⁴, A. Schoening ^{63b},
 P.G. Scholer ⁵⁴, E. Schopf ¹²⁵, M. Schott ⁹⁹, J. Schovancova ³⁶, S. Schramm ⁵⁶,

F. Schroeder ¹⁶⁹, H-C. Schultz-Coulon ^{63a}, M. Schumacher ⁵⁴, B.A. Schumm ¹³⁵, Ph. Schune ¹³⁴, A. Schwartzman ¹⁴², T.A. Schwarz ¹⁰⁵, Ph. Schwemling ¹³⁴, R. Schwiendorfer ¹⁰⁶, A. Sciandra ¹³⁵, G. Sciolla ²⁶, F. Scuri ^{73a}, F. Scutti ¹⁰⁴, C.D. Sebastiani ⁹¹, K. Sedlaczek ⁴⁹, P. Seema ¹⁸, S.C. Seidel ¹¹¹, A. Seiden ¹³⁵, B.D. Seidlitz ⁴¹, T. Seiss ³⁹, C. Seitz ⁴⁸, J.M. Seixas ^{81b}, G. Sekhniaidze ^{71a}, S.J. Sekula ⁴⁴, L. Selem ⁴, N. Semprini-Cesari ^{23b,23a}, S. Sen ⁵¹, D. Sengupta ⁵⁶, V. Senthilkumar ¹⁶¹, L. Serin ⁶⁶, L. Serkin ^{68a,68b}, M. Sessa ^{76a,76b}, H. Severini ¹¹⁹, S. Sevova ¹⁴², F. Sforza ^{57b,57a}, A. Sfyrta ⁵⁶, E. Shabalina ⁵⁵, R. Shaheen ¹⁴³, J.D. Shahinian ¹²⁷, N.W. Shaikh ^{47a,47b}, D. Shaked Renous ¹⁶⁷, L.Y. Shan ^{14a}, M. Shapiro ^{17a}, A. Sharma ³⁶, A.S. Sharma ¹⁶², P. Sharma ⁷⁹, S. Sharma ⁴⁸, P.B. Shatalov ³⁷, K. Shaw ¹⁴⁵, S.M. Shaw ¹⁰⁰, Q. Shen ^{62c,5}, P. Sherwood ⁹⁵, L. Shi ⁹⁵, C.O. Shimmin ¹⁷⁰, Y. Shimogama ¹⁶⁶, J.D. Shinner ⁹⁴, I.P.J. Shipsey ¹²⁵, S. Shirabe ⁶⁰, M. Shiyakova ³⁸, J. Shlomi ¹⁶⁷, M.J. Shochet ³⁹, J. Shojaii ¹⁰⁴, D.R. Shope ¹²⁴, S. Shrestha ^{118,ag}, E.M. Shrif ^{33g}, M.J. Shroff ¹⁶³, P. Sicho ¹³⁰, A.M. Sickles ¹⁶⁰, E. Sideras Haddad ^{33g}, A. Sidoti ^{23b}, F. Siegert ⁵⁰, Dj. Sijacki ¹⁵, R. Sikora ^{84a}, F. Sili ⁸⁹, J.M. Silva ²⁰, M.V. Silva Oliveira ³⁶, S.B. Silverstein ^{47a}, S. Simion ⁶⁶, R. Simoniello ³⁶, E.L. Simpson ⁵⁹, N.D. Simpson ⁹⁷, S. Simsek ^{21d}, S. Sindhu ⁵⁵, P. Sinervo ¹⁵⁴, V. Sinetckii ³⁷, S. Singh ¹⁴¹, S. Singh ¹⁵⁴, S. Sinha ⁴⁸, S. Sinha ^{33g}, M. Sioli ^{23b,23a}, I. Siral ³⁶, S.Yu. Sivoklov ^{37,*}, J. Sjölin ^{47a,47b}, A. Skaf ⁵⁵, E. Skorda ⁹⁷, P. Skubic ¹¹⁹, M. Slawinska ⁸⁵, V. Smakhtin ¹⁶⁷, B.H. Smart ¹³³, J. Smiesko ³⁶, S.Yu. Smirnov ³⁷, Y. Smirnov ³⁷, L.N. Smirnova ^{37,a}, O. Smirnova ⁹⁷, A.C. Smith ⁴¹, E.A. Smith ³⁹, H.A. Smith ¹²⁵, J.L. Smith ⁹¹, R. Smith ¹⁴², M. Smizanska ⁹⁰, K. Smolek ¹³¹, A. Smykiewicz ⁸⁵, A.A. Snegarev ³⁷, H.L. Snoek ¹¹³, S. Snyder ²⁹, R. Sobie ^{163,v}, A. Soffer ¹⁵⁰, C.A. Solans Sanchez ³⁶, E.Yu. Soldatov ³⁷, U. Soldevila ¹⁶¹, A.A. Solodkov ³⁷, S. Solomon ⁵⁴, A. Soloshenko ³⁸, K. Solovieva ⁵⁴, O.V. Solovyanov ³⁷, V. Solovyev ³⁷, P. Sommer ³⁶, A. Sonay ¹³, W.Y. Song ^{155b}, A. Sopczak ¹³¹, A.L. Soppio ⁹⁵, F. Sopkova ^{28b}, V. Sothilingam ^{63a}, S. Sottocornola ^{72a,72b}, R. Soualah ^{115c}, Z. Soumami ^{35e}, D. South ⁴⁸, S. Spagnolo ^{69a,69b}, M. Spalla ¹⁰⁹, F. Spanò ⁹⁴, D. Sperlich ⁵⁴, G. Spigo ³⁶, M. Spina ¹⁴⁵, S. Spinali ⁹⁰, D.P. Spiteri ⁵⁹, M. Spousta ¹³², E.J. Staats ³⁴, A. Stabile ^{70a,70b}, R. Stamen ^{63a}, M. Stamenkovic ¹¹³, A. Stampekis ²⁰, M. Standke ²⁴, E. Stanecka ⁸⁵, M.V. Stange ⁵⁰, B. Stanislaus ^{17a}, M.M. Stanitzki ⁴⁸, M. Stankaityte ¹²⁵, B. Stapf ⁴⁸, E.A. Starchenko ³⁷, G.H. Stark ¹³⁵, J. Stark ¹⁰¹, D.M. Starke ^{155b}, P. Staroba ¹³⁰, P. Starovoitov ^{63a}, S. Stärz ¹⁰³, R. Staszewski ⁸⁵, G. Stavropoulos ⁴⁶, J. Steentoft ¹⁵⁹, P. Steinberg ²⁹, A.L. Steinhebel ¹²², B. Stelzer ^{141,155a}, H.J. Stelzer ¹²⁸, O. Stelzer-Chilton ^{155a}, H. Stenzel ⁵⁸, T.J. Stevenson ¹⁴⁵, G.A. Stewart ³⁶, M.C. Stockton ³⁶, G. Stoica ^{27b}, M. Stolarski ^{129a}, S. Stonjek ¹⁰⁹, A. Straessner ⁵⁰, J. Strandberg ¹⁴³, S. Strandberg ^{47a,47b}, M. Strauss ¹¹⁹, T. Streblner ¹⁰¹, P. Strizenc ^{28b}, R. Ströhmer ¹⁶⁴, D.M. Strom ¹²², L.R. Strom ⁴⁸, R. Stroynowski ⁴⁴, A. Strubig ^{47a,47b}, S.A. Stucci ²⁹, B. Stugu ¹⁶, J. Stupak ¹¹⁹, N.A. Styles ⁴⁸, D. Su ¹⁴², S. Su ^{62a}, W. Su ^{62d,137,62c}, X. Su ^{62a,66}, K. Sugizaki ¹⁵², V.V. Sulim ³⁷, M.J. Sullivan ⁹¹, D.M.S. Sultan ^{77a,77b}, L. Sultanaliyeva ³⁷, S. Sultansoy ^{3b}, T. Sumida ⁸⁶, S. Sun ¹⁰⁵, S. Sun ¹⁶⁸, O. Sunneborn Gudnadottir ¹⁵⁹, M.R. Sutton ¹⁴⁵, M. Svatos ¹³⁰, M. Swiatlowski ^{155a}, T. Swirski ¹⁶⁴, I. Sykora ^{28a}, M. Sykora ¹³², T. Sykora ¹³², D. Ta ⁹⁹, K. Tackmann ^{48,u}, A. Taffard ¹⁵⁸, R. Tafirout ^{155a}, J.S. Tafuya Vargas ⁶⁶, R.H.M. Taibah ¹²⁶, R. Takashima ⁸⁷, K. Takeda ⁸³, E.P. Takeva ⁵², Y. Takubo ⁸², M. Talby ¹⁰¹, A.A. Talyshev ³⁷, K.C. Tam ^{64b}, N.M. Tamir ¹⁵⁰, A. Tanaka ¹⁵², J. Tanaka ¹⁵², R. Tanaka ⁶⁶, M. Tanasini ^{57b,57a}, J. Tang ^{62c}, Z. Tao ¹⁶², S. Tapia Araya ⁸⁰, S. Tapprogge ⁹⁹, A. Tarek Abouelfadl Mohamed ¹⁰⁶, S. Tarem ¹⁴⁹, K. Tariq ^{62b}, G. Tarna ^{101,27b}, G.F. Tartarelli ^{70a}, P. Tas ¹³², M. Tasevsky ¹³⁰, E. Tassi ^{43b,43a}, A.C. Tate ¹⁶⁰, G. Tateno ¹⁵², Y. Tayalati ^{35e}, G.N. Taylor ¹⁰⁴, W. Taylor ^{155b}, H. Teagle ⁹¹, A.S. Tee ¹⁶⁸, R. Teixeira De Lima ¹⁴², P. Teixeira-Dias ⁹⁴, J.J. Teoh ¹⁵⁴,

K. Terashi ¹⁵², J. Terron ⁹⁸, S. Terzo ¹³, M. Testa ⁵³, R.J. Teuscher ^{154,v}, A. Thaler ⁷⁸,
 O. Theiner ⁵⁶, N. Themistokleous ⁵², T. Thevenaux-Pelzer ¹⁸, O. Thielmann ¹⁶⁹, D.W. Thomas⁹⁴,
 J.P. Thomas ²⁰, E.A. Thompson ⁴⁸, P.D. Thompson ²⁰, E. Thomson ¹²⁷, E.J. Thorpe ⁹³,
 Y. Tian ⁵⁵, V. Tikhomirov ^{37,a}, Yu.A. Tikhonov ³⁷, S. Timoshenko³⁷, E.X.L. Ting ¹, P. Tipton ¹⁷⁰,
 S. Tisserant ¹⁰¹, S.H. Tlou ^{33g}, A. Tnourji ⁴⁰, K. Todome ^{23b,23a}, S. Todorova-Nova ¹³², S. Todt⁵⁰,
 M. Togawa ⁸², J. Tojo ⁸⁸, S. Tokár ^{28a}, K. Tokushuku ⁸², R. Tombs ³², M. Tomoto ^{82,110},
 L. Tompkins ¹⁴², K.W. Topolnicki ^{84b}, P. Tornambe ¹⁰², E. Torrence ¹²², H. Torres ⁵⁰,
 E. Torró Pastor ¹⁶¹, M. Toscani ³⁰, C. Toscirci ³⁹, M. Tost ¹¹, D.R. Tovey ¹³⁸, A. Traeet¹⁶,
 I.S. Trandafir ^{27b}, T. Trefzger ¹⁶⁴, A. Tricoli ²⁹, I.M. Trigger ^{155a}, S. Trincaz-Duvoid ¹²⁶,
 D.A. Trischuk ²⁶, B. Trocmé ⁶⁰, A. Trofymov ⁶⁶, C. Troncon ^{70a}, L. Truong ^{33c},
 M. Trzebinski ⁸⁵, A. Trzuppek ⁸⁵, F. Tsai ¹⁴⁴, M. Tsai ¹⁰⁵, A. Tsiamis ¹⁵¹, P.V. Tsiarehka³⁷,
 S. Tsigaridas ^{155a}, A. Tsigotis ^{151,s}, V. Tsiskaridze ¹⁴⁴, E.G. Tskhadadze^{148a}, M. Tsopoulou ¹⁵¹,
 Y. Tsujikawa ⁸⁶, I.I. Tsukerman ³⁷, V. Tsulaia ^{17a}, S. Tsuno ⁸², O. Tsur¹⁴⁹, D. Tsybychev ¹⁴⁴,
 Y. Tu ^{64b}, A. Tudorache ^{27b}, V. Tudorache ^{27b}, A.N. Tuna ³⁶, S. Turchikhin ³⁸, I. Turk Cakir ^{3a},
 R. Turra ^{70a}, T. Turtuvshin ^{38,w}, P.M. Tuts ⁴¹, S. Tzamarias ¹⁵¹, P. Tzanis ¹⁰, E. Tzovara ⁹⁹,
 K. Uchida¹⁵², F. Ukegawa ¹⁵⁶, P.A. Ulloa Poblete ^{136c}, G. Unal ³⁶, M. Unal ¹¹, A. Undrus ²⁹,
 G. Unel ¹⁵⁸, J. Urban ^{28b}, P. Urquijo ¹⁰⁴, G. Usai ⁸, R. Ushioda ¹⁵³, M. Usman ¹⁰⁷,
 Z. Uysal ^{21b}, L. Vacavant ¹⁰¹, V. Vacek ¹³¹, B. Vachon ¹⁰³, K.O.H. Vadla ¹²⁴, T. Vafeiadis ³⁶,
 A. Vaitkus ⁹⁵, C. Valderanis ¹⁰⁸, E. Valdes Santurio ^{47a,47b}, M. Valente ^{155a}, S. Valentinetti ^{23b,23a},
 A. Valero ¹⁶¹, A. Vallier ¹⁰¹, J.A. Valls Ferrer ¹⁶¹, T.R. Van Daalen ¹³⁷, P. Van Gemmeren ⁶,
 M. Van Rijnbach ^{124,36}, S. Van Stroud ⁹⁵, I. Van Vulpen ¹¹³, M. Vanadia ^{75a,75b}, W. Vandelli ³⁶,
 M. Vandenbroucke ¹³⁴, E.R. Vandewall ¹²⁰, D. Vannicola ¹⁵⁰, L. Vannoli ^{57b,57a}, R. Vari ^{74a},
 E.W. Varnes ⁷, C. Varni ^{17a}, T. Varol ¹⁴⁷, D. Varouchas ⁶⁶, L. Varriale ¹⁶¹, K.E. Varvell ¹⁴⁶,
 M.E. Vasile ^{27b}, L. Vaslin⁴⁰, G.A. Vasquez ¹⁶³, F. Vazeille ⁴⁰, T. Vazquez Schroeder ³⁶,
 J. Veatch ³¹, V. Vecchio ¹⁰⁰, M.J. Veen ¹⁰², I. Veliscek ¹²⁵, L.M. Veloce ¹⁵⁴, F. Veloso ^{129a,129c},
 S. Veneziano ^{74a}, A. Ventura ^{69a,69b}, A. Verbytskyi ¹⁰⁹, M. Verducci ^{73a,73b}, C. Vergis ²⁴,
 M. Verissimo De Araujo ^{81b}, W. Verkerke ¹¹³, J.C. Vermeulen ¹¹³, C. Vernieri ¹⁴²,
 P.J. Verschuuren ⁹⁴, M. Vessella ¹⁰², M.C. Vetterli ^{141,ac}, A. Vgenopoulos ¹⁵¹,
 N. Viaux Maira ^{136f}, T. Vickey ¹³⁸, O.E. Vickey Boeriu ¹³⁸, G.H.A. Viehhauser ¹²⁵, L. Vigani ^{63b},
 M. Villa ^{23b,23a}, M. Villaplana Perez ¹⁶¹, E.M. Villhauer⁵², E. Vilucchi ⁵³, M.G. Vincter ³⁴,
 G.S. Virdee ²⁰, A. Vishwakarma ⁵², C. Vittori ^{23b,23a}, I. Vivarelli ¹⁴⁵, V. Vladimirov¹⁶⁵,
 E. Voevodina ¹⁰⁹, F. Vogel ¹⁰⁸, P. Vokac ¹³¹, J. Von Ahnen ⁴⁸, E. Von Toerne ²⁴,
 B. Vormwald ³⁶, V. Vorobel ¹³², K. Vorobev ³⁷, M. Vos ¹⁶¹, J.H. Vosseveld ⁹¹, M. Vozak ¹¹³,
 L. Vozdecky ⁹³, N. Vranjes ¹⁵, M. Vranjes Milosavljevic ¹⁵, M. Vreeswijk ¹¹³, R. Vuillermet ³⁶,
 O. Vujanovic ⁹⁹, I. Vukotic ³⁹, S. Wada ¹⁵⁶, C. Wagner¹⁰², W. Wagner ¹⁶⁹, S. Wahdan ¹⁶⁹,
 H. Wahlberg ⁸⁹, R. Wakasa ¹⁵⁶, M. Wakida ¹¹⁰, V.M. Walbrecht ¹⁰⁹, J. Walder ¹³³,
 R. Walker ¹⁰⁸, W. Walkowiak ¹⁴⁰, A.M. Wang ⁶¹, A.Z. Wang ¹⁶⁸, C. Wang ^{62a}, C. Wang ^{62c},
 H. Wang ^{17a}, J. Wang ^{64a}, R.-J. Wang ⁹⁹, R. Wang ⁶¹, R. Wang ⁶, S.M. Wang ¹⁴⁷,
 S. Wang ^{62b}, T. Wang ^{62a}, W.T. Wang ⁷⁹, X. Wang ^{14c}, X. Wang ¹⁶⁰, X. Wang ^{62c},
 Y. Wang ^{62d}, Y. Wang ^{14c}, Z. Wang ¹⁰⁵, Z. Wang ^{62d,51,62c}, Z. Wang ¹⁰⁵, A. Warburton ¹⁰³,
 R.J. Ward ²⁰, N. Warrack ⁵⁹, A.T. Watson ²⁰, H. Watson ⁵⁹, M.F. Watson ²⁰, G. Watts ¹³⁷,
 B.M. Waugh ⁹⁵, A.F. Webb ¹¹, C. Weber ²⁹, H.A. Weber ¹⁸, M.S. Weber ¹⁹, S.M. Weber ^{63a},
 C. Wei^{62a}, Y. Wei ¹²⁵, A.R. Weidberg ¹²⁵, J. Weingarten ⁴⁹, M. Weirich ⁹⁹, C. Weiser ⁵⁴,
 C.J. Wells ⁴⁸, T. Wenaus ²⁹, B. Wendland ⁴⁹, T. Wengler ³⁶, N.S. Wenke¹⁰⁹, N. Wermes ²⁴,
 M. Wessels ^{63a}, K. Whalen ¹²², A.M. Wharton ⁹⁰, A.S. White ⁶¹, A. White ⁸, M.J. White ¹,
 D. Whiteson ¹⁵⁸, L. Wickremasinghe ¹²³, W. Wiedenmann ¹⁶⁸, C. Wiel ⁵⁰, M. Wielers ¹³³,
 N. Wieseotte⁹⁹, C. Wiglesworth ⁴², L.A.M. Wiik-Fuchs ⁵⁴, D.J. Wilbern¹¹⁹, H.G. Wilkens ³⁶,

D.M. Williams , H.H. Williams , S. Williams , S. Willocq , P.J. Windischhofer , F. Winklmeier , B.T. Winter , J.K. Winter , M. Wittgen , M. Wobisch , R. Wölker , J. Wollrath , M.W. Wolter , H. Wolters , V.W.S. Wong , A.F. Wongel , S.D. Worm , B.K. Wosiek , K.W. Woźniak , K. Wraight , J. Wu , M. Wu , M. Wu , S.L. Wu , X. Wu , Y. Wu , Z. Wu , J. Wuerzinger , T.R. Wyatt , B.M. Wynne , S. Xella , L. Xia , M. Xia , J. Xiang , X. Xiao , M. Xie , X. Xie , S. Xin , J. Xiong , I. Xiotidis , D. Xu , H. Xu , H. Xu , L. Xu , R. Xu , T. Xu , W. Xu , Y. Xu , Z. Xu , Z. Xu , B. Yabsley , S. Yacoob , N. Yamaguchi , Y. Yamaguchi , H. Yamauchi , T. Yamazaki , Y. Yamazaki , J. Yan , S. Yan , Z. Yan , H.J. Yang , H.T. Yang , S. Yang , T. Yang , X. Yang , X. Yang , Y. Yang , Z. Yang , W-M. Yao , Y.C. Yap , H. Ye , H. Ye , J. Ye , S. Ye , X. Ye , Y. Yeh , I. Yeletsikh , B.K. Yeo , M.R. Yexley , P. Yin , K. Yorita , C.J.S. Young , C. Young , M. Yuan , R. Yuan , L. Yue , X. Yue , M. Zaazoua , B. Zabinski , E. Zaid , T. Zakareishvili , N. Zakharchuk , S. Zambito , J.A. Zamora Saa , J. Zang , D. Zanzi , O. Zaplatilek , S.V. Zeißner , C. Zeitnitz , J.C. Zeng , D.T. Zenger Jr , O. Zenin , T. Ženiš , S. Zenz , S. Zerradi , D. Zerwas , B. Zhang , D.F. Zhang , G. Zhang , J. Zhang , J. Zhang , K. Zhang , L. Zhang , P. Zhang , R. Zhang , S. Zhang , T. Zhang , X. Zhang , X. Zhang , Y. Zhang , Z. Zhang , Z. Zhang , H. Zhao , P. Zhao , T. Zhao , Y. Zhao , Z. Zhao , A. Zhemchugov , X. Zheng , Z. Zheng , D. Zhong , B. Zhou , C. Zhou , H. Zhou , N. Zhou , Y. Zhou , C.G. Zhu , C. Zhu , H.L. Zhu , H. Zhu , J. Zhu , Y. Zhu , Y. Zhu , X. Zhuang , K. Zhukov , V. Zhulanov , N.I. Zimine , J. Zinsser , M. Ziolkowski , L. Živković , A. Zoccoli , K. Zoch , T.G. Zorbas , O. Zormpa , W. Zou , L. Zwalinski .

¹Department of Physics, University of Adelaide, Adelaide; Australia.

²Department of Physics, University of Alberta, Edmonton AB; Canada.

³(^a)Department of Physics, Ankara University, Ankara; (^b)Division of Physics, TOBB University of Economics and Technology, Ankara; Türkiye.

⁴LAPP, Univ. Savoie Mont Blanc, CNRS/IN2P3, Annecy; France.

⁵APC, Université Paris Cité, CNRS/IN2P3, Paris; France.

⁶High Energy Physics Division, Argonne National Laboratory, Argonne IL; United States of America.

⁷Department of Physics, University of Arizona, Tucson AZ; United States of America.

⁸Department of Physics, University of Texas at Arlington, Arlington TX; United States of America.

⁹Physics Department, National and Kapodistrian University of Athens, Athens; Greece.

¹⁰Physics Department, National Technical University of Athens, Zografou; Greece.

¹¹Department of Physics, University of Texas at Austin, Austin TX; United States of America.

¹²Institute of Physics, Azerbaijan Academy of Sciences, Baku; Azerbaijan.

¹³Institut de Física d'Altes Energies (IFAE), Barcelona Institute of Science and Technology, Barcelona; Spain.

¹⁴(^a)Institute of High Energy Physics, Chinese Academy of Sciences, Beijing; (^b)Physics Department, Tsinghua University, Beijing; (^c)Department of Physics, Nanjing University, Nanjing; (^d)University of Chinese Academy of Science (UCAS), Beijing; China.

¹⁵Institute of Physics, University of Belgrade, Belgrade; Serbia.

¹⁶Department for Physics and Technology, University of Bergen, Bergen; Norway.

¹⁷(^a)Physics Division, Lawrence Berkeley National Laboratory, Berkeley CA; (^b)University of California,

Berkeley CA; United States of America.

¹⁸Institut für Physik, Humboldt Universität zu Berlin, Berlin; Germany.

¹⁹Albert Einstein Center for Fundamental Physics and Laboratory for High Energy Physics, University of Bern, Bern; Switzerland.

²⁰School of Physics and Astronomy, University of Birmingham, Birmingham; United Kingdom.

²¹(^a)Department of Physics, Bogazici University, Istanbul; (^b)Department of Physics Engineering, Gaziantep University, Gaziantep; (^c)Department of Physics, Istanbul University, Istanbul; (^d)Istinye University, Sariyer, Istanbul; Türkiye.

²²(^a)Facultad de Ciencias y Centro de Investigaciones, Universidad Antonio Nariño, Bogotá; (^b)Departamento de Física, Universidad Nacional de Colombia, Bogotá; Colombia.

²³(^a)Dipartimento di Fisica e Astronomia A. Righi, Università di Bologna, Bologna; (^b)INFN Sezione di Bologna; Italy.

²⁴Physikalisches Institut, Universität Bonn, Bonn; Germany.

²⁵Department of Physics, Boston University, Boston MA; United States of America.

²⁶Department of Physics, Brandeis University, Waltham MA; United States of America.

²⁷(^a)Transilvania University of Brasov, Brasov; (^b)Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest; (^c)Department of Physics, Alexandru Ioan Cuza University of Iasi, Iasi; (^d)National Institute for Research and Development of Isotopic and Molecular Technologies, Physics Department, Cluj-Napoca; (^e)University Politehnica Bucharest, Bucharest; (^f)West University in Timisoara, Timisoara; (^g)Faculty of Physics, University of Bucharest, Bucharest; Romania.

²⁸(^a)Faculty of Mathematics, Physics and Informatics, Comenius University, Bratislava; (^b)Department of Subnuclear Physics, Institute of Experimental Physics of the Slovak Academy of Sciences, Kosice; Slovak Republic.

²⁹Physics Department, Brookhaven National Laboratory, Upton NY; United States of America.

³⁰Universidad de Buenos Aires, Facultad de Ciencias Exactas y Naturales, Departamento de Física, y CONICET, Instituto de Física de Buenos Aires (IFIBA), Buenos Aires; Argentina.

³¹California State University, CA; United States of America.

³²Cavendish Laboratory, University of Cambridge, Cambridge; United Kingdom.

³³(^a)Department of Physics, University of Cape Town, Cape Town; (^b)iThemba Labs, Western Cape; (^c)Department of Mechanical Engineering Science, University of Johannesburg, Johannesburg; (^d)National Institute of Physics, University of the Philippines Diliman (Philippines); (^e)University of South Africa, Department of Physics, Pretoria; (^f)University of Zululand, KwaDlangezwa; (^g)School of Physics, University of the Witwatersrand, Johannesburg; South Africa.

³⁴Department of Physics, Carleton University, Ottawa ON; Canada.

³⁵(^a)Faculté des Sciences Ain Chock, Réseau Universitaire de Physique des Hautes Energies - Université Hassan II, Casablanca; (^b)Faculté des Sciences, Université Ibn-Tofail, Kénitra; (^c)Faculté des Sciences Semlalia, Université Cadi Ayyad, LPHEA-Marrakech; (^d)LPMP, Faculté des Sciences, Université Mohamed Premier, Oujda; (^e)Faculté des sciences, Université Mohammed V, Rabat; (^f)Institute of Applied Physics, Mohammed VI Polytechnic University, Ben Guerir; Morocco.

³⁶CERN, Geneva; Switzerland.

³⁷Affiliated with an institute covered by a cooperation agreement with CERN.

³⁸Affiliated with an international laboratory covered by a cooperation agreement with CERN.

³⁹Enrico Fermi Institute, University of Chicago, Chicago IL; United States of America.

⁴⁰LPC, Université Clermont Auvergne, CNRS/IN2P3, Clermont-Ferrand; France.

⁴¹Nevis Laboratory, Columbia University, Irvington NY; United States of America.

⁴²Niels Bohr Institute, University of Copenhagen, Copenhagen; Denmark.

⁴³(^a)Dipartimento di Fisica, Università della Calabria, Rende; (^b)INFN Gruppo Collegato di Cosenza,

Laboratori Nazionali di Frascati; Italy.

⁴⁴Physics Department, Southern Methodist University, Dallas TX; United States of America.

⁴⁵Physics Department, University of Texas at Dallas, Richardson TX; United States of America.

⁴⁶National Centre for Scientific Research "Demokritos", Agia Paraskevi; Greece.

⁴⁷(^a) Department of Physics, Stockholm University; (^b) Oskar Klein Centre, Stockholm; Sweden.

⁴⁸Deutsches Elektronen-Synchrotron DESY, Hamburg and Zeuthen; Germany.

⁴⁹Fakultät Physik, Technische Universität Dortmund, Dortmund; Germany.

⁵⁰Institut für Kern- und Teilchenphysik, Technische Universität Dresden, Dresden; Germany.

⁵¹Department of Physics, Duke University, Durham NC; United States of America.

⁵²SUPA - School of Physics and Astronomy, University of Edinburgh, Edinburgh; United Kingdom.

⁵³INFN e Laboratori Nazionali di Frascati, Frascati; Italy.

⁵⁴Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, Freiburg; Germany.

⁵⁵II. Physikalisches Institut, Georg-August-Universität Göttingen, Göttingen; Germany.

⁵⁶Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève; Switzerland.

⁵⁷(^a) Dipartimento di Fisica, Università di Genova, Genova; (^b) INFN Sezione di Genova; Italy.

⁵⁸II. Physikalisches Institut, Justus-Liebig-Universität Giessen, Giessen; Germany.

⁵⁹SUPA - School of Physics and Astronomy, University of Glasgow, Glasgow; United Kingdom.

⁶⁰LPSC, Université Grenoble Alpes, CNRS/IN2P3, Grenoble INP, Grenoble; France.

⁶¹Laboratory for Particle Physics and Cosmology, Harvard University, Cambridge MA; United States of America.

⁶²(^a) Department of Modern Physics and State Key Laboratory of Particle Detection and Electronics, University of Science and Technology of China, Hefei; (^b) Institute of Frontier and Interdisciplinary Science and Key Laboratory of Particle Physics and Particle Irradiation (MOE), Shandong University, Qingdao; (^c) School of Physics and Astronomy, Shanghai Jiao Tong University, Key Laboratory for Particle Astrophysics and Cosmology (MOE), SKLPPC, Shanghai; (^d) Tsung-Dao Lee Institute, Shanghai; China.

⁶³(^a) Kirchhoff-Institut für Physik, Ruprecht-Karls-Universität Heidelberg, Heidelberg; (^b) Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg; Germany.

⁶⁴(^a) Department of Physics, Chinese University of Hong Kong, Shatin, N.T., Hong Kong; (^b) Department of Physics, University of Hong Kong, Hong Kong; (^c) Department of Physics and Institute for Advanced Study, Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong; China.

⁶⁵Department of Physics, National Tsing Hua University, Hsinchu; Taiwan.

⁶⁶IJCLab, Université Paris-Saclay, CNRS/IN2P3, 91405, Orsay; France.

⁶⁷Department of Physics, Indiana University, Bloomington IN; United States of America.

⁶⁸(^a) INFN Gruppo Collegato di Udine, Sezione di Trieste, Udine; (^b) ICTP, Trieste; (^c) Dipartimento Politecnico di Ingegneria e Architettura, Università di Udine, Udine; Italy.

⁶⁹(^a) INFN Sezione di Lecce; (^b) Dipartimento di Matematica e Fisica, Università del Salento, Lecce; Italy.

⁷⁰(^a) INFN Sezione di Milano; (^b) Dipartimento di Fisica, Università di Milano, Milano; Italy.

⁷¹(^a) INFN Sezione di Napoli; (^b) Dipartimento di Fisica, Università di Napoli, Napoli; Italy.

⁷²(^a) INFN Sezione di Pavia; (^b) Dipartimento di Fisica, Università di Pavia, Pavia; Italy.

⁷³(^a) INFN Sezione di Pisa; (^b) Dipartimento di Fisica E. Fermi, Università di Pisa, Pisa; Italy.

⁷⁴(^a) INFN Sezione di Roma; (^b) Dipartimento di Fisica, Sapienza Università di Roma, Roma; Italy.

⁷⁵(^a) INFN Sezione di Roma Tor Vergata; (^b) Dipartimento di Fisica, Università di Roma Tor Vergata, Roma; Italy.

⁷⁶(^a) INFN Sezione di Roma Tre; (^b) Dipartimento di Matematica e Fisica, Università Roma Tre, Roma; Italy.

⁷⁷(^a) INFN-TIFPA; (^b) Università degli Studi di Trento, Trento; Italy.

⁷⁸Universität Innsbruck, Department of Astro and Particle Physics, Innsbruck; Austria.

- ⁷⁹University of Iowa, Iowa City IA; United States of America.
- ⁸⁰Department of Physics and Astronomy, Iowa State University, Ames IA; United States of America.
- ⁸¹(^a)Departamento de Engenharia Elétrica, Universidade Federal de Juiz de Fora (UFJF), Juiz de Fora; (^b)Universidade Federal do Rio De Janeiro COPPE/EE/IF, Rio de Janeiro; (^c)Instituto de Física, Universidade de São Paulo, São Paulo; (^d)Rio de Janeiro State University, Rio de Janeiro; Brazil.
- ⁸²KEK, High Energy Accelerator Research Organization, Tsukuba; Japan.
- ⁸³Graduate School of Science, Kobe University, Kobe; Japan.
- ⁸⁴(^a)AGH University of Science and Technology, Faculty of Physics and Applied Computer Science, Krakow; (^b)Marian Smoluchowski Institute of Physics, Jagiellonian University, Krakow; Poland.
- ⁸⁵Institute of Nuclear Physics Polish Academy of Sciences, Krakow; Poland.
- ⁸⁶Faculty of Science, Kyoto University, Kyoto; Japan.
- ⁸⁷Kyoto University of Education, Kyoto; Japan.
- ⁸⁸Research Center for Advanced Particle Physics and Department of Physics, Kyushu University, Fukuoka ; Japan.
- ⁸⁹Instituto de Física La Plata, Universidad Nacional de La Plata and CONICET, La Plata; Argentina.
- ⁹⁰Physics Department, Lancaster University, Lancaster; United Kingdom.
- ⁹¹Oliver Lodge Laboratory, University of Liverpool, Liverpool; United Kingdom.
- ⁹²Department of Experimental Particle Physics, Jožef Stefan Institute and Department of Physics, University of Ljubljana, Ljubljana; Slovenia.
- ⁹³School of Physics and Astronomy, Queen Mary University of London, London; United Kingdom.
- ⁹⁴Department of Physics, Royal Holloway University of London, Egham; United Kingdom.
- ⁹⁵Department of Physics and Astronomy, University College London, London; United Kingdom.
- ⁹⁶Louisiana Tech University, Ruston LA; United States of America.
- ⁹⁷Fysiska institutionen, Lunds universitet, Lund; Sweden.
- ⁹⁸Departamento de Física Teórica C-15 and CIAFF, Universidad Autónoma de Madrid, Madrid; Spain.
- ⁹⁹Institut für Physik, Universität Mainz, Mainz; Germany.
- ¹⁰⁰School of Physics and Astronomy, University of Manchester, Manchester; United Kingdom.
- ¹⁰¹CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille; France.
- ¹⁰²Department of Physics, University of Massachusetts, Amherst MA; United States of America.
- ¹⁰³Department of Physics, McGill University, Montreal QC; Canada.
- ¹⁰⁴School of Physics, University of Melbourne, Victoria; Australia.
- ¹⁰⁵Department of Physics, University of Michigan, Ann Arbor MI; United States of America.
- ¹⁰⁶Department of Physics and Astronomy, Michigan State University, East Lansing MI; United States of America.
- ¹⁰⁷Group of Particle Physics, University of Montreal, Montreal QC; Canada.
- ¹⁰⁸Fakultät für Physik, Ludwig-Maximilians-Universität München, München; Germany.
- ¹⁰⁹Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), München; Germany.
- ¹¹⁰Graduate School of Science and Kobayashi-Maskawa Institute, Nagoya University, Nagoya; Japan.
- ¹¹¹Department of Physics and Astronomy, University of New Mexico, Albuquerque NM; United States of America.
- ¹¹²Institute for Mathematics, Astrophysics and Particle Physics, Radboud University/Nikhef, Nijmegen; Netherlands.
- ¹¹³Nikhef National Institute for Subatomic Physics and University of Amsterdam, Amsterdam; Netherlands.
- ¹¹⁴Department of Physics, Northern Illinois University, DeKalb IL; United States of America.
- ¹¹⁵(^a)New York University Abu Dhabi, Abu Dhabi; (^b)United Arab Emirates University, Al Ain; (^c)University of Sharjah, Sharjah; United Arab Emirates.

- ¹¹⁶Department of Physics, New York University, New York NY; United States of America.
- ¹¹⁷Ochanomizu University, Otsuka, Bunkyo-ku, Tokyo; Japan.
- ¹¹⁸Ohio State University, Columbus OH; United States of America.
- ¹¹⁹Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, Norman OK; United States of America.
- ¹²⁰Department of Physics, Oklahoma State University, Stillwater OK; United States of America.
- ¹²¹Palacký University, Joint Laboratory of Optics, Olomouc; Czech Republic.
- ¹²²Institute for Fundamental Science, University of Oregon, Eugene, OR; United States of America.
- ¹²³Graduate School of Science, Osaka University, Osaka; Japan.
- ¹²⁴Department of Physics, University of Oslo, Oslo; Norway.
- ¹²⁵Department of Physics, Oxford University, Oxford; United Kingdom.
- ¹²⁶LPNHE, Sorbonne Université, Université Paris Cité, CNRS/IN2P3, Paris; France.
- ¹²⁷Department of Physics, University of Pennsylvania, Philadelphia PA; United States of America.
- ¹²⁸Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh PA; United States of America.
- ¹²⁹(^a) Laboratório de Instrumentação e Física Experimental de Partículas - LIP, Lisboa; (^b) Departamento de Física, Faculdade de Ciências, Universidade de Lisboa, Lisboa; (^c) Departamento de Física, Universidade de Coimbra, Coimbra; (^d) Centro de Física Nuclear da Universidade de Lisboa, Lisboa; (^e) Departamento de Física, Universidade do Minho, Braga; (^f) Departamento de Física Teórica y del Cosmos, Universidad de Granada, Granada (Spain); (^g) Departamento de Física, Instituto Superior Técnico, Universidade de Lisboa, Lisboa; Portugal.
- ¹³⁰Institute of Physics of the Czech Academy of Sciences, Prague; Czech Republic.
- ¹³¹Czech Technical University in Prague, Prague; Czech Republic.
- ¹³²Charles University, Faculty of Mathematics and Physics, Prague; Czech Republic.
- ¹³³Particle Physics Department, Rutherford Appleton Laboratory, Didcot; United Kingdom.
- ¹³⁴IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette; France.
- ¹³⁵Santa Cruz Institute for Particle Physics, University of California Santa Cruz, Santa Cruz CA; United States of America.
- ¹³⁶(^a) Departamento de Física, Pontificia Universidad Católica de Chile, Santiago; (^b) Millennium Institute for Subatomic physics at high energy frontier (SAPHIR), Santiago; (^c) Instituto de Investigación Multidisciplinario en Ciencia y Tecnología, y Departamento de Física, Universidad de La Serena; (^d) Universidad Andres Bello, Department of Physics, Santiago; (^e) Instituto de Alta Investigación, Universidad de Tarapacá, Arica; (^f) Departamento de Física, Universidad Técnica Federico Santa María, Valparaíso; Chile.
- ¹³⁷Department of Physics, University of Washington, Seattle WA; United States of America.
- ¹³⁸Department of Physics and Astronomy, University of Sheffield, Sheffield; United Kingdom.
- ¹³⁹Department of Physics, Shinshu University, Nagano; Japan.
- ¹⁴⁰Department Physik, Universität Siegen, Siegen; Germany.
- ¹⁴¹Department of Physics, Simon Fraser University, Burnaby BC; Canada.
- ¹⁴²SLAC National Accelerator Laboratory, Stanford CA; United States of America.
- ¹⁴³Department of Physics, Royal Institute of Technology, Stockholm; Sweden.
- ¹⁴⁴Departments of Physics and Astronomy, Stony Brook University, Stony Brook NY; United States of America.
- ¹⁴⁵Department of Physics and Astronomy, University of Sussex, Brighton; United Kingdom.
- ¹⁴⁶School of Physics, University of Sydney, Sydney; Australia.
- ¹⁴⁷Institute of Physics, Academia Sinica, Taipei; Taiwan.
- ¹⁴⁸(^a) E. Andronikashvili Institute of Physics, Iv. Javakhishvili Tbilisi State University, Tbilisi; (^b) High

- Energy Physics Institute, Tbilisi State University, Tbilisi;^(c) University of Georgia, Tbilisi; Georgia.
- ¹⁴⁹ Department of Physics, Technion, Israel Institute of Technology, Haifa; Israel.
- ¹⁵⁰ Raymond and Beverly Sackler School of Physics and Astronomy, Tel Aviv University, Tel Aviv; Israel.
- ¹⁵¹ Department of Physics, Aristotle University of Thessaloniki, Thessaloniki; Greece.
- ¹⁵² International Center for Elementary Particle Physics and Department of Physics, University of Tokyo, Tokyo; Japan.
- ¹⁵³ Department of Physics, Tokyo Institute of Technology, Tokyo; Japan.
- ¹⁵⁴ Department of Physics, University of Toronto, Toronto ON; Canada.
- ¹⁵⁵ ^(a) TRIUMF, Vancouver BC; ^(b) Department of Physics and Astronomy, York University, Toronto ON; Canada.
- ¹⁵⁶ Division of Physics and Tomonaga Center for the History of the Universe, Faculty of Pure and Applied Sciences, University of Tsukuba, Tsukuba; Japan.
- ¹⁵⁷ Department of Physics and Astronomy, Tufts University, Medford MA; United States of America.
- ¹⁵⁸ Department of Physics and Astronomy, University of California Irvine, Irvine CA; United States of America.
- ¹⁵⁹ Department of Physics and Astronomy, University of Uppsala, Uppsala; Sweden.
- ¹⁶⁰ Department of Physics, University of Illinois, Urbana IL; United States of America.
- ¹⁶¹ Instituto de Física Corpuscular (IFIC), Centro Mixto Universidad de Valencia - CSIC, Valencia; Spain.
- ¹⁶² Department of Physics, University of British Columbia, Vancouver BC; Canada.
- ¹⁶³ Department of Physics and Astronomy, University of Victoria, Victoria BC; Canada.
- ¹⁶⁴ Fakultät für Physik und Astronomie, Julius-Maximilians-Universität Würzburg, Würzburg; Germany.
- ¹⁶⁵ Department of Physics, University of Warwick, Coventry; United Kingdom.
- ¹⁶⁶ Waseda University, Tokyo; Japan.
- ¹⁶⁷ Department of Particle Physics and Astrophysics, Weizmann Institute of Science, Rehovot; Israel.
- ¹⁶⁸ Department of Physics, University of Wisconsin, Madison WI; United States of America.
- ¹⁶⁹ Fakultät für Mathematik und Naturwissenschaften, Fachgruppe Physik, Bergische Universität Wuppertal, Wuppertal; Germany.
- ¹⁷⁰ Department of Physics, Yale University, New Haven CT; United States of America.
- ^a Also Affiliated with an institute covered by a cooperation agreement with CERN.
- ^b Also at Borough of Manhattan Community College, City University of New York, New York NY; United States of America.
- ^c Also at Bruno Kessler Foundation, Trento; Italy.
- ^d Also at Center for High Energy Physics, Peking University; China.
- ^e Also at Centro Studi e Ricerche Enrico Fermi; Italy.
- ^f Also at CERN, Geneva; Switzerland.
- ^g Also at Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève; Switzerland.
- ^h Also at Departament de Física de la Universitat Autònoma de Barcelona, Barcelona; Spain.
- ⁱ Also at Department of Financial and Management Engineering, University of the Aegean, Chios; Greece.
- ^j Also at Department of Physics and Astronomy, Michigan State University, East Lansing MI; United States of America.
- ^k Also at Department of Physics and Astronomy, University of Louisville, Louisville, KY; United States of America.
- ^l Also at Department of Physics, Ben Gurion University of the Negev, Beer Sheva; Israel.
- ^m Also at Department of Physics, California State University, East Bay; United States of America.
- ⁿ Also at Department of Physics, California State University, Sacramento; United States of America.
- ^o Also at Department of Physics, King's College London, London; United Kingdom.

- p* Also at Department of Physics, University of Fribourg, Fribourg; Switzerland.
- q* Also at Department of Physics, University of Thessaly; Greece.
- r* Also at Department of Physics, Westmont College, Santa Barbara; United States of America.
- s* Also at Hellenic Open University, Patras; Greece.
- t* Also at Institucio Catalana de Recerca i Estudis Avancats, ICREA, Barcelona; Spain.
- u* Also at Institut für Experimentalphysik, Universität Hamburg, Hamburg; Germany.
- v* Also at Institute of Particle Physics (IPP); Canada.
- w* Also at Institute of Physics and Technology, Ulaanbaatar; Mongolia.
- x* Also at Institute of Physics, Azerbaijan Academy of Sciences, Baku; Azerbaijan.
- y* Also at Institute of Theoretical Physics, Ilia State University, Tbilisi; Georgia.
- z* Also at Lawrence Livermore National Laboratory, Livermore; United States of America.
- aa* Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen; Germany.
- ab* Also at The Collaborative Innovation Center of Quantum Matter (CICQM), Beijing; China.
- ac* Also at TRIUMF, Vancouver BC; Canada.
- ad* Also at Università di Napoli Parthenope, Napoli; Italy.
- ae* Also at University of Chinese Academy of Sciences (UCAS), Beijing; China.
- af* Also at University of Colorado Boulder, Department of Physics, Colorado; United States of America.
- ag* Also at Washington College, Maryland; United States of America.
- ah* Also at Yeditepe University, Physics Department, Istanbul; Türkiye.
- * Deceased