


Evidence of a $J/\psi K_S^0$ Structure in $B^0 \rightarrow J/\psi \phi K_S^0$ DecaysR. Aaij *et al.**
(LHCb Collaboration) (Received 17 January 2023; accepted 21 March 2023; published 26 September 2023)

An amplitude analysis of $B^0 \rightarrow J/\psi \phi K_S^0$ decays is performed using proton-proton collision data, corresponding to an integrated luminosity of 9 fb^{-1} , collected with the LHCb detector at center-of-mass energies of 7, 8, and 13 TeV. Evidence with a significance of 4.0 standard deviations of a structure in the $J/\psi K_S^0$ system, named $T_{\psi s 1}^{\theta}(4000)^0$, is seen, with its mass and width measured to be $3991_{-10}^{+12} {}_{-17}^{+9} \text{ MeV}/c^2$ and $105_{-25}^{+29} {}_{-23}^{+17} \text{ MeV}$, respectively, where the first uncertainty is statistical and the second systematic. The $T_{\psi s 1}^{\theta}(4000)^0$ state is likely to be the isospin partner of the $T_{\psi s 1}^{\theta}(4000)^+$ state, previously observed in the $J/\psi K^+$ system of the $B^+ \rightarrow J/\psi \phi K^+$ decay. When isospin symmetry for the charged and neutral $T_{\psi s 1}^{\theta}(4000)$ states is assumed, the signal significance increases to 5.4 standard deviations.

DOI: [10.1103/PhysRevLett.131.131901](https://doi.org/10.1103/PhysRevLett.131.131901)

Hadrons formed by more than three quarks, namely exotic hadrons, can have complex color and flavor structures, and studies of the structures shed light on the nature of the strong interaction. Since the discovery of the $\chi_{c1}(3872)$ [1], several states compatible with a four or five quark composition have been observed [2], including the fully charmed tetraquark states [3–5] and the pentaquark states [6,7]. In 2020, the BESIII Collaboration reported the observation of the $Z_{cs}(3985)^+$ state in the $D_s^+ \bar{D}^{*0}$ and $D_s^{*+} \bar{D}^0$ mass spectra [8]. (Charge conjugation is implied throughout the Letter unless specified otherwise.) Two similar states, $T_{\psi s 1}^{\theta}(4000)^+$ and $T_{\psi s 1}(4220)^+$, were observed in $B^+ \rightarrow J/\psi \phi K^+$ decays by the LHCb experiment [9,10]. The observations of the $T_{\psi s 1}^{\theta}(4000)^+$, $T_{\psi s 1}(4220)^+$, and $Z_{cs}(3985)^+$ states have stimulated many theoretical studies to interpret their internal structures, such as hadronic molecules [12], compact tetraquarks [13], or threshold effects [14].

Searching for $T_{\psi s}^0$, the isospin partner of the $T_{\psi s}^+$, is an important step in identifying the full SU(3) nonet that involves these states. The quark contents for the $T_{\psi s}^+$ and $T_{\psi s}^0$ states are $c\bar{c}u\bar{s}$ and $c\bar{c}d\bar{s}$, respectively. Recently, the BESIII experiment reported evidence for the $Z_{cs}(3985)^0$ state ($c\bar{c}d\bar{s}$) in the $D_s^+ D^{*-}$ and $D_s^{*+} D^-$ mass spectra [15]. The $B^0 \rightarrow J/\psi \phi K_S^0$ decay [16–18] is the ideal candidate to search for such $T_{\psi s}^0$ states. The quark level processes of the

$B^0 \rightarrow J/\psi \phi K_S^0$ decay are similar to those of the $B^+ \rightarrow J/\psi \phi K^+$ decay, as shown in Figs. 1(c) and 1(d) of Ref. [17], and the two decays are related by isospin symmetry.

In this Letter, evidence of a $J/\psi K_S^0$ structure is reported by making use of an amplitude analysis of $B^0 \rightarrow J/\psi \phi K_S^0$ decays. The analysis is based on proton-proton (pp) collision data, collected with the LHCb detector at center-of-mass energies of 7, 8, and 13 TeV, corresponding to integrated luminosities of 1, 2, and 6 fb^{-1} , respectively.

The LHCb detector is a single-arm forward spectrometer covering the pseudorapidity range $2 < \eta < 5$, described in detail in Refs. [19,20]. Simulated samples of the signal decays are used to estimate the effect of reconstruction and event selections. The samples are generated with PYTHIA [21], EvtGen [22], and the GEANT4 toolkit [23] as described in Ref. [24].

For the $B^0 \rightarrow J/\psi \phi K_S^0$ decays, the J/ψ , ϕ , and K_S^0 candidates are reconstructed by combining two oppositely charged muons, kaons, and pions, respectively, forming vertices detached from any primary vertex (PV). The B^0 candidate is reconstructed by combining the J/ψ , ϕ , and K_S^0 candidates, and the resulting decay vertex must be of high quality. In order to improve the resolution of the mass of the B^0 candidates (referred to as $m_{J/\psi \phi K_S^0}$), a kinematic constraint [25] is applied, where the momenta of the final-state particles are recalculated constraining the masses of the J/ψ and K_S^0 candidates to their nominal values, and the momentum of B^0 candidate pointing back to its associated PV. The associated PV is the one with the smallest impact parameter χ^2 , defined as the difference in the vertex fit χ^2 of a given PV with and without the particle under consideration.

The B^0 candidates are selected by a multivariate classifier based on a multilayer perceptron [26,27].

*Full author list given at the end of the Letter.

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

The selection is similar to that used in Ref. [9], with the additional requirement on the significance of the flight distance of the K_S^0 from the B^0 decay vertex. Only candidates with the mass of the K^+K^- system within ± 15 MeV around the nominal ϕ mass [2] are kept, where the mass resolution is about 0.7 MeV. (Natural units with $\hbar = c = 1$ are used throughout the Letter.) The multivariate classifier uses 11 variables related to the decay-chain topology, particle transverse momentum, vertex fit quality, and charged particle identification information. The selection criterion on the classifier response is chosen by maximizing the figure of merit, $S^2/(S+B)^{3/2}$ [28], where S and B are the yields of signal and background in the signal region, respectively. The signal region in $m_{J/\psi\phi K_S^0}$ is defined to be ± 15 MeV around the nominal B^0 mass [2].

An extended maximum-likelihood fit is performed to the reconstructed B^0 mass, as is shown in Fig. 1(a). The B^0 signal is described by a Hypatia function [29] and the background is described by an exponential function. The B^0 yield is measured to be 1866 ± 47 in the signal region. The fraction of combinatorial background in the signal region is 6%. Roughly 4% of the B^0 yield corresponds to peaking background $B^0 \rightarrow J/\psi K^+ K^- K_S^0$ decays without an intermediate ϕ meson (referred to as non- ϕ). The non- ϕ contribution is neglected in the default amplitude model and is considered as a source of systematic uncertainty. A further kinematic fit is performed to improve the momentum resolution of the final-state particles by constraining the measured B^0 mass to its nominal value. Figure 1(b) shows the distribution of $m_{J/\psi K_S^0}^2$ versus $m_{J/\psi\phi}^2$ for candidates in the signal region.

An amplitude fit is performed to study the intermediate states. The $B^0 \rightarrow J/\psi\phi K_S^0$ sample size is small, while the intermediate contributions are complicated, making the fit difficult to converge. In order to solve this difficulty, a simultaneous fit is performed to the $B^0 \rightarrow J/\psi\phi K_S^0$ sample and the $B^+ \rightarrow J/\psi\phi K^+$ sample in Ref. [9], where the amplitudes in the two decay modes are related through

isospin symmetry. The likelihood for the B^0 decay is given by

$$\mathcal{L}(\vec{\omega}) = \prod_i [(1 - \beta)\mathcal{P}_{\text{sig}}(m_{\phi K_S^0}^i, \vec{\Omega}^i | \vec{\omega}) + \beta\mathcal{P}_{\text{bkg}}(m_{\phi K_S^0}^i, \vec{\Omega}^i)], \quad (1)$$

where $\mathcal{P}_{\text{sig}}(m_{\phi K_S^0}^i, \vec{\Omega}^i | \vec{\omega})$ and $\mathcal{P}_{\text{bkg}}(m_{\phi K_S^0}^i, \vec{\Omega}^i)$ refer to the probability density functions (PDFs) for the signal and background components, respectively. The superscript i refers to the i th candidate. The fraction of combinatorial background, β , is fixed to 6%. The likelihood of the B^+ decay is similar to Eq. (1), and is described in detail in Ref. [30]. The decay kinematics are described by a mass, chosen to be $m_{\phi K_S^0}$, and five angular variables $\vec{\Omega}$ as defined below. The signal PDF is proportional to the incoherent sum of the squared matrix elements for different final-state muon helicities $(\lambda_{\mu^+}, \lambda_{\mu^-})$, and depends on the set of fit parameters, $\vec{\omega}$, which includes masses, widths, and helicity couplings of intermediate states contributing to $B^0 \rightarrow J/\psi\phi K_S^0$ decays,

$$\mathcal{P}_{\text{sig}}(m_{\phi K_S^0}^i, \vec{\Omega}^i | \vec{\omega}) = \frac{1}{I(\vec{\omega})} \sum_{\lambda_{\mu^+}, \lambda_{\mu^-}} \sum_f |\mathcal{M}_f(m_{\phi K_S^0}^i, \vec{\Omega}^i | \vec{\omega})|^2 \times \Phi(m_{\phi K_S^0}^i, \vec{\Omega}^i) \epsilon(m_{\phi K_S^0}^i, \vec{\Omega}^i). \quad (2)$$

The term $\Phi(m_{\phi K_S^0}^i, \vec{\Omega}^i)$ represents the phase-space density, the term $\epsilon(m_{\phi K_S^0}^i, \vec{\Omega}^i)$ represents the efficiency, and the term $I(\vec{\omega})$ is a normalization factor. The efficiency is obtained from simulated samples. Since the $B^0 \rightarrow J/\psi\phi K_S^0$ decay is a self-charge-conjugated mode, the data sample contains both B^0 and \bar{B}^0 decays with unknown b flavor. Therefore, the signal PDF for the neutral B decay averages B^0 and \bar{B}^0 contributions, where the flavor of B candidates is denoted as f in Eq. (2).




FIG. 1. (a) Invariant-mass distribution of selected B^0 candidates and corresponding fit result. (b) Distribution of $m_{J/\psi K_S^0}^2$ versus $m_{J/\psi\phi}^2$ for candidates in the ± 15 MeV region around the nominal B^0 mass.




FIG. 2. Distributions of (left) $m_{\phi K}$, (middle) $m_{J/\psi\phi}$, and (right) $m_{J/\psi K}$, overlaid with the corresponding projections of the default fit model. The upper and lower rows correspond to the $B^+ \rightarrow J/\psi\phi K^+$ and $B^0 \rightarrow J/\psi\phi K_S^0$ decays, respectively.

The matrix element $\mathcal{M}_f(m_{\phi K_S^0}^i, \vec{\Omega}^i | \vec{\omega})$ is constructed based on the helicity formalism [31]. Three interfering decay sequences are considered: $B^0 \rightarrow (K^* \rightarrow \phi K_S^0)J/\psi$, $B^0 \rightarrow (X \rightarrow J/\psi\phi)K_S^0$, and $B^0 \rightarrow (T_{\psi s1} \rightarrow J/\psi K_S^0)\phi$. (The K^* denotes any excited K state. The X denotes $\chi_{c0,1}$, η_{c2} , and $T_{\psi\phi 1}^n$ states.) The decay sequences can be described by $m_{\phi K_S^0}$ and $\vec{\Omega} \equiv (\theta_{K^*}, \theta_{J/\psi}, \theta_\phi, \Delta\varphi_{K^*J/\psi}, \Delta\varphi_{K^*\phi})$, where the θ_{K^*} , $\theta_{J/\psi}$, and θ_ϕ are the helicity angles of the ϕ , μ , and K^+ particles in the K^* , J/ψ , and ϕ rest frames, respectively. The angles between the K^* and J/ψ decay planes, and between the K^* and ϕ decay planes in the B rest frame, are denoted as $\Delta\varphi_{K^*J/\psi}$, and $\Delta\varphi_{K^*\phi}$, respectively. The background PDF $\mathcal{P}_{\text{bkg}}(m_{\phi K_S^0}^i, \vec{\Omega}^i)$ is determined from studying the candidates in the sideband of the $m_{J/\psi\phi K_S^0}$ distribution with the methods outlined in Ref. [30].

The default model is taken from Ref. [9] and is composed of nine K^* components, seven X components, and two $T_{\psi s1}$ states [$T_{\psi s1}^\theta(4000)$ and $T_{\psi s1}(4220)$], and one $J/\psi\phi$ nonresonant component (NR). Relativistic Breit–Wigner (RBW) functions are used to model the line shapes of these resonances. The amplitudes of the B^0 and the B^+ decays are formed by intermediate states that are either identical (namely X) or isospin partners (K^* or $T_{\psi s1}$). Under the assumption of isospin symmetry, the mass, width, and helicity couplings for all the components except for the $T_{\psi s1}^\theta(4000)$ states are constrained to be identical. In order to test the significance of the $T_{\psi s1}^\theta(4000)^0$ signal without prior knowledge of its properties, a fit is performed with the masses, widths, and helicity couplings of $T_{\psi s1}^\theta(4000)^0$ and $T_{\psi s1}^\theta(4000)^+$ states allowed to vary independently. In an alternative model, isospin symmetry is imposed for the $T_{\psi s1}^\theta(4000)^0$ and $T_{\psi s1}^\theta(4000)^+$ components. The parameters for the $T_{\psi s1}(4220)^0$ state are always

constrained to be identical to the $T_{\psi s1}(4220)^+$ state due to the limited size of the B^0 sample.

Figure 2 shows the ϕK , $J/\psi\phi$, and $J/\psi K$ mass distributions and the corresponding fit projections of the default model for the two B decay modes. Table I summarizes measurements of the mass, width, and fit fraction of the $T_{\psi s1}^\theta(4000)^0$ state, and the mass difference between the $T_{\psi s1}^\theta(4000)^0$ and $T_{\psi s1}^\theta(4000)^+$ states, defined as $\Delta M \equiv M_{T_{\psi s1}^\theta(4000)^0} - M_{T_{\psi s1}^\theta(4000)^+}$. The fit value of ΔM is zero within uncertainties, consistent with the two states being isospin partners. The fit fraction of each component is defined as the integral of the signal PDF divided by the $I(\vec{\omega})$ term. All the fit parameters, including mass, width, and helicity couplings of the intermediate states, of the default model in this analysis are consistent with the corresponding parameters of the default model for the $B^+ \rightarrow J/\psi\phi K^+$ decay in Ref. [9].

The estimated systematic uncertainties on the mass, width, and fit fraction of the $T_{\psi s1}^\theta(4000)^0$ state, and on ΔM are summarized in Table II. Both the background PDF and efficiency function are described by an expansion with Legendre polynomials and a spherical harmonic function instead of interpolation. The Blatt-Weisskopf hadron size is varied between 1.5 and 4.5 GeV^{-1} from the default value 3 GeV^{-1} . The Flatté model [32] is used for the line shape of the $X(4140)$ state instead of the RBW function to account

TABLE I. Results for the $T_{\psi s1}^\theta(4000)^0$ state from the default model. The first uncertainty is statistical and the second systematic.

State	Mass (MeV)	Width (MeV)	Fit fraction (%)	ΔM (MeV)
$T_{\psi s1}^\theta(4000)^0$	$3991^{+12}_{-10} +9_{-17}$	$105^{+29}_{-25} +17_{-23}$	$7.9 \pm 2.5^{+3.0}_{-2.8}$	$-12^{+11}_{-10} +6_{-4}$

TABLE II. Systematic uncertainties associated to the mass, width, and fit fraction of the $T_{\psi s1}^{\theta}(4000)^0$ state, and the mass difference between the $T_{\psi s1}^{\theta}(4000)^0$ and $T_{\psi s1}^{\theta}(4000)^+$ states.

Source	Mass (MeV)	Width (MeV)	Fit fraction (%)	ΔM (MeV)
Efficiency and background models	+10.46	+1.91	-0.06	+1.02
Hadron radius	+0.29 -1.92	+2.33 -5.85	+0.05 -0.39	+0.92 -1.68
$X(4140)$ Flatté model	+0.61	-3.47	-0.31	-0.18
$J/\psi\phi$ NR representation	+1.72	+1.46	+0.38	+0.12
Simplified K-matrix model	-7.30	-17.54	-1.73	-3.38
RBW widths	+1.90	-3.41	-0.56	+1.70
Spin parity of the $T_{\psi s1}(4220)$ state	-16.32	-17.76	-2.33	-1.48
Non- ϕ contribution	+5.52	+7.98	+0.92	+5.49
Additional 1^+ $J/\psi\phi$ NR	+0.80	-3.66	+0.18	-1.28
Additional 2^+ $J/\psi\phi$ NR	+7.66	+2.57	+2.52	+0.92
Extended model	-6.36	-2.54	-0.20	-2.98
Additional X state	-5.58	+0.27	+0.34	-1.48
Fit fraction constraint	+0.07	+1.94	-0.48	-0.45
Impact parameter χ^2 modeling	+1.00	-2.49	-0.22	+0.72
Finite simulation sample size	+2.90	+11.20	+1.33	+2.17
Fixed mass and width of K^* states	-0.58	-1.30	-0.43	-0.58
Samples to construct the \mathcal{P}_{bkg}	+1.06 -1.38	+6.38 -7.25	+0.54 -0.59	+0.52 -0.68
Background fraction uncertainty	+0.37 -0.45	+1.38 -1.55	+0.06 -0.07	+0.22 -0.38
<i>Final</i>	+8.46 -16.71	+17.09 -23.33	+3.00 -2.84	+6.04 -4.23

for the possible additional decay channel $D_s^{*+}D_s^-$. The $J/\psi\phi$ NR shape is changed from constant to be a linear function of $m_{J/\psi\phi}$. The K^* states are described by a simplified one-channel K-matrix model [2] instead of a sum of RBW functions. The mass-dependent widths in the RBW function for the K^* resonances are calculated using the decay with the largest branching fraction, $K^* \rightarrow K\pi$ or $K^* \rightarrow K^*(892)\pi$, instead of using the $K^* \rightarrow \phi K$ decay. The spin parity of the $T_{\psi s1}(4220)$ state is changed from the default 1^+ to 1^- to account for its ambiguity between 1^+ and 1^- [9]. The uncertainty originating from the neglected non- ϕ contribution is evaluated by narrowing the ϕ mass window from ± 15 to ± 8 MeV. An additional $J/\psi\phi$ NR contribution with the spin parity equal to 1^+ or 2^+ is included. An extended model with more K^* states is also studied, which includes all the K^* resonances that are within the allowed phase space, as predicted in Ref. [33]. Possible additional X states, with spins ranging from 0 to 2, are checked based on the extended model. The total fit fraction of the default model is 165.2%, which indicates that the interference between the decay amplitudes is large. An alternative fit constraining the total fit fraction to be smaller than 140%, corresponding to a reduction by 3 times its uncertainty, is performed. The final positive (negative) systematic uncertainty is taken as the maximal positive (negative) deviation from the model uncertainties above summed in quadrature with the uncertainty from other sources. These other sources of systematic uncertainty

include mismodeling of the impact parameter χ^2 of B candidates, the finite size of the simulated samples, the fixed masses and widths of the known K^* states, the choice of the data sample used to construct the background PDF model, and the uncertainty on the background fraction.

The significance of the $T_{\psi s1}^{\theta}(4000)^0$ state is evaluated with a likelihood ratio test. The test statistic is defined as $t \equiv -2 \ln[\mathcal{L}(H_0)/\mathcal{L}(H_1)]$, with H_1 and H_0 denoting the default model with and without the $T_{\psi s1}^{\theta}(4000)^0$ state. Here, the term \mathcal{L} represents the likelihood of the B^0 decay. Five thousand pseudoexperiments are generated with the H_0 model, where the parameters of the model are fixed to the values determined from data. The sample size of each pseudoexperiment follows a Poisson distribution with its mean being equal to the number of candidates in the data sample. A fit is performed to each pseudoexperiment with the H_0 and H_1 models to determine the test statistic t . The t value from these pseudoexperiments should follow a χ^2 distribution, where the number of degrees of freedom is a freely varying parameter. Using the resulting χ^2 distribution, the probability of $t > t_{\text{data}}$ is taken as the p value. The corresponding significance is estimated to be 4.9σ , decreasing to 4.0σ after including systematic uncertainties, which provides evidence for the $T_{\psi s1}^{\theta}(4000)^0$ state. The significance for the $T_{\psi s1}^{\theta}(4000)^0$ state is taken as the smallest significance found when varying the sources of systematic uncertainty.

The significance is also evaluated assuming isospin symmetry for the $T_{\psi s 1}^{\theta}(4000)$ states, where the test statistic is defined as $t' \equiv -2 \ln[\mathcal{L}(H_0)/\mathcal{L}(H'_1)]$, with \mathcal{L} denoting the total likelihood of the B^+ and B^0 decays. The corresponding fit parameters for the $T_{\psi s 1}^{\theta}(4000)^+$ and $T_{\psi s 1}^{\theta}(4000)^0$ states, including mass, width, and helicity couplings, are constrained to be equal between the two states in the H'_1 model. A fit is performed to the t' distribution from the pseudoexperiments with a Gaussian function. The Gaussian function rather than a χ^2 distribution is used here because the number of degrees of freedom of the H_0 and H'_1 models is equal. The significance is estimated to be 7.2σ and decreases to 5.4σ after accounting for systematic uncertainties.

In conclusion, an amplitude analysis of the $B^0 \rightarrow J/\psi \phi K_S^0$ decay is performed. Evidence of a $J/\psi K_S^0$ structure, denoted the $T_{\psi s 1}^{\theta}(4000)^0$ state, is obtained with a significance of 4.0σ . The mass and width of this state are measured to be

$$M(T_{\psi s 1}^{\theta}(4000)^0) = 3991_{-10}^{+12} {}_{-17}^{+9} \text{ MeV},$$

$$\Gamma(T_{\psi s 1}^{\theta}(4000)^0) = 105_{-25}^{+29} {}_{-23}^{+17} \text{ MeV},$$

where the first uncertainty is statistical and the second systematic. The mass difference between the $T_{\psi s 1}^{\theta}(4000)^0$ and $T_{\psi s 1}^{\theta}(4000)^+$ states is measured to be

$$\Delta M = -12_{-10}^{+11} {}_{-4}^{+6} \text{ MeV},$$

which is consistent with the two states being isospin partners. With isospin symmetry imposed for the $T_{\psi s 1}^{\theta}(4000)$ states, the significance of the $T_{\psi s 1}^{\theta}(4000)^0$ structure is measured to be 5.4σ .

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); NWO (Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MICINN (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); DOE NP and NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (Netherlands), PIC (Spain), GridPP (United Kingdom), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), Polish WLCG (Poland), and NERSC (USA). We are indebted to the communities behind the multiple open-source software packages on which we depend. Individual groups or members have received support from ARC and ARDC (Australia); Minciencias (Colombia); AvH

Foundation (Germany); EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union); A*MIDEX, ANR, IPhU and Labex P2IO, and Région Auvergne-Rhône-Alpes (France); Key Research Program of Frontier Sciences of CAS, CAS PIFI, CAS CCEPP, Fundamental Research Funds for the Central Universities, and Science and Technology Program of Guangzhou (China); GVA, XuntaGal, GENCAT, and Prog. Atracción Talento, CM (Spain); SRC (Sweden); the Leverhulme Trust, the Royal Society, and UKRI (United Kingdom).

- [1] S. K. Choi *et al.* (Belle Collaboration), Observation of a Narrow Charmoniumlike State in Exclusive $B^{\pm} \rightarrow K^{\pm} \pi^+ \pi^- J/\psi$ Decays, *Phys. Rev. Lett.* **91**, 262001 (2003).
- [2] R. L. Workman *et al.* (Particle Data Group), Review of particle physics, *Prog. Theor. Exp. Phys.* **2022**, 083C01 (2022).
- [3] R. Aaij *et al.* (LHCb Collaboration), Observation of structure in the J/ψ -pair mass spectrum, *Sci. Bull.* **65**, 1983 (2020).
- [4] ATLAS Collaboration, Observation of an excess of di-charmonium events in the four-muon final state with the ATLAS detector, Report No. ATLAS-CONF-2022-040, CERN.
- [5] CMS Collaboration, Observation of new structures in the $J/\psi J/\psi$ mass spectrum in pp collisions at $\sqrt{s} = 13$ TeV, Report No. CMS-PAS-BPH-21-003, CERN.
- [6] R. Aaij *et al.* (LHCb Collaboration), Observation of a Narrow Pentaquark State, $P_c(4312)^+$, and of Two-Peak Structure of the $P_c(4450)^+$, *Phys. Rev. Lett.* **122**, 222001 (2019).
- [7] R. Aaij *et al.* (LHCb Collaboration), Observation of a $J/\psi \Lambda$ Resonance Consistent with a Strange Pentaquark Candidate in $B^- \rightarrow J/\psi \Lambda \bar{p}$ Decays *Phys. Rev. Lett.* **131**, 031901 (2023).
- [8] M. Ablikim *et al.* (BESIII Collaboration), Observation of a Near-Threshold Structure in the K^+ Recoil-Mass Spectra in $e^+ e^- \rightarrow K^+(D_s^- D^{*0} + D_s^{*-} D^0)$, *Phys. Rev. Lett.* **126**, 102001 (2021).
- [9] R. Aaij *et al.* (LHCb Collaboration), Observation of New Resonances Decaying to $J/\psi K^+$ and $J/\psi \phi$, *Phys. Rev. Lett.* **127**, 082001 (2021).
- [10] The new exotic hadron naming convention proposed by the LHCb Collaboration [11] is applied. The $T_{\psi s 1}^{\theta}(4000)^+$ and $T_{\psi s 1}(4220)^+$ states are named $Z_{cs}(4000)^+$ and $Z_{cs}(4220)^+$ in the original papers [8,9], respectively. The superscript θ refers to 1/2 isospin and positive parity.
- [11] T. Gershon *et al.* (LHCb Collaboration), Exotic hadron naming convention, [arXiv:2206.15233](https://arxiv.org/abs/2206.15233).
- [12] L. Meng, B. Wang, and S.-L. Zhu, $Z_{cs}(3985)^-$ as the U -spin partner of $Z_c(3900)^-$ and implication of other states in the $SU(3)_F$ symmetry and heavy quark symmetry, *Phys. Rev. D* **102**, 111502(R) (2020).
- [13] L. Maiani, A. D. Polosa, and V. Riquer, The new resonances $Z_{cs}(3985)$ and $Z_{cs}(4003)$ (almost) fill two tetraquark nonets of broken $SU(3)_F$, *Sci. Bull.* **66**, 1616 (2021).

- [14] N. Ikeno, R. Molina, and E. Oset, Z_{cs} states from the $D_s^* \bar{D}^*$ and $J/\psi K^*$ coupled channels: Signal in $B^+ \rightarrow J/\psi \phi K^+$ decay, *Phys. Rev. D* **105**, 014012 (2022); **106**, 099905(E) (2022).
- [15] M. Ablikim *et al.* (BESIII Collaboration), Evidence for a Neutral Near-Threshold Structure in the K_S^0 Recoil-Mass Spectra in $e^+e^- \rightarrow K_S^0 D_s^+ D^{*-}$ and $e^+e^- \rightarrow K_S^0 D_s^{*+} D^-$, *Phys. Rev. Lett.* **129**, 112003 (2022).
- [16] C. P. Jessop *et al.* (CLEO Collaboration), First Observation of the Decay $B \rightarrow J/\psi \phi K$, *Phys. Rev. Lett.* **84**, 1393 (2000).
- [17] B. Aubert *et al.* (BABAR Collaboration), Rare B Decays into States Containing a J/ψ Meson and a Meson with $s\bar{s}$ Quark Content, *Phys. Rev. Lett.* **91**, 071801 (2003).
- [18] J. P. Lees *et al.* (BABAR Collaboration), Study of $B^{\pm,0} \rightarrow J/\psi K^+ K^- K^{\pm,0}$ and search for $B^0 \rightarrow J/\psi \phi$ at BABAR, *Phys. Rev. D* **91**, 012003 (2015).
- [19] A. A. Alves Jr. *et al.* (LHCb Collaboration), The LHCb detector at the LHC, *J. Instrum.* **3**, S08005 (2008).
- [20] R. Aaij *et al.* (LHCb Collaboration), LHCb detector performance, *Int. J. Mod. Phys. A* **30**, 1530022 (2015).
- [21] T. Sjöstrand, S. Mrenna, and P. Skands, A brief introduction to PYTHIA 8.1, *Comput. Phys. Commun.* **178**, 852 (2008); T. Sjöstrand, S. Mrenna, and P. Skands, PYTHIA 6.4 physics and manual, *J. High Energy Phys.* **05** (2006) 026.
- [22] D. J. Lange, The EvtGen particle decay simulation package, *Nucl. Instrum. Methods Phys. Res., Sect. A* **462**, 152 (2001).
- [23] J. Allison *et al.* (GEANT4 Collaboration), GEANT4 developments and applications, *IEEE Trans. Nucl. Sci.* **53**, 270 (2006); S. Agostinelli *et al.* (GEANT4 Collaboration), GEANT4: A simulation toolkit, *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [24] M. Clemencic, G. Corti, S. Easo, C. R. Jones, S. Miglioranzi, M. Pappagallo, and P. Robbe, The LHCb simulation application, Gauss: Design, evolution and experience, *J. Phys. Conf. Ser.* **331**, 032023 (2011).
- [25] W. D. Hulsbergen, Decay chain fitting with a Kalman filter, *Nucl. Instrum. Methods Phys. Res., Sect. A* **552**, 566 (2005).
- [26] W. S. McCulloch and W. Pitts, A logical calculus of the ideas immanent in nervous activity, *Bull. Math. Biol.* **5**, 115 (1943).
- [27] F. Rosenblatt, The perceptron: A probabilistic model for information storage and organization in the brain, *Psy. Rev.* **65**, 386 (1958).
- [28] R. Aaij *et al.* (LHCb Collaboration), Amplitude analysis of $B_s^0 \rightarrow K_S^0 K^\pm \pi^\mp$ decays, *J. High Energy Phys.* **06** (2019) 114.
- [29] D. Martínez Santos and F. Dupertuis, Mass distributions marginalized over per-event errors, *Nucl. Instrum. Methods Phys. Res., Sect. A* **764**, 150 (2014).
- [30] R. Aaij *et al.* (LHCb Collaboration), Amplitude analysis of $B^+ \rightarrow J/\psi \phi K^+$ decays, *Phys. Rev. D* **95**, 012002 (2017).
- [31] S. U. Chung, Spin formalisms, Report No. CERN-71-08, CERN, 1971; J. D. Richman, An experimenter's guide to the helicity formalism, Report No. CALT-68-1148, CERN; M. Jacob and G. C. Wick, On the general theory of collisions for particles with spin, *Ann. Phys. (N.Y.)* **7**, 404 (1959).
- [32] S. M. Flatté, Coupled-channel analysis of the $\pi\eta$ and $K\bar{K}$ systems near $K\bar{K}$ threshold, *Phys. Lett.* **63B**, 224 (1976).
- [33] S. Godfrey and N. Isgur, Mesons in a relativized quark model with chromodynamics, *Phys. Rev. D* **32**, 189 (1985).

R. Aaij^{1b},³² A. S. W. Abdelmotteleb^{1b},⁵⁰ C. Abellan Beteta⁴⁴ F. Abudinén^{1b},⁵⁰ T. Ackernley^{1b},⁵⁴ B. Adeva^{1b},⁴⁰ M. Adinolfi^{1b},⁴⁸ P. Adlarson^{1b},⁷⁷ H. Afsharnia⁹ C. Agapopoulou^{1b},¹³ C. A. Aidala^{1b},⁷⁸ Z. Ajaltouni⁹ S. Akar^{1b},⁵⁹ K. Akiba^{1b},³² P. Albicocco^{1b},²³ J. Albrecht^{1b},¹⁵ F. Alessio^{1b},⁴² M. Alexander^{1b},⁵³ A. Alfonso Alberro^{1b},³⁹ Z. Aliouche^{1b},⁵⁶ P. Alvarez Cartelle^{1b},⁴⁹ R. Amalric^{1b},¹³ S. Amato^{1b},² J. L. Amey^{1b},⁴⁸ Y. Amhis^{1b},^{11,42} L. An^{1b},⁴² L. Anderlini^{1b},²² M. Andersson^{1b},⁴⁴ A. Andreianov^{1b},³⁸ M. Andreotti^{1b},²¹ D. Andreou^{1b},⁶² D. Ao^{1b},⁶ F. Archilli^{1b},^{31,b} A. Artamonov^{1b},³⁸ M. Artuso^{1b},⁶² E. Aslanides^{1b},¹⁰ M. Atzeni^{1b},⁴⁴ B. Audurier^{1b},¹² I. Bachiller Perea^{1b},⁸ S. Bachmann^{1b},¹⁷ M. Bachmayer^{1b},⁴³ J. J. Back^{1b},⁵⁰ A. Bailly-reyre^{1b},¹³ P. Baladron Rodriguez^{1b},⁴⁰ V. Balagura^{1b},¹² W. Baldini^{1b},^{21,42} J. Baptista de Souza Leite^{1b},¹ M. Barbetti^{1b},^{22,c} R. J. Barlow^{1b},⁵⁶ S. Barsuk^{1b},¹¹ W. Barter^{1b},⁵² M. Bartolini^{1b},⁴⁹ F. Baryshnikov^{1b},³⁸ J. M. Basels^{1b},¹⁴ G. Bassi^{1b},^{29,d} B. Batsukh^{1b},⁴ A. Battig^{1b},¹⁵ A. Bay^{1b},⁴³ A. Beck^{1b},⁵⁰ M. Becker^{1b},¹⁵ F. Bedeschi^{1b},²⁹ I. B. Bediaga^{1b},¹ A. Beiter^{1b},⁶² S. Belin^{1b},⁴⁰ V. Bellee^{1b},⁴⁴ K. Belous^{1b},³⁸ I. Belov^{1b},³⁸ I. Belyaev^{1b},³⁸ G. Benane^{1b},¹⁰ G. Bencivenni^{1b},²³ E. Ben-Haim^{1b},¹³ A. Berezhnoy^{1b},³⁸ R. Bernet^{1b},⁴⁴ S. Bernet Andres^{1b},⁷⁶ D. Berninghoff^{1b},¹⁷ H. C. Bernstein^{1b},⁶² C. Bertella^{1b},⁵⁶ A. Bertolin^{1b},²⁸ C. Betancourt^{1b},⁴⁴ F. Betti^{1b},⁴² Ia. Bezshyiko^{1b},⁴⁴ S. Bhasin^{1b},⁴⁸ J. Bhom^{1b},³⁵ L. Bian^{1b},⁶⁸ M. S. Bieker^{1b},¹⁵ N. V. Biesuz^{1b},²¹ P. Billoir^{1b},¹³ A. Biolchini^{1b},³² M. Birch^{1b},⁵⁵ F. C. R. Bishop^{1b},⁴⁹ A. Bitadze^{1b},⁵⁶ A. Bizzeti^{1b}, M. P. Blago^{1b},⁴⁹ T. Blake^{1b},⁵⁰ F. Blanc^{1b},⁴³ J. E. Blank^{1b},¹⁵ S. Blusk^{1b},⁶² D. Bobulska^{1b},⁵³ J. A. Boelhave^{1b},¹⁵ O. Boente Garcia^{1b},¹² T. Boettcher^{1b},⁵⁹ A. Boldyrev^{1b},³⁸ C. S. Bolognani^{1b},⁷⁴ R. Bolzonella^{1b},^{21,e} N. Bondar^{1b},^{38,42} F. Borgato^{1b},²⁸ S. Borghi^{1b},⁵⁶ M. Borsato^{1b},¹⁷ J. T. Borsuk^{1b},³⁵ S. A. Bouchiba^{1b},⁴³ T. J. V. Bowcock^{1b},⁵⁴ A. Boyer^{1b},⁴² C. Bozzi^{1b},²¹ M. J. Bradley^{1b},⁵⁵ S. Braun^{1b},⁶⁰ A. Brea Rodriguez^{1b},⁴⁰ J. Brodzicka^{1b},³⁵ A. Brossa Gonzalo^{1b},⁴⁰ J. Brown^{1b},⁵⁴ D. Brundu^{1b},²⁷ A. Buonaura^{1b},⁴⁴ L. Buonincontri^{1b},²⁸ A. T. Burke^{1b},⁵⁶ C. Burr^{1b},⁴² A. Bursche^{1b},⁶⁶ A. Butkevich^{1b},³⁸ J. S. Butter^{1b},³² J. Buytaert^{1b},⁴² W. Byczynski^{1b},⁴² S. Cadeddu^{1b},²⁷ H. Cai^{1b},⁶⁸ R. Calabrese^{1b},^{21,e} L. Calefice^{1b},¹⁵ S. Cali^{1b},²³

M. Calvi^{26,f} M. Calvo Gomez⁷⁶ P. Campana²³ D. H. Campora Perez⁷⁴ A. F. Campoverde Quezada⁶
 S. Capelli^{26,f} L. Capriotti²⁰ A. Carbone^{20,g} R. Cardinale^{24,h} A. Cardini²⁷ P. Carniti^{26,f} L. Carus¹⁴
 A. Casais Vidal⁴⁰ R. Caspary¹⁷ G. Casse⁵⁴ M. Cattaneo⁴² G. Cavallero^{55,42} V. Cavallini^{21,e} S. Celani⁴³
 J. Cerasoli¹⁰ D. Cervenkova⁵⁷ A. J. Chadwick⁵⁴ I. Chahrour⁷⁸ M. G. Chapman⁴⁸ M. Charles¹³
 Ph. Charpentier⁴² C. A. Chavez Barajas⁵⁴ M. Chefdeville⁸ C. Chen¹⁰ S. Chen⁴ A. Chernov³⁵
 S. Chernyshenko⁴⁶ V. Chobanova⁴⁰ S. Cholak⁴³ M. Chrzaszcz³⁵ A. Chubykin³⁸ V. Chulikov³⁸
 P. Ciambrone²³ M. F. Cicala⁵⁰ X. Cid Vidal⁴⁰ G. Ciezarek⁴² P. Cifra⁴² G. Ciullo^{21,e} P. E. L. Clarke⁵²
 M. Clemencic⁴² H. V. Cliff⁴⁹ J. Closier⁴² J. L. Cobbedick⁵⁶ V. Coco⁴² J. Cogan¹⁰ E. Cogneras⁹
 L. Cojocariu³⁷ P. Collins⁴² T. Colombo⁴² L. Congedo¹⁹ A. Contu²⁷ N. Cooke⁴⁷ I. Corredoira⁴⁰ G. Corti⁴²
 B. Couturier⁴² D. C. Craik⁴⁴ M. Cruz Torres^{1,i} R. Currie⁵² C. L. Da Silva⁶¹ S. Dadabaev³⁸ L. Dai⁶⁵
 X. Dai⁵ E. Dall'Occo¹⁵ J. Dalseno⁴⁰ C. D'Ambrosio⁴² J. Daniel⁹ A. Danilina³⁸ P. d'Argent¹⁹
 J. E. Davies⁵⁶ A. Davis⁵⁶ O. De Aguiar Francisco⁵⁶ J. de Boer⁴² K. De Bruyn⁷³ S. De Capua⁵⁶
 M. De Cian⁴³ U. De Freitas Carneiro Da Graca¹ E. De Lucia²³ J. M. De Miranda¹ L. De Paula² M. De Serio^{19,j}
 D. De Simone⁴⁴ P. De Simone²³ F. De Vellis¹⁵ J. A. de Vries⁷⁴ C. T. Dean⁶¹ F. Debernardis^{19,j} D. Decamp⁸
 V. Dedu¹⁰ L. Del Buono¹³ B. Delaney⁵⁸ H.-P. Dembinski¹⁵ V. Denysenko⁴⁴ O. Deschamps⁹ F. Dettori^{27,k}
 B. Dey⁷¹ P. Di Nezza²³ I. Diachkov³⁸ S. Didenko³⁸ L. Dieste Maronas⁴⁰ S. Ding⁶² V. Dobishuk⁴⁶
 A. Dolmatov³⁸ C. Dong³ A. M. Donohoe¹⁸ F. Dordei²⁷ A. C. dos Reis¹ L. Douglas⁵³ A. G. Downes⁸
 P. Duda⁷⁵ M. W. Dudek³⁵ L. Dufour⁴² V. Duk⁷² P. Durante⁴² M. M. Duras⁷⁵ J. M. Durham⁶¹ D. Dutta⁵⁶
 A. Dziurda³⁵ A. Dzyuba³⁸ S. Easo⁵¹ U. Egede⁶³ V. Egorychev³⁸ C. Eirea Orro⁴⁰ S. Eisenhardt⁵² E. Ejopu⁵⁶
 S. Ek-In⁴³ L. Eklund⁷⁷ M. Elashri⁵⁹ J. Ellbracht¹⁵ S. Ely⁵⁵ A. Ene³⁷ E. Eppe⁵⁹ S. Escher¹⁴ J. Eschle⁴⁴
 S. Esen⁴⁴ T. Evans⁵⁶ F. Fabiano^{27,k} L. N. Falcao¹ Y. Fan⁶ B. Fang^{11,68} L. Fantini^{72,1} M. Faria⁴³
 S. Farry⁵⁴ D. Fazzini^{26,f} L. Felkowski⁷⁵ M. Feo⁴² M. Fernandez Gomez⁴⁰ A. D. Femez⁶⁰ F. Ferrari²⁰
 L. Ferreira Lopes⁴³ F. Ferreira Rodrigues² S. Ferreres Sole³² M. Ferrillo⁴⁴ M. Ferro-Luzzi⁴² S. Filippov³⁸
 R. A. Fini¹⁹ M. Fiorini^{21,e} M. Firlej³⁴ K. M. Fischer⁵⁷ D. S. Fitzgerald⁷⁸ C. Fitzpatrick⁵⁶ T. Fiutowski³⁴
 F. Fleuret¹² M. Fontana¹³ F. Fontanelli^{24,h} R. Forty⁴² D. Foulds-Holt⁴⁹ V. Franco Lima⁵⁴
 M. Franco Sevilla⁶⁰ M. Frank⁴² E. Franzoso^{21,e} G. Frau¹⁷ C. Frei⁴² D. A. Friday⁵³ L. Frontini²⁵ J. Fu⁶
 Q. Fuehring¹⁵ T. Fulghesu¹³ E. Gabriel³² G. Galati^{19,j} M. D. Galati³² A. Gallas Torreira⁴⁰ D. Galli^{20,g}
 S. Gambetta^{52,42} M. Gandelman² P. Gandini²⁵ Y. Gao⁷ Y. Gao⁵ M. Garau^{27,k} L. M. Garcia Martin⁵⁰
 P. Garcia Moreno³⁹ J. Garcia Pardiñas^{26,f} B. Garcia Plana⁴⁰ F. A. Garcia Rosales¹² L. Garrido³⁹ C. Gaspar⁴²
 R. E. Geertsema³² D. Gerick¹⁷ L. L. Gerken¹⁵ E. Gersabeck⁵⁶ M. Gersabeck⁵⁶ T. Gershon⁵⁰
 L. Giambastiani²⁸ V. Gibson⁴⁹ H. K. Gienza³⁶ A. L. Gilman⁵⁷ M. Giovannetti^{23,b} A. Gioventù⁴⁰
 P. Gironella Gironell³⁹ C. Giugliano^{21,e} M. A. Giza³⁵ K. Gizdov⁵² E. L. Gkougkousis⁴² V. V. Gligorov^{13,42}
 C. Göbel⁶⁴ E. Golobardes⁷⁶ D. Golubkov³⁸ A. Golutvin^{55,38} A. Gomes^{1,m} S. Gomez Fernandez³⁹
 F. Goncalves Abrantes⁵⁷ M. Goncerz³⁵ G. Gong³ I. V. Gorelov³⁸ C. Gotti²⁶ J. P. Grabowski⁷⁰
 T. Grammatico¹³ L. A. Granado Cardoso⁴² E. Graugés³⁹ E. Graverini⁴³ G. Graziani⁴³ A. T. Grecu³⁷
 L. M. Greeven³² N. A. Grieser⁵⁹ L. Grillo⁵³ S. Gromov³⁸ B. R. Gruberg Cazon⁵⁷ C. Gu³ M. Guarise^{21,e}
 M. Guittiere¹¹ P. A. Günther¹⁷ E. Gushchin³⁸ A. Guth¹⁴ Y. Guz³⁸ T. Gys⁴² T. Hadavizadeh⁶³
 C. Hadjivasiliou⁶⁰ G. Haefeli⁴³ C. Haen⁴² J. Haimberger⁴² S. C. Haines⁴⁹ T. Halewood-leagas⁵⁴
 M. M. Halvorsen⁴² P. M. Hamilton⁶⁰ J. Hammerich⁵⁴ Q. Han⁷ X. Han¹⁷ E. B. Hansen⁵⁶
 S. Hansmann-Menzemer¹⁷ L. Hao⁶ N. Harnew⁵⁷ T. Harrison⁵⁴ C. Hasse⁴² M. Hatch⁴² J. He^{6,n}
 K. Heijhoff³² F. H. Hemmer⁴² C. Henderson⁵⁹ R. D. L. Henderson^{63,50} A. M. Hennequin⁵⁸ K. Hennessy⁵⁴
 L. Henry⁴² J. Herd⁵⁵ J. Heuel¹⁴ A. Hicheur² D. Hill⁴³ M. Hilton⁵⁶ S. E. Hollitt¹⁵ J. Horswill⁵⁶ R. Hou⁷
 Y. Hou⁸ J. Hu¹⁷ J. Hu⁶⁶ W. Hu⁵ X. Hu³ W. Huang⁶ X. Huang⁶⁸ W. Hulsbergen³² R. J. Hunter⁵⁰
 M. Hushchyn³⁸ D. Hutchcroft⁵⁴ P. Ibis¹⁵ M. Idzik³⁴ D. Ilin³⁸ P. Ilten⁵⁹ A. Inglessi³⁸ A. Iniukhin³⁸
 A. Ishteev³⁸ K. Ivshin³⁸ R. Jacobsson⁴² H. Jage¹⁴ S. J. Jaimes Elles⁴¹ S. Jakobsen⁴² E. Jans³²
 B. K. Jashal⁴¹ A. Jawahery⁶⁰ V. Jevtic¹⁵ E. Jiang⁶⁰ X. Jiang^{4,6} Y. Jiang⁶ M. John⁵⁷ D. Johnson⁵⁸
 C. R. Jones⁴⁹ T. P. Jones⁵⁰ B. Jost⁴² N. Jurik⁴² I. Juszczak³⁵ S. Kandybei⁴⁵ Y. Kang³ M. Karacson⁴²
 D. Karpenkov³⁸ M. Karpov³⁸ J. W. Kautz⁵⁹ F. Keizer⁴² D. M. Keller⁶² M. Kenzie⁵⁰ T. Ketel³² B. Khanji¹⁵
 A. Kharisova³⁸ S. Kholodenko³⁸ G. Khreich¹¹ T. Kim¹⁴ V. S. Kirsebom⁴³ O. Kitouni⁵⁸ S. Klaver³³

N. Kleijne^{29,d} K. Klimaszewski³⁶ M. R. Kmieć³⁶ S. Koliiev⁴⁶ L. Kolk¹⁵ A. Kondybayeva³⁸
A. Konoplyannikov³⁸ P. Kopciwicz³⁴ R. Kopečna¹⁷ P. Koppenburg³² M. Korolev³⁸ I. Kostiuik³² O. Kot,⁴⁶
S. Kotriakhova³⁸ A. Kozachuk³⁸ P. Kravchenko³⁸ L. Kravchuk³⁸ R. D. Krawczyk⁴² M. Kreps⁵⁰
S. Kretschmar¹⁴ P. Krokovny³⁸ W. Krupa³⁴ W. Krzemien³⁶ J. Kubat,¹⁷ S. Kubis⁷⁵ W. Kucewicz³⁵
M. Kucharczyk³⁵ V. Kudryavtsev³⁸ E. Kulikova³⁸ A. Kupsc⁷⁷ D. Lacarrere⁴² G. Lafferty⁵⁶ A. Lai²⁷
A. Lampis^{27,k} D. Lancierini⁴⁴ C. Landesa Gomez⁴⁰ J. J. Lane⁵⁶ R. Lane⁴⁸ C. Langenbruch¹⁴ J. Langer¹⁵
O. Lantwin³⁸ T. Latham⁵⁰ F. Lazzari^{29,o} M. Lazzaroni^{25,p} R. Le Gac¹⁰ S. H. Lee⁷⁸ R. Lefèvre⁹ A. Leflat³⁸
S. Legotin³⁸ P. Lenisa^{21,e} O. Leroy¹⁰ T. Lesiak³⁵ B. Leverington¹⁷ A. Li³ H. Li⁶⁶ K. Li⁷ P. Li⁴²
P.-R. Li⁶⁷ S. Li⁷ T. Li⁴ T. Li⁶⁶ Y. Li⁴ Z. Li⁶² X. Liang⁶² C. Lin⁶ T. Lin⁵¹ R. Lindner⁴²
V. Lisovskyi¹⁵ R. Litvinov^{27,k} G. Liu⁶⁶ H. Liu⁶ Q. Liu⁶ S. Liu^{4,6} A. Lobo Salvia³⁹ A. Loi²⁷ R. Lollini⁷²
J. Lomba Castro⁴⁰ I. Longstaff,⁵³ J. H. Lopes² A. Lopez Huertas³⁹ S. López Soliño⁴⁰ G. H. Lovell⁴⁹ Y. Lu^{4,q}
C. Lucarelli^{22,c} D. Lucchesi^{28,r} S. Luchuk³⁸ M. Lucio Martinez⁷⁴ V. Lukashenko^{32,46} Y. Luo³ A. Lupato⁵⁶
E. Luppi^{21,e} A. Lusiani^{29,d} K. Lynch¹⁸ X.-R. Lyu⁶ R. Ma⁶ S. Maccolini¹⁵ F. Machefert¹¹ F. Maciuc³⁷
I. Mackay⁵⁷ V. Macko⁴³ L. R. Madhan Mohan⁴⁸ A. Maevskiy³⁸ D. Maisuzenko³⁸ M. W. Majewski,³⁴
J. J. Malczewski³⁵ S. Malde⁵⁷ B. Malecki^{35,42} A. Malinin³⁸ T. Maltsev³⁸ G. Manca^{27,k} G. Mancinelli¹⁰
C. Mancuso^{11,25,p} R. Manera Escalero,³⁹ D. Manuzzi²⁰ C. A. Manzari⁴⁴ D. Marangotto^{25,p} J. F. Marchand⁸
U. Marconi²⁰ S. Mariani^{22,c} C. Marin Benito³⁹ J. Marks¹⁷ A. M. Marshall⁴⁸ P. J. Marshall,⁵⁴ G. Martelli^{72,1}
G. Martellotti³⁰ L. Martinazzoli^{42,f} M. Martinelli^{26,f} D. Martinez Santos⁴⁰ F. Martinez Vidal⁴¹ A. Massafferri¹
M. Materok¹⁴ R. Matev⁴² A. Mathad⁴⁴ V. Matiunin³⁸ C. Matteuzzi²⁶ K. R. Mattioli¹² A. Mauri³²
E. Maurice¹² J. Mauricio³⁹ M. Mazurek⁴² M. McCann⁵⁵ L. Mcconnell¹⁸ T. H. McGrath⁵⁶ N. T. McHugh⁵³
A. McNab⁵⁶ R. McNulty¹⁸ J. V. Mead⁵⁴ B. Meadows⁵⁹ G. Meier¹⁵ D. Melnychuk³⁶ S. Meloni^{26,f}
M. Merk^{32,74} A. Merli^{25,p} L. Meyer Garcia² D. Miao^{4,6} M. Mikhasenko^{70,s} D. A. Milanese⁶⁹ E. Millard,⁵⁰
M. Milovanovic⁴² M.-N. Minard,^{8,a} A. Minotti^{26,f} T. Miralles⁹ S. E. Mitchell⁵² B. Mitreska¹⁵ D. S. Mitzel¹⁵
A. Mödden¹⁵ R. A. Mohammed⁵⁷ R. D. Moise¹⁴ S. Mokhnenko³⁸ T. Mombächer⁴⁰ M. Monk^{50,63}
I. A. Monroy⁶⁹ S. Monteil⁹ G. Morello²³ M. J. Morello^{29,d} M. P. Morgenthaler¹⁷ J. Moron³⁴ A. B. Morris⁴²
A. G. Morris⁵⁰ R. Mountain⁶² H. Mu³ E. Muhammad⁵⁰ F. Muheim⁵² M. Mulder⁷³ K. Müller⁴⁴
C. H. Murphy⁵⁷ D. Murray⁵⁶ R. Murta⁵⁵ P. Muzzetto^{27,k} P. Naik⁴⁸ T. Nakada⁴³ R. Nandakumar⁵¹
T. Nanut⁴² I. Nasteva² M. Needham⁵² N. Neri^{25,p} S. Neubert⁷⁰ N. Neufeld⁴² P. Neustroev,³⁸ R. Newcombe,⁵⁵
J. Nicolini^{15,11} D. Nicotra⁷⁴ E. M. Niel⁴³ S. Nieswand,¹⁴ N. Nikitin³⁸ N. S. Nolte⁵⁸ C. Normand^{8,27,k}
J. Novoa Fernandez⁴⁰ G. Nowak⁵⁹ C. Nunez⁷⁸ A. Oblakowska-Mucha³⁴ V. Obraztsov³⁸ T. Oeser¹⁴
S. Okamura^{21,e} R. Oldeman^{27,k} F. Oliva⁵² C. J. G. Onderwater⁷³ R. H. O’Neil⁵² J. M. Otalora Goicochea²
T. Ovsianikova³⁸ P. Owen⁴⁴ A. Oyanguren⁴¹ O. Ozcelik⁵² K. O. Padeken⁷⁰ B. Pagare⁵⁰ P. R. Pais⁴²
T. Pajero⁵⁷ A. Palano¹⁹ M. Palutan²³ Y. Pan⁵⁶ G. Panshin³⁸ L. Paolucci⁵⁰ A. Papanestis⁵¹
M. Pappagallo^{19,j} L. L. Pappalardo^{21,e} C. Pappenheimer⁵⁹ W. Parker⁶⁰ C. Parkes⁵⁶ B. Passalacqua^{21,e}
G. Passaleva²² A. Pastore¹⁹ M. Patel⁵⁵ C. Patrignani^{20,g} C. J. Pawley⁷⁴ A. Pellegrino³² M. Pepe Altarelli⁴²
S. Perazzini²⁰ D. Pereima³⁸ A. Pereiro Castro⁴⁰ P. Perret⁹ K. Petridis⁴⁸ A. Petrolini^{24,h} A. Petrov,³⁸
S. Petrucci⁵² M. Petruzzo²⁵ H. Pham⁶² A. Philippov³⁸ R. Piandani⁶ L. Pica^{29,d} M. Piccini⁷² B. Pietrzyk⁸
G. Pietrzyk¹¹ M. Pili⁵⁷ D. Pinci³⁰ F. Pisani⁴² M. Pizzichemi^{26,42,f} V. Placinta³⁷ J. Plews⁴⁷
M. Plo Casaus⁴⁰ F. Polci^{13,42} M. Poli Lener²³ A. Poluektov¹⁰ N. Polukhina³⁸ I. Polyakov⁴² E. Polycarpo²
S. Ponce⁴² D. Popov^{6,42} S. Poslavskii³⁸ K. Prasanth³⁵ L. Promberger¹⁷ C. Prouve⁴⁰ V. Pugatch⁴⁶
V. Puill¹¹ G. Punzi^{29,o} H. R. Qi³ W. Qian⁶ N. Qin³ S. Qu³ R. Quagliani⁴³ N. V. Raab¹⁸ B. Rachwal³⁴
J. H. Rademacker⁴⁸ R. Rajagopalan,⁶² M. Rama²⁹ M. Ramos Pernas⁵⁰ M. S. Rangel² F. Ratnikov³⁸
G. Raven^{33,42} M. Rebollo De Miguel⁴¹ F. Redi⁴² J. Reich⁴⁸ F. Reiss⁵⁶ C. Remon Alepuz,⁴¹ Z. Ren³
P. K. Resmi⁵⁷ R. Ribatti^{29,d} A. M. Ricci²⁷ S. Ricciardi⁵¹ K. Richardson⁵⁸ M. Richardson-Slipper⁵²
K. Rinnert⁵⁴ P. Robbe¹¹ G. Robertson⁵² A. B. Rodrigues⁴³ E. Rodrigues⁵⁴ E. Rodriguez Fernandez⁴⁰
J. A. Rodriguez Lopez⁶⁹ E. Rodriguez Rodriguez⁴⁰ D. L. Rolf⁴² A. Rollings⁵⁷ P. Roloff⁴² V. Romanovskiy³⁸
M. Romero Lamas⁴⁰ A. Romero Vidal⁴⁰ J. D. Roth,^{78,a} M. Rotondo²³ M. S. Rudolph⁶² T. Ruf⁴²
R. A. Ruiz Fernandez⁴⁰ J. Ruiz Vidal,⁴¹ A. Ryzhikov³⁸ J. Ryzka³⁴ J. J. Saborido Silva⁴⁰ N. Sagidova³⁸
N. Sahoo⁴⁷ B. Saitta^{27,k} M. Salomoni⁴² C. Sanchez Gras³² I. Sanderswood⁴¹ R. Santacesaria³⁰

C. Santamarina Rios⁴⁰, M. Santimaria²³, E. Santovetti^{31,b}, D. Saranin³⁸, G. Sarpis¹⁴, M. Sarpis⁷⁰, A. Sarti³⁰,
 C. Satriano^{30,t}, A. Satta³¹, M. Saur¹⁵, D. Savrina³⁸, H. Sazak⁹, L. G. Scantlebury Smead⁵⁷, A. Scarabotto¹³,
 S. Schael¹⁴, S. Scherl⁵⁴, M. Schiller⁵³, H. Schindler⁴², M. Schmelling¹⁶, B. Schmidt⁴², S. Schmitt¹⁴,
 O. Schneider⁴³, A. Schopper⁴², M. Schubiger³², S. Schulte⁴³, M. H. Schune¹¹, R. Schwemmer⁴², B. Sciascia²³,
 A. Sciuccati⁴², S. Sellam⁴⁰, A. Semennikov³⁸, M. Senghi Soares³³, A. Sergi^{24,h}, N. Serra⁴⁴, L. Sestini²⁸,
 A. Seuthe¹⁵, Y. Shang⁵, D. M. Shangase⁷⁸, M. Shapkin³⁸, I. Shchemerov³⁸, L. Shchutska⁴³, T. Shears⁵⁴,
 L. Shekhtman³⁸, Z. Shen⁵, S. Sheng^{4,6}, V. Shevchenko³⁸, B. Shi⁶, E. B. Shields^{26,f}, Y. Shimizu¹¹,
 E. Shmanin³⁸, R. Shorkin³⁸, J. D. Shupperd⁶², B. G. Siddi^{21,e}, R. Silva Coutinho⁶², G. Simi²⁸, S. Simone^{19,j},
 M. Singla⁶³, N. Skidmore⁵⁶, R. Skuza¹⁷, T. Skwarnicki⁶², M. W. Slater⁴⁷, J. C. Smallwood⁵⁷, J. G. Smeaton⁴⁹,
 E. Smith⁴⁴, K. Smith⁶¹, M. Smith⁵⁵, A. Snoch³², L. Soares Lavra⁹, M. D. Sokoloff⁵⁹, F. J. P. Soler⁵³,
 A. Solomin^{38,48}, A. Solovev³⁸, I. Solovyev³⁸, R. Song⁶³, F. L. Souza De Almeida², B. Souza De Paula²,
 B. Spaan^{15,a}, E. Spadaro Norella^{25,p}, E. Spedicato²⁰, E. Spiridenkov³⁸, P. Spradlin⁵³, V. Sriskaran⁴², F. Stagni⁴²,
 M. Stahl⁴², S. Stahl⁴², S. Stanislaus⁵⁷, E. N. Stein⁴², O. Steinkamp⁴⁴, O. Stenyakin³⁸, H. Stevens¹⁵, S. Stone^{62,a},
 D. Strelakina³⁸, Y. Su⁶, F. Suljik⁵⁷, J. Sun²⁷, L. Sun⁶⁸, Y. Sun⁶⁰, P. Svihra⁵⁶, P. N. Swallow⁴⁷, K. Swientek³⁴,
 A. Szabelski³⁶, T. Szumlak³⁴, M. Szymanski⁴², Y. Tan³, S. Taneja⁵⁶, M. D. Tat⁵⁷, A. Terentev⁴⁴, F. Teubert⁴²,
 E. Thomas⁴², D. J. D. Thompson⁴⁷, K. A. Thomson⁵⁴, H. Tilquin⁵⁵, V. Tisserand⁹, S. T'Jampens⁸, M. Tobin⁴,
 L. Tomassetti^{21,e}, G. Tonani^{25,p}, X. Tong⁵, D. Torres Machado¹, D. Y. Tou³, S. M. Trilov⁴⁸, C. Trippl⁴³,
 G. Tuci⁶, N. Tuning³², A. Ukleja³⁶, D. J. Unverzagt¹⁷, A. Usachov³³, A. Ustyuzhanin³⁸, U. Uwer¹⁷,
 A. Vagner³⁸, V. Vagnoni²⁰, A. Valassi⁴², G. Valenti²⁰, N. Valls Canudas⁷⁶, M. Van Dijk⁴³, H. Van Hecke⁶¹,
 E. van Herwijnen⁵⁵, C. B. Van Hulse^{40,u}, M. van Veghel³², R. Vazquez Gomez³⁹, P. Vazquez Regueiro⁴⁰,
 C. Vázquez Sierra⁴², S. Vecchi²¹, J. J. Velthuis⁴⁸, M. Veltri^{22,v}, A. Venkateswaran⁴³, M. Veronesi³²,
 M. Vesterinen⁵⁰, D. Vieira⁵⁹, M. Vieites Diaz⁴³, X. Vilasis-Cardona⁷⁶, E. Vilella Figueras⁵⁴, A. Villa²⁰,
 P. Vincent¹³, F. C. Volle¹¹, D. vom Bruch¹⁰, A. Vorobyev³⁸, V. Vorobyev³⁸, N. Voropaev³⁸, K. Vos⁷⁴, C. Vrahas⁵²,
 J. Walsh²⁹, E. J. Walton⁶³, G. Wan⁵, C. Wang¹⁷, G. Wang⁷, J. Wang⁵, J. Wang⁴, J. Wang³, J. Wang⁶⁸,
 M. Wang²⁵, R. Wang⁴⁸, X. Wang⁶⁶, Y. Wang⁷, Z. Wang⁴⁴, Z. Wang³, Z. Wang⁶, J. A. Ward^{50,63},
 N. K. Watson⁴⁷, D. Websdale⁵⁵, Y. Wei⁵, B. D. C. Westhenry⁴⁸, D. J. White⁵⁶, M. Whitehead⁵³,
 A. R. Wiederhold⁵⁰, D. Wiedner¹⁵, G. Wilkinson⁵⁷, M. K. Wilkinson⁵⁹, I. Williams⁴⁹, M. Williams⁵⁸,
 M. R. J. Williams⁵², R. Williams⁴⁹, F. F. Wilson⁵¹, W. Wislicki³⁶, M. Witek³⁵, L. Witola¹⁷, C. P. Wong⁶¹,
 G. Wormser¹¹, S. A. Wotton⁴⁹, H. Wu⁶², J. Wu⁷, K. Wyllie⁴², Z. Xiang⁶, Y. Xie⁷, A. Xu⁵, J. Xu⁶, L. Xu³,
 L. Xu³, M. Xu⁵⁰, Q. Xu⁶, Z. Xu⁹, Z. Xu⁶, D. Yang³, S. Yang⁶, X. Yang⁵, Y. Yang⁶, Z. Yang⁵, Z. Yang⁶⁰,
 L. E. Yeomans⁵⁴, V. Yeroshenko¹¹, H. Yeung⁵⁶, H. Yin⁷, J. Yu⁶⁵, X. Yuan⁶², E. Zaffaroni⁴³, M. Zavertyaev¹⁶,
 M. Zdybal³⁵, M. Zeng³, C. Zhang⁵, D. Zhang⁷, L. Zhang³, S. Zhang⁶⁵, S. Zhang⁵, Y. Zhang⁵, Y. Zhang⁵⁷,
 Y. Zhao¹⁷, A. Zharkova³⁸, A. Zhelezov¹⁷, Y. Zheng⁶, T. Zhou⁵, X. Zhou⁶, Y. Zhou⁶, V. Zhovkovska¹¹,
 X. Zhu³, X. Zhu⁷, Z. Zhu⁶, V. Zhukov^{14,38}, Q. Zou^{4,6}, S. Zucchelli^{20,g}, D. Zuliani²⁸ and G. Zunica⁵⁶

(LHCb Collaboration)

¹Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

²Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

³Center for High Energy Physics, Tsinghua University, Beijing, China

⁴Institute of High Energy Physics (IHEP), Beijing, China

⁵School of Physics State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

⁶University of Chinese Academy of Sciences, Beijing, China

⁷Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China

⁸Université Savoie Mont Blanc, CNRS, IN2P3-LAPP, Annecy, France

⁹Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France

¹⁰Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France

¹¹Université Paris-Saclay, CNRS/IN2P3, IJCLab, Orsay, France

¹²Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

¹³LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France

¹⁴I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany

- ¹⁵*Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany*
- ¹⁶*Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany*
- ¹⁷*Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany*
- ¹⁸*School of Physics, University College Dublin, Dublin, Ireland*
- ¹⁹*INFN Sezione di Bari, Bari, Italy*
- ²⁰*INFN Sezione di Bologna, Bologna, Italy*
- ²¹*INFN Sezione di Ferrara, Ferrara, Italy*
- ²²*INFN Sezione di Firenze, Firenze, Italy*
- ²³*INFN Laboratori Nazionali di Frascati, Frascati, Italy*
- ²⁴*INFN Sezione di Genova, Genova, Italy*
- ²⁵*INFN Sezione di Milano, Milano, Italy*
- ²⁶*INFN Sezione di Milano-Bicocca, Milano, Italy*
- ²⁷*INFN Sezione di Cagliari, Monserrato, Italy*
- ²⁸*Università degli Studi di Padova, Università e INFN, Padova, Padova, Italy*
- ²⁹*INFN Sezione di Pisa, Pisa, Italy*
- ³⁰*INFN Sezione di Roma La Sapienza, Roma, Italy*
- ³¹*INFN Sezione di Roma Tor Vergata, Roma, Italy*
- ³²*Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands*
- ³³*Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, Netherlands*
- ³⁴*AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland*
- ³⁵*Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*
- ³⁶*National Center for Nuclear Research (NCBJ), Warsaw, Poland*
- ³⁷*Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*
- ³⁸*Affiliated with an institute covered by a cooperation agreement with CERN*
- ³⁹*ICCUB, Universitat de Barcelona, Barcelona, Spain*
- ⁴⁰*Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain*
- ⁴¹*Instituto de Física Corpuscular, Centro Mixto Universidad de Valencia - CSIC, Valencia, Spain*
- ⁴²*European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- ⁴³*Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- ⁴⁴*Physik-Institut, Universität Zürich, Zürich, Switzerland*
- ⁴⁵*NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- ⁴⁶*Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- ⁴⁷*University of Birmingham, Birmingham, United Kingdom*
- ⁴⁸*H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- ⁴⁹*Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- ⁵⁰*Department of Physics, University of Warwick, Coventry, United Kingdom*
- ⁵¹*STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- ⁵²*School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- ⁵³*School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- ⁵⁴*Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- ⁵⁵*Imperial College London, London, United Kingdom*
- ⁵⁶*Department of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- ⁵⁷*Department of Physics, University of Oxford, Oxford, United Kingdom*
- ⁵⁸*Massachusetts Institute of Technology, Cambridge, 02139 Massachusetts, USA*
- ⁵⁹*University of Cincinnati, Cincinnati, 45221 Ohio, USA*
- ⁶⁰*University of Maryland, College Park, 20742 Maryland, USA*
- ⁶¹*Los Alamos National Laboratory (LANL), Los Alamos, 87545 New Mexico, USA*
- ⁶²*Syracuse University, Syracuse, 13244 New York, USA*
- ⁶³*School of Physics and Astronomy, Monash University, Melbourne, Australia*
(associated with Institution Department of Physics, University of Warwick, Coventry, United Kingdom)
- ⁶⁴*Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil*
(associated with Institution Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil)
- ⁶⁵*Physics and Micro Electronic College, Hunan University, Changsha City, China*
(associated with Institution Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China)
- ⁶⁶*Guangdong Provincial Key Laboratory of Nuclear Science, Guangdong-Hong Kong Joint Laboratory of Quantum Matter, Institute of Quantum Matter, South China Normal University, Guangzhou, China*
(associated with Institution Center for High Energy Physics, Tsinghua University, Beijing, China)
- ⁶⁷*Lanzhou University, Lanzhou, China*
(associated with Institution Institute Of High Energy Physics (IHEP), Beijing, China)

- ⁶⁸*School of Physics and Technology, Wuhan University, Wuhan, China*
(associated with Institution Center for High Energy Physics, Tsinghua University, Beijing, China)
- ⁶⁹*Departamento de Física, Universidad Nacional de Colombia, Bogota, Colombia*
(associated with Institution LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France)
- ⁷⁰*Universität Bonn - Helmholtz-Institut für Strahlen und Kernphysik, Bonn, Germany*
(associated with Institution Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany)
- ⁷¹*Eotvos Lorand University, Budapest, Hungary*
(associated with Institution European Organization for Nuclear Research (CERN), Geneva, Switzerland)
- ⁷²*INFN Sezione di Perugia, Perugia, Italy*
(associated with Institution INFN Sezione di Ferrara, Ferrara, Italy)
- ⁷³*Van Swinderen Institute, University of Groningen, Groningen, Netherlands*
(associated with Institution Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands)
- ⁷⁴*Universiteit Maastricht, Maastricht, Netherlands*
(associated with Institution Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands)
- ⁷⁵*Tadeusz Kosciuszko Cracow University of Technology, Cracow, Poland*
(associated with Institution Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland)
- ⁷⁶*DS4DS, La Salle, Universitat Ramon Llull, Barcelona, Spain*
(associated with Institution ICCUB, Universitat de Barcelona, Barcelona, Spain)
- ⁷⁷*Department of Physics and Astronomy, Uppsala University, Uppsala, Sweden*
(associated with Institution School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom)
- ⁷⁸*University of Michigan, Ann Arbor, 48109 Michigan, USA*
(associated with Institution Syracuse University, Syracuse, New York, USA)

^aDeceased.

^bAlso at Università di Roma Tor Vergata, Roma, Italy.

^cAlso at Università di Firenze, Firenze, Italy.

^dAlso at Scuola Normale Superiore, Pisa, Italy.

^eAlso at Università di Ferrara, Ferrara, Italy.

^fAlso at Università di Milano Bicocca, Milano, Italy.

^gAlso at Università di Bologna, Bologna, Italy.

^hAlso at Università di Genova, Genova, Italy.

ⁱAlso at Universidad Nacional Autónoma de Honduras, Tegucigalpa, Honduras.

^jAlso at Università di Bari, Bari, Italy.

^kAlso at Università di Cagliari, Cagliari, Italy.

^lAlso at Università di Perugia, Perugia, Italy.

^mAlso at Universidade de Brasília, Brasília, Brazil.

ⁿAlso at Hangzhou Institute for Advanced Study, UCAS, Hangzhou, China.

^oAlso at Università di Pisa, Pisa, Italy.

^pAlso at Università degli Studi di Milano, Milano, Italy.

^qAlso at Central South University, Changsha, China.

^rAlso at Università di Padova, Padova, Italy.

^sAlso at Excellence Cluster ORIGINS, Munich, Germany.

^tAlso at Università della Basilicata, Potenza, Italy.

^uAlso at Universidad de Alcalá, Alcalá de Henares, Spain.

^vAlso at Università di Urbino, Urbino, Italy.