

Observation of new structure in the $J/\psi J/\psi$ mass spectrum in proton-proton collisions at $\sqrt{s} = 13$ TeV

The CMS Collaboration*

Abstract

A search is reported for near-threshold structures in the $J/\psi J/\psi$ invariant mass spectrum produced in proton-proton collisions at $\sqrt{s} = 13$ TeV from data collected by the CMS experiment, corresponding to an integrated luminosity of 135 fb^{-1} . A new structure is observed with a significance above 5 standard deviations at a mass of 6552 ± 10 (stat) ± 12 (syst) MeV. Another structure with even higher significance is found at a mass of 6927 ± 9 (stat) ± 4 (syst) MeV, which is consistent with the $X(6900)$ resonance reported by the LHCb experiment and confirmed by the ATLAS experiment. Evidence for another new structure, with a local significance of 4.1 standard deviations, is found at a mass of 7287^{+20}_{-18} (stat) ± 5 (syst) MeV. The masses and significances are obtained in a model without considering possible quantum mechanical interference between the resonances. Incorporating this interference provides a better description of the mass spectrum between the resonances and shifts the measured masses by up to 150 MeV.

Submitted to Physical Review Letters

The prospect of exotic hadrons—states other than $q\bar{q}$ or qqq —dates back to Gell-Mann’s seminal 1964 paper [1]. However, interest, both theoretical and experimental, in exotics wavered for decades. The discovery of the charmonium-like $X(3872)$ state by the Belle Collaboration in 2003 [2] propelled exotics from speculative chimeras to the frontier of hadronic physics [3]. While many tetraquark candidates containing heavy quarks (c , b) are now known, questions still abound, including which of these are truly exotic hadrons [3], and if they are bound states, what is their internal structure (e.g., molecules, bound states of diquarks, etc. [4–6]).

Beginning with the J/ψ discovery in 1974 [7, 8], heavy quarkonia brought clarity to the quark model. Similarly, exotic states composed entirely of heavy quarks may offer analogous insights. Exotic states decaying into the $J/\psi J/\psi$ and YY channels are particularly promising experimentally because of the efficient triggering and reconstruction of muonic channels. The possibility of structure appearing in these channels is enhanced by an empirical pattern first observed with light mesons: the mass spectra of two vector states with the same isospin (e.g., $\phi\phi$ and $\omega\phi$) have near-threshold enhancements, whereas systems of two vector mesons of different isospin (e.g., $\phi\rho$ and $\omega\rho$) have no structure [9]. Such a behavior has been corroborated by the enhancements observed in $J/\psi\omega$ [10] and $J/\psi\phi$ [11] spectra.

The first searches for all heavy quark states were in the all-bottom sector, and were inconclusive [12, 13]. In 2020, the LHCb Collaboration reported structure in the $J/\psi J/\psi$ channel, which consisted of a broad enhancement just above the $J/\psi J/\psi$ mass and a narrow peak designated as $X(6900)$ [14]. The nature of the broad structure was uncertain, but the narrow one was interpreted to be an all-charm tetraquark state [15–22]. Considerations of such states date back to 1976 [23, 24] and have recently become the object of renewed interest [25–35]. However, non-tetraquark interpretations of this new structure have also been proposed [36–38].

Here we report on the $J/\psi J/\psi$ invariant mass spectrum from proton-proton (pp) collisions as recorded with the CMS detector, where the J/ψ mesons are reconstructed from $\mu^+\mu^-$ pairs. The data sample corresponds to an integrated luminosity of 135 fb^{-1} [39–41] at a center-of-mass energy of 13 TeV.

The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the solenoid volume are a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter, and a brass and scintillator hadron calorimeter, each composed of a barrel and two endcap sections. Muons are detected in gas-ionization chambers embedded in the steel flux-return yoke outside the solenoid. A more detailed description of the CMS detector can be found in Ref. [42].

Several Monte Carlo (MC) samples are used to model the detector mass resolution and efficiency of signals. As simulations of tetraquarks states are not available and the analysis is independent of signal kinematic details, we use the production of other particles to simulate the signal, with the masses defined appropriately. We use $J^P = 0^+$ mesons for a variety of masses decaying to $J/\psi J/\psi$ as the signal, and simulate their production via gluon-gluon interactions. We model signal particles by $\chi_{b0}(1P)$ production using the PYTHIA event generator (version 8.230) [43]. As an alternate production model, signal is also simulated by generating a sample with ‘Higgs bosons’ decaying to $ZZ^{(*)}$, where the $Z^{(*)}$ is redefined as a J/ψ , using the JHUGEN event generator (version 7.40) [44, 45]. To understand the detector mass resolution, the samples are generated with a negligible natural width.

Background events, i.e., selected events that are not from a resonant decay to two J/ψ candidates, are expected to be dominated by two genuine J/ψ decays arising either from a single parton-parton collision, i.e., nonresonant single-parton scattering (NRSPS) or a pair of parton-

parton interactions in a single pp collision, i.e., double-parton scattering (DPS). The NRSPS contribution is expected to dominate the DPS contribution near the $J/\psi/\psi$ threshold, with the DPS contribution dominating at masses above 11 GeV. To simulate these backgrounds we use the PYTHIA generator. The next-to-leading order CASCADE generator [46] and the next-to-next-to-leading order HELAC-ONIA generator [47–50] are also used to simulate alternative NRSPS shapes for the estimation of the systematic uncertainty in the background. Direct production, as well as feed-down processes where J/ψ mesons arise as decay products, are included. Feed-down processes from double-charmonium resonances [38] are generated using the PYTHIA generator separately. Generated events are processed through the full CMS detector simulation, which utilizes the GEANT4 toolkit [51].

Events of interest are selected using a two-tiered trigger system [52]. The first level (L1), composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events. The second level, known as the high-level trigger (HLT), consists of a farm of processors running a faster version of the full event reconstruction software, which reduces the event rate to around 1 kHz. The data collected in 2016 correspond to an integrated luminosity of 36.3 fb^{-1} and were obtained with an L1 trigger that required at least three muon candidates. The HLT required that the pseudorapidity of each muon satisfies $|\eta| < 2.5$, and a pair of oppositely charged muon candidates with an invariant mass between 2.95 and 3.25 GeV, a distance of closest approach between the two muons less than 0.5 cm, and a fit to a common vertex with a χ^2 probability greater than 0.5%. The data collected in 2017–2018 correspond to an integrated luminosity of 98.6 fb^{-1} and utilized an L1 trigger requiring at least three muons with the highest and second-highest transverse momentum (p_T) muons having $p_T > 5$ and 3 GeV, respectively, and at least one oppositely charged pair of muons with invariant mass below 9 GeV. The HLT criteria replicated those of 2016 with the additional requirement that the two muons with invariant mass between 2.95 and 3.25 GeV each have $p_T > 3.5$ GeV.

The offline selection criteria were finalized before unblinding the signal region (defined as invariant mass below 7.8 GeV) and are based in part on an examination of data collected in 2011 and 2012. It requires at least four muons, each of them having $p_T > 2$ GeV, $|\eta| < 2.4$, and satisfying the “soft muon identification” requirements [53]. Pairs of oppositely charged muons are used to form J/ψ candidates, and these are required to have $p_T > 3.5$ GeV. To confirm the HLT requirement, each muon pair is fit to a vertex, requiring a fit probability greater than 0.5%. Combinations of four muons from two J/ψ candidates are fit to a common vertex, which must have a fit probability greater than 0.5%. A mass constrained vertex fit is applied to the oppositely-charged muon pairs, in which the muon pair’s mass is constrained to the known J/ψ mass $m_{J/\psi}$ [54], and the probability is required to be greater than 0.1%. The two constrained J/ψ candidates are then fit to a common vertex and its fit probability must exceed 0.1%. An event may have multiple J/ψ -pair candidates. If both charge permutations of the same four muons have masses within three standard deviations (3σ) of the known J/ψ mass, we select the pairing that minimizes $(\Delta m_1/\sigma_{m_1})^2 + (\Delta m_2/\sigma_{m_2})^2$, where Δm is the difference between the reconstructed dimuon mass and the J/ψ mass and σ_m is the calculated uncertainty of the reconstructed dimuon mass (affects 0.2% of events). All $J/\psi/J/\psi$ candidates are kept if they arise from more than four muons (affects 0.2% of events).

The $J/\psi/J/\psi$ invariant mass spectrum of the 8651 selected candidate pairs below 9 GeV is shown in Fig. 1. To better constrain the smooth background continuum, the full analysis range extends up to masses of 15 GeV (the corresponding mass spectrum is shown in Fig. A.1 of Appendix A).

Using an unbinned extended likelihood method, we fit the data from the threshold, $2m_{J/\psi}$, to 15 GeV. Signals are represented by relativistic S-wave Breit–Wigner (BW) functions [54, 56, 57]




Figure 1: The $J/\psi J/\psi$ invariant mass spectrum in the range up to 9 GeV, with fits consisting of three signal functions (BW_1 , BW_2 and BW_3) and a background model (see text). The left plot shows the fit without interference. The right plot shows the fit that includes interference, where “Interfering BWs” refers to the total contribution of all the interfering amplitudes, and their cross-terms. For clarity, only the sum of the three background components (NRSPS+DPS+ BW_0) is shown on the plots. The lower portion of the plots shows the pulls, i.e., the number of standard deviations (statistical uncertainties only) that the binned data differ from the fit.

convolved with resolution functions, which are sums of two Gaussian functions with the same mean value. The detector mass resolution varies from about 10 MeV at a mass of 6500 MeV to 18 MeV at 7300 MeV. The BW functions are not modified by acceptance or efficiency corrections because these vary only slowly, by less than 1.3% over the natural width of the $X(6900)$.

The NRSPS and DPS backgrounds are parameterizations of distributions of the corresponding simulated events. The NRSPS component includes an exponential function where one of its parameters is a free parameter in the fit. The effect of feed-down from heavier mass states such as the $\psi(2S)$ is accounted for as a systematic uncertainty. Combinatorial backgrounds arise when one or more muon candidates are not from a J/ψ meson decay, and from hadrons that are misidentified as a muon or misreconstructed. The shapes of combinatorial backgrounds are well modeled by the NRSPS + DPS parameterizations, and the residual effects are accounted for in the systematic uncertainties.

Aside from the NRSPS and DPS background components, a component to model a threshold enhancement is included, as was done in Ref. [14]. This excess may be due to a resonance, but other processes could be responsible: coupled-channel interactions [58], triangle singularities [59], Pomeron exchange [36], or simply an inadequate NRSPS model. This region may also include feed-down from potential higher mass tetraquarks that are only partially reconstructed. Because of these uncertainties, we regard this enhancement as an additional background component, and model it with a BW function (BW_0) with free mass and width, which provides a good *ad hoc* description of this feature.

We begin with a background-only fit (NRSPS + DPS + BW_0) and then add signal contributions one at a time to the fit. A signal BW is kept as long as its local significance exceeds 3σ , with the local significance calculated from the difference in log-likelihood between including and not including the component, and only accounting for statistical uncertainties. Three resonance structures, labeled BW_1 , BW_2 , and BW_3 in order of increasing mass, are found to be statistically significant. The local significance of these peaks, calculated from the log-likelihood differences between the full fit and the fit with the BW of interest removed, are 6.5σ , 9.4σ , and 4.1σ , respectively. Figure 1 (left) shows this fit, with numerical results in Table 1. The BW widths are

all much larger than the detector mass resolution.

Table 1: Summary of the fit results for the $J/\psi/\psi$ invariant mass distribution. The mass m and natural width Γ for both the no-interference model and the interference model, and the signal yields N for the no-interference model, are given for the three signal structures. The dual uncertainties are the statistical followed by the systematic components, and single uncertainties are only statistical.

		BW ₁	BW ₂	BW ₃
No-interference	m [MeV]	$6552 \pm 10 \pm 12$	$6927 \pm 9 \pm 4$	$7287^{+20}_{-18} \pm 5$
	Γ [MeV]	$124^{+32}_{-26} \pm 33$	$122^{+24}_{-21} \pm 18$	$95^{+59}_{-40} \pm 19$
	N	470^{+120}_{-110}	492^{+78}_{-73}	156^{+64}_{-51}
Interference	m [MeV]	6638^{+43+16}_{-38-31}	6847^{+44+48}_{-28-20}	7134^{+48+41}_{-25-15}
	Γ [MeV]	$440^{+230+110}_{-200-240}$	191^{+66+25}_{-49-17}	97^{+40+29}_{-29-26}

We quantify the goodness of fit by the probability of the fit's χ^2 (over the full mass range with the bin size of Fig. 1) and the number of degrees of freedom. Because the signal region is only a small fraction of the fit range, the impact of deviations from the model in this region is diluted. As an alternative, we also compute a probability using only the bins below 7.8 GeV, which we refer to as the signal-region χ^2 probability.

The full χ^2 fit probability is 98%. However, the dips in the data around 6750 and 7150 MeV are poorly described: the signal-region χ^2 probability is 9%. Agreement may be improved by introducing interference, as was done by LHCb [14].

Our construction of interference models is based on the three structures found in the no-interference fit. We consider interference between pairs of (“two-way”), or all three (“three-way”), components. The three-way interference is implemented with a term proportional to $|r_1 \exp(i\phi_1)BW_1 + BW_2 + r_3 \exp(i\phi_3)BW_3|^2$, where $r_{1,3}$ and $\phi_{1,3}$ are the relative magnitudes and phases of $BW_{1,3}$ with respect to BW_2 [54]. We choose this model that shows a signal-region χ^2 probability of 65%—whereas all two-way models are below 30%.

The three-way interference fit is shown in Fig. 1 (right), and the fit parameters are listed in Table 1. In contrast, the interference model utilized by LHCb [14] consisted of a BW for the $X(6900)$ signal plus an auxiliary BW that interfered with the NRSPS background (Model II). This model by itself (without including interference between resonances) does not fit as well and adding this interference to the three-way interference does not improve upon the three-way interference fit.

The interference fit assumes that the unknown J^{PC} quantum numbers of these states are identical, and they are formed from a coherent production process. Other models may also be able to describe the dips. For example, inspired by predictions of dense spectroscopy of tetra-charm states [15–21], representing individual peaks by multiple overlapping narrow resonances could provide good fits by enabling the BW functions to fall more deeply into the dips. Because of these uncertainties we adopt the no-interference case as our main result.

Systematic uncertainties for masses and widths are determined by varying aspects of, and inputs to, the fits. For a given source, the largest deviation from a parameter's nominal value is taken as its systematic uncertainty. The sources of systematic uncertainty that have been considered are: different BW shapes (P - and D -wave, and alternative values for the “interaction radius” in the Blatt–Weisskopf barrier factor [56, 57]); alternative DPS parameterizations, such

as those obtained by mixing data events (artificial $J/\psi J/\psi$ mass spectrum formed by two J/ψ s from different events); NRSPS parameterizations from different simulations (CASCADE and HELAC-ONIA) or floating otherwise fixed parameters individually; momentum scale (based on the shift in the unconstrained fitted J/ψ meson mass from the world average value [54]); detector mass resolution (slightly different resolution models based on the PYTHIA or JHUGEN generators); combinatorial background shape (the result of altering the parameterization of the combinatorial background); efficiency corrections (the difference between not applying and applying efficiency corrections based on the PYTHIA or JHUGEN generator); and the effects of including various feed-down components from hypothetical heavier tetraquarks [38]. The effects of feed-down components in the interference model can produce asymmetric uncertainties, so an asymmetric uncertainty is assigned for this source. For other sources, symmetric uncertainties are assigned. The principal systematic effects are summarized in Table 2. The total uncertainties are their sum in quadrature and also appear in Table 1. Including feed-down in the fit affects signal structures, especially for the BW_1 , because of their overlap in the $J/\psi J/\psi$ mass distribution. Compared to the no-interference model, the greater complexity and increased parameter correlations of the interference model results in larger systematic uncertainties.

Table 2: Dominant contributions to the systematic uncertainties in masses and widths, in MeV, for the two fits. The “Total uncertainty” is the quadratic sum of all individual components, including the unlisted non-dominant contributions.

Fit	Dominant sources	M_{BW_1}	M_{BW_2}	M_{BW_3}	Γ_{BW_1}	Γ_{BW_2}	Γ_{BW_3}
No-interference	Signal shape	3	3	3	10	5	5
	NRSPS shape	3	1	1	18	15	17
	Feed-down	11	1	1	25	8	6
	Total uncertainty	12	4	5	33	18	19
Interference	Signal shape	7	12	7	56	8	7
	DPS shape	1	3	2	18	6	2
	NRSPS shape	9	14	13	85	9	20
	Mass resolution	8	4	1	24	7	13
	Combinatorial bkg.	7	2	<1	5	3	2
	Feed-down	+0 -27	+44 -0	+38 -0	+0 -210	+19 -0	+12 -0
	Total uncertainty	+16 -31	+48 -20	+41 -15	+110 -240	+25 -17	+29 -26

The impact of systematic uncertainties on the significances of the structures was checked in the no-interference fit model by using discrete sets of individual alternative hypotheses and re-computing the significances. The only appreciable reduction is for BW_1 , where the significance is reduced from 6.5 to 5.7 σ .

The BW_2 parameters are within 2 σ , when comparing the analogous no-interference or interference models, of the $X(6900)$ values reported by LHCb. For their Model I, consisting of a BW for the $X(6900)$ signal plus two auxiliary threshold BWs, they reported an $X(6900)$ mass of $6905 \pm 11 \pm 7$ MeV with a natural width of $80 \pm 19 \pm 33$ MeV, and $6886 \pm 11 \pm 11$ MeV and $168 \pm 33 \pm 69$ MeV for Model II [14]. However, because the two experiments use different fit models, quantitative comparisons between them may be distorted, and thus we also fit our data with the two LHCb models. Neither LHCb model provides a good description of the CMS data, but nevertheless our $X(6900)$ parameters are again compatible. More information about these comparisons, including where some small discrepancies are present, is summarized in

Appendix B.

Our measured masses appear compatible with recent calculations of the $c\bar{c}\bar{c}$ spectrum [21, 60], which would indicate that these three structures may be a family of radial excitations of the same J^{PC} . This is the case for both no-interference and interference masses, albeit for different theoretical models.

In summary, the $J/\psi J/\psi$ invariant mass spectrum has been presented. The data were collected with the CMS detector from pp collisions at $\sqrt{s} = 13$ TeV and correspond to an integrated luminosity of 135 fb^{-1} . Two new structures, tentatively named $X(6600)$ and $X(7300)$, are found with masses of 6552 ± 10 (stat) ± 12 (syst) MeV and 7287_{-18}^{+20} (stat) ± 5 (syst) MeV, and the $X(6900)$ structure observed by LHCb is confirmed with a mass of 6927 ± 9 (stat) ± 4 (syst) MeV. The local significances of these peaks are, for increasing mass, 6.5, 9.4, and 4.1 standard deviations. These results are from a fit that does not include interference effects. Adding interference terms between the three signals results in better agreement to the data in the regions between the resonances and results in shifts of the resonance parameters. Numeric results are provided in the HEPData record [61].

As this Letter was being finalized the ATLAS Collaboration released a paper [62] confirming the $X(6900)$ structure in the $J/\psi J/\psi$ spectrum, and also reported near-threshold excesses, including a possible feature in the $\psi(2S)J/\psi$ channel.

Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid and other centers for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, NSFC, Nanjing Normal University research start-up funding project, Tsinghua University Initiative Scientific Research and Dushi Programs, and Natural Science Foundation of China under Grants 11975010, 12075123, 12061141002 (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); MoER, ERC PUT and ERDF (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); BMBF, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LAS (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES and NSC (Poland); FCT (Portugal); MESTD (Serbia); MCIN/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI and NSTDA (Thailand); TUBITAK and TENMAK (Turkey); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (USA).

References

- [1] M. Gell-Mann, “A schematic model of baryons and mesons”, *Phys. Lett.* **8** (1964) 214, doi:10.1016/S0031-9163(64)92001-3.
- [2] Belle Collaboration, “Observation of a narrow charmoniumlike state in exclusive $B^\pm \rightarrow K^\pm \pi^+ \pi^- J/\psi$ decays”, *Phys. Rev. Lett.* **91** (2003) 262001, doi:10.1103/PhysRevLett.91.262001, arXiv:hep-ex/0309032.
- [3] A. Ali, L. Maiani, and A. D. Polosa, “Multiquark hadrons”. Cambridge University Press, Cambridge, 2019. doi:10.1017/9781316761465.
- [4] N. Brambilla et al., “The XYZ states: experimental and theoretical status and perspectives”, *Phys. Rept.* **873** (2020) 1, doi:10.1016/j.physrep.2020.05.001, arXiv:1907.07583.
- [5] M. Nielsen et al., “Supersymmetry in the double-heavy hadronic spectrum”, *Phys. Rev. D* **98** (2018) 034002, doi:10.1103/PhysRevD.98.034002, arXiv:1805.11567.
- [6] M. Nielsen and S. J. Brodsky, “Hadronic superpartners from a superconformal and supersymmetric algebra”, *Phys. Rev. D* **97** (2018) 114001, doi:10.1103/PhysRevD.97.114001, arXiv:1802.09652.
- [7] J. J. Aubert et al., “Experimental observation of a heavy particle j ”, *Phys. Rev. Lett.* **33** (1974) 1404, doi:10.1103/PhysRevLett.33.1404.
- [8] J. E. Augustin et al., “Discovery of a narrow resonance in e^+e^- annihilation”, *Phys. Rev. Lett.* **33** (1974) 1406, doi:10.1103/PhysRevLett.33.1406.
- [9] K. Yi, “Experimental review of structures in the $J/\psi\phi$ mass spectrum”, *Int. J. Mod. Phys. A* **28** (2013) 1330020, doi:10.1142/S0217751X13300202, arXiv:1308.0772.
- [10] BaBar Collaboration, “Evidence for the decay $X(3872) \rightarrow J/\psi\omega$ ”, *Phys. Rev. D* **82** (2010) 011101, doi:10.1103/PhysRevD.82.011101, arXiv:1005.5190.
- [11] LHCb Collaboration, “Observation of new resonances decaying to $J/\psi K^+$ and $J/\psi\phi$ ”, *Phys. Rev. Lett.* **127** (2021) 082001, doi:10.1103/PhysRevLett.127.082001, arXiv:2103.01803.
- [12] LHCb Collaboration, “Search for beautiful tetraquarks in the $Y(1S)\mu^+\mu^-$ invariant-mass spectrum”, *JHEP* **10** (2018) 086, doi:10.1007/JHEP10(2018)086, arXiv:1806.09707.
- [13] CMS Collaboration, “Measurement of the $Y(1S)$ pair production cross section and search for resonances decaying to $Y(1S)\mu^+\mu^-$ in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *Phys. Lett. B* **808** (2020) 135578, doi:10.1016/j.physletb.2020.135578, arXiv:2002.06393.
- [14] LHCb Collaboration, “Observation of structure in the J/ψ -pair mass spectrum”, *Sci. Bull.* **65** (2020) 1983, doi:10.1016/j.scib.2020.08.032, arXiv:2006.16957.
- [15] J.-Z. Wang, D.-Y. Chen, X. Liu, and T. Matsuki, “Producing fully charm structures in the J/ψ -pair invariant mass spectrum”, *Phys. Rev. D* **103** (2021) 071503, doi:10.1103/PhysRevD.103.L071503, arXiv:2008.07430.

-
- [16] H.-X. Chen, W. Chen, X. Liu, and S.-L. Zhu, “Strong decays of fully-charm tetraquarks into di-charmonia”, *Sci. Bull.* **65** (2020) 1994, doi:10.1016/j.scib.2020.08.038, arXiv:2006.16027.
- [17] M. A. Bedolla, J. Ferretti, C. D. Roberts, and E. Santopinto, “Spectrum of fully-heavy tetraquarks from a diquark+antidiquark perspective”, *Eur. Phys. J. C* **80** (2020) 1004, doi:10.1140/epjc/s10052-020-08579-3, arXiv:1911.00960.
- [18] X. Jin, Y. Xue, H. Huang, and J. Ping, “Full-heavy tetraquarks in constituent quark models”, *Eur. Phys. J. C* **80** (2020) 1083, doi:10.1140/epjc/s10052-020-08650-z, arXiv:2006.13745.
- [19] M.-S. Liu, F.-X. Liu, X.-H. Zhong, and Q. Zhao, “Full-heavy tetraquark states and their evidences in the LHCb di- J/ψ spectrum”, 2020. arXiv:2006.11952.
- [20] F.-X. Liu, M.-S. Liu, X.-H. Zhong, and Q. Zhao, “Higher mass spectra of the fully-charmed and fully-bottom tetraquarks”, *Phys. Rev. D* **104** (2021) 116029, doi:10.1103/PhysRevD.104.116029, arXiv:2110.09052.
- [21] R. Zhu, “Fully-heavy tetraquark spectra and production at hadron colliders”, *Nucl. Phys. B* **966** (2021) 115393, doi:10.1016/j.nuclphysb.2021.115393, arXiv:2010.09082.
- [22] J. F. Giron and R. F. Lebed, “Simple spectrum of $c\bar{c}c\bar{c}$ states in the dynamical diquark model”, *Phys. Rev. D* **102** (2020) 074003, doi:10.1103/PhysRevD.102.074003, arXiv:2008.01631.
- [23] Y. Iwasaki, “Is a state $c\bar{c}c\bar{c}$ found at 6.0 GeV?”, *Phys. Rev. Lett.* **36** (1976) 1266, doi:10.1103/PhysRevLett.36.1266.
- [24] K.-T. Chao, “The $(cc) - (\bar{c}\bar{c})$ (diquark-antidiquark) states in e^+e^- annihilation”, *Z. Phys. C* **7** (1981) 317, doi:10.1007/BF01431564.
- [25] A. V. Berezhnoy, A. V. Luchinsky, and A. A. Novoselov, “Heavy tetraquarks production at the LHC”, *Phys. Rev. D* **86** (2012) 034004, doi:10.1103/PhysRevD.86.034004, arXiv:1111.1867.
- [26] J. Wu et al., “Heavy-flavored tetraquark states with the $QQ\bar{Q}\bar{Q}$ configuration”, *Phys. Rev. D* **97** (2018) 094015, doi:10.1103/PhysRevD.97.094015, arXiv:1605.01134.
- [27] Y. Bai, S. Lu, and J. Osborne, “Beauty-full tetraquarks”, *Phys. Lett. B* **798** (2019) 134930, doi:10.1016/j.physletb.2019.134930, arXiv:1612.00012.
- [28] Z.-G. Wang, “Analysis of the $QQ\bar{Q}\bar{Q}$ tetraquark states with QCD sum rules”, *Eur. Phys. J. C* **77** (2017) 432, doi:10.1140/epjc/s10052-017-4997-0, arXiv:1701.04285.
- [29] M. N. Anwar et al., “Spectroscopy and decays of the fully-heavy tetraquarks”, *Eur. Phys. J. C* **78** (2018) 647, doi:10.1140/epjc/s10052-018-6073-9, arXiv:1710.02540.
- [30] J.-M. Richard, A. Valcarce, and J. Vijande, “String dynamics and metastability of all-heavy tetraquarks”, *Phys. Rev. D* **95** (2017) 054019, doi:10.1103/PhysRevD.95.054019, arXiv:1703.00783.
- [31] A. Esposito and A. D. Polosa, “A $bb\bar{b}\bar{b}$ di-bottomonium at the LHC?”, *Eur. Phys. J. C* **78** (2018) 782, doi:10.1140/epjc/s10052-018-6269-z, arXiv:1807.06040.

- [32] M. Karliner, S. Nussinov, and J. L. Rosner, “ $QQ\bar{Q}\bar{Q}$ states: Masses, production, and decays”, *Phys. Rev. D* **95** (2017) 034011, doi:10.1103/PhysRevD.95.034011, arXiv:1611.00348.
- [33] M.-S. Liu, Q.-F. Lü, X.-H. Zhong, and Q. Zhao, “All-heavy tetraquarks”, *Phys. Rev. D* **100** (2019) 016006, doi:10.1103/PhysRevD.100.016006, arXiv:1901.02564.
- [34] W. Chen et al., “Hunting for exotic doubly hidden-charm/bottom tetraquark states”, *Phys. Lett. B* **773** (2017) 247, doi:10.1016/j.physletb.2017.08.034, arXiv:1605.01647.
- [35] G.-J. Wang, L. Meng, and S.-L. Zhu, “Spectrum of the fully-heavy tetraquark state $QQ\bar{Q}'\bar{Q}'$ ”, *Phys. Rev. D* **100** (2019) 096013, doi:10.1103/PhysRevD.100.096013, arXiv:1907.05177.
- [36] C. Gong et al., “Nature of $X(6900)$ and its production mechanism at LHCb”, *Phys. Lett. B* **824** (2022) 136794, doi:10.1016/j.physletb.2021.136794, arXiv:2011.11374.
- [37] B.-D. Wan and C.-F. Qiao, “Gluonic tetracharm configuration of $X(6900)$ ”, *Phys. Lett. B* **817** (2021) 136339, doi:10.1016/j.physletb.2021.136339, arXiv:2012.00454.
- [38] J.-Z. Wang, X. Liu, and T. Matsuki, “Fully-heavy structures in the invariant mass spectrum of $J/\psi\psi(3686)$, $J/\psi\psi(3770)$, $\psi(3686)\psi(3686)$, and $J/\psi Y(1S)$ at hadron colliders”, *Phys. Lett. B* **816** (2021) 136209, doi:10.1016/j.physletb.2021.136209, arXiv:2012.03281.
- [39] CMS Collaboration, “Precision luminosity measurement in proton-proton collisions at $\sqrt{s} = 13$ TeV in 2015 and 2016 at CMS”, *Eur. Phys. J. C* **81** (2021) 800, doi:10.1140/epjc/s10052-021-09538-2, arXiv:2104.01927.
- [40] CMS Collaboration, “CMS luminosity measurement for the 2017 data-taking period at $\sqrt{s} = 13$ TeV”, CMS Physics Analysis Summary CMS-PAS-LUM-17-004, 2018.
- [41] CMS Collaboration, “CMS luminosity measurement for the 2018 data-taking period at $\sqrt{s} = 13$ TeV”, CMS Physics Analysis Summary CMS-PAS-LUM-18-002, 2019.
- [42] CMS Collaboration, “The CMS experiment at the CERN LHC”, *JINST* **3** (2008) S08004, doi:10.1088/1748-0221/3/08/S08004.
- [43] T. Sjöstrand et al., “An introduction to PYTHIA 8.2”, *Comput. Phys. Commun.* **191** (2015) 159, doi:10.1016/j.cpc.2015.01.024, arXiv:1410.3012.
- [44] Y. Gao et al., “Spin determination of single-produced resonances at hadron colliders”, *Phys. Rev. D* **81** (2010) 075022, doi:10.1103/PhysRevD.81.075022, arXiv:1001.3396.
- [45] A. V. Gritsan et al., “New features in the JHU generator framework: Constraining Higgs boson properties from on-shell and off-shell production”, *Phys. Rev. D* **102** (2020) 056022, doi:10.1103/PhysRevD.102.056022, arXiv:2002.09888.
- [46] H. Jung et al., “The CCFM Monte Carlo generator CASCADE version 2.2.03”, *Eur. Phys. J. C* **70** (2010) 1237, doi:10.1140/epjc/s10052-010-1507-z, arXiv:1008.0152.
- [47] H.-S. Shao, “HELAC-ONIA: An automatic matrix element generator for heavy quarkonium physics”, *Comput. Phys. Commun.* **184** (2013) 2562, doi:10.1016/j.cpc.2013.05.023, arXiv:1212.5293.

-
- [48] J.-P. Lansberg and H.-S. Shao, “ J/ψ -pair production at large momenta: Indications for double parton scatterings and large α_S^5 contributions”, *Phys. Lett. B* **751** (2015) 479, doi:10.1016/j.physletb.2015.10.083, arXiv:1410.8822.
- [49] J.-P. Lansberg and H.-S. Shao, “Production of $J/\psi + \eta_c$ versus $J/\psi + J/\psi$ at the LHC: Importance of real α_S^5 corrections”, *Phys. Rev. Lett.* **111** (2013) 122001, doi:10.1103/PhysRevLett.111.122001, arXiv:1308.0474.
- [50] H.-S. Shao, “HELAC-ONIA 2.0: An upgraded matrix-element and event generator for heavy quarkonium physics”, *Comput. Phys. Commun.* **198** (2016) 238, doi:10.1016/j.cpc.2015.09.011, arXiv:1507.03435.
- [51] GEANT4 Collaboration, “GEANT4—a simulation toolkit”, *Nucl. Instrum. Meth. A* **506** (2003) 250, doi:10.1016/S0168-9002(03)01368-8.
- [52] CMS Collaboration, “The CMS trigger system”, *JINST* **12** (2017) P01020, doi:10.1088/1748-0221/12/01/P01020, arXiv:1609.02366.
- [53] CMS Collaboration, “Performance of CMS muon reconstruction in pp collision events at $\sqrt{s} = 7$ TeV”, *JINST* **7** (2012) P10002, doi:10.1088/1748-0221/7/10/P10002, arXiv:1206.4071.
- [54] Particle Data Group Collaboration, “Review of Particle Physics”, *PTEP* **2022** (2022) 083C01, doi:10.1093/ptep/ptac097.
- [55] See Supplemental Material at [URL will be inserted by publisher] for the fit projections in full range and the fit results with LHCb models.
- [56] F. Von Hippel and C. Quigg, “Centrifugal-barrier effects in resonance partial decay widths, shapes, and production amplitudes”, *Phys. Rev. D* **5** (1972) 624, doi:10.1103/PhysRevD.5.624.
- [57] S. U. Chung, “Helicity-coupling amplitudes in tensor formalism”, *Phys. Rev. D* **48** (1993) 1225, doi:10.1103/PhysRevD.48.1225. [Erratum: doi:10.1103/PhysRevD.56.4419].
- [58] X.-K. Dong et al., “Coupled-channel interpretation of the LHCb double- J/ψ spectrum and hints of a new state near the $J/\psi J/\psi$ threshold”, *Phys. Rev. Lett.* **126** (2021) 132001, doi:10.1103/PhysRevLett.126.132001, arXiv:2009.07795. [Erratum: doi:10.1103/PhysRevLett.127.119901].
- [59] F.-K. Guo, X.-H. Liu, and S. Sakai, “Threshold cusps and triangle singularities in hadronic reactions”, *Prog. Part. Nucl. Phys.* **112** (2020) 103757, doi:10.1016/j.ppnp.2020.103757, arXiv:1912.07030.
- [60] R. Tiwari, D. P. Rathaud, and A. K. Rai, “Spectroscopy of all charm tetraquark states”, *Indian J. Phys.* **97** (2023) 943, doi:10.1007/s12648-022-02427-8, arXiv:2108.04017.
- [61] HEPData record for this analysis, 2023. doi:10.17182/hepdata.141028.
- [62] ATLAS Collaboration, “Observation of an excess of di-charmonium events in the four-muon final state with the ATLAS detector”, 2023. arXiv:2304.08962. Submitted to *Phys. Rev. Lett.*

A Fit projections in full range

The $J/\psi J/\psi$ invariant mass spectrum for the full range of the fit is shown in Fig. A.1 for the no-interference and the interference models.




Figure A.1: The $J/\psi J/\psi$ invariant mass spectrum with the no-interference fit (upper) and the interference fit (lower) in the full fit range (see text for model details). The “Interference BWs” curve is the total contribution of all the interference amplitudes and their cross terms. The lower portion of the plots shows the pulls, i.e., the number of standard deviations (statistical uncertainties only) that the binned data differ from the fit.

B Fitting with the LHCb models

In order to make a direct comparison to the results from the LHCb Collaboration [14] we also used their two principal models (I and II) to fit the CMS data. The details are provided here.

Model I consists of two auxiliary relativistic Breit–Wigner (BW) functions below the $X(6900)$ BW. The parameter values for these two BWs are not reported in Ref. [14], but the second auxiliary BW is in the same mass region as the CMS “ BW_1 ” and so we extend this labeling to the LHCb models.

The Model I fit is shown in Fig. B.1 (upper), with numerical results listed in Table B.1. The fit quality is poor with a signal-region χ^2 probability of 0.9%. The no-interference fit using the CMS model with three structures—and roughly the same number of $J/\psi J/\psi$ candidates after the final selection as LHCb—has a signal-region χ^2 probability of 9%.

Model II has, aside from the signal BW, a single auxiliary BW (here labeled as ‘ BW_1' ’), which




Figure B.1: The $J/\psi/\psi$ invariant mass spectrum with fits using LHCb models [14]: Model I (no-interference) (upper) and Model II (interference) (lower). Model details are given in the text. The “Interfering components” is the total contribution of BW_1 , NRSPS, and their interference. The lower portion of the plots shows the pulls between the binned data and the fit.


Table B.1: Summary of results for the two LHCb fit models from LHCb and CMS: the mass m and natural width Γ , in MeV, are given for the principal structures (the Breit–Wigner function at the threshold is not included in Model I), and LHCb values not reported in Ref. [14] are marked as “—”. Single uncertainties are statistical only.

Exp.	Fit	M_{BW_1}	Γ_{BW_1}	$M_{X(6900)}$	$\Gamma_{X(6900)}$
LHCb [14]	Model I	—	—	$6905 \pm 11 \pm 7$	$80 \pm 19 \pm 33$
CMS	Model I	6550 ± 10	112 ± 27	6927 ± 10	117 ± 24
LHCb [14]	Model II	6741 ± 6	288 ± 16	$6886 \pm 11 \pm 11$	$168 \pm 33 \pm 69$
CMS	Model II	6736 ± 38	439 ± 65	6918 ± 10	187 ± 40



















interferes with the NRSPS background. This interference was introduced to better model the dip around 6750 MeV. Our fit results for Model II are shown in Fig. B.1 (lower) and Table B.1. The 6750 MeV dip is now well described: the pulls in Fig. B.1 (lower) are well behaved in the vicinity of the $X(6900)$ peak. However, systematic deviations are evident elsewhere in the pulls, and the signal-region χ^2 probability of 0.8% for this fit is actually lower than for Model I.

C The CMS Collaboration




Yerevan Physics Institute, Yerevan, Armenia

A. Hayrapetyan, A. Tumasyan¹ 

Institut für Hochenergiephysik, Vienna, Austria

W. Adam , J.W. Andrejkovic, T. Bergauer , S. Chatterjee , K. Damanakis , M. Dragicevic , A. Escalante Del Valle , P.S. Hussain , M. Jeitler² , N. Krammer , D. Liko , I. Mikulec , J. Schieck² , R. Schöffbeck , D. Schwarz , M. Sonawane , S. Templ , W. Waltenberger , C.-E. Wulz² 














Universiteit Antwerpen, Antwerpen, Belgium

M.R. Darwish³ , T. Janssen , P. Van Mechelen 

Vrije Universiteit Brussel, Brussel, Belgium

E.S. Bols , J. D'Hondt , S. Dansana , A. De Moor , M. Delcourt , H. El Faham , S. Lowette , I. Makarenko , D. Müller , A.R. Sahasransu , S. Tavernier , M. Tytgat⁴ , S. Van Putte , D. Vannerom 

Université Libre de Bruxelles, Bruxelles, Belgium

B. Clerbaux , G. De Lentdecker , L. Favart , D. Hohov , J. Jaramillo , A. Khalilzadeh, K. Lee , M. Mahdavihorrami , A. Malara , S. Paredes , L. Pétré , N. Postiau, L. Thomas , M. Vanden Bemden, C. Vander Velde , P. Vanlaer 






Ghent University, Ghent, Belgium

M. De Coen , D. Dobur , Y. Hong , J. Knolle , L. Lambrecht , G. Mestdach, C. Rendón, A. Samalan, K. Skovpen , N. Van Den Bossche , L. Wezenbeek 




















Université Catholique de Louvain, Louvain-la-Neuve, Belgium

A. Benecke , G. Bruno , C. Caputo , C. Delaere , I.S. Donertas , A. Giammanco , K. Jaffel , Sa. Jain , V. Lemaitre, J. Lidrych , P. Mastrapasqua , K. Mondal , T.T. Tran , S. Wertz 

Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil

G.A. Alves , E. Coelho , C. Hensel , T. Menezes De Oliveira, A. Moraes , P. Rebello Teles , M. Soeiro

Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W.L. Aldá Júnior , M. Alves Gallo Pereira , M. Barroso Ferreira Filho , H. Brandao Malbouisson , W. Carvalho , J. Chinellato⁵, E.M. Da Costa , G.G. Da Silveira⁶ , D. De Jesus Damiao , S. Fonseca De Souza , J. Martins⁷ , C. Mora Herrera , K. Mota Amarilo , L. Mundim , H. Nogima , A. Santoro , S.M. Silva Do Amaral , A. Sznajder , M. Thiel , A. Vilela Pereira 

Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

C.A. Bernardes⁶ , L. Calligaris , T.R. Fernandez Perez Tomei , E.M. Gregores , P.G. Mercadante , S.F. Novaes , B. Orzari , Sandra S. Padula 

Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Aleksandrov , G. Antchev , R. Hadjiiska , P. Iaydjiev , M. Misheva , M. Shopova , G. Sultanov 





University of Sofia, Sofia, Bulgaria

A. Dimitrov , T. Ivanov , L. Litov , B. Pavlov , P. Petkov , A. Petrov , E. Shumka 




Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile

S. Keshri , S. Thakur 












Beihang University, Beijing, China

T. Cheng , Q. Guo, T. Javaid , M. Mittal , L. Yuan 









Department of Physics, Tsinghua University, Beijing, China

G. Bauer^{8,9}, J. Chen⁹, J. Gu⁹, Z. Hu , J. Liu, X. Wang¹⁰, H. Wen⁹, K. Yi^{9,11} , J. Zhang⁹ 

Institute of High Energy Physics, Beijing, China

G.M. Chen¹² , H.S. Chen¹² , M. Chen¹² , F. Iemmi , C.H. Jiang, A. Kapoor , H. Liao , Z.-A. Liu¹³ , F. Monti , R. Sharma , J.N. Song¹³, J. Tao , C. Wang¹², J. Wang , Z. Wang, H. Zhang 

State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

A. Agapitos , Y. Ban , A. Levin , C. Li , Q. Li , X. Lyu, Y. Mao, S.J. Qian , X. Sun , D. Wang , H. Yang, C. Zhou 




Sun Yat-Sen University, Guangzhou, China

Z. You 

University of Science and Technology of China, Hefei, China

N. Lu 

Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) - Fudan University, Shanghai, China

X. Gao¹⁴ , D. Leggat, H. Okawa , Y. Zhang 





Zhejiang University, Hangzhou, Zhejiang, China

Z. Lin , C. Lu , M. Xiao 





Universidad de Los Andes, Bogota, Colombia

C. Avila , D.A. Barbosa Trujillo, A. Cabrera , C. Florez , J. Fraga , J.A. Reyes Vega

Universidad de Antioquia, Medellin, Colombia

J. Mejia Guisao , F. Ramirez , M. Rodriguez , J.D. Ruiz Alvarez 

University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia

D. Giljanovic , N. Godinovic , D. Lelas , A. Sculac 









University of Split, Faculty of Science, Split, Croatia

M. Kovac , T. Sculac 




Institute Rudjer Boskovic, Zagreb, Croatia

P. Bargassa , V. Brigljevic , B.K. Chitroda , D. Ferencek , S. Mishra , A. Starodumov¹⁵ , T. Susa 

University of Cyprus, Nicosia, Cyprus




















































































































A. Attikis , K. Christoforou , S. Konstantinou , J. Mousa , C. Nicolaou, F. Ptochos , P.A. Razis , H. Rykaczewski, H. Saka , A. Stepennov 

Charles University, Prague, Czech Republic

M. Finger , M. Finger Jr. , A. Kveton 









Escuela Politecnica Nacional, Quito, Ecuador

E. Ayala 

Universidad San Francisco de Quito, Quito, EcuadorE. Carrera Jarrin **Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt**A.A. Abdelalim^{16,17} , E. Salama^{18,19} **Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt**M. Abdullah Al-Mashad , M.A. Mahmoud **National Institute of Chemical Physics and Biophysics, Tallinn, Estonia**R.K. Dewanjee²⁰ , K. Ehataht , M. Kadastik , T. Lange , S. Nandan , C. Nielsen , J. Pata , M. Raidal , L. Tani , C. Veelken **Department of Physics, University of Helsinki, Helsinki, Finland**H. Kirschenmann , K. Osterberg , M. Voutilainen **Helsinki Institute of Physics, Helsinki, Finland**S. Bharthuar , E. Brücken , F. Garcia , J. Havukainen , K.T.S. Kallonen , M.S. Kim , R. Kinnunen , T. Lampén , K. Lassila-Perini , S. Lehti , T. Lindén , M. Lotti , L. Martikainen , M. Myllymäki , M.m. Rantanen , H. Siikonen , E. Tuominen , J. Tuominiemi **Lappeenranta-Lahti University of Technology, Lappeenranta, Finland**P. Luukka , H. Petrow , T. Tuuva[†]**IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France**M. Besancon , F. Couderc , M. Dejardin , D. Denegri , J.L. Faure , F. Ferri , S. Ganjour , P. Gras , G. Hamel de Monchenault , V. Lohezic , J. Malcles , J. Rander , A. Rosowsky , M.Ö. Sahin , A. Savoy-Navarro²¹ , P. Simkina , M. Titov **Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France**C. Baldenegro Barrera , F. Beaudette , A. Buchot Perraguin , P. Busson , A. Cappati , C. Charlot , F. Damas , O. Davignon , G. Falmagne , B.A. Fontana Santos Alves , S. Ghosh , A. Gilbert , R. Granier de Cassagnac , A. Hakimi , B. Harikrishnan , L. Kalipoliti , G. Liu , J. Motta , M. Nguyen , C. Ochando , L. Portales , R. Salerno , U. Sarkar , J.B. Sauvan , Y. Sirois , A. Tarabini , E. Vernazza , A. Zabi , A. Zghiche **Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France**J.-L. Agram²² , J. Andrea , D. Apparú , D. Bloch , J.-M. Brom , E.C. Chabert , C. Collard , S. Falke , U. Goerlach , C. Grimault , R. Haeberle , A.-C. Le Bihan , M.A. Sessini , P. Van Hove **Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France**S. Beauceron , B. Blancon , G. Boudoul , N. Chanon , J. Choi , D. Contardo , P. Depasse , C. Dozen²³ , H. El Mamouni , J. Fay , S. Gascon , M. Gouzevitch , C. Greenberg , G. Grenier , B. Ille , I.B. Laktineh , M. Lethuillier , L. Mirabito , S. Perries , M. Vander Donckt , P. Verdier , J. Xiao **Georgian Technical University, Tbilisi, Georgia**I. Lomidze , T. Toriashvili²⁴ , Z. Tsamalaidze¹⁵ **RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany**V. Botta , L. Feld , K. Klein , M. Lipinski , D. Meuser , A. Pauls , N. Röwert 

M. Teroerde 









RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

S. Diekmann , A. Dodonova , N. Eich , D. Eliseev , F. Engelke , M. Erdmann , P. Fackeldey , B. Fischer , T. Hebbeker , K. Hoepfner , F. Ivone , A. Jung , M.y. Lee , L. Mastrolorenzo, M. Merschmeyer , A. Meyer , S. Mukherjee , D. Noll , A. Novak , F. Nowotny, A. Pozdnyakov , Y. Rath, W. Redjeb , F. Rehm, H. Reithler , V. Sarkisovi , A. Schmidt , S.C. Schuler, A. Sharma , A. Stein , F. Torres Da Silva De Araujo²⁵ , L. Vigilante, S. Wiedenbeck , S. Zaleski









RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany

C. Dziwok , G. Flügge , W. Haj Ahmad²⁶ , T. Kress , A. Nowack , O. Pooth , A. Stahl , T. Ziemons , A. Zotz









Deutsches Elektronen-Synchrotron, Hamburg, Germany

H. Aarup Petersen , M. Aldaya Martin , J. Alimena , S. Amoroso, Y. An , S. Baxter , M. Bayatmakou , H. Becerril Gonzalez , O. Behnke , A. Belvedere , S. Bhattacharya , F. Blekman²⁷ , K. Borras²⁸ , D. Brunner , A. Campbell , A. Cardini , C. Cheng, F. Colombina , S. Consuegra Rodríguez , G. Correia Silva , M. De Silva , G. Eckerlin, D. Eckstein , L.I. Estevez Banos , O. Filatov , E. Gallo²⁷ , A. Geiser , A. Giraldi , G. Greau, V. Guglielmi , M. Guthoff , A. Hinzmann , A. Jafari²⁹ , L. Jeppe , N.Z. Jomhari , B. Kaech , M. Kasemann , H. Kaveh , C. Kleinwort , R. Kogler , M. Komm , D. Krücker , W. Lange, D. Leyva Pernia , K. Lipka³⁰ , W. Lohmann³¹ , R. Mankel , I.-A. Melzer-Pellmann , M. Mendizabal Morentin , J. Metwally, A.B. Meyer , G. Milella , A. Mussgiller , A. Nürnberg , Y. Otarid, D. Pérez Adán , E. Ranken , A. Raspereza , B. Ribeiro Lopes , J. Rübenach, A. Saggio , M. Scham^{32,28} , V. Scheurer, S. Schnake²⁸ , P. Schütze , C. Schwanenberger²⁷ , M. Shchedrolosiev , R.E. Sosa Ricardo , L.P. Sreelatha Pramod , D. Stafford, F. Vazzoler , A. Ventura Barroso , R. Walsh , Q. Wang , Y. Wen , K. Wichmann, L. Wiens²⁸ , C. Wissing , S. Wuchterl , Y. Yang , A. Zimmermann Castro Santos

University of Hamburg, Hamburg, Germany

A. Albrecht , S. Albrecht , M. Antonello , S. Bein , L. Benato , M. Bonanomi , P. Connor , M. Eich, K. El Morabit , Y. Fischer , A. Fröhlich, C. Garbers , E. Garutti , A. Grohsjean , M. Hajheidari, J. Haller , H.R. Jabusch , G. Kasieczka , P. Keicher, R. Klanner , W. Korcari , T. Kramer , V. Kutzner , F. Labe , J. Lange , A. Lobanov , C. Matthies , A. Mehta , L. Moureaux , M. Mrowietz, A. Nigamova , Y. Nissan, A. Paasch , K.J. Pena Rodriguez , T. Quadfasel , B. Raciti , M. Rieger , D. Savoie , J. Schindler , P. Schleper , M. Schröder , J. Schwandt , M. Sommerhalder , H. Stadie , G. Steinbrück , A. Tews, M. Wolf

Karlsruher Institut fuer Technologie, Karlsruhe, Germany

S. Brommer , M. Burkart, E. Butz , T. Chwalek , A. Dierlamm , A. Droll, N. Faltermann , M. Giffels , A. Gottmann , F. Hartmann³³ , M. Horzela , U. Husemann , M. Klute , R. Koppenhöfer , M. Link, A. Lintuluoto , S. Maier , S. Mitra , M. Mormile , Th. Müller , M. Neukum, M. Oh , G. Quast , K. Rabbertz , I. Shvetsov , H.J. Simonis , N. Trevisani , R. Ulrich , J. van der Linden , R.F. Von Cube , M. Wassmer , S. Wieland , F. Wittig, R. Wolf , S. Wunsch, X. Zuo

Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

G. Anagnostou, P. Assiouras , G. Daskalakis , A. Kyriakis, A. Papadopoulos³³, A. Stakia 

National and Kapodistrian University of Athens, Athens, Greece

D. Karasavvas, P. Kontaxakis , G. Melachroinos, A. Panagiotou, I. Papavergou , I. Paraskevas , N. Saoulidou , K. Theofilatos , E. Tziaferi , K. Vellidis , I. Zisopoulos 




National Technical University of Athens, Athens, Greece

G. Bakas , T. Chatzistavrou, G. Karapostoli , K. Kousouris , I. Papakrivopoulos , E. Siamarkou, G. Tsipolitis, A. Zacharopoulou






University of Ioánnina, Ioánnina, Greece

K. Adamidis, I. Bestintzanos, I. Evangelou , C. Foudas, P. Gianneios , C. Kamtsikis, P. Katsoulis, P. Kokkas , P.G. Kosmoglou Kioseoglou , N. Manthos , I. Papadopoulos , J. Strologas 

MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

M. Csanád , K. Farkas , M.M.A. Gadallah³⁴ , Á. Kadlecik , P. Major , K. Mandal , G. Pásztor , A.J. Rádli³⁵ , G.I. Veres 



Wigner Research Centre for Physics, Budapest, Hungary

M. Bartók³⁶ , C. Hajdu , D. Horvath^{37,38} , F. Sikler , V. Veszpremi 




Institute of Nuclear Research ATOMKI, Debrecen, Hungary

G. Bencze, S. Czellar, J. Karancsi³⁶ , J. Molnar, Z. Szillasi

















Institute of Physics, University of Debrecen, Debrecen, Hungary

P. Raics, B. Ujvari³⁹ , G. Zilizi 

Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary

T. Csorgo³⁵ , F. Nemes³⁵ , T. Novak 

Panjab University, Chandigarh, India

J. Babbar , S. Bansal , S.B. Beri, V. Bhatnagar , G. Chaudhary , S. Chauhan , N. Dhingra⁴⁰ , R. Gupta, A. Kaur , A. Kaur , H. Kaur , M. Kaur , S. Kumar , P. Kumari , M. Meena , K. Sandeep , T. Sheokand, J.B. Singh⁴¹ , A. Singla 
















University of Delhi, Delhi, India

A. Ahmed , A. Bhardwaj , A. Chhetri , B.C. Choudhary , A. Kumar , M. Naimuddin , K. Ranjan , S. Saumya 


Saha Institute of Nuclear Physics, HBNI, Kolkata, India

S. Baradia , S. Barman⁴² , S. Bhattacharya , D. Bhowmik, S. Dutta , S. Dutta, B. Gomber⁴³ , P. Palit , G. Saha , B. Sahu⁴³ , S. Sarkar

Indian Institute of Technology Madras, Madras, India

P.K. Behera , S.C. Behera , S. Chatterjee , P. Jana , P. Kalbhor , J.R. Komaragiri⁴⁴ , D. Kumar⁴⁴ , M. Mohammad Mobassir Ameen , L. Panwar⁴⁴ , R. Pradhan , P.R. Pujahari , N.R. Saha , A. Sharma , A.K. Sikdar , S. Verma 

Tata Institute of Fundamental Research-A, Mumbai, India

T. Aziz, I. Das , S. Dugad, M. Kumar , G.B. Mohanty , P. Suryadevara

Tata Institute of Fundamental Research-B, Mumbai, India

A. Bala , S. Banerjee , R.M. Chatterjee, M. Guchait , S. Karmakar , S. Kumar , G. Majumder , K. Mazumdar , S. Mukherjee , A. Thachayath 

National Institute of Science Education and Research, An OCC of Homi Bhabha National



Institute, Bhubaneswar, Odisha, India

S. Bahinipati⁴⁵ , A.K. Das, C. Kar , D. Maity⁴⁶ , P. Mal , T. Mishra , V.K. Muraleedharan Nair Bindhu⁴⁶ , K. Naskar⁴⁶ , A. Nayak⁴⁶ , P. Sadangi, P. Saha , S.K. Swain, S. Varghese⁴⁶ , D. Vats⁴⁶ 

Indian Institute of Science Education and Research (IISER), Pune, India

A. Alpana , S. Dube , B. Kansal , A. Laha , A. Rastogi , S. Sharma 

Isfahan University of Technology, Isfahan, Iran

H. Bakhshiansohi⁴⁷ , E. Khazaie⁴⁸ , M. Zeinali⁴⁹ 
































Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

S. Chenarani⁵⁰ , S.M. Etesami , M. Khakzad , M. Mohammadi Najafabadi 




























University College Dublin, Dublin, Ireland

M. Grunewald 

INFN Sezione di Bari^a, Università di Bari^b, Politecnico di Bari^c, Bari, Italy

M. Abbrescia^{a,b} , R. Aly^{a,c,16} , A. Colaleo^a , D. Creanza^{a,c} , B. D' Anzi^{a,b} , N. De Filippis^{a,c} , M. De Palma^{a,b} , A. Di Florio^{a,c} , W. Elmetenawee^{a,b,16} , L. Fiore^a , G. Iaselli^{a,c} , G. Maggi^{a,c} , M. Maggi^a , I. Margjeka^{a,b} , V. Mastrapasqua^{a,b} , S. My^{a,b} , S. Nuzzo^{a,b} , A. Pellecchia^{a,b} , A. Pompili^{a,b} , G. Pugliese^{a,c} , R. Radogna^a , G. Ramirez-Sanchez^{a,c} , D. Ramos^a , A. Ranieri^a , L. Silvestris^a , F.M. Simone^{a,b} , Ü. Sözbilir^a , A. Stamerra^a , R. Venditti^a , P. Verwilligen^a , A. Zaza^{a,b} 

















INFN Sezione di Bologna^a, Università di Bologna^b, Bologna, Italy

G. Abbiendi^a , C. Battilana^{a,b} , D. Bonacorsi^{a,b} , L. Borgonovi^a , R. Campanini^{a,b} , P. Capiluppi^{a,b} , A. Castro^{a,b} , F.R. Cavallo^a , M. Cuffiani^{a,b} , G.M. Dallavalle^a , T. Diotallevi^{a,b} , F. Fabbri^a , A. Fanfani^{a,b} , D. Fasanella^{a,b} , P. Giacomelli^a , L. Giommi^{a,b} , C. Grandi^a , L. Guiducci^{a,b} , S. Lo Meo^{a,51} , L. Lunerti^{a,b} , S. Marcellini^a , G. Masetti^a , F.L. Navarra^{a,b} , A. Perrotta^a , F. Primavera^{a,b} , A.M. Rossi^{a,b} , T. Rovelli^{a,b} 

INFN Sezione di Catania^a, Università di Catania^b, Catania, Italy

S. Costa^{a,b,52} , A. Di Mattia^a , R. Potenza^{a,b}, A. Tricomi^{a,b,52} , C. Tuve^{a,b} 

INFN Sezione di Firenze^a, Università di Firenze^b, Firenze, Italy

G. Barbagli^a , G. Bardelli^{a,b} , B. Camaiani^{a,b} , A. Cassese^a , R. Ceccarelli^a , V. Ciulli^{a,b} , C. Civinini^a , R. D'Alessandro^{a,b} , E. Focardi^{a,b} , G. Latino^{a,b} , P. Lenzi^{a,b} , M. Lizzo^{a,b} , M. Meschini^a , S. Paoletti^a , A. Papanastassiou^{a,b}, G. Sguazzoni^a , L. Viliani^a 
























INFN Laboratori Nazionali di Frascati, Frascati, Italy

L. Benussi , S. Bianco , S. Meola⁵³ , D. Piccolo 












INFN Sezione di Genova^a, Università di Genova^b, Genova, Italy

P. Chatagnon^a , F. Ferro^a , E. Robutti^a , S. Tosi^{a,b} 































INFN Sezione di Milano-Bicocca^a, Università di Milano-Bicocca^b, Milano, Italy

A. Benaglia^a , G. Boldrini^a , F. Brivio^a , F. Cetorelli^a , F. De Guio^{a,b} , M.E. Dinardo^{a,b} , P. Dini^a , S. Gennai^a , A. Ghezzi^{a,b} , P. Govoni^{a,b} , L. Guzzi^a , M.T. Lucchini^{a,b} , M. Malberti^a , S. Malvezzi^a , A. Massironi^a , D. Menasce^a , L. Moroni^a , M. Paganoni^{a,b} , D. Pedrini^a , B.S. Pinolini^a, S. Ragazzi^{a,b} , N. Redaelli^a , T. Tabarelli de Fatis^{a,b} , D. Zuolo^a 












INFN Sezione di Napoli^a, Università di Napoli 'Federico II'^b, Napoli, Italy; Università della Basilicata^c, Potenza, Italy; Università G. Marconi^d, Roma, Italy

S. Buontempo^a , A. Cagnotta^{a,b} , F. Carnevali^{a,b} , N. Cavallo^{a,c} , A. De Iorio^{a,b} , F. Fabozzi^{a,c} , A.O.M. Iorio^{a,b} , L. Lista^{a,b,54} , P. Paolucci^{a,33} , B. Rossi^a , C. Sciacca^{a,b} 

















INFN Sezione di Padova^a, Università di Padova^b, Padova, Italy; Università di Trento^c, Trento, Italy

R. Ardino^a , P. Azzi^a , N. Bacchetta^{a,55} , P. Bortignon^a , A. Bragagnolo^{a,b} , R. Carlin^{a,b} , P. Checchia^a , T. Dorigo^a , F. Gasparini^{a,b} , U. Gasparini^{a,b} , G. Grosso^a , L. Layer^{a,56} , E. Lusiani^a , M. Margoni^{a,b} , A.T. Meneguzzo^{a,b} , M. Migliorini^{a,b} , F. Montecassiano^a , J. Pazzini^{a,b} , P. Ronchese^{a,b} , R. Rossin^{a,b} , F. Simonetto^{a,b} , G. Strong^a , M. Tosi^{a,b} , A. Triossi^{a,b} , S. Ventura^a , H. Yarar^{a,b} , M. Zanetti^{a,b} , P. Zotto^{a,b} , A. Zucchetta^{a,b} , G. Zumerle^{a,b} 
































INFN Sezione di Pavia^a, Università di Pavia^b, Pavia, Italy

S. Abu Zeid^{a,19} , C. Aimè^{a,b} , A. Braghieri^a , S. Calzaferri^{a,b} , D. Fiorina^{a,b} , P. Montagna^{a,b} , V. Re^a , C. Riccardi^{a,b} , P. Salvini^a , I. Vai^{a,b} , P. Vitulo^{a,b} 





















INFN Sezione di Perugia^a, Università di Perugia^b, Perugia, Italy

S. Ajmal^{a,b} , P. Asenov^{a,57} , G.M. Bilei^a , D. Ciangottini^{a,b} , L. Fanò^{a,b} , M. Magherini^{a,b} , G. Mantovani^{a,b} , V. Mariani^{a,b} , M. Menichelli^a , F. Moscatelli^{a,57} , A. Piccinelli^{a,b} , M. Presilla^{a,b} , A. Rossi^{a,b} , A. Santocchia^{a,b} , D. Spiga^a , T. Tedeschi^{a,b} 









































INFN Sezione di Pisa^a, Università di Pisa^b, Scuola Normale Superiore di Pisa^c, Pisa, Italy; Università di Siena^d, Siena, Italy

P. Azzurri^a , G. Bagliesi^a , R. Bhattacharya^a , L. Bianchini^{a,b} , T. Boccali^a , E. Bossini^a , D. Bruschini^{a,c} , R. Castaldi^a , M.A. Ciocci^{a,b} , M. Cipriani^{a,b} , V. D'Amante^{a,d} , R. Dell'Orso^a , S. Donato^a , A. Giassi^a , F. Ligabue^{a,c} , D. Matos Figueiredo^a , A. Messineo^{a,b} , M. Musich^{a,b} , F. Palla^a , S. Parolia^a , A. Rizzi^{a,b} , G. Rolandi^{a,c} , S. Roy Chowdhury^a , T. Sarkar^a , A. Scribano^a , P. Spagnolo^a , R. Tenchini^{a,b} , G. Tonelli^{a,b} , N. Turini^{a,d} , A. Venturi^a , P.G. Verdini^a 







INFN Sezione di Roma^a, Sapienza Università di Roma^b, Roma, Italy

P. Barria^a , M. Campana^{a,b} , F. Cavallari^a , L. Cunqueiro Mendez^{a,b} , D. Del Re^{a,b} , E. Di Marco^a , M. Diemoz^a , F. Errico^{a,b} , E. Longo^{a,b} , P. Meridiani^a , J. Mijuskovic^{a,b} , G. Organtini^{a,b} , F. Pandolfi^a , R. Paramatti^{a,b} , C. Quaranta^{a,b} , S. Rahatlou^{a,b} , C. Rovelli^a , F. Santanastasio^{a,b} , L. Soffi^a , R. Tramontano^{a,b} 













INFN Sezione di Torino^a, Università di Torino^b, Torino, Italy; Università del Piemonte Orientale^c, Novara, Italy

N. Amapane^{a,b} , R. Arcidiacono^{a,c} , S. Argiro^{a,b} , M. Arneodo^{a,c} , N. Bartosik^a , R. Bellan^{a,b} , A. Bellora^{a,b} , C. Biino^a , N. Cartiglia^a , M. Costa^{a,b} , R. Covarelli^{a,b} , N. Demaria^a , L. Finco^a , M. Grippo^{a,b} , B. Kiani^{a,b} , F. Legger^a , F. Luongo^{a,b} , C. Mariotti^a , S. Maselli^a , A. Mecca^{a,b} , E. Migliore^{a,b} , M. Monteno^a , R. Mulargia^a , M.M. Obertino^{a,b} , G. Ortona^a , L. Pacher^{a,b} , N. Pastrone^a , M. Pelliccioni^a , M. Ruspa^{a,c} , F. Siviero^{a,b} , V. Sola^{a,b} , A. Solano^{a,b} , D. Soldi^{a,b} , A. Staiano^a , C. Tarricone^{a,b} , M. Tornago^{a,b} , D. Trocino^a , G. Umoret^{a,b} , A. Vagnerini^{a,b} , E. Vlasov^{a,b} 

INFN Sezione di Trieste^a, Università di Trieste^b, Trieste, Italy

S. Belforte^a , V. Candelise^{a,b} , M. Casarsa^a , F. Cossutti^a , K. De Leo^{a,b} ,
G. Della Ricca^{a,b} 

Kyungpook National University, Daegu, Korea

S. Dogra , J. Hong , C. Huh , B. Kim , D.H. Kim , J. Kim, H. Lee, S.W. Lee ,
C.S. Moon , Y.D. Oh , S.I. Pak , M.S. Ryu , S. Sekmen , Y.C. Yang 




Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

G. Bak , P. Gwak , H. Kim , D.H. Moon 

Hanyang University, Seoul, Korea

E. Asilar , D. Kim , T.J. Kim , J.A. Merlin, J. Park 

Korea University, Seoul, Korea

S. Choi , S. Han, B. Hong , K. Lee, K.S. Lee , J. Park, S.K. Park, J. Yoo 

Kyung Hee University, Department of Physics, Seoul, Korea

J. Goh 









Sejong University, Seoul, Korea

H. S. Kim , Y. Kim, S. Lee



Seoul National University, Seoul, Korea

J. Almond, J.H. Bhyun, J. Choi , S. Jeon , W. Jun , J. Kim , J.S. Kim, S. Ko , H. Kwon ,
H. Lee , J. Lee , J. Lee , S. Lee, B.H. Oh , S.B. Oh , H. Seo , U.K. Yang, I. Yoon 

University of Seoul, Seoul, Korea

W. Jang , D.Y. Kang, Y. Kang , S. Kim , B. Ko, J.S.H. Lee , Y. Lee , I.C. Park , Y. Roh,
I.J. Watson , S. Yang 


Yonsei University, Department of Physics, Seoul, Korea

S. Ha , H.D. Yoo 

Sungkyunkwan University, Suwon, Korea

M. Choi , M.R. Kim , H. Lee, Y. Lee , I. Yu 


College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait

T. Beyrouthy, Y. Maghrbi 

Riga Technical University, Riga, Latvia

K. Dreimanis , A. Gaile , G. Pikurs, A. Potrebko , M. Seidel , V. Veckalns⁵⁸ 

University of Latvia (LU), Riga, Latvia

N.R. Strautnieks 






Vilnius University, Vilnius, Lithuania

M. Ambrozias , A. Juodagalvis , A. Rinkevicius , G. Tamulaitis 

National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia

N. Bin Norjoharuddeen , I. Yusuff⁵⁹ , Z. Zolkapli

Universidad de Sonora (UNISON), Hermosillo, Mexico

J.F. Benitez , A. Castaneda Hernandez , H.A. Encinas Acosta, L.G. Gallegos Maríñez,
M. León Coello , J.A. Murillo Quijada , A. Sehrawat , L. Valencia Palomo 

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

G. Ayala , H. Castilla-Valdez , E. De La Cruz-Burelo , I. Heredia-De La Cruz⁶⁰ ,
R. Lopez-Fernandez , C.A. Mondragon Herrera, A. Sánchez Hernández 


Universidad Iberoamericana, Mexico City, Mexico

C. Oropeza Barrera , M. Ramírez García 

Benemerita Universidad Autonoma de Puebla, Puebla, Mexico

I. Bautista , I. Pedraza , H.A. Salazar Ibarguen , C. Uribe Estrada 

University of Montenegro, Podgorica, Montenegro

I. Bubanja, N. Raicevic 

University of Canterbury, Christchurch, New Zealand

P.H. Butler 








National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan

A. Ahmad , M.I. Asghar, A. Awais , M.I.M. Awan, H.R. Hoorani , W.A. Khan 







AGH University of Science and Technology Faculty of Computer Science, Electronics and Telecommunications, Krakow, Poland

V. Avati, L. Grzanka , M. Malawski 

National Centre for Nuclear Research, Swierk, Poland

H. Bialkowska , M. Bluj , B. Boimska , M. Górski , M. Kazana , M. Szeper ,
P. Zalewski 



Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland

K. Bunkowski , K. Doroba , A. Kalinowski , M. Konecki , J. Krolikowski ,
A. Muhammad 



Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal

M. Araujo , D. Bastos , C. Beirão Da Cruz E Silva , A. Boletti , M. Bozzo , P. Faccioli ,
M. Gallinaro , J. Hollar , N. Leonardo , T. Niknejad , M. Pisano , J. Seixas , J. Varela 




























Faculty of Physics, University of Belgrade, Belgrade, Serbia

P. Adzic , P. Milenovic 


VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia

M. Dordevic , J. Milosevic , V. Rekovic

Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain

M. Aguilar-Benitez, J. Alcaraz Maestre , M. Barrio Luna, Cristina F. Bedoya , M. Cepeda ,
M. Cerrada , N. Colino , B. De La Cruz , A. Delgado Peris , D. Fernández Del Val ,
J.P. Fernández Ramos , J. Flix , M.C. Fouz , O. Gonzalez Lopez , S. Goy Lopez ,
J.M. Hernandez , M.I. Josa , J. León Holgado , D. Moran , C. M. Morcillo Perez ,
Á. Navarro Tobar , C. Perez Dengra , A. Pérez-Calero Yzquierdo , J. Puerta Pelayo ,
I. Redondo , D.D. Redondo Ferrero , L. Romero, S. Sánchez Navas , L. Urda Gómez ,
J. Vazquez Escobar , C. Willmott

Universidad Autónoma de Madrid, Madrid, Spain


















J.F. de Trocóniz 

Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain

B. Alvarez Gonzalez , J. Cuevas , J. Fernandez Menendez , S. Folgueras , I. Gonza-

lez Caballero , J.R. González Fernández , E. Palencia Cortezon , C. Ramón Álvarez , V. Rodríguez Bouza , A. Soto Rodríguez , A. Trapote , C. Vico Villalba , P. Vischia 

Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

S. Blanco Fernández , J.A. Brochero Cifuentes , I.J. Cabrillo , A. Calderon , J. Duarte Campderros , M. Fernandez , C. Fernandez Madrazo , G. Gomez , C. Lasasa García , C. Martinez Rivero , P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas , E. Navarrete Ramos, J. Piedra Gomez , C. Prieels, L. Scodellaro , I. Vila , J.M. Vizan Garcia 

University of Colombo, Colombo, Sri Lanka

M.K. Jayananda , B. Kailasapathy⁶¹ , D.U.J. Sonnadara , D.D.C. Wickramarathna 









University of Ruhuna, Department of Physics, Matara, Sri Lanka

W.G.D. Dharmaratna , K. Liyanage , N. Perera , N. Wickramage 





































CERN, European Organization for Nuclear Research, Geneva, Switzerland

D. Abbaneo , C. Amendola , E. Auffray , G. Auzinger , J. Baechler, D. Barney , A. Bermúdez Martínez , M. Bianco , B. Bilin , A.A. Bin Anuar , A. Bocci , E. Brondolin , C. Caillol , T. Camporesi , G. Cerminara , N. Chernyavskaya , D. d'Enterria , A. Dabrowski , A. David , A. De Roeck , M.M. Defranchis , M. Deile , M. Dobson , F. Fallavollita⁶², L. Forthomme , G. Franzoni , W. Funk , S. Giani, D. Gigi, K. Gill , F. Glege , L. Gouskos , M. Haranko , J. Hegeman , V. Innocente , T. James , P. Janot , J. Kieseler , S. Laurila , P. Lecoq , E. Leutgeb , C. Lourenço , B. Maier , L. Malgeri , M. Mannelli , A.C. Marini , F. Meijers , S. Mersi , E. Meschi , V. Milosevic , F. Moortgat , M. Mulders , S. Orfanelli, F. Pantaleo , M. Peruzzi , A. Petrilli , G. Petrucciani , A. Pfeiffer , M. Pierini , D. Piparo , H. Qu , D. Rabadý , G. Reales Gutiérrez, M. Rovere , H. Sakulin , S. Scarfi , M. Selvaggi , A. Sharma , K. Shchelina , P. Silva , P. Sphicas⁶³ , A.G. Stahl Leitner , A. Steen , S. Summers , D. Treille , P. Tropea , A. Tsiros, D. Walter , J. Wanczyk⁶⁴ , K.A. Wozniak⁶⁵ , P. Zehetner, P. Zexid , W.D. Zeuner
























Paul Scherrer Institut, Villigen, Switzerland

T. Bevilacqua⁶⁶ , L. Caminada⁶⁶ , A. Ebrahimi , W. Erdmann , R. Horisberger , Q. Ingram , H.C. Kaestli , D. Kotlinski , C. Lange , M. Missiroli⁶⁶ , L. Noehte⁶⁶ , T. Rohe 




ETH Zurich - Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

T.K. Aarrestad , K. Androsov⁶⁴ , M. Backhaus , A. Calandri , C. Cazzaniga , K. Datta , A. De Cosa , G. Dissertori , M. Dittmar, M. Donegà , F. Eble , M. Galli , K. Gedia , F. Glessgen , C. Grab , D. Hits , W. Lustermann , A.-M. Lyon , R.A. Manzoni , M. Marchegiani , L. Marchese , C. Martin Perez , A. Mascellani⁶⁴ , F. Nessi-Tedaldi , F. Pauss , V. Perovic , S. Pigazzini , M.G. Ratti , M. Reichmann , C. Reissel , T. Reitenspiess , B. Ristic , F. Riti , D. Ruini, D.A. Sanz Becerra , R. Seidita , J. Steggemann⁶⁴ , D. Valsecchi , R. Wallny 










Universität Zürich, Zurich, Switzerland

C. Amsler⁶⁷ , P. Bäertschi , C. Botta , D. Brzhechko, M.F. Canelli , K. Cormier , A. De Wit , R. Del Burgo, J.K. Heikkilä , M. Huwiler , W. Jin , A. Jofrehei , B. Kilminster , S. Leontsinis , S.P. Liechi , A. Macchiolo , P. Meiring , V.M. Mikuni , U. Molinatti , I. Neutelings , A. Reimers , P. Robmann, S. Sanchez Cruz , K. Schweiger , M. Senger , Y. Takahashi 




National Central University, Chung-Li, Taiwan

C. Adloff⁶⁸, C.M. Kuo, W. Lin, P.K. Rout , P.C. Tiwari⁴⁴ , S.S. Yu 




















National Taiwan University (NTU), Taipei, Taiwan

L. Ceard, Y. Chao , K.F. Chen , P.s. Chen, Z.g. Chen, W.-S. Hou , T.h. Hsu, Y.w. Kao, R. Khurana, G. Kole , Y.y. Li , R.-S. Lu , E. Paganis , A. Psallidas, X.f. Su, J. Thomas-Wilsker , H.y. Wu, E. Yazgan 

Chulalongkorn University, Faculty of Science, Department of Physics, Bangkok, Thailand

C. Asawatrangkuldee , N. Srimanobhas , V. Wachirapusanand 

Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

D. Agyel , F. Boran , Z.S. Demiroglu , F. Dolek , I. Dumanoglu⁶⁹ , E. Eskut , Y. Guler⁷⁰ , E. Gurpınar Guler⁷⁰ , C. Isik , O. Kara, A. Kayis Topaksu , U. Kiminsu , G. Onengut , K. Ozdemir⁷¹ , A. Polatoz , B. Tali⁷² , U.G. Tok , S. Turkcapar , E. Uslan , I.S. Zorbakir 

Middle East Technical University, Physics Department, Ankara, Turkey

K. Ocalan⁷³ , M. Yalvac⁷⁴ 














Bogazici University, Istanbul, Turkey

B. Akgun , I.O. Atakisi , E. Gülmez , M. Kaya⁷⁵ , O. Kaya⁷⁶ , S. Tekten⁷⁷ 

Istanbul Technical University, Istanbul, Turkey

A. Cakir , K. Cankocak⁶⁹ , Y. Komurcu , S. Sen⁷⁸ 

Istanbul University, Istanbul, Turkey

O. Aydilek , S. Cerci⁷² , V. Epshteyn , B. Hacıahinoglu , I. Hos⁷⁹ , B. Isildak⁸⁰ , B. Kaynak , S. Ozkorucuklu , O. Potok , H. Sert , C. Simsek , D. Sunar Cerci⁷² , C. Zorbilmez 
















Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine

A. Boyaryntsev , B. Grynyov 



















National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine

L. Levchuk 



















University of Bristol, Bristol, United Kingdom

D. Anthony , J.J. Brooke , A. Bundock , F. Bury , E. Clement , D. Cussans , H. Flacher , M. Glowacki, J. Goldstein , H.F. Heath , L. Kreczko , B. Krikler , S. Paramesvaran , S. Seif El Nasr-Storey, V.J. Smith , N. Stylianou⁸¹ , K. Walkingshaw Pass, R. White 

Rutherford Appleton Laboratory, Didcot, United Kingdom

A.H. Ball, K.W. Bell , A. Belyaev⁸² , C. Brew , R.M. Brown , D.J.A. Cockerill , C. Cooke , K.V. Ellis, K. Harder , S. Harper , M.-L. Holmberg⁸³ , Sh. Jain , J. Linacre , K. Manolopoulos, D.M. Newbold , E. Olaiya, D. Petyt , T. Reis , G. Salvi , T. Schuh, C.H. Shepherd-Themistocleous , I.R. Tomalin , T. Williams 

Imperial College, London, United Kingdom














R. Bainbridge , P. Bloch , C.E. Brown , O. Buchmuller, V. Cacchio, C.A. Carrillo Montoya , G.S. Chahal⁸⁴ , D. Colling , J.S. Dancu, P. Dauncey , G. Davies , J. Davies, M. Della Negra , S. Fayer, G. Fedi , G. Hall , M.H. Hassanshahi , A. Howard, G. Iles , M. Knight , J. Langford , L. Lyons , A.-M. Magnan , S. Malik, A. Martelli 

M. Mieskolainen , J. Nash⁸⁵ , M. Pesaresi, B.C. Radburn-Smith , A. Richards, A. Rose , C. Seez , R. Shukla , A. Tapper , K. Uchida , G.P. Uttley , L.H. Vage, T. Virdee³³ , M. Vojinovic , N. Wardle , D. Winterbottom 





Brunel University, Uxbridge, United Kingdom

K. Coldham, J.E. Cole , A. Khan, P. Kyberd , I.D. Reid 

Baylor University, Waco, Texas, USA

S. Abdullin , A. Brinkerhoff , B. Caraway , J. Dittmann , K. Hatakeyama , J. Hiltbrand , A.R. Kanuganti , B. McMaster , M. Saunders , S. Sawant , C. Sutantawibul , M. Toms⁸⁶ , J. Wilson 
















Catholic University of America, Washington, DC, USA

R. Bartek , A. Dominguez , C. Huerta Escamilla, A.E. Simsek , R. Uniyal , A.M. Vargas Hernandez 


















The University of Alabama, Tuscaloosa, Alabama, USA

R. Chudasama , S.I. Cooper , S.V. Gleyzer , C.U. Perez , P. Rumerio⁸⁷ , E. Usai , C. West , R. Yi 




















Boston University, Boston, Massachusetts, USA

A. Akpınar , A. Albert , D. Arcaro , C. Cosby , Z. Demiragli , C. Erice , E. Fontanesi , D. Gastler , J. Rohlf , K. Salyer , D. Sperka , D. Spitzbart , I. Suarez , A. Tsatsos , S. Yuan 













Brown University, Providence, Rhode Island, USA

G. Benelli , X. Coubez²⁸, D. Cutts , M. Hadley , U. Heintz , J.M. Hogan⁸⁸ , T. Kwon , G. Landsberg , K.T. Lau , D. Li , J. Luo , S. Mondal , M. Narain[†] , N. Pervan , S. Sagir⁸⁹ , F. Simpson , M. Stamenkovic , W.Y. Wong, X. Yan , W. Zhang





University of California, Davis, Davis, California, USA

S. Abbott , J. Bonilla , C. Brainerd , R. Breedon , M. Calderon De La Barca Sanchez , M. Chertok , M. Citron , J. Conway , P.T. Cox , R. Erbacher , G. Haza , F. Jensen , O. Kukral , G. Mocellin , M. Mulhearn , D. Pellett , B. Regnery , W. Wei, Y. Yao , F. Zhang 
























University of California, Los Angeles, California, USA

M. Bachtis , R. Cousins , A. Datta , J. Hauser , M. Ignatenko , M.A. Iqbal , T. Lam , E. Manca , W.A. Nash , D. Saltzberg , B. Stone , V. Valuev 

















University of California, Riverside, Riverside, California, USA

R. Clare , M. Gordon, G. Hanson , W. Si , S. Wimpenny[†] 











University of California, San Diego, La Jolla, California, USA

J.G. Branson , S. Cittolin , S. Cooperstein , D. Diaz , J. Duarte , R. Gerosa , L. Gianini , J. Guiang , R. Kansal , V. Krutelyov , R. Lee , J. Letts , M. Masciovecchio , F. Mokhtar , M. Pieri , M. Quinnan , B.V. Sathia Narayanan , V. Sharma , M. Tadel , E. Vourliotis , F. Würthwein , Y. Xiang , A. Yagil 















University of California, Santa Barbara - Department of Physics, Santa Barbara, California, USA

A. Barzdukas , L. Brennan, C. Campagnari , G. Collura , A. Dorsett , J. Incandela , M. Kilpatrick , J. Kim , A.J. Li , P. Masterson , H. Mei , M. Oshiro , J. Richman , U. Sarica , R. Schmitz , F. Setti , J. Sheplock , D. Stuart , S. Wang 














California Institute of Technology, Pasadena, California, USA

A. Bornheim , O. Cerri, A. Latorre, J.M. Lawhorn , J. Mao , H.B. Newman , T. Q. Nguyen , M. Spiropulu , J.R. Vlimant , C. Wang , S. Xie , R.Y. Zhu 

Carnegie Mellon University, Pittsburgh, Pennsylvania, USA

J. Alison , S. An , M.B. Andrews , P. Bryant , V. Dutta , T. Ferguson , A. Harilal , C. Liu , T. Mudholkar , S. Murthy , M. Paulini , A. Roberts , A. Sanchez , W. Terrill 








































































University of Colorado Boulder, Boulder, Colorado, USA

J.P. Cumalat , W.T. Ford , A. Hassani , G. Karathanasis , E. MacDonald, N. Manganello , F. Marini , A. Perloff , C. Savard , N. Schonbeck , K. Stenson , K.A. Ulmer , S.R. Wagner , N. Zipper 




















Cornell University, Ithaca, New York, USA

J. Alexander , S. Bright-Thonney , X. Chen , D.J. Cranshaw , J. Fan , X. Fan , D. Gadkari , S. Hogan , J. Monroy , J.R. Patterson , J. Reichert , M. Reid , A. Ryd , J. Thom , P. Wittich , R. Zou 















Fermi National Accelerator Laboratory, Batavia, Illinois, USA

M. Albrow , M. Alyari , O. Amram , G. Apollinari , A. Apresyan , L.A.T. Bauerdick , D. Berry , J. Berryhill , P.C. Bhat , K. Burkett , J.N. Butler , A. Canepa , G.B. Cerati , H.W.K. Cheung , F. Chlebana , G. Cummings , J. Dickinson , I. Dutta , V.D. Elvira , Y. Feng , J. Freeman , A. Gandrakota , Z. Gecse , L. Gray , D. Green, S. Grünendahl , D. Guerrero , O. Gutsche , R.M. Harris , R. Heller , T.C. Herwig , J. Hirschauer , L. Horyn , B. Jayatilaka , S. Jindariani , M. Johnson , U. Joshi , T. Klijnsma , B. Klima , K.H.M. Kwok , S. Lammel , D. Lincoln , R. Lipton , T. Liu , C. Madrid , K. Maeshima , C. Mantilla , D. Mason , P. McBride , P. Merkel , S. Mrenna , S. Nahn , J. Ngadiuba , D. Noonan , V. Papadimitriou , N. Pastika , K. Pedro , C. Pena⁹⁰ , F. Ravera , A. Reinsvold Hall⁹¹ , L. Ristori , E. Sexton-Kennedy , N. Smith , A. Soha , L. Spiegel , S. Stoynev , L. Taylor , S. Tkaczyk , N.V. Tran , L. Uplegger , E.W. Vaandering , I. Zoi 






University of Florida, Gainesville, Florida, USA

C. Aruta , P. Avery , D. Bourilkov , L. Cadamuro , P. Chang , V. Cherepanov , R.D. Field, E. Koenig , M. Kolosova , J. Konigsberg , A. Korytov , K.H. Lo, K. Matchev , N. Menendez , G. Mitselmakher , A. Muthirakalayil Madhu , N. Rawal , D. Rosenzweig , S. Rosenzweig , K. Shi , J. Wang 























Florida State University, Tallahassee, Florida, USA

T. Adams , A. Al Kadhimi , A. Askew , N. Bower , R. Habibullah , V. Hagopian , R. Hashmi , R.S. Kim , S. Kim , T. Kolberg , G. Martinez, H. Prosper , P.R. Prova, O. Viazlo , M. Wulansatiti , R. Yohay , J. Zhang












Florida Institute of Technology, Melbourne, Florida, USA

B. Alsufyani, M.M. Baarmand , S. Butalla , T. Elkafrawy¹⁹ , M. Hohlmann , R. Kumar Verma , M. Rahmani













University of Illinois at Chicago (UIC), Chicago, Illinois, USA

M.R. Adams , C. Bennett, R. Cavanaugh , S. Dittmer , R. Escobar Franco , O. Evdokimov , C.E. Gerber , D.J. Hofman , J.h. Lee , D. S. Lemos , A.H. Merrit , C. Mills , S. Nanda , G. Oh , B. Ozek , D. Pilipovic , T. Roy , S. Rudrabhatla , M.B. Tonjes , N. Varelas , X. Wang , Z. Ye , J. Yoo 
























The University of Iowa, Iowa City, Iowa, USA

M. Alhusseini , D. Blend, K. Dilsiz⁹² , L. Emediato , G. Karaman , O.K. Köseyan , J.-P. Merlo, A. Mestvirishvili⁹³ , J. Nachtman , O. Neogi, H. Ogul⁹⁴ , Y. Onel , A. Penzo , C. Snyder, E. Tiras⁹⁵ 









Johns Hopkins University, Baltimore, Maryland, USA

B. Blumenfeld , L. Corcodilos , J. Davis , A.V. Gritsan , L. Kang , S. Kyriacou , P. Maksimovic , M. Roguljic , J. Roskes , S. Sekhar , M. Swartz , T.Á. Vámi 

The University of Kansas, Lawrence, Kansas, USA

A. Abreu , L.F. Alcerro Alcerro , J. Anguiano , P. Baringer , A. Bean , Z. Flowers , D. Grove, J. King , G. Krintiras , M. Lazarovits , C. Le Mahieu , C. Lindsey, J. Marquez , N. Minafra , M. Murray , M. Nickel , M. Pitt , S. Popescu⁹⁶ , C. Rogan , C. Royon , R. Salvatico , S. Sanders , C. Smith , Q. Wang , G. Wilson 














Kansas State University, Manhattan, Kansas, USA

B. Allmond , A. Ivanov , K. Kaadze , A. Kalogeropoulos , D. Kim, Y. Maravin , K. Nam, J. Natoli , D. Roy , G. Sorrentino 

























Lawrence Livermore National Laboratory, Livermore, California, USA

F. Rebasoo , D. Wright 















University of Maryland, College Park, Maryland, USA

E. Adams , A. Baden , O. Baron, A. Belloni , A. Bethani , Y.M. Chen , S.C. Eno , N.J. Hadley , S. Jabeen , R.G. Kellogg , T. Koeth , Y. Lai , S. Lascio , A.C. Mignerey , S. Nabili , C. Palmer , C. Papageorgakis , M.M. Paranjpe, L. Wang , K. Wong 

Massachusetts Institute of Technology, Cambridge, Massachusetts, USA

J. Bendavid , W. Busza , I.A. Cali , Y. Chen , M. D'Alfonso , J. Eysermans , C. Freer , G. Gomez-Ceballos , M. Goncharov, P. Harris, D. Hoang, D. Kovalskiy , J. Krupa , L. Lavezzo , Y.-J. Lee , K. Long , C. Mironov , C. Paus , D. Rankin , C. Roland , G. Roland , S. Rothman , Z. Shi , G.S.F. Stephans , J. Wang, Z. Wang , B. Wyslouch , T. J. Yang 















University of Minnesota, Minneapolis, Minnesota, USA

B. Crossman , B.M. Joshi , C. Kapsiak , M. Krohn , D. Mahon , J. Mans , B. Marzocchi , S. Pandey , M. Revering , R. Rusack , R. Saradhy , N. Schroeder , N. Strobbe , M.A. Wadud 

University of Mississippi, Oxford, Mississippi, USA

L.M. Cremaldi 

















University of Nebraska-Lincoln, Lincoln, Nebraska, USA

K. Bloom , M. Bryson, D.R. Claes , C. Fangmeier , F. Golf , J. Hossain , C. Joo , I. Kravchenko , I. Reed , J.E. Siado , G.R. Snow[†], W. Tabb , A. Wightman , F. Yan , D. Yu , A.G. Zecchinelli 









State University of New York at Buffalo, Buffalo, New York, USA

G. Agarwal , H. Bandyopadhyay , L. Hay , I. Iashvili , A. Kharchilava , C. McLean , M. Morris , D. Nguyen , J. Pekkanen , S. Rappoccio , H. Rejeb Sfar, A. Williams 

























Northeastern University, Boston, Massachusetts, USA

G. Alverson , E. Barberis , Y. Haddad , Y. Han , A. Krishna , J. Li , M. Lu , G. Madigan , D.M. Morse , V. Nguyen , T. Orimoto , A. Parker , L. Skinnari , A. Tishelman-Charny , B. Wang , D. Wood 










Northwestern University, Evanston, Illinois, USA

S. Bhattacharya , J. Bueghly, Z. Chen , K.A. Hahn , Y. Liu , Y. Miao , D.G. Monk , M.H. Schmitt , A. Taliercio , M. Velasco



















University of Notre Dame, Notre Dame, Indiana, USA

R. Band , R. Bucci, S. Castells , M. Cremonesi, A. Das , R. Goldouzian , M. Hildreth , K.W. Ho , K. Hurtado Anampa , C. Jessop , K. Lannon , J. Lawrence , N. Loukas , L. Lutton , J. Mariano, N. Marinelli, I. Mcalister, T. McCauley , C. Mcgrady , K. Mohrman , C. Moore , Y. Musienko¹⁵ , H. Nelson , M. Osherson , R. Ruchti , A. Townsend , M. Wayne , H. Yockey, M. Zarucki , L. Zygala 

The Ohio State University, Columbus, Ohio, USA

A. Basnet , B. Bylsma, M. Carrigan , L.S. Durkin , C. Hill , M. Joyce , A. Lesauvage , M. Nunez Ornelas , K. Wei, B.L. Winer , B. R. Yates 






















Princeton University, Princeton, New Jersey, USA

F.M. Addesa , H. Bouchamaoui , P. Das , G. Dezoort , P. Elmer , A. Frankenthal , B. Greenberg , N. Haubrich , S. Higginbotham , G. Kopp , S. Kwan , D. Lange , A. Loeliger , D. Marlow , I. Ojalvo , J. Olsen , D. Stickland , C. Tully 




University of Puerto Rico, Mayaguez, Puerto Rico, USA

S. Malik 















Purdue University, West Lafayette, Indiana, USA

A.S. Bakshi , V.E. Barnes , S. Chandra , R. Chawla , S. Das , A. Gu , L. Gutay, M. Jones , A.W. Jung , D. Kondratyev , A.M. Koshy, M. Liu , G. Negro , N. Neumeister , G. Paspalaki , S. Piperov , A. Purohit , J.F. Schulte , M. Stojanovic , J. Thieman , A. K. Viridi , F. Wang , W. Xie 












Purdue University Northwest, Hammond, Indiana, USA

J. Dolen , N. Parashar , A. Pathak 


Rice University, Houston, Texas, USA

D. Acosta , A. Baty , T. Carnahan , S. Dildick , K.M. Ecklund , P.J. Fernández Manteca , S. Freed, P. Gardner, F.J.M. Geurts , A. Kumar , W. Li , O. Miguel Colin , B.P. Padley , R. Redjimi, J. Rotter , E. Yigitbasi , Y. Zhang 

















University of Rochester, Rochester, New York, USA

A. Bodek , P. de Barbaro , R. Demina , J.L. Dulemba , C. Fallon, A. Garcia-Bellido , O. Hindrichs , A. Khukhunaishvili , P. Parygin⁸⁶ , E. Popova⁸⁶ , R. Taus , G.P. Van Onsem 









The Rockefeller University, New York, New York, USA

K. Goulianos 















Rutgers, The State University of New Jersey, Piscataway, New Jersey, USA

B. Chiarito, J.P. Chou , Y. Gershtein , E. Halkiadakis , A. Hart , M. Heindl , D. Jaroslowski , O. Karacheban³¹ , I. Laflotte , A. Lath , R. Montalvo, K. Nash, H. Routray , S. Salur , S. Schnetzer, S. Somalwar , R. Stone , S.A. Thayil , S. Thomas, J. Vora , H. Wang 












University of Tennessee, Knoxville, Tennessee, USA

H. Acharya, D. Ally , A.G. Delannoy , S. Fiorendi , T. Holmes , N. Karunarathna , L. Lee , E. Nibigira , S. Spanier 

Texas A&M University, College Station, Texas, USA

D. Aebi , M. Ahmad , O. Bouhali⁹⁷ , M. Dalchenko , R. Eusebi , J. Gilmore , T. Huang , T. Kamon⁹⁸ , H. Kim , S. Luo , S. Malhotra, R. Mueller , D. Overton , D. Rathjens , A. Safonov 









Texas Tech University, Lubbock, Texas, USA

N. Akchurin , J. Damgov , V. Hegde , A. Hussain , Y. Kazhykarim, K. Lamichhane , S.W. Lee , A. Mankel , T. Mengke, S. Muthumuni , T. Peltola , I. Volobouev , A. Whitbeck 

Vanderbilt University, Nashville, Tennessee, USA

E. Appelt , S. Greene, A. Gurrola , W. Johns , R. Kunnawalkam Elayavalli , A. Melo , F. Romeo , P. Sheldon , S. Tuo , J. Velkovska , J. Viinikainen 






















University of Virginia, Charlottesville, Virginia, USA

B. Cardwell , B. Cox , J. Hakala , R. Hirosky , A. Ledovskoy , A. Li , C. Neu , C.E. Perez Lara 

Wayne State University, Detroit, Michigan, USA

P.E. Karchin 

University of Wisconsin - Madison, Madison, Wisconsin, USA

A. Aravind, S. Banerjee , K. Black , T. Bose , S. Dasu , I. De Bruyn , P. Everaerts , C. Galloni, H. He , M. Herndon , A. Herve , C.K. Koraka , A. Lanaro, R. Loveless , J. Madhusudanan Sreekala , A. Mallampalli , A. Mohammadi , S. Mondal, G. Parida , D. Pinna, A. Savin, V. Shang , V. Sharma , W.H. Smith , D. Teague, H.F. Tsoi , W. Vetens , A. Warden 

Authors affiliated with an institute or an international laboratory covered by a cooperation agreement with CERN

S. Afanasiev , V. Andreev , Yu. Andreev , T. Aushev , M. Azarkin , A. Babaev , A. Belyaev , V. Blinov⁹⁹, E. Boos , V. Borshch , D. Budkouski , V. Chekhovsky, R. Chistov⁹⁹ , M. Danilov⁹⁹ , A. Dermenev , T. Dimova⁹⁹ , D. Druzhkin¹⁰⁰ , M. Dubinin⁹⁰ , L. Dudko , A. Ershov , G. Gavrilo , V. Gavrilo , S. Gninenko , V. Golovtsov , N. Golubev , I. Golutvin , I. Gorbunov , A. Gribushin , Y. Ivanov , V. Kachanov , L. Kardapoltsev⁹⁹ , V. Karjavine , A. Karneyeu , V. Kim⁹⁹ , M. Kirakosyan, D. Kirpichnikov , M. Kirsanov , V. Klyukhin , O. Kodolova¹⁰¹ , D. Konstantinov , V. Korenkov , V. Korotkikh, A. Kozyrev⁹⁹ , N. Krasnikov , A. Lanev , P. Levchenko¹⁰² , N. Lychkovskaya , V. Makarenko , A. Malakhov , V. Matveev⁹⁹ , V. Murzin , A. Nikitenko^{103,101} , S. Obraztsov , V. Oreshkin , A. Oskin, V. Palichik , V. Perelygin , N. Petrov, S. Petrushanko , S. Polikarpov⁹⁹ , V. Popov, O. Radchenko⁹⁹ , M. Savina , V. Savrin , V. Shalaev , S. Shmatov , S. Shulha , Y. Skovpen⁹⁹ , S. Slabospitskii , V. Smirnov , A. Snigirev , D. Sosnov , V. Sulimov , E. Tcherniaev , A. Terkulov , O. Teryaev , I. Tlisova , A. Toropin , L. Uvarov , A. Uzunian , A. Vorobyev[†], N. Voytishin , B.S. Yuldashev¹⁰⁴, A. Zarubin , I. Zhizhin , A. Zhokin 

†: Deceased

¹Also at Yerevan State University, Yerevan, Armenia

²Also at TU Wien, Vienna, Austria

³Also at Institute of Basic and Applied Sciences, Faculty of Engineering, Arab Academy for Science, Technology and Maritime Transport, Alexandria, Egypt

⁴Also at Ghent University, Ghent, Belgium

-
- ⁵Also at Universidade Estadual de Campinas, Campinas, Brazil
- ⁶Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil
- ⁷Also at UFMS, Nova Andradina, Brazil
- ⁸Also at Henan Normal University, Xinxiang, China
- ⁹Now at Nanjing Normal University Department of Physics, Nanjing, China
- ¹⁰Also at Sun Yat-Sen University, Guangzhou, China
- ¹¹Now at The University of Iowa, Iowa City, Iowa, USA
- ¹²Also at University of Chinese Academy of Sciences, Beijing, China
- ¹³Also at University of Chinese Academy of Sciences, Beijing, China
- ¹⁴Also at Université Libre de Bruxelles, Bruxelles, Belgium
- ¹⁵Also at an institute or an international laboratory covered by a cooperation agreement with CERN
- ¹⁶Also at Helwan University, Cairo, Egypt
- ¹⁷Now at Zewail City of Science and Technology, Zewail, Egypt
- ¹⁸Also at British University in Egypt, Cairo, Egypt
- ¹⁹Now at Ain Shams University, Cairo, Egypt
- ²⁰Also at Birla Institute of Technology, Mesra, Mesra, India
- ²¹Also at Purdue University, West Lafayette, Indiana, USA
- ²²Also at Université de Haute Alsace, Mulhouse, France
- ²³Also at Department of Physics, Tsinghua University, Beijing, China
- ²⁴Also at Tbilisi State University, Tbilisi, Georgia
- ²⁵Also at The University of the State of Amazonas, Manaus, Brazil
- ²⁶Also at Erzincan Binali Yildirim University, Erzincan, Turkey
- ²⁷Also at University of Hamburg, Hamburg, Germany
- ²⁸Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany
- ²⁹Also at Isfahan University of Technology, Isfahan, Iran
- ³⁰Also at Bergische University Wuppertal (BUW), Wuppertal, Germany
- ³¹Also at Brandenburg University of Technology, Cottbus, Germany
- ³²Also at Forschungszentrum Jülich, Juelich, Germany
- ³³Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland
- ³⁴Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt
- ³⁵Also at Wigner Research Centre for Physics, Budapest, Hungary
- ³⁶Also at Institute of Physics, University of Debrecen, Debrecen, Hungary
- ³⁷Also at Institute of Nuclear Research ATOMKI, Debrecen, Hungary
- ³⁸Now at Universitatea Babeş-Bolyai - Facultatea de Fizica, Cluj-Napoca, Romania
- ³⁹Also at Faculty of Informatics, University of Debrecen, Debrecen, Hungary
- ⁴⁰Also at Punjab Agricultural University, Ludhiana, India
- ⁴¹Also at UPES - University of Petroleum and Energy Studies, Dehradun, India
- ⁴²Also at University of Visva-Bharati, Santiniketan, India
- ⁴³Also at University of Hyderabad, Hyderabad, India
- ⁴⁴Also at Indian Institute of Science (IISc), Bangalore, India
- ⁴⁵Also at IIT Bhubaneswar, Bhubaneswar, India
- ⁴⁶Also at Institute of Physics, Bhubaneswar, India
- ⁴⁷Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany
- ⁴⁸Also at Department of Physics, Isfahan University of Technology, Isfahan, Iran
- ⁴⁹Also at Sharif University of Technology, Tehran, Iran
- ⁵⁰Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran
- ⁵¹Also at Italian National Agency for New Technologies, Energy and Sustainable Economic

Development, Bologna, Italy

⁵²Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy

⁵³Also at Università degli Studi Guglielmo Marconi, Roma, Italy

⁵⁴Also at Scuola Superiore Meridionale, Università di Napoli 'Federico II', Napoli, Italy

⁵⁵Also at Fermi National Accelerator Laboratory, Batavia, Illinois, USA

⁵⁶Also at Università di Napoli 'Federico II', Napoli, Italy

⁵⁷Also at Consiglio Nazionale delle Ricerche - Istituto Officina dei Materiali, Perugia, Italy

⁵⁸Also at Riga Technical University, Riga, Latvia

⁵⁹Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia

⁶⁰Also at Consejo Nacional de Ciencia y Tecnología, Mexico City, Mexico

⁶¹Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka

⁶²Also at INFN Sezione di Pavia, Università di Pavia, Pavia, Italy

⁶³Also at National and Kapodistrian University of Athens, Athens, Greece

⁶⁴Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland

⁶⁵Also at University of Vienna Faculty of Computer Science, Vienna, Austria

⁶⁶Also at Universität Zürich, Zurich, Switzerland

⁶⁷Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria

⁶⁸Also at Laboratoire d'Annecy-le-Vieux de Physique des Particules, IN2P3-CNRS, Annecy-le-Vieux, France

⁶⁹Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey

⁷⁰Also at Konya Technical University, Konya, Turkey

⁷¹Also at Izmir Bakircay University, Izmir, Turkey

⁷²Also at Adiyaman University, Adiyaman, Turkey

⁷³Also at Necmettin Erbakan University, Konya, Turkey

⁷⁴Also at Bozok Universitetesi Rektörlüğü, Yozgat, Turkey

⁷⁵Also at Marmara University, Istanbul, Turkey

⁷⁶Also at Milli Savunma University, Istanbul, Turkey

⁷⁷Also at Kafkas University, Kars, Turkey

⁷⁸Also at Hacettepe University, Ankara, Turkey

⁷⁹Also at Istanbul University - Cerrahpasa, Faculty of Engineering, Istanbul, Turkey

⁸⁰Also at Yildiz Technical University, Istanbul, Turkey

⁸¹Also at Vrije Universiteit Brussel, Brussel, Belgium

⁸²Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom

⁸³Also at University of Bristol, Bristol, United Kingdom

⁸⁴Also at IPPP Durham University, Durham, United Kingdom

⁸⁵Also at Monash University, Faculty of Science, Clayton, Australia

⁸⁶Now at an institute or an international laboratory covered by a cooperation agreement with CERN

⁸⁷Also at Università di Torino, Torino, Italy

⁸⁸Also at Bethel University, St. Paul, Minnesota, USA

⁸⁹Also at Karamanoğlu Mehmetbey University, Karaman, Turkey

⁹⁰Also at California Institute of Technology, Pasadena, California, USA

⁹¹Also at United States Naval Academy, Annapolis, Maryland, USA

⁹²Also at Bingol University, Bingol, Turkey

⁹³Also at Georgian Technical University, Tbilisi, Georgia

⁹⁴Also at Sinop University, Sinop, Turkey

⁹⁵Also at Erciyes University, Kayseri, Turkey

⁹⁶Also at Horia Hulubei National Institute of Physics and Nuclear Engineering (IFIN-HH), Bucharest, Romania

⁹⁷Also at Texas A&M University at Qatar, Doha, Qatar

⁹⁸Also at Kyungpook National University, Daegu, Korea

⁹⁹Also at another institute or international laboratory covered by a cooperation agreement with CERN

¹⁰⁰Also at Universiteit Antwerpen, Antwerpen, Belgium

¹⁰¹Also at Yerevan Physics Institute, Yerevan, Armenia

¹⁰²Also at Northeastern University, Boston, Massachusetts, USA

¹⁰³Also at Imperial College, London, United Kingdom

¹⁰⁴Also at Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan