



# Search for flavour-changing neutral-current couplings between the top quark and the Higgs boson in multi-lepton final states in 13 TeV $pp$ collisions with the ATLAS detector

The ATLAS Collaboration

A search is presented for flavour-changing neutral-current interactions involving the top quark, the Higgs boson and an up-type quark ( $q = u, c$ ) with the ATLAS detector at the Large Hadron Collider. The analysis considers leptonic decays of the top quark along with Higgs boson decays into two  $W$  bosons, two  $Z$  bosons or a  $\tau^+\tau^-$  pair. It focuses on final states containing either two leptons (electrons or muons) of the same charge or three leptons. The considered processes are  $t\bar{t}$  and  $Ht$  production. For the  $t\bar{t}$  production, one top quark decays via  $t \rightarrow Hq$ . The proton–proton collision data set analysed amounts to  $140 \text{ fb}^{-1}$  at  $\sqrt{s} = 13 \text{ TeV}$ . No significant excess beyond Standard Model expectations is observed and upper limits are set on the  $t \rightarrow Hq$  branching ratios at 95% confidence level, amounting to observed (expected) limits of  $\mathcal{B}(t \rightarrow Hu) < 2.8 (3.0) \times 10^{-4}$  and  $\mathcal{B}(t \rightarrow Hc) < 3.3 (3.8) \times 10^{-4}$ . Combining this search with other searches for  $tHq$  flavour-changing neutral-current interactions previously conducted by ATLAS, considering  $H \rightarrow b\bar{b}$  and  $H \rightarrow \gamma\gamma$  decays, as well as  $H \rightarrow \tau^+\tau^-$  decays with one or two hadronically decaying  $\tau$ -leptons, yields observed (expected) upper limits on the branching ratios of  $\mathcal{B}(t \rightarrow Hu) < 2.6 (1.8) \times 10^{-4}$  and  $\mathcal{B}(t \rightarrow Hc) < 3.4 (2.3) \times 10^{-4}$ .

# Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>The ATLAS detector</b>	<b>5</b>
<b>3</b>	<b>Samples of data and simulated events</b>	<b>6</b>
3.1	Simulation of $tHq$ FCNC signal samples	7
3.2	Simulation of background processes	7
<b>4</b>	<b>Object reconstruction and event preselection</b>	<b>9</b>
<b>5</b>	<b>Background estimate</b>	<b>11</b>
5.1	Non-prompt lepton background	12
5.2	Charge misidentification	12
5.3	Prompt-lepton background	13
<b>6</b>	<b>Event categorisation</b>	<b>13</b>
6.1	Definition of signal regions	13
6.2	Control regions for non-prompt-lepton backgrounds	15
6.3	Control regions for $t\bar{t}W$ and $t\bar{t}Z$ backgrounds	15
<b>7</b>	<b>Separation of signal and background</b>	<b>16</b>
7.1	Event reconstruction in signal regions	16
7.2	Training of feed-forward neural networks	18
<b>8</b>	<b>Systematic uncertainties</b>	<b>19</b>
8.1	Experimental uncertainties	19
8.2	Modelling uncertainties	20
<b>9</b>	<b>Statistical analysis</b>	<b>22</b>
<b>10</b>	<b>Results</b>	<b>23</b>
10.1	Cross-checks on the modelling of the neural network input variables	23
10.2	Full fit to data	23
<b>11</b>	<b>Combination of results with other searches</b>	<b>25</b>
<b>12</b>	<b>Conclusion</b>	<b>31</b>
	<b>Appendix</b>	<b>33</b>

## 1 Introduction

Following the discovery of the Higgs boson at the Large Hadron Collider (LHC) by the ATLAS and CMS experiments in 2012 [1, 2], several measurements have probed properties of the particle. In addition to

Higgs boson couplings to fermions and gauge bosons, which agree well with the Standard Model (SM) predictions [3, 4], further interactions can be investigated to search for evidence of possible new physics in the Higgs sector. A possibility is that the Higgs boson has an interaction involving two up-type quarks of different generations, denoted by  $tHq$  where  $q = (u, c)$ , leading to  $gq \rightarrow Ht$  production and  $t \rightarrow Hq$  decay. These are flavour-changing neutral-current (FCNC) interactions, forbidden at tree-level in the SM and suppressed at higher orders due to the Glashow-Iliopoulos-Maiani (GIM) mechanism [5]. The SM predictions for the  $t \rightarrow Hq$  branching ratios  $\mathcal{B}(t \rightarrow Hq)$  are very small, around  $10^{-15}$  [6–10], and beyond the sensitivity of the LHC. However, in the framework of new physics models such as two-Higgs-doublet models (2HDMs) [11], the minimal supersymmetric SM (MSSM) [12–15], supersymmetric models with R-parity violation [16], quark-singlet models [17], and warped extra dimensions [18], the branching ratios of these processes are modified making them large enough to be measured at the LHC. For example, branching ratios of up to  $10^{-3}$  are possible in 2HDMs without explicit flavour symmetry, since there are no symmetries explicitly forbidding tree-level FCNC interactions [19–26] in this model. The FCNC interaction can also mediate the production of a Higgs boson with a single top quark ( $pp \rightarrow tH$ ) [27]. The SM prediction of the  $pp \rightarrow tH$  cross-section is  $74.3_{-0.3}^{+0.4}$  fb at  $\sqrt{s} = 13$  TeV [28].

The FCNC interactions are introduced using an effective field theory (EFT) framework, which is used for indirect searches for new physics [29]. Here the SM is regarded as a low-energy approximation of an ultraviolet complete theory containing new particles, whose masses are characterised by an energy scale  $\Lambda = 1$  TeV. The new physics contributions are parameterised in terms of operators with mass dimension greater than four containing only the SM fields, scaled by dimensionless Wilson coefficients and inverse powers of  $\Lambda$ . In the case where only the  $tHu$  and  $tHc$  interactions are considered, the relevant operators are

$$O_{u\phi}^{qt} = \left( \phi^\dagger \phi - \frac{v^2}{2} \right) (\bar{q}_L t_R) \tilde{\phi}, \quad O_{u\phi}^{tq} = \left( \phi^\dagger \phi - \frac{v^2}{2} \right) (\bar{t}_L q_R) \tilde{\phi}, \quad (1)$$

where  $q$  corresponds to an up or charm quark, depending on the FCNC coupling. The index  $u$  is the coupling to any up-type quark,  $t$  is the top-quark,  $\phi$  denotes the Higgs boson field with  $v$  corresponding to the absolute value of its vacuum expectation value. The two left-handed quark doublet fields are  $\bar{q}_L$  and  $\bar{t}_L$ , with  $q_R$  and  $t_R$  being the corresponding right-handed singlets. The operators are scaled with Wilson coefficients  $C_{u\phi}^{qt}$  and  $C_{u\phi}^{tq}$ , and  $1/\Lambda^2$  to give the relevant Lagrangian:

$$\mathcal{L}_{\text{EFT}} = \mathcal{L}_{\text{SM}} + \sum_{q=u,c} \left[ \frac{C_{u\phi}^{qt}}{\Lambda^2} O_{u\phi}^{qt} + \frac{C_{u\phi}^{tq}}{\Lambda^2} O_{u\phi}^{tq} \right]. \quad (2)$$

In the case of  $t\bar{t}$  production with a FCNC  $t \rightarrow Hq$  decay, no kinematic differences between  $O_{u\phi}^{qt}$  and  $O_{u\phi}^{tq}$  are expected because the top quarks are produced unpolarised and the Higgs boson is a scalar particle. For  $t \rightarrow Hq$  production, comparisons of simulations using either  $O_{u\phi}^{qt}$  or  $O_{u\phi}^{tq}$  have shown that differences between production rates and kinematic distributions are negligible in the phase space considered. Thus, the mean of both the couplings  $C_{u\phi}^{qt,tq} = \frac{C_{u\phi}^{qt} + C_{u\phi}^{tq}}{2}$  is considered.

Both the ATLAS and CMS Collaborations have undertaken various searches for  $tHq$  FCNC processes, split by Higgs boson final state [30–38]. In this paper, a search is conducted using final states containing either two




Figure 1: Example Feynman diagrams of the  $t\bar{t}(t \rightarrow Hq)$  decay signal process resulting in (a) the  $2\ell$ SS final state and (b) the  $3\ell$  final state. The FCNC vertex is highlighted in red.

leptons (electrons or muons) of the same charge ( $2\ell$ SS,  $\ell = e, \mu$ ) or three leptons, exactly two of which have the same charge ( $3\ell$ ,  $\ell = e, \mu$ ). The ATLAS Collaboration searched for  $tHq$  FCNC couplings in  $2\ell$ SS and  $3\ell$  final states using a partial Run 2 data sample of  $36.1 \text{ fb}^{-1}$  [32], resulting in observed (expected) 95% CL upper limits of  $\mathcal{B}(t \rightarrow Hu) < 19(15) \times 10^{-4}$  and  $\mathcal{B}(t \rightarrow Hc) < 16(15) \times 10^{-4}$ . The strongest limits set by the ATLAS Collaboration come from a combination of searches in the  $H \rightarrow b\bar{b}$ ,  $H \rightarrow \gamma\gamma$  and  $H \rightarrow \tau^+\tau^-$  channels using the full Run 2 data sample [35]. The obtained limits are  $\mathcal{B}(t \rightarrow Hu) < 4.0(2.4) \times 10^{-4}$  and  $\mathcal{B}(t \rightarrow Hc) < 5.8(3.0) \times 10^{-4}$ . For the  $tHc$  coupling, these are the most stringent limits published to date. However, for the  $tHu$  coupling the strongest limits are set by a search conducted with the CMS detector in the  $H \rightarrow \gamma\gamma$  channel [38], amounting to  $\mathcal{B}(t \rightarrow Hu) < 1.9(3.1) \times 10^{-4}$ .

The dominant Higgs-boson decay mode resulting in  $2\ell$ SS and  $3\ell$  final states across production and decay processes is  $H \rightarrow WW^*$ , with  $WW^* \rightarrow \ell\nu jj$  or  $WW^* \rightarrow \ell\nu\ell\nu$  for the  $2\ell$ SS and  $3\ell$  final state respectively. The corresponding Feynman diagrams are shown in Figures 1 and 2. Other Higgs-boson decay modes such as  $H \rightarrow ZZ^*$  and  $H \rightarrow \tau^+\tau^-$  can also meet the selection criteria. Based on the SM branching ratios of the Higgs boson and the top quark, approximately 73% of all  $2\ell$ SS signal events are expected from the  $H \rightarrow WW^*$  decay mode, while the rest originate from leptonic  $H \rightarrow \tau^+\tau^-$  decays. Less than 1% of events are attributable to the  $H \rightarrow ZZ^*$  decay mode. In the  $3\ell$  final state, the latter contributes more significantly, being responsible for 14% of all events. The majority, accounting for 54% of the  $3\ell$  signal events, originate from  $H \rightarrow WW^*$  decays, while the remaining 32% arise from  $H \rightarrow \tau^+\tau^-$  events. These numbers do not take into account detector acceptance effects or selection requirements. Although the total number of events meeting the  $2\ell$ SS and  $3\ell$  criteria is modest, the low statistical precision is balanced by minimal background contributions.

The strategy of the analysis is to first identify kinematic regions enriched in the signal process. In these regions, various reconstruction algorithms are implemented to create variables that can distinguish between the signal and several background processes. These reconstructed variables are combined into a single discriminant using a feed-forward neural network [39]. The same kinematic regions are used for the  $tHu$  and the  $tHc$  channel of the analysis, while individual neural networks are trained for each of the channels. Aside from this, the analysis strategy is identical for the  $tHu$  and the  $tHc$  channel. The distribution of the neural network output is then used as input to a maximum-likelihood fit, which considers statistical and systematic uncertainties. In addition to the signal-enriched regions, several regions enriched in specific background processes are included in the fit, to constrain the normalisation of these processes. If the fit




Figure 2: Example Feynman diagrams of the  $gq \rightarrow Ht$  production signal process resulting in (a) the  $2\ell SS$  final state and (b) the  $3\ell$  final state. The FCNC vertex is highlighted in red.

shows evidence of a signal, the corresponding significance is determined. Otherwise, upper limits are set on the FCNC branching ratios and the Wilson coefficients of the EFT dimension-6 operators. Finally, the results of this analysis are statistically combined with those from other ATLAS searches for  $tHq$  FCNC interactions in different final states.

## 2 The ATLAS detector

The ATLAS detector [40] at the LHC covers nearly the entire solid angle around the collision point.<sup>1</sup> It consists of an inner tracking detector surrounded by a thin superconducting solenoid, electromagnetic and hadron calorimeters, and a muon spectrometer incorporating three large superconducting air-core toroidal magnets.

The inner-detector system (ID) is immersed in a 2 T axial magnetic field and provides charged-particle tracking in the range of  $|\eta| < 2.5$ . The high-granularity silicon pixel detector covers the vertex region and typically provides four measurements per track, the first hit normally being in the insertable B-layer (IBL) installed before Run 2 [41, 42]. It is followed by the silicon microstrip tracker (SCT), which usually provides eight measurements per track. These silicon detectors are complemented by the transition radiation tracker (TRT), which enables radially extended track reconstruction up to  $|\eta| = 2.0$ . The TRT also provides electron identification information based on the fraction of hits (typically 30 in total) above a higher energy-deposit threshold corresponding to transition radiation.

The calorimeter system covers the pseudorapidity range  $|\eta| < 4.9$ . Within the region  $|\eta| < 3.2$ , electromagnetic calorimetry is provided by barrel and endcap high-granularity lead/liquid-argon (LAr) calorimeters, with an additional thin LAr presampler covering  $|\eta| < 1.8$  to correct for energy loss in material upstream of the calorimeters. Hadron calorimetry is provided by the steel/scintillator-tile calorimeter,

<sup>1</sup> ATLAS uses a right-handed coordinate system with its origin at the nominal interaction point (IP) in the centre of the detector and the  $z$ -axis along the beam pipe. The  $x$ -axis points from the IP to the centre of the LHC ring, and the  $y$ -axis points upwards. Cylindrical coordinates  $(r, \phi)$  are used in the transverse plane,  $\phi$  being the azimuthal angle around the  $z$ -axis. The pseudorapidity is defined in terms of the polar angle  $\theta$  as  $\eta = -\ln \tan(\theta/2)$ . Angular distance is measured in units of  $\Delta R \equiv \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$ .

segmented into three barrel structures within  $|\eta| = 1.7$ , and two copper/LAr hadron endcap calorimeters. The solid angle coverage is completed with forward copper/LAr and tungsten/LAr calorimeter modules optimised for electromagnetic and hadronic energy measurements respectively.

The muon spectrometer (MS) comprises separate trigger and high-precision tracking chambers measuring the deflection of muons in a magnetic field generated by the superconducting air-core toroidal magnets. The field integral of the toroids ranges between 2.0 and 6.0 T m across most of the detector. Three layers of precision chambers, each consisting of layers of monitored drift tubes, cover the region  $|\eta| < 2.7$  and are complemented by cathode-strip chambers in the forward region, where the detector occupancy is highest. The muon trigger system covers the range of  $|\eta| < 2.4$  with resistive-plate chambers in the barrel, and thin-gap chambers in the endcap regions.

Recorded events are selected by the first-level trigger system implemented in custom hardware, followed by selections made by algorithms implemented in software in the high-level trigger [43]. The first-level trigger accepts events from the 40 MHz bunch crossings at a rate below 100 kHz, which the high-level trigger further reduces to record events to disk at about 1 kHz.

An extensive software suite [44] is used in data simulation, in the reconstruction and analysis of real and simulated data, in detector operations, and in the trigger and data acquisition systems of the experiment.

### 3 Samples of data and simulated events

Proton–proton ( $pp$ ) collision data are used recorded with the ATLAS detector in the years 2015 to 2018 at a centre-of-mass energy of 13 TeV. After applying data-quality requirements [45], the data sample corresponds to an integrated luminosity of  $(140.1 \pm 1.2) \text{ fb}^{-1}$  [46]. The LUCID-2 detector [47] was used for the primary luminosity measurements, complemented by measurements using the inner detector and calorimeters.

Events were selected online during data taking by single-electron or single-muon triggers [48, 49]. Multiple unprescaled triggers were combined in a logical OR to increase the selection efficiency. The lowest-threshold triggers utilised isolation requirements to reduce the trigger rate. The higher-level lepton triggers had transverse momentum ( $p_T$ ) thresholds of 20 GeV for muons and 24 GeV for electrons in 2015 data, and 26 GeV for both the lepton types in 2016, 2017 and 2018 data. They were complemented by other triggers with higher  $p_T$  thresholds but no isolation requirements to increase the trigger efficiency.

Large sets of simulated events from signal and background processes were produced with Monte Carlo (MC) event generator programs to model the selected data. After event simulation, the response of the ATLAS detector was simulated using the GEANT4 toolkit [50] with a full detector model [51] or a fast simulation [51, 52] which employed a parameterisation of the calorimeter response. Samples relying on fast simulation were used to evaluate systematic uncertainties in the event generators and for the modelling of  $t\bar{t}$  production with two bosons.

To account for additional inelastic  $pp$  collisions in the same and neighbouring bunch crossings (pile-up), minimum-bias interactions were overlaid on the hard-scattering events at the level of energy depositions simulated using GEANT4. The minimum-bias events were simulated using PYTHIA 8.186 [53] with the A3 [54] set of tuned parameters and the NNPDF2.3LO set of parton distribution functions (PDF) [55]. The resulting events were weighted to reproduce the observed pile-up distribution. The average number of interactions per bunch crossing during the entire data-taking period from 2015 to 2018 is 33.7.

Finally, the simulated events were reconstructed using the same software as applied to the collision data. The same event selection requirements were applied and the selected events were passed through the same analysis chain. Small corrections were applied to simulated events such that the efficiencies of the utilised reconstruction methods are in better agreement with the response observed in data. More details of the simulated event samples are provided in the following subsections. Except for the events simulated with the SHERPA generator [56, 57], the EVTGEN [58] program was used to simulate bottom and charm hadron decays. If not mentioned otherwise, the top-quark mass was set to  $m_t = 172.5$  GeV and the Higgs-boson mass to  $m_H = 125$  GeV. All processes for which the parton shower is simulated using PYTHIA 8 use the A14 set of tuned parameters [59] and the NNPDF2.3LO PDF set.

### 3.1 Simulation of $tHq$ FCNC signal samples

Two different signal processes are studied: the  $t\bar{t}(t \rightarrow Hq)$  decay and the  $gq \rightarrow Ht$  production process. The EFT operators parametrising the FCNC coupling are implemented in the TopFCNC [60] model using the FEYNRULES 2.0 framework [61], allowing for next-to-leading-order (NLO) calculations of the postulated processes. The decay process events were simulated using the NLO matrix-element generator POWHEG BOX v2 [62–68], with the NNPDF3.0<sub>NLO</sub> PDF set [69]. The production of a top quark–antiquark pair is performed according to SM calculations in the five-flavour scheme, setting the masses of all quarks except for the top quark to zero. One top quark is required to decay via the SM decay mode  $t \rightarrow Wb$ , while the other performs the FCNC decay  $t \rightarrow Hq$ . Both top-quark decays are modelled using MADSPIN [70, 71]. Only leptonic final states for the SM decay of the top quark are considered. The Higgs-boson decay, and parton showers, hadronisation, and the underlying event, were modelled using PYTHIA 8.308 [72]. The matrix-element-to-parton-shower matching is steered by the  $h_{\text{damp}}$  parameter, which controls the  $p_T$  of the first additional gluon emission beyond the leading-order (LO) Feynman diagram in the parton shower and therefore regulates the high- $p_T$  emission against which the  $t\bar{t}$  system recoils. The event generation was performed with  $h_{\text{damp}} = 1.5m_t$  [73]. Four separate samples are produced, modelling the  $tHu/tHc$  coupling with either the top quark or the top antiquark decaying via the FCNC interaction. For consistency across searches, all  $tHq$  FCNC samples are normalised to the cross-section corresponding to a branching ratio of  $\mathcal{B}(t \rightarrow Hq) = 0.1\%$ . As explained previously, kinematic differences between the left- and right-handed couplings are negligible in the case of  $t \rightarrow Hq$  decay process. Therefore, the same samples are used to model the left-handed and the right-handed FCNC couplings. The matrix element generator MADGRAPH5\_AMC@NLO 2.9.9 [74] was used to model the  $gq \rightarrow Ht$  production signal using the TopFCNC model with the NNPDF3.0<sub>NLO</sub> PDF set and the five-flavour scheme. Separate samples are produced, each with exactly one of the Wilson coefficients set to 1.0 and the others set to zero, all of them at NLO in QCD.

The SM top-quark and Higgs-boson decays are simulated using MADSPIN. The three relevant Higgs-boson decays ( $H \rightarrow WW^*$ ,  $H \rightarrow ZZ^*$ ,  $H \rightarrow \tau^+\tau^-$ ) are modelled in separate samples. All Higgs-boson decays, parton shower, hadronisation, and the underlying event, were modelled using PYTHIA 8.307. The samples are normalised to the cross-section calculated with MADGRAPH5\_AMC@NLO.

### 3.2 Simulation of background processes

Samples of simulated events from SM  $t\bar{t}$  and single-top-quark production were simulated using the NLO matrix-element generator POWHEG BOX v2. For  $t\bar{t}$  and  $tW$  production and  $s$ -channel single-top-quark production ( $t\bar{b}$  production) the NNPDF3.0<sub>NLO</sub> PDF set was used with the five-flavour scheme. Following

a recommendation given in Ref. [68], single top-quark production in the  $t$ -channel ( $tq$  production) was simulated with the NNPDF3.0<sub>NLO\_NF4</sub> PDF set, which implements the four-flavour scheme. Parton showers, hadronisation, and the underlying event were modelled using PYTHIA 8.230. The event simulation used a value of  $h_{\text{damp}} = 1.5m_t$ .

The  $t\bar{t}$  production cross-section was scaled to  $\sigma(t\bar{t}) = 832$  pb, the value obtained from next-to-next-to-leading-order predictions from the TOP++2.0 program (see Ref. [75] and references therein), and it includes the resummation of next-to-next-to-leading logarithmic soft-gluon terms. The predicted cross-sections of  $tq$  and  $\bar{t}q$  production used to scale the corresponding samples of simulated events are  $\sigma(tq) = 136$  pb and  $\sigma(\bar{t}q) = 81$  pb and were calculated at NLO in QCD with the HATHOR 2.1 program [76, 77]. The total cross-section for  $t\bar{b}$  production was also computed at NLO in QCD with the HATHOR 2.1 program and the corresponding sample of simulated events was scaled to  $\sigma(t\bar{b} + \bar{t}b) = 10.32$  pb. The cross-section used for normalising the  $tW$  sample is  $\sigma(tW + \bar{t}W) = 71.7$  pb [78].

The production of a vector boson in association with jets ( $V$ +jets,  $V = W, Z$ ), including  $b$ - and  $c$ -jets, was simulated with the SHERPA 2.2.11 generator. NLO-accurate matrix elements for up to two partons and LO-accurate matrix elements for three to five partons are calculated in the five-flavour scheme using the COMIX [79] and OPENLOOPS [80–82] libraries. The default SHERPA parton shower [83] based on Catani–Seymour dipole factorisation and the cluster hadronisation model [84] are used. The samples are simulated using a dedicated set of tuned parameters developed by the SHERPA authors and use the NNPDF3.0<sub>NNLO</sub> PDF set.

The  $t\bar{t}W$  process was simulated using the SHERPA 2.2.10 generator. The matrix elements were calculated for up to one additional parton at NLO and up to two partons at LO using COMIX and OPENLOOPS, and merged with the SHERPA parton shower using the MEPS@NLO prescription [85]. In addition to the nominal prediction at NLO in QCD, higher-order corrections related to electroweak (EWK) contributions are also included. Event-by-event correction factors are applied that provide virtual NLO EWK corrections to  $\mathcal{O}(\alpha^2\alpha_s^2)$  and LO corrections to  $\mathcal{O}(\alpha^3)$  [57, 86, 87]. An independent SHERPA 2.2.10 sample was produced at LO to account for the sub-leading EWK corrections to  $\mathcal{O}(\alpha^3\alpha_s)$  [88]. The NLO QCD and NLO EWK contributions from SHERPA are combined following the method of Ref. [89]. The  $t\bar{t}W$  samples are normalised using a cross-section of 722 fb computed at NLO including the hard non-logarithmically enhanced radiation at NLO in QCD [89].

The modelling of the production of a top quark–antiquark pair with a  $Z$  boson decaying leptonically is done using the MADGRAPH5\_AMC@NLO 2.8.1 generator. It provides matrix elements at NLO in QCD, employing the NNPDF2.3<sub>LO</sub> PDF set. The generated events are interfaced with PYTHIA 8.244 for the parton shower and hadronisation stages. The  $t\bar{t}Z/\gamma^*(\rightarrow \ell^+\ell^-)$  prediction was normalised to the calculation at NLO QCD and NLO EWK accuracy based on Ref. [28] with the additional inclusion of off-shell effects, corresponding to 162 fb.

The production of  $t\bar{t}H$  events was modelled using the POWHEG BOX v2 generator at NLO in QCD with the NNPDF3.0<sub>NLO</sub> PDF set. The events were interfaced to PYTHIA 8.230. The cross-section was calculated at NLO QCD and NLO EWK using MADGRAPH5\_AMC@NLO and amounts to 507 fb.

The production of two or three weak vector bosons ( $VV, VVV$ ) in  $2\ell, 3\ell$  and  $4\ell$  final states was simulated using the SHERPA 2.2.12 generator including the off-shell contributions. Matrix elements including additional partons were calculated at NLO in QCD for up to one parton and at LO accuracy for two or three additional partons using COMIX and OPENLOOPS. For the parton shower simulation, the same method as in the  $W$ +jets and  $Z$ +jets samples is used. The  $VV$  and  $VVV$  samples are normalised to the total cross-sections provided by SHERPA.



The  $tZq$  process was modelled at NLO with the MADGRAPH5\_AMC@NLO 2.9.5 generator and the NNPDF3.0<sub>NLO</sub> PDF. The generated events were interfaced with PYTHIA 8.230. Matrix elements of the  $tWZ$  process were also calculated with MADGRAPH5\_AMC@NLO 2.2.2, using the same PDF set. The events were interfaced with PYTHIA 8.212 using the same tune and PDF set as used for  $tZq$  production. The samples were normalised to the theoretical cross-section at NLO QCD.

Some rare processes are considered:  $t\bar{t}$ ,  $t\bar{t}t$ ,  $tHW$ ,  $tHq$ ,  $t\bar{t}WW$ ,  $t\bar{t}HH$ ,  $t\bar{t}WH$ ,  $t\bar{t}ZZ$ ,  $t\bar{t}WZ$  and  $VH$  production. The processes involving at least one top quark were simulated with MADGRAPH5\_AMC@NLO, while  $VH$  was produced with POWHEG BOX v2. Simulated events of all processes were interfaced to PYTHIA 8 to simulate parton showers and hadronisation. The processes are normalised to their predicted NLO cross-sections. Their combined contribution is nearly negligible, ranging from 0.1 to 1%, depending on the kinematic region considered.

## 4 Object reconstruction and event preselection

Events are required to have at least one vertex reconstructed from at least two ID tracks with transverse momenta of  $p_T > 0.5$  GeV. The primary vertex of an event is defined as the vertex with the highest  $p_T^2$  summed over the ID tracks [90] matched to it.

Electron candidates are reconstructed by matching a track in the ID to clusters of energy deposits in the electromagnetic calorimeter [91]. The pseudorapidity of clusters,  $\eta_{\text{cluster}}$ , is required to be in the range of  $|\eta_{\text{cluster}}| < 2.47$ . However, clusters are excluded if they are in the transition region  $1.37 < |\eta_{\text{cluster}}| < 1.52$  between the barrel and endcap electromagnetic calorimeters. Electron candidates must have  $p_T > 10$  GeV. A likelihood-based discriminant is constructed to simultaneously evaluate several properties of electron candidates, including shower shapes in the electromagnetic calorimeter, track quality, and the detection of transition radiation produced in the TRT [92]. Applying a discriminant requirement enhances the selection of prompt electrons produced in  $W$ -,  $Z$ - or Higgs-boson decays and leptonic  $\tau$ -lepton decays, while effectively rejecting photon conversions and hadrons misidentified as electrons. All selected electrons must meet the *Tight* selection criteria as defined in Ref. [92]. Muon candidates are reconstructed by combining tracks in the MS with tracks in the ID [93]. The tracks must be in the range of  $|\eta| < 2.5$  and have  $p_T > 10$  GeV. Similarly to electrons, likelihood-based identification criteria are applied [93]. Muons are required to satisfy the *Medium* selection defined in Ref. [93].

The tracks matched to electron and muon candidates must point to the primary vertex, which is ensured by requirements imposed on the transverse impact-parameter significance,  $|d_0/\sigma(d_0)| < 5.0$  for electrons and  $|d_0/\sigma(d_0)| < 3.0$  for muons, and on the longitudinal impact parameter,  $z_0$ , for which  $|z_0 \sin(\theta)| < 0.5$  mm has to be satisfied for both lepton flavours. Non-prompt electrons and muons are leptons produced by mechanisms other than  $W$ -,  $Z$ - or Higgs-boson decays and leptonic  $\tau$ -lepton decays. They are effectively rejected by using multivariate discriminants computed with boosted decision trees (BDT), which integrate electromagnetic shower shapes and track information from the ID [94]. Separate BDTs are trained for electrons in the barrel ( $|\eta| < 1.37$ ) and endcap ( $|\eta| > 1.37$ ) regions, while a single BDT is employed for muons. The efficiency for correctly identifying prompt muons (electrons) ranges from approximately 80% to 95% (65% to 90%) for  $p_T$  values between 20 GeV and 45 GeV, with the efficiency plateauing beyond 45 GeV. Overall, the BDTs achieve a 71% (90%) rejection rate for muons (electrons) from  $B$ -hadron decay. Additionally, the contribution of electrons reconstructed with an incorrectly reconstructed charge is significantly reduced by using an additional BDT [91], which consolidates information about an electron candidate's charge, impact parameter, energy, and ID track into a single discriminant. This achieves a

95% efficiency for electrons with a correct charge assignment while rejecting 94% of charge-misidentified electrons.

Scale factors are used to correct the efficiencies in simulation to match the efficiencies measured for the electron [48] and muon [49] triggers, and the reconstruction, identification and isolation criteria [91, 93].

Jets are reconstructed from particle-flow objects [95] with the anti- $k_t$  clustering algorithm [96, 97] using a radius parameter of 0.4. This algorithm matches topological energy clusters [98] in the calorimeters to selected tracks in the ID. The energy of tracks is subtracted from the matched topological clusters and both the tracks and the energy-subtracted topological clusters are used as input to the jet reconstruction. The jet energy is calibrated by applying several simulation-based corrections and techniques correcting for differences between simulation and data [99]. The jets must satisfy  $p_T > 20$  GeV and  $|\eta| < 2.5$ .

To suppress jets originating from pile-up collisions, several track-based variables are combined with a multivariate technique to form the jet-vertex-tagger (JVT) discriminant [100]. Jets with  $p_T < 60$  GeV and  $|\eta| < 2.4$  are required to have a JVT discriminant above 0.5, which corresponds to an efficiency of 92% for jets from the primary vertex, while 98% of jets from pile-up events are rejected.

Identification of jets containing  $B$  hadrons ( $b$ -tagging) is performed with the DL1r algorithm, which uses a deep feed-forward neural network with several  $b$ -tagging algorithms as inputs [101]. These input algorithms exploit the impact parameters of charged-particle tracks, the properties of reconstructed secondary vertices and the topology of  $b$ - and  $c$ -hadron decays inside the jets. The requirement on the DL1r discriminant is chosen such that the average tagging efficiency of  $b$ -jets from simulated dileptonic  $t\bar{t}$  events is 70%. The corresponding  $p_T$ -dependent  $c$ -jet rejection factors range from 10 to 14, while those for light-flavour jets range from 100 to 900. Differences between the  $b$ -tagging efficiency between collision data and simulation are corrected using simulation-to-data scale factors derived from  $t\bar{t}$  events. The scale factors depend on the  $p_T$  of the jets and are consistent with unity within the uncertainties. The obtained scale factors depend on the parton-shower generator used to produce the  $t\bar{t}$  samples. When using samples produced with a different parton-shower generator, for example SHERPA, to model  $W$ +jets events, or when evaluating systematic uncertainties with a setup based on HERWIG, additional correction factors called MC-to-MC scale factors are applied. Since the DL1r algorithm uses measurements from the ID, the identification of  $b$ -jets is limited to the region with  $|\eta| < 2.5$ .

With the above definitions of physical objects it is possible for ambiguities to arise, since all objects are reconstructed independently of each other. To avoid the double-counting of candidates which satisfy more than one selection criterion, a procedure called *overlap removal* is applied. Reconstructed physical objects are removed in the following order: electrons sharing an ID track with a muon; jets within  $\Delta R = 0.2$  of an electron, thereby avoiding double-counting electron energy deposits as jets; electrons within  $\Delta R = 0.4$  of a remaining jet, for reducing the impact of non-prompt electrons; jets within  $\Delta R = 0.2$  of a muon if they have two or fewer matched tracks; muons within  $\Delta R = 0.4$  of a remaining jet, reducing the rate of non-prompt muons.

The missing transverse momentum  $\vec{p}_T^{\text{miss}}$  is reconstructed as the negative vector sum of the transverse momentum of the reconstructed leptons and jets, and ID tracks that point to the primary vertex but are not matched to a reconstructed object [102]. The magnitude of  $\vec{p}_T^{\text{miss}}$  is denoted by  $E_T^{\text{miss}}$ .

Events selected by this analysis are required to have at least one charged lepton with  $p_T > 28$  GeV. Additional leptons must satisfy a  $p_T$  threshold of 10 GeV. In the  $2\ell\text{SS}$  final state, exactly two leptons of the same charge are required, while in the  $3\ell$  final state the requirement is three leptons with a total charge of  $\pm 1e$ . The lepton with the highest  $p_T$  is referred to as  $\ell_0$ . The lepton with the second highest  $p_T$  is denoted

Table 1: Overview of the preselections applied in the analysis. These selections ensure jet and  $b$ -tag multiplicities, lepton momenta, and define the  $2\ell\text{SS}$  and  $3\ell$  final states.

Preselection		
$N_{\text{jets}}$	$\geq 1$	
$N_{b\text{-tags}}$	$\geq 1$	
$p_{\text{T}}(\text{jet})$	$\geq 20 \text{ GeV}$	
$p_{\text{T}}(\ell)$	$\geq 10 \text{ GeV}$	
$p_{\text{T}}(\ell_0)$	$\geq 28 \text{ GeV}$	
	$2\ell\text{SS}$	$3\ell$
$N_{\ell}$	$= 2$	$= 3$
$\sum q(\ell_i)$	$= \pm 2e$	$= \pm 1e$

by  $\ell_1$  and, for events in the  $3\ell$  final state, the lepton with the lowest  $p_{\text{T}}$  is called  $\ell_2$ . In the  $3\ell$  final state an alternative labelling of leptons is introduced based on their charge. In any given  $3\ell$  event two leptons have the same charge while the remaining lepton has an opposite charge. The latter is labelled  $\ell_{\text{OS}}$ , the former  $\ell_{\text{SS},0}$  and  $\ell_{\text{SS},1}$  with the number indexing the lepton in order of decreasing  $p_{\text{T}}$ . In addition to the requirements on leptons, considered events need to contain at least one jet, at least one of which is  $b$ -tagged. A summary of the preselection applied to events is given in Table 1.

## 5 Background estimate

Several SM background processes are present in the signal regions. Many of these processes can be simulated by using MC, others require dedicated treatment. Particularly the modelling of background from non-prompt leptons must be validated and potentially corrected. This background is separated into various categories, encompassing electrons from prompt muon decay, electrons from photon conversion and leptons produced in a hadronic jet, where further differentiation depending on the jet primary particle are made. Jets initiated by  $b$ -quarks, by other quarks or gluons, and by  $\tau$ -leptons are considered, where leptons from  $b$ -jets are further categorised into electrons and muons. The production of these non-prompt leptons is fully accounted for by the simulated SM samples. By navigating through the generator-level parent particles of a given lepton, this lepton is sorted into one of the previously discussed lepton-origin categories. Based on this, several MC templates are defined containing the events corresponding to exactly one non-prompt lepton-origin category. The remaining events include only prompt leptons and remain in their original SM background template. All background estimate methods discussed in the following are based on these prompt and non-prompt MC templates. Several of the processes discussed in this section are assigned unconstrained (*free-floating*) normalisation factors in the final maximum-likelihood fit. These normalisation factors are constrained in the fit through dedicated control regions, enriched in the respective processes. It should be noted that the QCD multijet background is negligible in this analysis, owing to the rarity of the considered  $2\ell\text{SS}$  and  $3\ell$  final states.

## 5.1 Non-prompt lepton background

The primary source of non-prompt lepton background is the decay of  $B$ -hadrons. These particular objects are also referred to as *heavy-flavour (HF) decay leptons*. They mainly consist of events from  $t\bar{t}$  production (75%), while the remaining events are split evenly between  $V$ +jets and single-top-quark production. The *Template fit method* [103] is used to determine the rates of HF-decay electrons and HF-decay muons. It assumes the kinematic distributions of the corresponding processes are well described, while their normalisation may require corrections. The normalisation factors are determined in the final maximum-likelihood fit, which is performed simultaneously in the  $2\ell$ SS and the  $3\ell$  final states.

As explained above, various other non-prompt lepton processes contribute to this analysis. Their contribution is minor compared with the HF-decay  $e$  and HF-decay  $\mu$  processes. In the fit, they are assigned normalisation uncertainties of 50%.

## 5.2 Charge misidentification

In addition to the processes discussed above, the phase space of the  $2\ell$ SS final state is contaminated by events containing prompt electrons with a misidentified charge. The modelling of such objects is strongly dependent on their  $p_T$ . Furthermore, electrons can radiate hard bremsstrahlung through interaction with the detector material, which then undergoes asymmetric conversion into an electron-positron-pair. The probability for electrons to interact with the detector material increases with the absolute value of the pseudorapidity  $|\eta|$ . Both, charge misidentified electrons and electrons from bremsstrahlung, are considered as a single background labelled Q-misID, which is modelled using a data-driven method based on the one described in Ref. [103]. To avoid double-counting Q-misID events, MC events featuring an electron from any of the considered processes are identified using generator level information and are excluded from the analysis.

The data-driven Q-misID background estimate is performed as follows. Events containing two isolated electrons with an invariant mass around the mass of the  $Z$  boson are considered, where the electrons can have either an opposite-sign charge (OS) or the same-sign charge (SS). To maintain orthogonality to the kinematic regions of the main analysis, there is an  $N_{b\text{-tags}} = 0$  requirement. The  $m_{ee}$  distribution in the vicinity of  $m_Z = 91.19$  GeV [104] is fitted separately for OS and SS events using Breit–Wigner functions. The fitted centres  $m_{\text{peak}}$  and widths  $\sigma$  are used to define the  $Z$ -windows up to  $4\sigma$  from the centres, and sideband regions  $[m_{\text{peak}} - 8\sigma, m_{\text{peak}} - 4\sigma] \cup [m_{\text{peak}} + 4\sigma, m_{\text{peak}} + 8\sigma]$ . The assumption is made, that events in the  $Z$ -window consist of prompt  $Z \rightarrow e^+e^-$  events and a uniform background. The uniform background is estimated from the average yield in the sideband regions, separately for the OS and SS events, and are subtracted in the  $Z$ -window.

Q-misID efficiencies are extracted in bins of  $|\eta|$  and  $p_T$ , corresponding to the probability for an electron in a given bin to be charge-misidentified. These relate the number of background-corrected SS and OS events in the  $Z$ -window and can thus be determined from the obtained event yields. The Q-misID events in a given region of the  $2\ell$ SS phase space are then modelled by selecting data events with two opposite-charge leptons while keeping all other selection requirements of the region unmodified, and weighting these data events with the obtained efficiencies. It should be noted that due to the high accuracy of the ATLAS muon spectrometer, the muon charge misidentification is negligible in this analysis.

### 5.3 Prompt-lepton background

The  $t\bar{t}W$  and  $t\bar{t}Z$  processes represent the primary background sources from prompt leptons. While measurements of  $t\bar{t}W$  production deviate from theoretical predictions by  $1.4\sigma$  [105], measurements of  $t\bar{t}Z$  production align closely with SM expectations. However, it's worth noting that the latest measurement of  $t\bar{t}Z$  production is conducted in regions with a jet multiplicity of at least four [106]. This analysis extends to kinematic regions with fewer jets, which may not be adequately modelled by MC simulations. To address this discrepancy, the normalisation of  $t\bar{t}W$  and  $t\bar{t}Z$  production is left unconstrained in the maximum-likelihood fit. Similar to the non-prompt templates, a single normalisation factor is applied per process across the entire phase space, encompassing both the  $2\ell$ SS and the  $3\ell$  final states.

The  $2\ell$ SS and  $3\ell$  event selection requires at least one  $b$ -jet in the final state. In the  $VV$  MC samples used in the analysis these quarks must be produced by the shower generator. To account for potential mismodeling of the jet flavour composition, the  $VV$  background is split according to the number of leptons ( $2\ell$ ,  $3\ell$ ,  $4\ell$ ) and the generator-level flavour of additional jets. If the event contains at least one jet originating from the decay of either a bottom or a charm hadron is sorted into the  $b/c$  category. If it exclusively contains jets from light quarks or  $\tau$ -leptons it is included in the  $l/\tau$  category. The  $VV3\ell + b/c$  template has the largest contribution in the phase space and its normalisation is left free-floating in the maximum-likelihood fit. The remaining templates ( $VV2\ell + l/\tau$ ,  $VV2\ell + b/c$ ,  $VV3\ell + l/\tau$ ,  $VV4\ell + b/c$  and  $VV4\ell + l/\tau$ ) are assigned individual 50% normalisation uncertainties. These minor  $VV$  templates, alongside the minor non-prompt templates and several SM processes are combined into one *Others* category in the following. The rates of the majority of processes in this category are similar in the analysis phase space.

## 6 Event categorisation

Preselected events are categorised into several kinematic regions. These regions are either enriched in signal events, to improve the overall sensitivity of the analysis, or in certain background events, to control the normalisation of specific processes.

### 6.1 Definition of signal regions

After event preselection, four signal regions are defined in this analysis, two in the  $2\ell$ SS final state and two in the  $3\ell$  final state. One region per final state is enriched in the  $t \rightarrow Hq$  decay signal. These two regions are referred to as SR $2\ell$ Dec and SR $3\ell$ Dec. The other two signal regions contain a larger fraction of  $gq \rightarrow tH$  production signal and are correspondingly named SR $2\ell$ Prod and SR $3\ell$ Prod.

Both the  $2\ell$ SS signal regions require exactly one  $b$ -tagged jet. Given that a higher jet multiplicity is anticipated for the decay signal compared to the production signal, the SR $2\ell$ Dec is defined by requiring  $N_{\text{jets}} \geq 4$ . A cut of  $N_{\text{jets}} \leq 3$  in the SR $2\ell$ Prod ensures the orthogonality of the two regions. To minimise the contribution from HF-decay leptons, the  $p_T$  of  $\ell_1$  is required to be larger than 12 GeV in the SR $2\ell$ Dec and 16 GeV in the SR $2\ell$ Prod. As Q-misID events strongly contaminate the  $2\ell$ SS final state, a cut on the invariant mass of the two leptons is imposed in both the regions. If both the leptons are identified as electrons,  $|m(e, e) - m_Z| \geq 10$  GeV must be satisfied.

Similar to the  $2\ell$ SS final state, both the signal regions in the  $3\ell$  final state require exactly one  $b$ -tagged jet. The primary difference between the two regions is again the jet multiplicity. Overall fewer jets are

Table 2: Overview of the event selections applied to define the signal regions of the analysis.

	SR2 $\ell$ Dec	SR2 $\ell$ Prod	SR3 $\ell$ Dec	SR3 $\ell$ Prod
$N_{\text{jets}}$	$\geq 4$	$\leq 3$	$\geq 3$	$\leq 2$
$N_{b\text{-tags}}$	$= 1$	$= 1$	$= 1$	$= 1$
$p_{\text{T}}(\ell_1)$	$\geq 12 \text{ GeV}$	$\geq 16 \text{ GeV}$	$\geq 20 \text{ GeV}$	$\geq 20 \text{ GeV}$
$p_{\text{T}}(\ell_2)$	–	–	$\geq 16 \text{ GeV}$	$\geq 16 \text{ GeV}$
$ m(e, e) - m_Z $	$\geq 10 \text{ GeV}$	$\geq 10 \text{ GeV}$	–	–




Figure 3: The composition of background processes in the (a) SR2 $\ell$ Dec, (b) SR2 $\ell$ Prod, (c) SR3 $\ell$ Dec and (d) SR3 $\ell$ Prod. The *Others* template contains various minor processes, which individually only have small contributions. No correction to the normalisation of any process is applied.

expected for the signal in the 3 $\ell$  final state. Hence,  $N_{\text{jets}} \geq 3$  is required for the SR3 $\ell$ Dec, while  $N_{\text{jets}} \leq 2$  is required for the SR3 $\ell$ Prod. To reduce the HF-decay contribution, the definition of both the regions require cuts on the  $p_{\text{T}}$  of the two sub-leading leptons:  $p_{\text{T}}(\ell_1) \geq 20 \text{ GeV}$  and  $p_{\text{T}}(\ell_2) \geq 16 \text{ GeV}$ . As the Q-misID process does not contribute to the 3 $\ell$  final state, no further restrictions to the event selection are made. The definition of all signal regions is summarised in Table 2, while their respective composition in terms of background processes is shown in Figure 3.

Table 3: Overview of the event selections applied to define the non-prompt lepton background control regions of the analysis.

	CR2 $\ell$ HF $e$	CR2 $\ell$ HF $\mu$	CR3 $\ell$ HF $e$	CR3 $\ell$ HF $\mu$
$N_{\text{jets}}$	$\leq 3$	$\leq 3$	$\geq 1$	$\geq 1$
$N_{b\text{-tags}}$	$\geq 1$	$\geq 1$	$= 1$	$= 1$
$\ell_0$ flavour	$\mu$	$\mu$	–	–
$\ell_1$ flavour	$e$	$\mu$	–	–
$p_{\text{T}}(\ell_1)$	$< 16 \text{ GeV}$	$< 16 \text{ GeV}$	$\geq 20 \text{ GeV}$	$\geq 20 \text{ GeV}$
$\ell_2$ flavour	–	–	$e$	$\mu$
$p_{\text{T}}(\ell_2)$	–	–	$< 16 \text{ GeV}$	$< 16 \text{ GeV}$

## 6.2 Control regions for non-prompt-lepton backgrounds

The template-fit method assumes that corrective normalisation factors are applicable independently of the considered final state. To ensure this assumption is correct, two control regions are defined for each of the HF-decay processes: one in the  $2\ell$ SS final state (CR2 $\ell$ HF $e$  and CR2 $\ell$ HF $\mu$ ) and one in the  $3\ell$  final state (CR3 $\ell$ HF $e$  and CR3 $\ell$ HF $\mu$ ).

Overall, the average  $p_{\text{T}}$  of leptons originating from a HF decay is smaller than that of prompt leptons. Thus, all HF-decay control regions are defined by imposing upper boundaries on the  $p_{\text{T}}$  of the lowest- $p_{\text{T}}$  lepton. The boundaries are chosen such that an orthogonality to the signal regions of the respective final state is ensured. Requiring the flavour of the lowest- $p_{\text{T}}$  lepton to be either  $e$  or  $\mu$  ensures that the region is exclusively enriched in HF-decay  $e$  or HF-decay  $\mu$  events. In the CR2 $\ell$ HF $e$  and the CR2 $\ell$ HF $\mu$  the  $p_{\text{T}}$ -leading lepton is required to be a muon, reducing the contamination of Q-misID events. Due to high statistical uncertainty, only the event yields of the CR2 $\ell$ HF $e$  and the CR2 $\ell$ HF $\mu$  enter the maximum-likelihood fit. For the CR3 $\ell$ HF $e$  and the CR3 $\ell$ HF $\mu$ , the distributions of the  $p_{\text{T}}(\ell_1)$  variable are included. A summary of the definition of all non-prompt control regions can be found in Table 3.

## 6.3 Control regions for $t\bar{t}W$ and $t\bar{t}Z$ backgrounds

Three additional control regions are defined to determine the normalisation of the background from  $t\bar{t}W$  and  $t\bar{t}Z$  production. Studies on various kinematic distributions in the signal regions of the  $2\ell$ SS final state show a strong similarity between the two processes. Therefore, in this final state only one control region (denoted CR2 $\ell t\bar{t}V$ ) is defined to constrain the sum of  $t\bar{t}W$  and  $t\bar{t}Z$ . This region requires at least four jets, exactly two of which are  $b$ -tagged, because both the top quarks in the process are expected to produce a  $b$ -quark. An additional requirement of  $p_{\text{T}}(\ell_1) \geq 18 \text{ GeV}$  further increases the purity of the region in  $t\bar{t}V$  events by reducing contamination from HF-decay events. The  $p_{\text{T}}(\ell_1)$  variable is used as input for the maximum-likelihood fit.

In the  $3\ell$  final state, distinct shape differences between  $t\bar{t}W$  and  $t\bar{t}Z$  production are observed due to the absence of any requirement on the invariant mass of lepton pairs. Therefore, two control regions, CR3 $\ell t\bar{t}W$  and CR3 $\ell t\bar{t}Z$ , are defined. Both the regions require at least two jets, two of which are  $b$ -tagged. Events are then assigned to the CR3 $\ell t\bar{t}Z$  if at least one pair of leptons with the same flavour and opposite charge can be found satisfying  $|m(\ell^+, \ell^-) - m_Z| < 10 \text{ GeV}$ . In cases where this criterion is not met, events are assigned to CR3 $\ell t\bar{t}W$ . In both the control regions, the transverse momentum of the  $p_{\text{T}}$ -leading  $b$ -tagged jet

Table 4: Overview of the event selections applied to define the  $t\bar{t}W$  and  $t\bar{t}Z$  control regions of the analysis.

	CR2 $t\bar{t}V$	CR3 $t\bar{t}W$	CR3 $t\bar{t}Z$
$N_{\text{jets}}$	$\geq 4$	$\geq 2$	$\geq 2$
$N_{b\text{-tags}}$	$= 2$	$= 2$	$= 2$
$\ell_0$ flavour	$\mu$	–	–
$p_{\text{T}}(\ell_1)$	$\geq 18 \text{ GeV}$	$\geq 20 \text{ GeV}$	$\geq 20 \text{ GeV}$
$p_{\text{T}}(\ell_2)$	–	$\geq 16 \text{ GeV}$	$\geq 16 \text{ GeV}$
$ m(\ell^+, \ell^-) - m_Z $	–	$\geq 10 \text{ GeV}$	$< 10 \text{ GeV}$

$p_{\text{T}}(b\text{-jet}_0)$  is used as input for the maximum-likelihood fit. A summary of the definition of all  $t\bar{t}V$  control regions can be found in Table 4.

## 7 Separation of signal and background

Artificial neural networks (NNs) are used to separate signal and background in all four signal regions combining several kinematic variables into optimised NN discriminants. In addition to variables derived from the reconstructed objects the NNs build on a diverse range of reconstructed kinematic variables. Detailed information regarding the reconstruction algorithms is provided in Section 7.1, while a comprehensive description of the NN training process can be found in Section 7.2.

### 7.1 Event reconstruction in signal regions

The reconstruction algorithms employed are described below. In  $tHq$  FCNC interactions, multiple decay modes of the Higgs boson that contribute to the  $2\ell\text{SS}$  and  $3\ell$  final states are considered. However, the  $H \rightarrow WW^*$  decay mode has by far the largest contribution with  $\geq 75\%$  in all signal regions. Thus, all reconstruction efforts presented here focus on this specific Higgs-boson decay mode. Most algorithms are customised to fit one of the two considered final states. The relevant final states for each algorithm are given in parentheses.

**Recursive Jigsaw Reconstruction ( $2\ell\text{SS}$  and  $3\ell$ )** The Recursive Jigsaw Reconstruction (RJR) technique [107] is a method for resolving combinatorial and kinematic ambiguities which arise in hadron collider events. RJR relies on decay trees which describe the topologies of interest, and relates the reference frames of the intermediate particles in the decay. Lepton and jet assignment, along with the splitting of  $E_{\text{T}}^{\text{miss}}$ , is governed by a set of jigsaw rules which rely on known properties of the decay tree. For instance the top-quark and Higgs-boson masses can be used to constrain the assignment of the jets, or the  $W$ -boson mass can be used to facilitate  $E_{\text{T}}^{\text{miss}}$  splitting and matching to leptons.

This process leads to the computation of four-vectors of the reconstructed intermediate states, accessible in any of the frames defined in the decay tree. For each of the four topologies ( $2\ell 1b 2j$ ,  $2\ell 1b 3j$ ,  $3\ell 1b 0j$ ,  $3\ell 1b 1j$ , where  $b$  refers to  $N_{b\text{-jets}}$  and  $j$  to  $N_{\text{jets}}$ ) there are different intermediate states, split by production versus decay and again by final state. The particles that are reconstructed for each individual region are listed in Table 5. These include  $W$  and Higgs bosons, and top quarks. Figure 4 depicts the probability



Table 5: Particles from the  $tHq$  processes which are reconstructed using the Recursive Jigsaw Reconstruction (RJR) in the respective signal regions.

Regions	Particle name	Description
SR2 $\ell$ Dec/ SR3 $\ell$ Dec	$t_{\text{SM}}$	The top quark decaying via $t \rightarrow Wb$
	$W_t$	The $W$ boson from the SM top-quark decay
	$t_{\text{FCNC}}$	The top quark decaying via $t \rightarrow Hq$
	$H$	The Higgs boson originating from th FCNC top-quark decay
SR2 $\ell$ Dec	$W_{\text{had}}$	The hadronically decaying $W$ boson from the $H \rightarrow WW^*$ decay
SR2 $\ell$ Prod/ SR3 $\ell$ Prod	$t_{\text{SM}}$	The top quark produced in the $gq \rightarrow Ht$ process
	$W_t$	The $W$ boson from the top-quark decay
	$H$	The Higgs boson produced in the $gq \rightarrow Ht$ process




Figure 4: Probability densities of (a)  $m(\ell_1, t_{\text{FCNC}})$  in the SR2 $\ell$ Dec and (b)  $m(W_t)$  in the SR3 $\ell$ Dec for the  $tHu$  signal process and the sum of all background processes. The vertical lines on bins represent MC-statistical uncertainties. The last bin includes the overflow.

densities of two RJR reconstructed variables for the  $tHu$  signal and the combination of all background processes, showing clear shape differences between the two.

**Neutrino Independent Combinatorics Estimator (3 $\ell$ )** The considered decay scenario of a signal event in the 3 $\ell$  final state involves one top quark decaying into a  $b$ -quark, a lepton and a neutrino, while the Higgs-boson decay is expected to yield two leptons and two neutrinos. Studies of the MC signal samples show that on average the two leptons from the Higgs-boson decay have the smallest angular separation out of any lepton pair in the event. Based on this information, the two leptons with the smallest angular separation  $\Delta R$  are labelled  $\ell_{H,0}$  and  $\ell_{H,1}$ , ordered by their transverse momentum. The remaining lepton is labelled  $\ell_t$ , while the  $b$ -tagged jet is denoted  $b_t$ . Various kinematic parameters are calculated for these objects, two of which are shown as examples in Figure 5 for the full  $tHu$  signal and the combination of all background processes, demonstrating their high separation power. However, these parameters are discarded, if the angular separation of  $b_t$  from either  $\ell_{H,i}$  is smaller than the separation from  $\ell_t$ . Conversely, when




Figure 5: Probability densities of (a)  $m(\ell_{H,0}, \ell_{H,1})$  and (b)  $m(\ell_t, b_t)$  in the SR3lDec for the  $tHu$  signal process and the sum of all background processes. The vertical lines on bins represent MC-statistical uncertainties. The last bin includes the overflow.

the angular separation of  $b_t$  from  $\ell_t$  is found to be smaller than from any other lepton, and further, when  $\ell_{H,0}$  and  $\ell_{H,1}$  have opposite charges the *Neutrino independent combinatorics estimator* reconstruction (NICE Reco) condition is defined as satisfied. In this scenario, the calculated kinematic parameters have a particularly high separation power between signal and background processes. This information is provided as additional input to the NNs.

## 7.2 Training of feed-forward neural networks

The NNs are implemented using the NeuroBayes package [108, 109], which combines a three-layer feed-forward NN with a complex and robust preprocessing of the input variables before they are presented to the NN. The preprocessing produces a ranking of the input variables based on an algorithm employing the total correlation of a set of variables to the target function which assumes the value 1 for signal and 0 for background events. The input variable selection is performed once per signal region.

The selected variables of all signal regions are listed in Tables 11–14. Overall, invariant masses of reconstructed objects, including RJR objects, compose the largest fraction. The most significant variables in both the  $3\ell$  signal regions are the invariant mass of the opposite charge and either of the two same charge leptons  $m(\ell_{OS}, \ell_{SS,0})$  and  $m(\ell_{OS}, \ell_{SS,1})$ . In the SR2lProd, the invariant mass of the subleading- $p_T$  lepton and the RJR Higgs boson  $m(\ell_1, H)$  is the most important variable, followed by the number of jets  $N_{\text{jets}}$ . The most significant variable of the SR2lDec is the scalar  $p_T$  sum of all jets  $H_T(\text{jets})$ , a quantity that is also highly ranked in other trainings. The invariant mass of the leading- $p_T$  lepton and the event's  $b$ -jet  $m(\ell_0, b\text{-jet})$  is the second-highest ranked in the training.

Individual NNs are created for both the signal processes and every signal region, summing up to a total of eight NN architectures. The training uses the data-driven Q-misID estimate together with unmodified MC templates as input, before any corrections to their normalisation by the maximum-likelihood fit. Cross-training is employed to prevent overfitting, where the signal and background MC events are split

into half. The splitting is performed in a pseudo-random way based on the parity of the unique event identification number. For each half, a separate NN model is trained, which is then applied to the other half of MC events in the final analysis.

During the training process of all the NNs, the signal is trained against all considered backgrounds with a fraction of 50% signal events and 50% background events. The different background processes are weighted relative to each other according to their expected number of events. NeuroBayes uses Bayesian regularisation techniques for the training process to improve the generalisation performance and to avoid overfitting. The network infrastructure consists of one input node for each input variable, a single hidden layer, and one output node giving a continuous output in the interval  $(-1, +1)$ . The number of nodes in the hidden layer is optimised individually for each NN model. As a non-linear activation function NeuroBayes uses the symmetric sigmoid function. In the region close to zero, the sigmoid function has a linear response. The final discriminant  $D_{\text{NN}}$  is obtained by linearly scaling the output of the NNs to the interval  $(0, 1)$ .

## 8 Systematic uncertainties

Several sources of systematic uncertainty affect the expected event yield from signal and background processes and the shape of the NN discriminants used in the maximum-likelihood fits. The systematic uncertainties are divided into two major categories. There are experimental uncertainties in the reconstruction of the four-momenta of the final-state objects: electrons, muons, jets,  $b$ -tagged jets, and  $E_{\text{T}}^{\text{miss}}$  as a sign of a high- $p_{\text{T}}$  neutrino. The second category of uncertainties is related to the modelling of scattering processes with event generators and of the interaction of particles with the detector. All uncertainties are propagated through the analysis and their effects on the expected event yields and discriminant distributions are accounted for by including corresponding nuisance parameters in the fit. In the following, the estimate of experimental and modelling uncertainties is explained in more detail.

### 8.1 Experimental uncertainties

The uncertainty in the integrated luminosity of the combined 2015–2018 data set is 0.83% and is based on a calibration of the luminosity scale using  $x$ - $y$  beam-separation scans [46]. The luminosity uncertainty is applied to the expected signal and background event yields.

Scale factors are applied to simulated events to correct for reconstruction, identification, isolation and trigger performance differences between data and detector simulation for electrons and muons. These scale factors and their systematic uncertainties, and the lepton momentum scale and resolution, were assessed using  $Z \rightarrow e^+e^-$  and  $Z \rightarrow \mu^+\mu^-$  events in simulation and data [91, 93].

The jet energy scale (JES) was calibrated using a combination of test-beam data, simulation and in situ techniques [99]. The JES is parameterised in bins of jet  $p_{\text{T}}$  and  $\eta$ . Its uncertainty is decomposed into a set of 30 uncorrelated components, of which 29 are non-zero in a given event depending on the type of simulation used. Sources of uncertainty contributing to the JES uncertainties include the  $\eta$  intercalibration of forward jets in the range of  $0.8 < |\eta_{\text{det}}| < 4.5$  with those in the central barrel region ( $|\eta_{\text{det}}| < 0.8$ ), pile-up modelling, jet flavour composition and response, differences between jets induced by  $b$ -quarks and those from gluons or light-quarks, single-particle response, detector modelling, non-closure, and effects of jets not fully contained in the calorimeter.

The uncertainty of the jet energy resolution (JER) is evaluated by smearing jet energies according to a Gaussian function [99]. Thirteen orthogonal components account for jet- $p_T$  and  $\eta$ -dependent differences between simulation and data which were determined using dijet events and noise measurements based on random cones. The smearing is applied to simulated events if the resolution in data is larger than in MC simulation, and to pseudo-data when the resolution is larger in simulation than in collision data. The JER uncertainties are defined by comparing both types of smearing and thereby taking the anti-correlation between different components into account. The nominal data remains unchanged. The uncertainty in the efficiency to satisfy the JVT requirement for pile-up suppression was derived in  $Z(\rightarrow \mu^+\mu^-)+\text{jets}$  events and is also considered [100]. The uncertainty in  $E_T^{\text{miss}}$  due to a possible miscalibration of its soft-track component was derived from data–simulation comparisons of the  $p_T$  balance between the hard and soft  $E_T^{\text{miss}}$  components [102].

The  $b$ -tagging requirement made in the measurement has uncertainties in the  $b$ -tagging efficiency of true  $b$ -jets and in the mistagging rates of light-quark jets and  $c$ -jets. The  $b$ -tagging efficiency is measured in dileptonic  $t\bar{t}$  events. Differences between data and detector simulation are corrected by  $p_T$ -dependent scale factors applied to simulated events. The uncertainty in the scale factors is decomposed into 45 orthogonal components [110]. The uncertainties are propagated through the analysis as weights. The set of uncertainties used covers both the 70% working point used in this analysis, as well as other working points of the DL1r discriminant. The rate of mistagging  $c$ -jets as  $b$ -jets was measured in semileptonic  $t\bar{t}$  events, where one of the  $W$  bosons decays into an electron or a muon and a neutrino and the other decays into a quark–antiquark pair [111]. This event sample allows to utilise the relatively large and known  $W \rightarrow cs$  contribution. The mistagging rate of  $c$ -jets depends on the jet  $p_T$  and has a total uncertainty in the range of 3 to 17%. The uncertainties are decomposed into 20 orthogonal components. The misidentification rate of light-quark jets was evaluated based on the techniques described in Ref. [112]. The resulting calibration factors are in the range of about 1.5 – 3.0 with uncertainties up to 50%. The uncertainties are decomposed into 20 independent eigenvectors.

To account for differences between simulation and data in the pile-up distribution, the pile-up profile in the simulation is corrected to match the one in data. The uncertainty in the correction factor is applied in the measurement as a variation of the event weight.

## 8.2 Modelling uncertainties

Normalisation uncertainties are applied to all processes, except for the ones whose normalisation is a free parameter of the maximum-likelihood fit, specifically for the HF-decay templates, for  $t\bar{t}W$  and  $t\bar{t}Z$  production, and for the largest  $VV$  template  $VV3\ell + b/c$ . Thus, no normalisation uncertainty is applied to  $t\bar{t}$ , single-top-quark,  $Z+\text{jets}$  and  $W+\text{jets}$  production, all of which exclusively contribute to non-prompt templates. Minor non-prompt templates are assigned 50% normalisation uncertainties. The same is true for the minor  $VV$  templates and all rare processes discussed in Section 3.2. For  $tZq$  and  $tWZ$  production, a respective uncertainty of 30% is assigned based on a previous ATLAS measurement of the processes [113]. Asymmetric uncertainties of  $^{+7\%}_{-10\%}$  are assigned to  $t\bar{t}H$  production, estimated by varying the renormalisation and factorisation scales, and the combined PDF+ $\alpha_s$  uncertainties [28].

Uncertainties in modelling parton showers and hadronisation are assigned to the  $tHq$  signal samples, and the background originating from  $t\bar{t}$ ,  $t\bar{t}H$ ,  $t\bar{t}Z$  and  $t\bar{t}W$  production. For the signal and  $t\bar{t}$ ,  $t\bar{t}H$  and  $t\bar{t}Z$  production the nominal samples are compared with alternative samples, for which the individual matrix element generators were interfaced to HERWIG 7.1.3 [114, 115] (for  $t\bar{t}$  production), HERWIG 7.0.4 (for  $t\bar{t}H$

production) and HERWIG 7.2.1 (for  $t\bar{t}Z$  production) instead of the respective versions of PYTHIA 8. For  $t\bar{t}W$  production, two alternative samples are generated. The matrix element of both the samples is generated with POWHEG BOX v2, which is interfaced to PYTHIA 8 for one sample and to HERWIG 7 for the other. The relative differences between the two samples is determined for all bins in the analysis and applied to the nominal SHERPA 2.2.10  $t\bar{t}W$  sample.

The impact of the generator choice for  $t\bar{t}W$  is evaluated using an alternative sample and is applied as systematic uncertainties for both the normalisation and shape. This alternative sample is generated with up to one additional parton in the final state at NLO accuracy using MADGRAPH5\_AMC@NLO 2.9.3, with different parton multiplicities being merged using the FxFx NLO matrix-element and parton-shower merging prescription [89] with a merging scale of 30 GeV. The events are interfaced with PYTHIA 8.245 using the A14 set of tuned parameters and the NNPDF2.3LO PDF set to simulate hadronisation and showering. The uncertainty is split into three components: a shape uncertainty in each region, a region migration uncertainty and a global normalisation uncertainty.

Uncertainties related to the choice of the renormalisation scale  $\mu_r$  and the factorisation scale  $\mu_f$  for the matrix-element calculations are evaluated by varying the scales independently by factors of 2 and 0.5, separately for  $VV$  production and each of the top-quark production processes, including production with an additional boson and the two signal processes. The scale variations are implemented as generator weights in the nominal sample. The uncertainties are treated as uncorrelated across individual processes.

The uncertainty in matching the NLO matrix elements to the parton shower when generating  $t\bar{t}$  events is evaluated by comparing the nominal samples of simulated events to samples with an alternative setting of the  $p_T^{\text{hard}}$  parameter in the matching code, using one instead of the default setting of zero. This parameter regulates the definition of the vetoed region of the parton shower, which is needed to avoid overlap in the phase space filled by POWHEG and PYTHIA. This estimate of the uncertainty follows the description in Ref. [116]. The uncertainty in the choice of the  $h_{\text{damp}}$  parameter for the  $t\bar{t}$  event generation is estimated by using an additional  $t\bar{t}$  sample produced with the  $h_{\text{damp}}$  parameter set to  $3m_t$ , while keeping all other generator settings the same as used for the nominal sample of events.

Three uncertainties are defined for the modelling of the Q-misID template. The statistical uncertainty in the Q-misID efficiencies obtained from the likelihood fit is taken into account. An uncertainty in the selection of the  $Z$ -window is calculated by redefining the  $Z$ -window with  $3\sigma$  and  $5\sigma$ , where  $\sigma$  corresponds to the width of the fitted Breit–Wigner distribution. The width of the  $Z$ -window is varied and the efficiencies are determined again, with the symmetrised difference to the nominal values comprising the uncertainty, which is then propagated through the analysis. Finally, a non-closure uncertainty is implemented by applying the Q-misID estimation method to a  $Z \rightarrow ee$  MC sample and comparing the resulting misidentification rates to the MC prediction. The symmetrised difference applied to the nominal Q-misID rates constitutes the uncertainty.

Uncertainties in the amount of initial-state radiation are defined for the top-quark production processes  $t\bar{t}$ ,  $t\bar{t}Z$  and  $t\bar{t}H$ . They are assessed by varying the eigentune Var3c of the A14 PYTHIA 8 tune consistent with the uncertainties of the tune. An uncertainty in the final-state radiation is introduced by varying the renormalisation scale  $\mu_r$  in the parton shower, at which the strong coupling constant  $\alpha_s$  is evaluated, by factors of 0.5 and 2.0.

In all uncertainty evaluations mentioned above the alternative samples or reweighted samples are normalised to the total cross-section of the nominal samples.

Uncertainties in the PDFs are evaluated for the background processes simulated using POWHEG BOX, meaning  $t\bar{t}$ , single- $t$  and  $t\bar{t}H$  production, using the PDF4LHC15 prescription with 30 eigenvectors [117]. Simulated events are reweighted to the central value and the eigenvectors of the combined PDF set. Systematically varied templates are constructed by taking the differences between the samples reweighted to the central value and those reweighted to the eigenvectors. In the likelihood fit, the PDF uncertainties are treated as correlated across the top-quark production processes.

The uncertainties due to the finite number of simulated events, also called the MC statistical uncertainty, is accounted for by adding a nuisance parameter for each bin of the  $D_{\text{NN}}$  distributions and the distributions in the CRs, implementing the Barlow–Beeston approach [118].

## 9 Statistical analysis

The normalisation  $\mu$  of the  $tHu$  and  $tHc$  signal couplings is determined in binned profile maximum-likelihood fits. The fits are performed simultaneously in all signal and control regions. In the signal regions, the respective  $D_{\text{NN}}$  distributions optimised for the considered signal coupling are used. For the CR2 $\ell$ HFe and CR2 $\ell$ HF $\mu$  only event yields are employed in the fit, whereas for the CR3 $\ell$ HFe and CR3 $\ell$ HF $\mu$  the transverse momentum of the third-leading- $p_{\text{T}}$  lepton is fitted. In the CR2 $\ell t\bar{t}V$ , the distribution of the subleading- $p_{\text{T}}$  lepton’s transverse momentum is used, as it provides separation between prompt and non-prompt leptons. The CR3 $\ell t\bar{t}W$  and CR3 $\ell t\bar{t}Z$  are incorporated into the fit with the distribution of the transverse momentum of the leading- $p_{\text{T}}$   $b$ -tagged jet. This distribution separates the  $t\bar{t}Z$  background from remaining  $VV$  events in the CR3 $\ell t\bar{t}Z$  and is also adopted in the CR3 $\ell t\bar{t}W$  for reasons of consistency. In the fits, each of the signal processes are treated as maximally anti-correlated, implying that the presence of one signal automatically precludes the presence of the other signal.

The likelihood function is constructed as a product of Poisson probability terms over all considered bins. The fitted event yields in the bins depend on nuisance parameters  $\theta$  which include the effects of systematic uncertainties. Each nuisance parameter, except those representing the MC statistical uncertainties, is constrained by a Gaussian distribution term in the likelihood function. Some systematically varied discriminant distributions are smoothed and nuisance parameters of systematic uncertainties with negligible impact on the parameter of interest are entirely removed to reduce spurious effects in minimisation, improve convergence of the fit, and reduce the computing time. Normalisation and shape effects from a source of systematic uncertainty are treated separately in this removal process. Single-sided systematic variations are turned into symmetric variations by taking the difference between the nominal model and the alternative model and mirroring this difference in the opposite direction. This is done for the event yield and for the event shape, where the distributions are subtracted from the nominal one. For most sources with two variations, their effects are made symmetric by using the average deviation from the nominal prediction. Exceptions are the uncertainties in the JER, for which the asymmetric variations are kept because the underlying effects are known to be asymmetric. Free-floating normalisation factors are assigned to the templates modelling the HF-decay  $e$  and HF-decay  $\mu$  background, as well as the  $VV3\ell+b/c$ ,  $t\bar{t}W$ , and  $t\bar{t}Z$  templates.

The test statistic  $q_{\mu}$  is defined as the profile likelihood ratio

$$q_{\mu} = -2 \ln \frac{\mathcal{L}(\mu, \hat{\theta})}{\mathcal{L}(\hat{\mu}, \hat{\theta})},$$

where  $\hat{\mu}$  and  $\hat{\theta}$  are the values of the parameters that maximise the likelihood function and  $\hat{\hat{\theta}}$  are the values of the nuisance parameters that maximise the likelihood function for a given value of  $\mu$ . The test statistic is evaluated with the RooFit package [119]. If the observed signal normalisation  $\mu$  is compatible with the background hypothesis, that is  $\mu = 0$ , the test statistic  $q_\mu$  is used in the CL<sub>S</sub> method [120] to obtain exclusion limits on the signal normalisation. For a given signal scenario, values of  $\mu$  yielding CL<sub>S</sub> ≤ 0.05, where CL<sub>S</sub> is computed using the asymptotic approximation [121], are excluded at ≥ 95% CL. The obtained upper limits on the signal strength  $\mu$  are then transformed into limits on the respective Wilson coefficient  $C_{u\phi}$  and the corresponding branching ratio  $\mathcal{B}(t \rightarrow Hq)$  through their dimension-6 operators [60].

## 10 Results

This section presents the results of the profile-likelihood fit. Prior to the full fit to extract the signal normalisation, a background-only fit using only regions with little signal contribution is performed to check the modelling of NN input variables. Once the modelling is checked and validity of the fit model is established, the signal normalisation is determined in a full fit to data using the entire phase space of the analysis.

### 10.1 Cross-checks on the modelling of the neural network input variables

Before the application of the NNs to the full collision data, modelling of the input variables is checked in parts of the signal regions. For this purpose, the events populating the respective signal regions up to a threshold value of  $D_{\text{NN}} = 0.5$  are selected. The threshold of  $D_{\text{NN}} = 0.5$  is chosen to maximise the available statistics while avoiding a signal contamination of more than 10%. The full fit is performed with both statistical and systematic uncertainties considered in all control regions and the low- $D_{\text{NN}}$  signal regions. The corrections made by the fit are then applied to the input variables of the various NNs, using only events with  $D_{\text{NN}} < 0.5$ , to check the variable modelling. The resulting distributions of the most important NN input variables in each signal region are shown in Figure 6. Overall, no discrepancies between the post-fit distributions and data could be observed in any of the input variables.

### 10.2 Full fit to data

Figure 7 shows the  $D_{\text{NN}}(tHc)$  distributions in all four signal regions after the fit. A summary plot of all  $2\ell\text{SS}$  and  $3\ell$  control regions is shown in Figure 8. Overall, a good agreement between MC and data is observed. The event yields shown in Tables 6 and 7 confirm this. The best fit value of the normalisation of the  $tHu$  ( $tHc$ ) signal is found to be  $\mu_{tHq} = -0.03 \pm 0.15$  ( $-0.08 \pm 0.19$ ). No pulls beyond  $1\sigma$  or strong constraints are observed for any of the nuisance parameters. The observed post-fit values of all normalisation factors and nuisance parameters are in agreement with the background-only fit consistent with the statistical uncertainties, suggesting a good modelling of the most signal-sensitive  $D_{\text{NN}}$  bins. The obtained normalisation factors for free-floating background processes are listed in Table 8. For the HF-decay  $e$  and HF-decay  $\mu$  processes they are compatible with one. A 40% increase can be observed for the  $VV3\ell + b/c$  template, owing to the poorer modelling of  $b$ -jets by the shower generator. The normalisations of the  $t\bar{t}W$  and  $t\bar{t}Z$  processes are increased by approximately 15%, which is compatible at a  $1\sigma$  level. The increased normalisation of  $t\bar{t}W$  production is expected based on dedicated measurements of the process, while for  $t\bar{t}Z$ , the slightly higher normalisation comes from the inclusion of events with low jet




Figure 6: Distributions of the most important NN input variable for the signal regions (a) SR2ℓDec, (b) SR2ℓProd, (c) SR3ℓDec and (d) SR3ℓProd. The corrections from a background-only fit are applied. Only events with  $D_{NN} < 0.5$  were used in the fit and in the plots. The first bin contains all events below its lower boundary, while the last bin contains all events falling above its upper boundary. The *Others* template contains various minor processes, which individually only have small contributions. The total statistical and systematic uncertainty is indicated by the hatched band. The dashed line depicts the sum of all background processes prior to the fit. The last bin includes the overflow.



multiplicity. Studies on the process confirm that for higher values of  $N_{\text{jets}}$  the normalisation is compatible with one.

As the best fit value of the signal strength is compatible with  $\mu = 0$ , upper limits are determined with the  $\text{CL}_S$  method. In addition to the nominal limits of the analysis, expected limits are determined for a fit considering only statistical uncertainties, as well as for fits in the  $2\ell\text{SS}$  and  $3\ell$  channels separately. A comparison of the expected upper limits on the branching ratio  $\mathcal{B}(t \rightarrow Hq)$  for each configuration is shown in Table 9. A degradation in the expected upper limit of approximately 20% can be observed for both the signal processes when systematic uncertainties are included. This shows that the sensitivity of this analysis is primarily limited by available statistics. When comparing the two final states with each other, a clear dominance of the  $2\ell\text{SS}$  final state can be observed. Including the  $3\ell$  final states yields an improvement of 20% in the  $tHu$  channel and 13% in the  $tHc$  channel.

The observed upper limits on the branching ratio  $\mathcal{B}(t \rightarrow Hq)$  together with a reinterpretation as limits on the absolute value of the Wilson coefficient  $|C_{u\phi}|$  for the EFT dimension-6 operators can be found in Table 10. The signal does not interfere with other SM processes, meaning that the signal normalisation is proportional to  $|C_{u\phi}|^2$  and limits can only be set on the coefficient's absolute value. Additionally, owing to the averaging of the left-handed and the right-handed signal component, these limits are set on the average of the two respective coefficients  $|C_{u\phi}^{tq}|$  and  $|C_{u\phi}^{qt}|$ . The final observed (expected) upper limits on the branching ratio are  $\mathcal{B}(t \rightarrow Hu) < 2.8 (3.0) \times 10^{-4}$  and  $\mathcal{B}(t \rightarrow Hc) < 3.3 (3.8) \times 10^{-4}$  for a cutoff scale  $\Lambda = 1 \text{ TeV}$ . The limits on the Wilson coefficients amount to  $|C_{u\phi}^{ut,tu}| < 0.71 (0.73)$  and  $|C_{u\phi}^{ct,tc}| < 0.76 (0.82)$ . Limits for each of the two signal couplings are determined with the assumption that the other signal coupling do not contribute.

Compared with a previous ATLAS search for  $tHq$  FCNC couplings in  $2\ell\text{SS}/3\ell$  final states using a partial Run 2 data sample of  $36 \text{ fb}^{-1}$  [32], a luminosity-adjusted improvement by a factor of two for the  $tHc$  channel and three for the  $tHu$  channel can be observed. This is determined by creating an Asimov-data sample, scaling its luminosity to that of the previous analysis, and comparing the resulting expected upper limits to the original analysis. This improvement primarily results from using improved analysis techniques such as new methods for event reconstruction and a more thorough multivariate analysis, and more precise detector calibrations. The additional improvement in the  $tHu$  channel is achieved because of the inclusion of the  $gq \rightarrow Ht$  production process in the analysis. This process primarily plays a role in the  $tHu$  channel because the up quark is a valence quark of the proton.

## 11 Combination of results with other searches

The reported results are combined with the corresponding ATLAS searches that use  $H \rightarrow \tau^+\tau^-$  decays with one or two hadronically decaying  $\tau$ -leptons [33], and  $H \rightarrow b\bar{b}$  [34] and  $H \rightarrow \gamma\gamma$  [35] decays. For clarity, this analysis is referred to as the  $H \rightarrow VV^*$  channel in the following, as the  $H \rightarrow WW^*/ZZ^*$  decay modes are by far the most dominant ones contributing to  $2\ell\text{SS}$  and  $3\ell$  final states. Orthogonality among all analyses is ensured by their respective event selections. The  $H \rightarrow \tau^+\tau^-$  and the  $H \rightarrow VV^*$  analysis, specifically, are orthogonal to each other, because the  $H \rightarrow \tau^+\tau^-$  analysis exclusively uses events with exactly one or zero leptons. The combination is performed through a simultaneous profile-likelihood fit of all four analyses. The correlations between the uncertainties in the different channels are assessed. In each search, the dominant systematic uncertainties are different. In addition, the  $H \rightarrow \tau^+\tau^-$ ,  $H \rightarrow \gamma\gamma$  and  $H \rightarrow VV^*$  channels are dominated by the statistical uncertainty in the data. Therefore, the combination exhibits




Figure 7: The  $D_{NN}$  distributions in the (a) SR2 $l$ Dec, (b) SR2 $l$ Prod, (c) SR3 $l$ Dec and (d) SR3 $l$ Prod, obtained from the signal-plus-background fit to data in the  $tHc$  channel. The *Others* template contains various minor processes, which individually only have small contributions. The total statistical and systematic uncertainty is indicated by the hatched band. The dotted line represents the distribution of the signal, scaled to the number of background events. The dashed line depicts the sum of all background processes prior to the fit.




Figure 8: Summary plot of the fitted distributions in all control regions obtained from the signal-plus-background fit to data in the  $tHc$  channel. The bin boundaries for each region are depicted by the parenthesised numbers. The *Others* template contains various minor processes, which individually only have small contributions. The total statistical and systematic uncertainty is indicated by the hatched band. The dashed line depicts the sum of all background processes prior to the fit.

minimal sensitivity to possible correlations of uncertainties across channels. Specifically, uncertainties pertaining to luminosity, pile-up modelling, and jet energy scale and resolution are correlated among the four channels. The uncertainties related to  $b$ -tagging are correlated between the  $H \rightarrow \tau^+\tau^-$ , the  $H \rightarrow b\bar{b}$  and the  $H \rightarrow VV^*$  analyses, but uncorrelated with the  $H \rightarrow \gamma\gamma$  analysis, which uses a simplified  $b$ -tagging scheme. The remaining uncertainties (mostly from experimental sources, and signal and background modelling) are taken as uncorrelated. Some of the sources of systematic uncertainties (especially related to electron and muon identification and signal modelling) are common to the three search channels, but are treated as uncorrelated due to slight differences in their treatment by individual analyses. The combined  $p$ -value for the  $tHu$  fit is 0.134, while that for the  $tHc$  fit it is 0.117, showing an overall good agreement among the four analyses.

The observed (expected) 95% CL combined upper limits on the branching ratios are  $\mathcal{B}(t \rightarrow Hu) < 2.6 (1.8) \times 10^{-4}$  and  $\mathcal{B}(t \rightarrow Hc) < 3.4 (2.3) \times 10^{-4}$ . For the  $tHc$  coupling the observed limit is larger than that of the individual  $H \rightarrow VV^*$  analysis, because of the strong upward-fluctuation in the  $H \rightarrow \tau^+\tau^-$  analysis. A summary of the upper limits on the branching ratios from the individual searches, and their combination, is given in Figure 9. In the EFT framework, the limits observed for the respective branching ratios translate to a limit on the Wilson coefficients of the  $tHu$  dimension-6 operators of  $|C_{u\phi}^{ut,tu}| < 0.68(0.56)$  at 95% CL, assuming  $C_{u\phi}^{ct,tc} = 0$ , and for a mass scale  $\Lambda = 1$  TeV. The analogous limit on the Wilson coefficient of the  $tHc$  dimension-6 operator amounts to  $|C_{u\phi}^{ct,tc}| < 0.78(0.64)$  assuming  $C_{u\phi}^{ut,tu} = 0$ .

Table 6: The predicted and observed yields in all  $2\ell$ SS regions of the analysis from the signal-plus-background fit to signal and control regions. The pre-fit predictions for the two signal components are presented also, scaled to a cross-section equivalent to a branching ratio  $\mathcal{B}(t \rightarrow Hq) = 0.1\%$ . The *Others* template contains various minor processes, which individually only have small contributions. The uncertainties in non-signal MC yields reflect the sum in quadrature of all post-fit systematic and statistical uncertainties. The row labelled *Post-fit BG* shows the sum of all above background (BG) processes with post-fit uncertainties, while the row labelled *Pre-fit BG* shows the total pre-fit background yield with the corresponding pre-fit uncertainties. Hyphens signify that the corresponding process does not contribute to the given region.

Process	SR2 $\ell$ Dec	SR2 $\ell$ Prod	CR2 $\ell$ HFe	CR2 $\ell$ HFe $\mu$	CR2 $\ell$ t $\bar{t}$ V
HF-decay $e$	$122 \pm 27$	$113 \pm 25$	$66 \pm 13$	–	$2.9 \pm 0.9$
HF-decay $\mu$	$201 \pm 36$	$192 \pm 35$	$0.1 \pm 0.02$	$120 \pm 22$	$5.6 \pm 1.2$
Q-misID	$204 \pm 16$	$457 \pm 35$	$2.4 \pm 0.2$	–	$15.5 \pm 1.4$
$t\bar{t}H$	$132 \pm 20$	$27 \pm 5$	$0.6 \pm 0.1$	$1.0 \pm 0.2$	$51 \pm 8$
$t\bar{t}W$	$512 \pm 61$	$285 \pm 42$	$4.8 \pm 0.9$	$7.5 \pm 1.4$	$216 \pm 24$
$t\bar{t}Z$	$210 \pm 21$	$66 \pm 9$	$1.5 \pm 0.2$	$2.4 \pm 0.4$	$70 \pm 6$
VV $3\ell+b/c$	$104 \pm 20$	$192 \pm 32$	$4.7 \pm 1.0$	$6.6 \pm 1.4$	$6.0 \pm 1.2$
$tWZ$	$23 \pm 7$	$12 \pm 4$	$0.11 \pm 0.04$	$0.17 \pm 0.06$	$3.6 \pm 1.1$
$tZq$	$26 \pm 8$	$63 \pm 18$	$0.7 \pm 0.2$	$1.1 \pm 0.3$	$5.8 \pm 1.7$
Others	$340 \pm 64$	$322 \pm 46$	$36 \pm 8$	$59 \pm 20$	$79 \pm 14$
Pre-fit BG	$1845 \pm 91$	$1585 \pm 70$	$111 \pm 11$	$210 \pm 32$	$424 \pm 22$
Post-fit BG	$1874 \pm 38$	$1729 \pm 36$	$117 \pm 10$	$198 \pm 12$	$455 \pm 17$
$t\bar{t}(t \rightarrow Hu)$	$207 \pm 22$	$181 \pm 10$	$3.4 \pm 0.3$	$5.4 \pm 0.7$	$6.8 \pm 0.6$
$ug \rightarrow Ht$	$31 \pm 4$	$68 \pm 2$	$1.2 \pm 0.1$	$2.1 \pm 0.2$	$1.1 \pm 0.1$
$t\bar{t}(t \rightarrow Hc)$	$196 \pm 22$	$180 \pm 10$	$3.5 \pm 0.4$	$5.9 \pm 0.7$	$13.4 \pm 1.5$
$cg \rightarrow Ht$	$5 \pm 1$	$11 \pm 1$	$0.2 \pm 0.1$	$0.4 \pm 0.1$	$0.2 \pm 0.1$
Data	1847	1723	116	193	443

Table 7: The predicted and observed yields in all  $3\ell$  regions of the analysis from the signal-plus-background fit to signal and control regions. The pre-fit predictions for the two signal components are presented as well, scaled to a cross-section equivalent to a branching ratio  $\mathcal{B}(t \rightarrow Hq) = 0.1\%$ . The *Others* template contains various minor processes, which individually only have small contributions. The uncertainties in non-signal MC yields reflect the sum in quadrature of all post-fit systematic and statistical uncertainties. The row labelled *Post-fit BG* shows the sum of all above background (BG) processes with post-fit uncertainties, while the row labelled *Pre-fit BG* shows the total pre-fit background yield with the corresponding pre-fit uncertainties. Hyphens signify that the corresponding process does not contribute to the given region.

Process	SR3 $\ell$ Prod	SR3 $\ell$ Dec	CR3 $\ell t\bar{t}W$	CR3 $\ell t\bar{t}Z$	CR3 $\ell HFe$	CR3 $\ell HF\mu$
HF-decay $e$	38 $\pm$ 9	14 $\pm$ 3	1.3 $\pm$ 0.3	0.28 $\pm$ 0.09	53 $\pm$ 11	–
HF-decay $\mu$	63 $\pm$ 11	22 $\pm$ 4	1.6 $\pm$ 0.3	0.37 $\pm$ 0.08	0.2 $\pm$ 0.1	122 $\pm$ 19
Q-misID	–	–	–	–	–	–
$t\bar{t}H$	10 $\pm$ 2	47 $\pm$ 7	32 $\pm$ 5	6.7 $\pm$ 1.1	3.0 $\pm$ 0.5	5.2 $\pm$ 0.9
$t\bar{t}W$	77 $\pm$ 12	80 $\pm$ 10	98 $\pm$ 16	12.5 $\pm$ 1.6	5.8 $\pm$ 1.0	9.5 $\pm$ 1.4
$t\bar{t}Z$	75 $\pm$ 11	438 $\pm$ 40	78 $\pm$ 7	261 $\pm$ 20	14.7 $\pm$ 1.8	28 $\pm$ 3
VV 3 $\ell$ + $b/c$	296 $\pm$ 49	215 $\pm$ 39	4.8 $\pm$ 0.9	27 $\pm$ 5	15 $\pm$ 3	30 $\pm$ 5
$tWZ$	19 $\pm$ 6	57 $\pm$ 18	2.9 $\pm$ 0.9	16 $\pm$ 5	1.9 $\pm$ 0.6	3.6 $\pm$ 1.1
$tZq$	134 $\pm$ 38	69 $\pm$ 20	3.5 $\pm$ 1.0	35 $\pm$ 10	6.1 $\pm$ 1.8	12 $\pm$ 3
Others	171 $\pm$ 32	119 $\pm$ 23	43 $\pm$ 7	11.7 $\pm$ 1.5	59 $\pm$ 8	48 $\pm$ 9
Pre-fit BG	710 $\pm$ 48	941 $\pm$ 42	228 $\pm$ 17	312 $\pm$ 19	148 $\pm$ 11	248 $\pm$ 16
Post-fit BG	882 $\pm$ 28	1061 $\pm$ 28	265 $\pm$ 14	371 $\pm$ 17	159 $\pm$ 10	258 $\pm$ 14
$t\bar{t}(t \rightarrow Hu)$	26 $\pm$ 2	39 $\pm$ 3	1.2 $\pm$ 0.2	0.7 $\pm$ 0.1	4.6 $\pm$ 0.5	8.0 $\pm$ 0.8
$ug \rightarrow Ht$	14 $\pm$ 1	7 $\pm$ 1	0.4 $\pm$ 0.1	0.2 $\pm$ 0.1	1.2 $\pm$ 0.1	2.1 $\pm$ 0.2
$t\bar{t}(t \rightarrow Hc)$	27 $\pm$ 2	37 $\pm$ 3	4.3 $\pm$ 0.4	1.5 $\pm$ 0.1	4.4 $\pm$ 0.5	7.8 $\pm$ 0.7
$cg \rightarrow Ht$	2 $\pm$ 1	1 $\pm$ 1	0.1 $\pm$ 0.1	0.1 $\pm$ 0.1	0.2 $\pm$ 0.1	0.4 $\pm$ 0.1
Data	896	1046	268	381	159	263

Table 8: Normalisation factors for various backgrounds determined from the signal-plus-background fit to signal and control regions. The results for the  $tHu$  fit and the  $tHc$  fit are shown.

Process	$tHu$ fit	$tHc$ fit
HF-decay $e$	1.05 $\pm$ 0.24	1.02 $\pm$ 0.23
HF-decay $\mu$	0.94 $\pm$ 0.18	0.92 $\pm$ 0.18
VV3 $\ell$ + $b/c$	1.41 $\pm$ 0.23	1.37 $\pm$ 0.24
$t\bar{t}W$	1.15 $\pm$ 0.14	1.19 $\pm$ 0.14
$t\bar{t}Z$	1.16 $\pm$ 0.11	1.17 $\pm$ 0.11

Table 9: The expected 95% upper limits on the branching ratio  $\mathcal{B}(t \rightarrow Hq)$  for the nominal and alternative fit configurations. One fit is performed in the full phase space considering only statistical uncertainties. Two other fits are performed using the full set of uncertainties in only the  $2\ell$ SS or  $3\ell$  final states. The  $\pm 1\sigma$  interval of the expected limit is indicated by the upper and lower indices. For the  $tHu$  channel, the assumption of  $\mathcal{B}(t \rightarrow Hc) = 0$  is made and vice versa.

Fit configuration	Expected 95% CL upper limits / $10^{-4}$	
	$\mathcal{B}(t \rightarrow Hu)$	$\mathcal{B}(t \rightarrow Hc)$
Nominal fit	$3.0^{+1.2}_{-0.8}$	$3.8^{+1.5}_{-1.1}$
Statistical uncertainties only	$2.6^{+1.1}_{-0.7}$	$3.3^{+1.2}_{-1.0}$
$2\ell$ SS final state only	$3.6^{+1.5}_{-1.0}$	$4.3^{+1.9}_{-1.2}$
$3\ell$ final state only	$6.5^{+2.7}_{-1.9}$	$8.9^{+3.7}_{-2.6}$

Table 10: The observed (expected) 95% upper limits on the branching ratio  $\mathcal{B}(t \rightarrow Hq)$  and the absolute value of the Wilson coefficient  $|C_{u\phi}|$  under the assumption of a cutoff-scale of  $\Lambda = 1$  TeV. For the  $tHu$  channel, the assumption of  $\mathcal{B}(t \rightarrow Hc) = 0$  is made and vice versa.

Signal	Observed (expected) 95% CL upper limits	
	$\mathcal{B}(t \rightarrow Hq)$	$ C_{u\phi}^{qt,tq} $
$tHu$	$2.8 (3.0) \times 10^{-4}$	0.71 (0.73)
$tHc$	$3.3 (3.8) \times 10^{-4}$	0.76 (0.82)




Figure 9: The 95% CL upper limits on (a)  $\mathcal{B}(t \rightarrow Hu)$  assuming  $\mathcal{B}(t \rightarrow Hc) = 0$  and (b)  $\mathcal{B}(t \rightarrow Hc)$  assuming  $\mathcal{B}(t \rightarrow Hu) = 0$  for the individual searches and their combination. The observed limits (solid lines) are compared with the expected (median) limits under the background-only hypothesis (dotted lines). The surrounding shaded bands correspond to the 68% and 95% CL intervals around the expected limits, denoted by  $\pm 1\sigma$  and  $\pm 2\sigma$ , respectively. The  $H \rightarrow VV^*$  also includes events from leptonic  $H \rightarrow \tau^+\tau^-$  decays.

## 12 Conclusion

A search is reported for FCNC couplings between the top quark, the Higgs boson and a second up-type quark in final states containing either two leptons of the same charge or three leptons. This search uses  $140 \text{ fb}^{-1}$  of proton–proton collision data collected with the ATLAS detector at the LHC between 2015 and 2018. The FCNC processes are considered in  $t\bar{t}$  production where either the top quark or the top antiquark decays via  $t \rightarrow Hq$ , and in the production of a single top quark with a Higgs boson via the  $q \rightarrow tH$  FCNC process. The results are compatible with the SM and no evidence of FCNC couplings is observed. Upper limits at 95% CL are set on the branching ratio  $\mathcal{B}(t \rightarrow Hq)$ . The observed (expected) upper limits on the branching ratio are  $\mathcal{B}(t \rightarrow Hu) < 2.8 (3.0) \times 10^{-4}$  and  $\mathcal{B}(t \rightarrow Hc) < 3.3 (3.8) \times 10^{-4}$ . They are reinterpreted as limits on the average of the Wilson coefficients for the left-handed and the right-handed dimension-6 operators modelling the effective  $tHq$  couplings with a new physics scale at  $\Lambda = 1 \text{ TeV}$ . These limits amount to  $|C_{u\phi}^{ut,tu}| < 0.71 (0.73)$  and  $|C_{u\phi}^{ct,tc}| < 0.76 (0.82)$ . These are the most stringent upper limits reported by any analysis searching for  $tHq$  FCNC couplings with the ATLAS detector. They are also lower than any other search for these couplings in  $2\ell\text{SS}$  and  $3\ell$  final states.

The results are combined with three other searches for  $tHq$  FCNC couplings with the ATLAS detector. The combined observed (expected) limits set by all four analyses, considering correlations between the individual searches, are  $\mathcal{B}(t \rightarrow Hu) < 2.6 (1.8) \times 10^{-4}$  and  $\mathcal{B}(t \rightarrow Hc) < 3.4 (2.3) \times 10^{-4}$  on the branching ratios, and  $|C_{u\phi}^{ut,tu}| < 0.68(0.56)$  and  $|C_{u\phi}^{ct,tc}| < 0.78(0.64)$  on the dimension-6 Wilson coefficients.

## Acknowledgements

We thank CERN for the very successful operation of the LHC and its injectors, as well as the support staff at CERN and at our institutions worldwide without whom ATLAS could not be operated efficiently.

The crucial computing support from all WLCG partners is acknowledged gratefully, in particular from CERN, the ATLAS Tier-1 facilities at TRIUMF/SFU (Canada), NDGF (Denmark, Norway, Sweden), CC-IN2P3 (France), KIT/GridKA (Germany), INFN-CNAF (Italy), NL-T1 (Netherlands), PIC (Spain), RAL (UK) and BNL (USA), the Tier-2 facilities worldwide and large non-WLCG resource providers. Major contributors of computing resources are listed in Ref. [122].

We gratefully acknowledge the support of ANPCyT, Argentina; YerPhI, Armenia; ARC, Australia; BMWFW and FWF, Austria; ANAS, Azerbaijan; CNPq and FAPESP, Brazil; NSERC, NRC and CFI, Canada; CERN; ANID, Chile; CAS, MOST and NSFC, China; Minciencias, Colombia; MEYS CR, Czech Republic; DNRF and DNSRC, Denmark; IN2P3-CNRS and CEA-DRF/IRFU, France; SRNSFG, Georgia; BMBF, HGF and MPG, Germany; GSRI, Greece; RGC and Hong Kong SAR, China; ISF and Benoziyo Center, Israel; INFN, Italy; MEXT and JSPS, Japan; CNRST, Morocco; NWO, Netherlands; RCN, Norway; MEiN, Poland; FCT, Portugal; MNE/IFA, Romania; MESTD, Serbia; MSSR, Slovakia; ARRS and MIZŠ, Slovenia; DSI/NRF, South Africa; MICINN, Spain; SRC and Wallenberg Foundation, Sweden; SERI, SNSF and Cantons of Bern and Geneva, Switzerland; MOST, Taipei; TENMAK, Türkiye; STFC, United Kingdom; DOE and NSF, United States of America.

Individual groups and members have received support from BCKDF, CANARIE, CRC and DRAC, Canada; CERN-CZ, PRIMUS 21/SCI/017 and UNCE SCI/013, Czech Republic; COST, ERC, ERDF, Horizon 2020, ICSC-NextGenerationEU and Marie Skłodowska-Curie Actions, European Union; Investissements d’Avenir

Labex, Investissements d'Avenir Idex and ANR, France; DFG and AvH Foundation, Germany; Herakleitos, Thales and Aristeia programmes co-financed by EU-ESF and the Greek NSRF, Greece; BSF-NSF and MINERVA, Israel; Norwegian Financial Mechanism 2014-2021, Norway; NCN and NAWA, Poland; La Caixa Banking Foundation, CERCA Programme Generalitat de Catalunya and PROMETEO and GenT Programmes Generalitat Valenciana, Spain; Göran Gustafssons Stiftelse, Sweden; The Royal Society and Leverhulme Trust, United Kingdom.

In addition, individual members wish to acknowledge support from CERN: European Organization for Nuclear Research (CERN PJAS); Chile: Agencia Nacional de Investigación y Desarrollo (FONDECYT 1190886, FONDECYT 1210400, FONDECYT 1230812, FONDECYT 1230987); China: Chinese Ministry of Science and Technology (MOST-2023YFA1605700), National Natural Science Foundation of China (NSFC - 12175119, NSFC 12275265, NSFC-12075060); Czech Republic: PRIMUS Research Programme (PRIMUS/21/SCI/017); EU: H2020 European Research Council (ERC - 101002463); European Union: European Research Council (ERC - 948254, ERC 101089007), Horizon 2020 Framework Programme (MUCCA - CHIST-ERA-19-XAI-00), European Union, Future Artificial Intelligence Research (FAIR-NextGenerationEU PE00000013), Italian Center for High Performance Computing, Big Data and Quantum Computing (ICSC, NextGenerationEU); France: Agence Nationale de la Recherche (ANR-20-CE31-0013, ANR-21-CE31-0013, ANR-21-CE31-0022), Investissements d'Avenir Labex (ANR-11-LABX-0012); Germany: Baden-Württemberg Stiftung (BW Stiftung-Postdoc Eliteprogramme), Deutsche Forschungsgemeinschaft (DFG - 469666862, DFG - CR 312/5-2); Italy: Istituto Nazionale di Fisica Nucleare (ICSC, NextGenerationEU); Japan: Japan Society for the Promotion of Science (JSPS KAKENHI JP21H05085, JSPS KAKENHI JP22H01227, JSPS KAKENHI JP22H04944, JSPS KAKENHI JP22KK0227); Netherlands: Netherlands Organisation for Scientific Research (NWO Veni 2020 - VI.Veni.202.179); Norway: Research Council of Norway (RCN-314472); Poland: Polish National Agency for Academic Exchange (PPN/PPO/2020/1/00002/U/00001), Polish National Science Centre (NCN 2021/42/E/ST2/00350, NCN OPUS nr 2022/47/B/ST2/03059, NCN UMO-2019/34/E/ST2/00393, UMO-2020/37/B/ST2/01043, UMO-2021/40/C/ST2/00187, UMO-2022/47/O/ST2/00148); Slovenia: Slovenian Research Agency (ARIS grant J1-3010); Spain: BBVA Foundation (LEO22-1-603), Generalitat Valenciana (Artemisa, FEDER, IDIFEDER/2018/048), Ministry of Science and Innovation (MCIN & NextGenEU PCI2022-135018-2, MICIN & FEDER PID2021-125273NB, RYC2019-028510-I, RYC2020-030254-I, RYC2021-031273-I, RYC2022-038164-I), PROMETEO and GenT Programmes Generalitat Valenciana (CIDEGENT/2019/023, CIDEGENT/2019/027); Sweden: Swedish Research Council (VR 2018-00482, VR 2022-03845, VR 2022-04683, VR grant 2021-03651), Knut and Alice Wallenberg Foundation (KAW 2017.0100, KAW 2018.0157, KAW 2018.0458, KAW 2019.0447, KAW 2022.0358); Switzerland: Swiss National Science Foundation (SNSF - PCEFP2\_194658); United Kingdom: Leverhulme Trust (Leverhulme Trust RPG-2020-004), Royal Society (NIF-R1-231091); United States of America: U.S. Department of Energy (ECA DE-AC02-76SF00515), Neubauer Family Foundation.



## Appendix

Tables 11 to 14 list the input variables to the NNs used in the four signal regions, approximately ordered by the increase in significance provided by each variable. The exact order differs among the two signal processes.

Table 11: List of input variables to the NN in the SR3 $\ell$ Prod, approximately ordered by the increase in significance provided by each variable. The exact order differs among the two signal processes.

Variable	Description
$m(\ell_{OS}, \ell_{SS,1})$	Invariant mass of the opposite-charge and the subleading- $p_T$ same-charge lepton
$m(\ell_{OS}, \ell_{SS,0})$	Invariant mass of the opposite-charge and the leading- $p_T$ same-charge lepton
$m(\ell_t, b_t)$	Invariant mass of the $b$ -tagged jet and the lepton assigned to the top-quark decay
$N_{\text{jets}}$	Number of jets
$H_T(\text{jets})$	Scalar sum of the $p_T$ of all jets
$m(t_{SM}, H)$	Invariant mass of the RJR top quark decaying via $t \rightarrow Wb$ and the Higgs boson
$\Delta R(\ell_{SS,0}, \ell_{SS,1})$	Angular separation between the leading and subleading- $p_T$ same-charge lepton
$m(\ell_{H,0}, \ell_{H,1})$	Invariant mass of the two leptons assigned to the Higgs-boson decay
$m(b\text{-jet}, \ell_{SS,0})$	Invariant mass of the $b$ -tagged jet and the leading- $p_T$ same-charge lepton
$\Delta R(\ell_t, b_t)$	Angular separation between the $b$ -tagged jet and the lepton assigned to the top-quark decay
$p_T(t_{SM})$	Transverse momentum of the RJR top quark decaying via $t \rightarrow Wb$
$p_T(b\text{-jet})$	Transverse momentum of the $b$ -tagged jet
$\eta(\ell_{SS,1})$	Pseudorapidity of the subleading- $p_T$ same-charge lepton
$p_T(\ell_{SS,1})$	Transverse momentum of the subleading- $p_T$ same-charge lepton
$m(H, \ell_{SS,1})$	Invariant mass of the RJR Higgs boson and the subleading- $p_T$ same-charge lepton
$\Delta R(t_{SM}, \ell_{OS})$	Angular separation between the RJR top quark decaying via $t \rightarrow Wb$ and the opposite-charge lepton
$\Delta R(H, \ell_{OS})$	Angular separation between the RJR Higgs boson and the opposite-charge lepton
$\Delta R(\ell_{OS}, \ell_{SS,1})$	Angular separation between the opposite-charge and the subleading- $p_T$ same-charge lepton

Table 12: List of input variables to the NN in the SR3 $\ell$ Dec, approximately ordered by the increase in significance provided by each variable. The exact order differs among the two signal processes. Variables labelled NICE were reconstructed with a fulfilled NICE Reco condition.

Variable	Description
$m(\ell_{\text{OS}}, \ell_{\text{SS},1})$	Invariant mass of the opposite-charge and the subleading- $p_{\text{T}}$ same-charge lepton
$m(\ell_{\text{OS}}, \ell_{\text{SS},0})$	Invariant mass of the opposite-charge and the leading- $p_{\text{T}}$ same-charge lepton
NICE $m(\ell_t, b_t)$	Invariant mass of the $b$ -tagged jet and the lepton assigned to the top-quark decay with a fulfilled NICE Reco condition
$H_{\text{T}}(\text{jets})$	Scalar sum of the $p_{\text{T}}$ of all jets
$m(b\text{-jet}, \ell_{\text{SS},0})$	Invariant mass of the $b$ -tagged jet and the leading- $p_{\text{T}}$ same-charge lepton
$m(t_{\text{SM}}, H)$	Invariant mass of the RJR top quark decaying via $t \rightarrow Wb$ and the RJR Higgs boson
$m(\ell_{H,0}, \ell_{H,1})$	Invariant mass of the two leptons assigned to the Higgs-boson decay
$m(H, \ell_{\text{SS},1})$	Invariant mass of the RJR Higgs boson and the subleading- $p_{\text{T}}$ same-charge lepton
$\Delta R(b\text{-jet}, t_{\text{SM}})$	Angular separation between the $b$ -tagged jet and the RJR top quark decaying via $t \rightarrow Wb$
$m(\ell_0, t_{\text{SM}})$	Invariant mass of the leading- $p_{\text{T}}$ lepton and the RJR top quark decaying via $t \rightarrow Wb$
$p_{\text{T}}(t_{\text{SM}})$	Transverse momentum of the RJR top quark decaying via $t \rightarrow Wb$
$m(t_{\text{SM}}, \ell_{\text{SS},1})$	Invariant mass of the RJR top quark decaying via $t \rightarrow Wb$ and the subleading- $p_{\text{T}}$ same-charge lepton
$\Delta R(\ell_{\text{OS}}, \ell_{\text{SS},0})$	Angular separation between the opposite-charge and the leading- $p_{\text{T}}$ same-charge lepton
$p_{\text{T}}(\ell_{\text{OS}})$	Transverse momentum of the opposite-charge lepton
$m(b\text{-jet}, \ell_{\text{OS}})$	Invariant mass of the $b$ -tagged jet and the opposite-charge lepton
$m(b\text{-jet}, H)$	Invariant mass of the $b$ -tagged jet and the RJR Higgs boson
$p_{\text{T}}(\ell_2)$	Transverse momentum of the third-leading- $p_{\text{T}}$ lepton
$\eta(\ell_0)$	Pseudorapidity of the leading- $p_{\text{T}}$ lepton
$m(W_t)$	Mass of the RJR $W$ boson from the top-quark decay
$m(\ell_t, b_t)$	Invariant mass of the $b$ -tagged jet and the lepton assigned to the top-quark decay

Table 13: List of input variables to the NN in the SR2 $\ell$ Prod, approximately ordered by the increase in significance provided by each variable. The exact order differs among the various signal processes.

Variable	Description
$m(\ell_1, H)$	Invariant mass of the subleading- $p_T$ lepton and the RJR Higgs boson
$N_{\text{jets}}$	Number of jets
$m(b\text{-jet}, t_{\text{SM}})$	Invariant mass of the $b$ -tagged jet and the RJR top quark decaying via $t \rightarrow Wb$
$m(H, b\text{-jet})$	Invariant mass of the RJR Higgs boson and the $b$ -tagged jet
$p_T(W_{\text{had}})$	Transverse momentum of the hadronically decaying RJR $W$ boson
$\Delta R(\ell_1, H)$	Angular separation between the subleading- $p_T$ lepton and the RJR Higgs boson
$m(W_{\text{had}})$	Mass of the hadronically decaying RJR $W$ boson
$p_T(\ell_1)$	Transverse momentum of the subleading- $p_T$ lepton
$\eta(\ell_1)$	Pseudorapidity of the subleading- $p_T$ lepton
$\Delta R(H, W_t)$	Angular separation between the RJR Higgs boson and the RJR $W$ boson from the top-quark decay
$\Delta R(\ell_0, \ell_1)$	Angular separation between leading and subleading- $p_T$ lepton
$m(\ell_1, b\text{-jet})$	Invariant mass of the subleading- $p_T$ lepton and the $b$ -tagged jet
$\eta(b\text{-jet})$	Pseudorapidity of the $b$ -tagged jet
$\Delta R(\ell_0, t_{\text{SM}})$	Angular separation between the leading- $p_T$ lepton and the RJR top quark decaying via $t \rightarrow Wb$
$E_{\text{T}}^{\text{miss}}$	Missing transverse momentum
$\text{fl.}(\ell_0)$	Flavour of the leading- $p_T$ lepton
$\eta(\ell_0)$	Pseudorapidity of the leading- $p_T$ lepton
$p_T(\ell_0)$	Transverse momentum of the leading- $p_T$ lepton
$\Delta R(\ell_1, t_{\text{SM}})$	Angular separation between the subleading- $p_T$ lepton and the RJR top quark decaying via $t \rightarrow Wb$
$m(H, W_t)$	Invariant mass of the RJR Higgs boson and the RJR $W$ boson from the top-quark decay
$\Delta R(\ell_1, W_t)$	Angular separation between the subleading- $p_T$ lepton and the RJR $W$ boson from the top-quark decay
$m(\ell_0, H)$	Invariant mass of the leading- $p_T$ lepton and the RJR Higgs boson
$p_T(b\text{-jet})$	Transverse momentum of the $b$ -tagged jet

Table 14: List of input variables to the NN in the SR2 $\ell$ Dec, approximately ordered by the increase in significance provided by each variable. The exact order differs among the various signal processes.

Variable	Description
$H_T(\text{jets})$	Scalar sum of the $p_T$ of all jets
$m(\ell_0, b\text{-jet})$	Invariant mass of the leading- $p_T$ lepton and the $b$ -tagged jet
$\Delta R(\ell_1, H)$	Angular separation between the subleading- $p_T$ lepton and the RJR Higgs boson
$p_T(\ell_1)$	Transverse momentum of the subleading- $p_T$ lepton
$m(\text{jets}_{\min\Delta R})$	Invariant mass of the two non- $b$ -tagged jets with the smallest $\Delta R$
$m(t_{\text{SM}}, l\text{-jet}_0)$	Invariant mass of the RJR top quark decaying via $t \rightarrow Wb$ and the leading- $p_T$ non- $b$ -tagged jet
$\eta(\ell_1)$	Pseudorapidity of the subleading- $p_T$ lepton
$\Delta R(\ell_0, l\text{-jet}_1)$	Angular separation between the leading- $p_T$ lepton and the subleading- $p_T$ non- $b$ -tagged jet
$m(\ell_1, l\text{-jet}_0)$	Invariant mass of the subleading- $p_T$ lepton and the leading- $p_T$ non- $b$ -tagged jet
$m(\ell_0, l\text{-jet}_0)$	Invariant mass of the leading- $p_T$ lepton and the leading- $p_T$ non- $b$ -tagged jet
$\Delta R(\ell_0, l\text{-jet}_2)$	Angular separation between the leading- $p_T$ lepton and the third-leading- $p_T$ non- $b$ -tagged jet
$\Delta R(\ell_1, l\text{-jet}_2)$	Angular separation between the subleading- $p_T$ lepton and the third-leading- $p_T$ non- $b$ -tagged jet
$m(t_{\text{FCNC}}, l\text{-jet}_0)$	Invariant mass of the RJR top quark decaying via $t \rightarrow Hq$ and the leading- $p_T$ non- $b$ -tagged jet
$m(\ell_1, l\text{-jet}_1)$	Invariant mass of the subleading- $p_T$ lepton and the subleading- $p_T$ non- $b$ -tagged jet
$m(\ell_1, t_{\text{FCNC}})$	Invariant mass of the subleading- $p_T$ lepton and the RJR top quark decaying via $t \rightarrow Hq$
$m(W_t, W_{\text{had}})$	Invariant mass of the RJR $W$ boson from the top-quark decay and the hadronically decaying RJR $W$ boson
$\Delta R(\ell_0, l\text{-jet}_0)$	Angular separation between the leading- $p_T$ lepton and the leading- $p_T$ non- $b$ -tagged jet
$m(\ell_1, b\text{-jet})$	Invariant mass of the subleading- $p_T$ lepton and the $b$ -tagged jet
$N_{\text{jets}}$	Number of jets
$m(H, b\text{-jet})$	Invariant mass of the RJR Higgs boson and the $b$ -tagged jet
$H_T(\ell_0, \ell_1)$	Scalar sum of the $p_T$ of all leptons
$p_T(\ell_0)$	Transverse momentum of the leading- $p_T$ lepton
$m(W_t, t_{\text{FCNC}})$	Invariant mass of the RJR $W$ boson from the top-quark decay and the RJR top quark decaying via $t \rightarrow Hq$

## References

- [1] ATLAS Collaboration, *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, *Phys. Lett. B* **716** (2012) 1, arXiv: [1207.7214 \[hep-ex\]](#).
- [2] CMS Collaboration, *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, *Phys. Lett. B* **716** (2012) 30, arXiv: [1207.7235 \[hep-ex\]](#).
- [3] ATLAS Collaboration, *A detailed map of Higgs boson interactions by the ATLAS experiment ten years after the discovery*, *Nature* **607** (2022) 52, arXiv: [2207.00092 \[hep-ex\]](#), Erratum: *Nature* **612** (2022) E24.
- [4] CMS Collaboration, *A portrait of the Higgs boson by the CMS experiment ten years after the discovery*, *Nature* **607** (2022) 60, arXiv: [2207.00043 \[hep-ex\]](#), Erratum: *Nature* **623** (2023) E4.
- [5] S. L. Glashow, J. Iliopoulos and L. Maiani, *Weak Interactions with Lepton-Hadron Symmetry*, *Phys. Rev. D* **2** (1970) 1285.
- [6] G. Eilam, J. L. Hewett and A. Soni, *Rare decays of the top quark in the standard and two-Higgs-doublet models*, *Phys. Rev. D* **44** (1991) 1473, Erratum: *Phys. Rev. D* **59** (1998) 039901.
- [7] B. Mele, S. Petrarca and A. Soddu, *A new evaluation of the  $t \rightarrow cH$  decay width in the standard model*, *Phys. Lett. B* **435** (1998) 401, arXiv: [hep-ph/9805498 \[hep-ph\]](#).
- [8] J. A. Aguilar-Saavedra, *Top flavor-changing neutral interactions: theoretical expectations and experimental detection*, *Acta Phys. Polon. B* **35** (2004) 2695, arXiv: [hep-ph/0409342 \[hep-ph\]](#).
- [9] C. Zhang and F. Maltoni, *Top-quark decay into Higgs boson and a light quark at next-to-leading order in QCD*, *Phys. Rev. D* **88** (2013) 054005, arXiv: [1305.7386 \[hep-ph\]](#).
- [10] W. Altmannshofer, B. Maddock and D. Tuckler, *Rare top decays as probes of flavorful Higgs bosons*, *Phys. Rev. D* **100** (2019) 015003, arXiv: [1904.10956 \[hep-ph\]](#).
- [11] G. C. Branco et al., *Theory and phenomenology of two-Higgs-doublet models*, *Phys. Rep.* **516** (2012) 1, arXiv: [1106.0034 \[hep-ph\]](#).
- [12] S. B ejar, J. Guasch and J. Sol a, *Loop induced flavor changing neutral decays of the top quark in a general two-Higgs-doublet model*, *Nucl. Phys. B* **600** (2001) 21, arXiv: [hep-ph/0011091 \[hep-ph\]](#).
- [13] J. Guasch and J. Sol a, *FCNC top quark decays in the MSSM: a door to SUSY physics in high luminosity colliders?*, *Nucl. Phys. B* **562** (1999) 3, arXiv: [hep-ph/9906268 \[hep-ph\]](#).
- [14] J. J. Cao et al., *Supersymmetry-induced flavor-changing neutral-current top-quark processes at the CERN Large Hadron Collider*, *Phys. Rev. D* **75** (2007) 075021, arXiv: [hep-ph/0702264 \[hep-ph\]](#).

- [15] J. Cao, C. Han, L. Wu, J. M. Yang and M. Zhang, *SUSY induced top quark FCNC decay  $t \rightarrow ch$  after Run I of LHC*, *Eur. Phys. J. C* **74** (2014) 3058, arXiv: [1404.1241 \[hep-ph\]](#).
- [16] G. Eilam, A. Gemintern, T. Han, J. M. Yang and X. Zhang, *Top-quark rare decay  $t \rightarrow ch$  in R-parity-violating SUSY*, *Phys. Lett. B* **510** (2001) 227, arXiv: [hep-ph/0102037 \[hep-ph\]](#).
- [17] J. A. Aguilar-Saavedra, *Effects of mixing with quark singlets*, *Phys. Rev. D* **67** (2003) 035003, Erratum: *Phys. Rev. D* **69** (2004) 099901, arXiv: [hep-ph/0210112 \[hep-ph\]](#).
- [18] A. Azatov, M. Toharia and L. Zhu, *Higgs mediated flavor changing neutral currents in warped extra dimensions*, *Phys. Rev. D* **80** (2009) 035016, arXiv: [0906.1990 \[hep-ph\]](#).
- [19] T. P. Cheng and M. Sher, *Mass-matrix ansatz and flavor nonconservation in models with multiple Higgs doublets*, *Phys. Rev. D* **35** (1987) 3484.
- [20] I. Baum, G. Eilam and S. Bar-Shalom, *Scalar flavor changing neutral currents and rare top quark decays in a two Higgs doublet model "for the top quark"*, *Phys. Rev. D* **77** (2008) 113008, arXiv: [0802.2622 \[hep-ph\]](#).
- [21] K.-F. Chen, W.-S. Hou, C. Kao and M. Kohda, *When the Higgs meets the top: search for  $t \rightarrow ch^0$  at the LHC*, *Phys. Lett. B* **725** (2013) 378, arXiv: [1304.8037 \[hep-ph\]](#).
- [22] C.-W. Chiang, H. Fukuda, M. Takeuchi and T. T. Yanagida, *Flavor-changing neutral-current decays in top-specific variant axion model*, *JHEP* **11** (2015) 057, arXiv: [1507.04354 \[hep-ph\]](#).
- [23] A. Crivellin, J. Heeck and P. Stoffer, *Perturbed Lepton-Specific Two-Higgs-Doublet Model Facing Experimental Hints for Physics beyond the Standard Model*, *Phys. Rev. Lett.* **116** (2016) 081801, arXiv: [1507.07567 \[hep-ph\]](#).
- [24] F. J. Botella, G. C. Branco, M. Nebot and M. N. Rebelo, *Flavour-changing Higgs couplings in a class of two Higgs doublet models*, *Eur. Phys. J. C* **76** (2016) 161, arXiv: [1508.05101 \[hep-ph\]](#).
- [25] S. Gori, C. Grojean, A. Juste and A. Paul, *Heavy Higgs searches: flavour matters*, *JHEP* **01** (2018) 108, arXiv: [1710.03752 \[hep-ph\]](#).
- [26] C.-W. Chiang, H. Fukuda, M. Takeuchi and T. T. Yanagida, *Current status of top-specific variant axion model*, *Phys. Rev. D* **97** (2018) 035015, arXiv: [1711.02993 \[hep-ph\]](#).
- [27] A. Greljo, J. F. Kamenik and J. Kopp, *Disentangling flavor violation in the top-Higgs sector at the LHC*, *JHEP* **07** (2014) 046, arXiv: [1404.1278 \[hep-ph\]](#).
- [28] D. de Florian et al., *Handbook of LHC Higgs Cross Sections: 4. Deciphering the Nature of the Higgs Sector*, (2017), arXiv: [1610.07922 \[hep-ph\]](#).
- [29] G. Durieux, F. Maltoni and C. Zhang, *Global approach to top-quark flavor-changing interactions*, *Phys. Rev. D* **91** (2015) 074017, arXiv: [1412.7166 \[hep-ph\]](#).

- [30] ATLAS Collaboration, *Search for top-quark decays  $t \rightarrow Hq$  with  $36 \text{ fb}^{-1}$  of  $pp$  collision data at  $\sqrt{s} = 13 \text{ TeV}$  with the ATLAS detector*, *JHEP* **05** (2019) 123, arXiv: [1812.11568 \[hep-ex\]](#).
- [31] ATLAS Collaboration, *Search for top quark decays  $t \rightarrow qH$ , with  $H \rightarrow \gamma\gamma$ , in  $\sqrt{s} = 13 \text{ TeV}$   $pp$  collisions using the ATLAS detector*, *JHEP* **10** (2017) 129, arXiv: [1707.01404 \[hep-ex\]](#).
- [32] ATLAS Collaboration, *Search for flavor-changing neutral currents in top quark decays  $t \rightarrow Hc$  and  $t \rightarrow Hu$  in multilepton final states in proton–proton collisions at  $\sqrt{s} = 13 \text{ TeV}$  with the ATLAS detector*, *Phys. Rev. D* **98** (2018) 032002, arXiv: [1805.03483 \[hep-ex\]](#).
- [33] ATLAS Collaboration, *Search for flavour-changing neutral current interactions of the top quark and the Higgs boson in events with a pair of  $\tau$ -leptons in  $pp$  collisions at  $\sqrt{s} = 13 \text{ TeV}$  with the ATLAS detector*, *JHEP* **06** (2023) 155, arXiv: [2208.11415 \[hep-ex\]](#).
- [34] ATLAS Collaboration, *Search for a new scalar resonance in flavour-changing neutral-current top-quark decays  $t \rightarrow qX$  ( $q = u, c$ ), with  $X \rightarrow b\bar{b}$ , in proton–proton collisions at  $\sqrt{s} = 13 \text{ TeV}$  with the ATLAS detector*, *JHEP* **07** (2023) 199, arXiv: [2301.03902 \[hep-ex\]](#).
- [35] ATLAS Collaboration, *Search for flavour-changing neutral  $tqH$  interactions with  $H \rightarrow \gamma\gamma$  in  $pp$  collisions at  $\sqrt{s} = 13 \text{ TeV}$  using the ATLAS detector*, *JHEP* **12** (2023) 195, arXiv: [2309.12817 \[hep-ex\]](#).
- [36] CMS Collaboration, *Search for the flavor-changing neutral current interactions of the top quark and the Higgs boson which decays into a pair of  $b$  quarks at  $\sqrt{s} = 13 \text{ TeV}$* , *JHEP* **06** (2018) 102, arXiv: [1712.02399 \[hep-ex\]](#).
- [37] CMS Collaboration, *Search for flavor-changing neutral current interactions of the top quark and the Higgs boson decaying to a bottom quark-antiquark pair at  $\sqrt{s} = 13 \text{ TeV}$* , *JHEP* **02** (2022) 169, arXiv: [2112.09734 \[hep-ex\]](#).
- [38] CMS Collaboration, *Search for flavor-changing neutral current interactions of the top quark and Higgs boson in final states with two photons in proton–proton collisions at  $\sqrt{s} = 13 \text{ TeV}$* , *JHEP* **02** (2022) 169, arXiv: [2111.02219 \[hep-ex\]](#).
- [39] D. E. Rumelhart, G. E. Hinton and R. J. Williams, ‘Learning Internal Representations by Error Propagation’, *Parallel Distributed Processing: Explorations in the Microstructure of Cognition, Volume 1: Foundations*, ed. by D. E. Rumelhart and J. L. McClelland, Cambridge, MA: MIT Press, 1986 318.
- [40] ATLAS Collaboration, *The ATLAS Experiment at the CERN Large Hadron Collider*, *JINST* **3** (2008) S08003.
- [41] ATLAS Collaboration, *ATLAS Insertable B-Layer: Technical Design Report*, ATLAS-TDR-19; CERN-LHCC-2010-013, 2010, URL: <https://cds.cern.ch/record/1291633>, Addendum: ATLAS-TDR-19-ADD-1; CERN-LHCC-2012-009, 2012, URL: <https://cds.cern.ch/record/1451888>.
- [42] B. Abbott et al., *Production and integration of the ATLAS Insertable B-Layer*, *JINST* **13** (2018) T05008, arXiv: [1803.00844 \[physics.ins-det\]](#).
- [43] ATLAS Collaboration, *Performance of the ATLAS trigger system in 2015*, *Eur. Phys. J. C* **77** (2017) 317, arXiv: [1611.09661 \[hep-ex\]](#).
- [44] ATLAS Collaboration, *The ATLAS Collaboration Software and Firmware*, ATL-SOFT-PUB-2021-001, 2021, URL: <https://cds.cern.ch/record/2767187>.

- [45] ATLAS Collaboration, *ATLAS data quality operations and performance for 2015–2018 data-taking*, [JINST \*\*15\*\* \(2020\) P04003](#), arXiv: [1911.04632 \[physics.ins-det\]](#).
- [46] ATLAS Collaboration, *Luminosity determination in  $pp$  collisions at  $\sqrt{s} = 13$  TeV using the ATLAS detector at the LHC*, [Eur. Phys. J. C \*\*83\*\* \(2023\) 982](#), arXiv: [2212.09379 \[hep-ex\]](#).
- [47] G. Avoni et al., *The new LUCID-2 detector for luminosity measurement and monitoring in ATLAS*, [JINST \*\*13\*\* \(2018\) P07017](#).
- [48] ATLAS Collaboration, *Performance of electron and photon triggers in ATLAS during LHC Run 2*, [Eur. Phys. J. C \*\*80\*\* \(2020\) 47](#), arXiv: [1909.00761 \[hep-ex\]](#).
- [49] ATLAS Collaboration, *Performance of the ATLAS muon triggers in Run 2*, [JINST \*\*15\*\* \(2020\) P09015](#), arXiv: [2004.13447 \[physics.ins-det\]](#).
- [50] S. Agostinelli et al., *GEANT4 – a simulation toolkit*, [Nucl. Instrum. Meth. A \*\*506\*\* \(2003\) 250](#).
- [51] ATLAS Collaboration, *The ATLAS Simulation Infrastructure*, [Eur. Phys. J. C \*\*70\*\* \(2010\) 823](#), arXiv: [1005.4568 \[physics.ins-det\]](#).
- [52] ATLAS Collaboration, *The simulation principle and performance of the ATLAS fast calorimeter simulation FastCaloSim*, ATL-PHYS-PUB-2010-013, 2010, URL: <https://cds.cern.ch/record/1300517>.
- [53] T. Sjöstrand et al., *An introduction to PYTHIA 8.2*, [Comput. Phys. Commun. \*\*191\*\* \(2015\) 159](#), arXiv: [1410.3012 \[hep-ph\]](#).
- [54] ATLAS Collaboration, *The Pythia 8 A3 tune description of ATLAS minimum bias and inelastic measurements incorporating the Donnachie–Landshoff diffractive model*, ATL-PHYS-PUB-2016-017, 2016, URL: <https://cds.cern.ch/record/2206965>.
- [55] NNPDF Collaboration, R. D. Ball et al., *Parton distributions with LHC data*, [Nucl. Phys. B \*\*867\*\* \(2013\) 244](#), arXiv: [1207.1303 \[hep-ph\]](#).
- [56] T. Gleisberg et al., *Event generation with SHERPA 1.1*, [JHEP \*\*02\*\* \(2009\) 007](#), arXiv: [0811.4622 \[hep-ph\]](#).
- [57] E. Bothmann et al., *Event generation with Sherpa 2.2*, [SciPost Phys. \*\*7\*\* \(2019\) 034](#), arXiv: [1905.09127 \[hep-ph\]](#).
- [58] D. J. Lange, *The EvtGen particle decay simulation package*, [Nucl. Instrum. Meth. A \*\*462\*\* \(2001\) 152](#).
- [59] ATLAS Collaboration, *ATLAS Pythia 8 tunes to 7 TeV data*, ATL-PHYS-PUB-2014-021, 2014, URL: <https://cds.cern.ch/record/1966419>.
- [60] C. Degrande, F. Maltoni, J. Wang and C. Zhang, *Automatic computations at next-to-leading order in QCD for top-quark flavor-changing neutral processes*, [Phys. Rev. D \*\*91\*\* \(2015\) 034024](#), arXiv: [1412.5594 \[hep-ex\]](#).
- [61] A. Alloul, N. D. Christensen, C. Degrande, C. Duhr and B. Fuks, *FeynRules 2.0 — A complete toolbox for tree-level phenomenology*, [Comput. Phys. Commun. \*\*185\*\* \(2014\) 2250](#), arXiv: [1310.1921 \[hep-ph\]](#).
- [62] P. Nason, *A new method for combining NLO QCD with shower Monte Carlo algorithms*, [JHEP \*\*11\*\* \(2004\) 040](#), arXiv: [hep-ph/0409146](#).



- [63] S. Frixione, G. Ridolfi and P. Nason,  
*A positive-weight next-to-leading-order Monte Carlo for heavy flavour hadroproduction*,  
[JHEP \*\*09\*\* \(2007\) 126](#), arXiv: [0707.3088 \[hep-ph\]](#).
- [64] S. Frixione, P. Nason and C. Oleari,  
*Matching NLO QCD computations with parton shower simulations: the POWHEG method*,  
[JHEP \*\*11\*\* \(2007\) 070](#), arXiv: [0709.2092 \[hep-ph\]](#).
- [65] S. Alioli, P. Nason, C. Oleari and E. Re,  
*NLO single-top production matched with shower in POWHEG: s- and t-channel contributions*,  
[JHEP \*\*09\*\* \(2009\) 111](#), arXiv: [0907.4076 \[hep-ph\]](#), Erratum: [JHEP \*\*02\*\* \(2010\) 011](#).
- [66] S. Alioli, P. Nason, C. Oleari and E. Re, *A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX*, [JHEP \*\*06\*\* \(2010\) 043](#), arXiv: [1002.2581 \[hep-ph\]](#).
- [67] E. Re,  
*Single-top Wt-channel production matched with parton showers using the POWHEG method*,  
[Eur. Phys. J. C \*\*71\*\* \(2011\) 1547](#), arXiv: [1009.2450 \[hep-ph\]](#).
- [68] R. Frederix, E. Re and P. Torrielli,  
*Single-top t-channel hadroproduction in the four-flavour scheme with POWHEG and aMC@NLO*,  
[JHEP \*\*09\*\* \(2012\) 130](#), arXiv: [1207.5391 \[hep-ph\]](#).
- [69] NNPDF Collaboration, R. D. Ball et al., *Parton distributions for the LHC run II*,  
[JHEP \*\*04\*\* \(2015\) 040](#), arXiv: [1410.8849 \[hep-ph\]](#).
- [70] S. Frixione, E. Laenen, P. Motylinski and B. R. Webber, *Angular correlations of lepton pairs from vector boson and top quark decays in Monte Carlo simulations*, [JHEP \*\*04\*\* \(2007\) 081](#), arXiv: [hep-ph/0702198](#).
- [71] P. Artoisenet, R. Frederix, O. Mattelaer and R. Rietkerk,  
*Automatic spin-entangled decays of heavy resonances in Monte Carlo simulations*,  
[JHEP \*\*03\*\* \(2013\) 015](#), arXiv: [1212.3460 \[hep-ph\]](#).
- [72] C. Bierlich et al., *A comprehensive guide to the physics and usage of PYTHIA 8.3*,  
[SciPost Phys. Codeb. \*\*2022\*\* \(2022\) 8](#), arXiv: [2203.11601 \[hep-ph\]](#).
- [73] ATLAS Collaboration, *Studies on top-quark Monte Carlo modelling for Top2016*,  
ATL-PHYS-PUB-2016-020, 2016, URL: <https://cds.cern.ch/record/2216168>.
- [74] J. Alwall et al., *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, [JHEP \*\*07\*\* \(2014\) 079](#), arXiv: [1405.0301 \[hep-ph\]](#).
- [75] M. Czakon and A. Mitov,  
*Top++: A program for the calculation of the top-pair cross-section at hadron colliders*,  
[Comput. Phys. Commun. \*\*185\*\* \(2014\) 2930](#), arXiv: [1112.5675 \[hep-ph\]](#).
- [76] M. Aliev et al., *HATHOR – HAdronic Top and Heavy quarks crOss section calculatoR*,  
[Comput. Phys. Commun. \*\*182\*\* \(2011\) 1034](#), arXiv: [1007.1327 \[hep-ph\]](#).
- [77] P. Kant et al., *HatHor for single top-quark production: Updated predictions and uncertainty estimates for single top-quark production in hadronic collisions*,  
[Comput. Phys. Commun. \*\*191\*\* \(2015\) 74](#), arXiv: [1406.4403 \[hep-ph\]](#).

- [78] N. Kidonakis, *Theoretical results for electroweak boson and single-top production*, [PoS DIS2015 \(2016\) 170](#), arXiv: [1506.04072 \[hep-ph\]](#).
- [79] T. Gleisberg and S. Höche, *Comix, a new matrix element generator*, [JHEP 12 \(2008\) 039](#), arXiv: [0808.3674 \[hep-ph\]](#).
- [80] F. Buccioni et al., *OpenLoops 2*, [Eur. Phys. J. C 79 \(2019\) 866](#), arXiv: [1907.13071 \[hep-ph\]](#).
- [81] F. Cascioli, P. Maierhöfer and S. Pozzorini, *Scattering Amplitudes with Open Loops*, [Phys. Rev. Lett. 108 \(2012\) 111601](#), arXiv: [1111.5206 \[hep-ph\]](#).
- [82] A. Denner, S. Dittmaier and L. Hofer, *COLLIER: A fortran-based complex one-loop library in extended regularizations*, [Comput. Phys. Commun. 212 \(2017\) 220](#), arXiv: [1604.06792 \[hep-ph\]](#).
- [83] S. Schumann and F. Krauss, *A parton shower algorithm based on Catani–Seymour dipole factorisation*, [JHEP 03 \(2008\) 038](#), arXiv: [0709.1027 \[hep-ph\]](#).
- [84] J.-C. Winter, F. Krauss and G. Soff, *A modified cluster-hadronisation model*, [Eur. Phys. J. C 36 \(2004\) 381](#), arXiv: [hep-ph/0311085](#).
- [85] S. Höche, F. Krauss, M. Schönherr and F. Siegert, *QCD matrix elements + parton showers. The NLO case*, [JHEP 04 \(2013\) 027](#), arXiv: [1207.5030 \[hep-ph\]](#).
- [86] S. Kallweit, J. M. Lindert, P. Maierhöfer, S. Pozzorini and M. Schönherr, *NLO QCD+EW predictions for  $V + jets$  including off-shell vector-boson decays and multijet merging*, [JHEP 04 \(2016\) 021](#), arXiv: [1511.08692 \[hep-ph\]](#).
- [87] C. Gütschow, J. M. Lindert and M. Schönherr, *Multi-jet merged top-pair production including electroweak corrections*, [Eur. Phys. J. C 78 \(2018\) 317](#), arXiv: [1803.00950 \[hep-ph\]](#).
- [88] R. Frederix, D. Pagani and M. Zaro, *Large NLO corrections in  $t\bar{t}W^\pm$  and  $t\bar{t}\bar{t}$  hadroproduction from supposedly subleading EW contributions*, [JHEP 02 \(2018\) 031](#), arXiv: [1711.02116 \[hep-ph\]](#).
- [89] R. Frederix and I. Tsinikos, *On improving NLO merging for  $t\bar{t}W$  production*, [JHEP 11 \(2021\) 029](#), arXiv: [2108.07826 \[hep-ph\]](#).
- [90] ATLAS Collaboration, *Vertex Reconstruction Performance of the ATLAS Detector at  $\sqrt{s} = 13$  TeV*, ATL-PHYS-PUB-2015-026, 2015, URL: <https://cds.cern.ch/record/2037717>.
- [91] ATLAS Collaboration, *Electron and photon performance measurements with the ATLAS detector using the 2015–2017 LHC proton–proton collision data*, [JINST 14 \(2019\) P12006](#), arXiv: [1908.00005 \[hep-ex\]](#).
- [92] ATLAS Collaboration, *Electron and photon efficiencies in LHC Run 2 with the ATLAS experiment*, (2023), arXiv: [2308.13362 \[hep-ex\]](#).
- [93] ATLAS Collaboration, *Muon reconstruction and identification efficiency in ATLAS using the full Run 2  $pp$  collision data set at  $\sqrt{s} = 13$  TeV*, [Eur. Phys. J. C 81 \(2021\) 578](#), arXiv: [2012.00578 \[hep-ex\]](#).
- [94] ATLAS Collaboration, *Evidence for the associated production of the Higgs boson and a top quark pair with the ATLAS detector*, [Phys. Rev. D 97 \(2018\) 072003](#), arXiv: [1712.08891 \[hep-ex\]](#).

- [95] ATLAS Collaboration, *Jet reconstruction and performance using particle flow with the ATLAS Detector*, [Eur. Phys. J. C \*\*77\*\* \(2017\) 466](#), arXiv: [1703.10485 \[hep-ex\]](#).
- [96] M. Cacciari, G. P. Salam and G. Soyez, *The anti- $k_t$  jet clustering algorithm*, [JHEP \*\*04\*\* \(2008\) 063](#), arXiv: [0802.1189 \[hep-ph\]](#).
- [97] M. Cacciari, G. P. Salam and G. Soyez, *FastJet user manual*, [Eur. Phys. J. C \*\*72\*\* \(2012\) 1896](#), arXiv: [1111.6097 \[hep-ph\]](#).
- [98] ATLAS Collaboration, *Topological cell clustering in the ATLAS calorimeters and its performance in LHC Run 1*, [Eur. Phys. J. C \*\*77\*\* \(2017\) 490](#), arXiv: [1603.02934 \[hep-ex\]](#).
- [99] ATLAS Collaboration, *Jet energy scale and resolution measured in proton–proton collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector*, [Eur. Phys. J. C \*\*81\*\* \(2021\) 689](#), arXiv: [2007.02645 \[hep-ex\]](#).
- [100] ATLAS Collaboration, *Performance of pile-up mitigation techniques for jets in pp collisions at  $\sqrt{s} = 8$  TeV using the ATLAS detector*, [Eur. Phys. J. C \*\*76\*\* \(2016\) 581](#), arXiv: [1510.03823 \[hep-ex\]](#).
- [101] ATLAS Collaboration, *ATLAS flavour-tagging algorithms for the LHC Run 2 pp collision dataset*, [Eur. Phys. J. C \*\*83\*\* \(2023\) 681](#), arXiv: [2211.16345 \[physics.data-an\]](#).
- [102] ATLAS Collaboration, *The performance of missing transverse momentum reconstruction and its significance with the ATLAS detector using  $140\text{fb}^{-1}$  of  $\sqrt{s} = 13$  TeV pp collisions*, (2024), arXiv: [2402.05858 \[hep-ex\]](#).
- [103] ATLAS Collaboration, *Observation of four-top-quark production in the multilepton final state with the ATLAS detector*, [Eur. Phys. J. C \*\*83\*\* \(2023\) 496](#), arXiv: [2303.15061 \[hep-ex\]](#).
- [104] Particle Data Group, M. Tanabashi et al., *Review of Particle Physics*, [Phys. Rev. D \*\*98\*\* \(2018\) 030001](#).
- [105] ATLAS Collaboration, *Measurement of the total and differential cross-sections of  $t\bar{t}W$  production in pp collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector*, 2024, arXiv: [2401.05299 \[hep-ex\]](#).
- [106] ATLAS Collaboration, *Inclusive and differential cross-section measurements of  $t\bar{t}Z$  production in pp collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector, including EFT and spin-correlation interpretations*, 2023, arXiv: [2312.04450 \[hep-ex\]](#).
- [107] P. Jackson and C. Rogan, *Recursive Jigsaw Reconstruction: HEP event analysis in the presence of kinematic and combinatoric ambiguities*, [Phys. Rev. D \*\*96\*\* \(2017\) 112007](#), arXiv: [1705.10733 \[hep-ph\]](#).
- [108] M. Feindt, *A Neural Bayesian Estimator for Conditional Probability Densities*, 2004, arXiv: [physics/0402093 \[physics.data-an\]](#).
- [109] M. Feindt and U. Kerzel, *The NeuroBayes neural network package*, [Nucl. Instrum. Meth. A \*\*559\*\* \(2006\) 190](#).
- [110] ATLAS Collaboration, *ATLAS b-jet identification performance and efficiency measurement with  $t\bar{t}$  events in pp collisions at  $\sqrt{s} = 13$  TeV*, [Eur. Phys. J. C \*\*79\*\* \(2019\) 970](#), arXiv: [1907.05120 \[hep-ex\]](#).

- [111] ATLAS Collaboration, *Measurement of the  $c$ -jet mistagging efficiency in  $t\bar{t}$  events using  $pp$  collision data at  $\sqrt{s} = 13$  TeV collected with the ATLAS detector*, *Eur. Phys. J. C* **82** (2022) 95, arXiv: [2109.10627](https://arxiv.org/abs/2109.10627) [[hep-ex](#)].
- [112] ATLAS Collaboration, *Calibration of the light-flavour jet mistagging efficiency of the  $b$ -tagging algorithms with Z+jets events using  $139\text{ fb}^{-1}$  of ATLAS proton–proton collision data at  $\sqrt{s} = 13$  TeV*, *Eur. Phys. J. C* **83** (2023) 728, arXiv: [2301.06319](https://arxiv.org/abs/2301.06319) [[hep-ex](#)].
- [113] ATLAS Collaboration, *Measurement of the production cross-section of a single top quark in association with a Z boson in proton–proton collisions at 13 TeV with the ATLAS detector*, *Phys. Lett. B* **780** (2018) 557, arXiv: [1710.03659](https://arxiv.org/abs/1710.03659) [[hep-ex](#)].
- [114] M. Bähr et al., *Herwig++ physics and manual*, *Eur. Phys. J. C* **58** (2008) 639, arXiv: [0803.0883](https://arxiv.org/abs/0803.0883) [[hep-ph](#)].
- [115] J. Bellm et al., *Herwig 7.0/Herwig++ 3.0 release note*, *Eur. Phys. J. C* **76** (2016) 196, arXiv: [1512.01178](https://arxiv.org/abs/1512.01178) [[hep-ph](#)].
- [116] S. Höche, S. Mrenna, S. Payne, C. T. Preuss and P. Skands, *A Study of QCD Radiation in VBF Higgs Production with Vincia and Pythia*, *SciPost Phys.* **12** (2022) 010, arXiv: [2106.10987](https://arxiv.org/abs/2106.10987) [[hep-ph](#)].
- [117] J. Butterworth et al., *PDF4LHC recommendations for LHC Run II*, *J. Phys. G* **43** (2016) 023001, arXiv: [1510.03865](https://arxiv.org/abs/1510.03865) [[hep-ph](#)].
- [118] R. Barlow and C. Beeston, *Fitting using finite Monte Carlo samples*, *Comput. Phys. Commun.* **77** (1993) 219.
- [119] W. Verkerke and D. Kirkby, *The RooFit toolkit for data modeling*, 2003, arXiv: [physics/0306116](https://arxiv.org/abs/physics/0306116) [[physics.data-an](#)].
- [120] A. L. Read, *Presentation of search results: the  $CL_S$  technique*, *J. Phys. G* **28** (2002) 2693.
- [121] G. Cowan, K. Cranmer, E. Gross and O. Vitells, *Asymptotic formulae for likelihood-based tests of new physics*, *Eur. Phys. J. C* **71** (2011) 1554, arXiv: [1007.1727](https://arxiv.org/abs/1007.1727) [[physics.data-an](#)], Erratum: *Eur. Phys. J. C* **73** (2013) 2501.
- [122] ATLAS Collaboration, *ATLAS Computing Acknowledgements*, ATL-SOFT-PUB-2023-001, 2023, URL: <https://cds.cern.ch/record/2869272>.

## The ATLAS Collaboration

G. Aad <sup>103</sup>, E. Aakvaag <sup>16</sup>, B. Abbott <sup>121</sup>, K. Abeling <sup>55</sup>, N.J. Abicht <sup>49</sup>, S.H. Abidi <sup>29</sup>, M. Aboeela <sup>44</sup>, A. Aboulhorma <sup>35e</sup>, H. Abramowicz <sup>152</sup>, H. Abreu <sup>151</sup>, Y. Abulaiti <sup>118</sup>, B.S. Acharya <sup>69a,69b,1</sup>, A. Ackermann <sup>63a</sup>, C. Adam Bourdarios <sup>4</sup>, L. Adamczyk <sup>86a</sup>, S.V. Addepalli <sup>26</sup>, M.J. Addison <sup>102</sup>, J. Adelman <sup>116</sup>, A. Adiguzel <sup>21c</sup>, T. Adaye <sup>135</sup>, A.A. Affolder <sup>137</sup>, Y. Afik <sup>39</sup>, M.N. Agaras <sup>13</sup>, J. Agarwala <sup>73a,73b</sup>, A. Aggarwal <sup>101</sup>, C. Agheorghiesei <sup>27c</sup>, A. Ahmad <sup>36</sup>, F. Ahmadov <sup>38,y</sup>, W.S. Ahmed <sup>105</sup>, S. Ahuja <sup>96</sup>, X. Ai <sup>62e</sup>, G. Aielli <sup>76a,76b</sup>, A. Aikot <sup>164</sup>, M. Ait Tamlihat <sup>35e</sup>, B. Aitbenchikh <sup>35a</sup>, I. Aizenberg <sup>170</sup>, M. Akbiyik <sup>101</sup>, T.P.A. Åkesson <sup>99</sup>, A.V. Akimov <sup>37</sup>, D. Akiyama <sup>169</sup>, N.N. Akolkar <sup>24</sup>, S. Aktas <sup>21a</sup>, K. Al Houry <sup>41</sup>, G.L. Alberghi <sup>23b</sup>, J. Albert <sup>166</sup>, P. Albicocco <sup>53</sup>, G.L. Albouy <sup>60</sup>, S. Alderweireldt <sup>52</sup>, Z.L. Alegria <sup>122</sup>, M. Aleksa <sup>36</sup>, I.N. Aleksandrov <sup>38</sup>, C. Alexa <sup>27b</sup>, T. Alexopoulos <sup>10</sup>, F. Alfonsi <sup>23b</sup>, M. Algren <sup>56</sup>, M. Alhroob <sup>142</sup>, B. Ali <sup>133</sup>, H.M.J. Ali <sup>92</sup>, S. Ali <sup>149</sup>, S.W. Alibocus <sup>93</sup>, M. Aliev <sup>33c</sup>, G. Alimonti <sup>71a</sup>, W. Alkakhri <sup>55</sup>, C. Allaire <sup>66</sup>, B.M.M. Allbrooke <sup>147</sup>, J.F. Allen <sup>52</sup>, C.A. Allendes Flores <sup>138f</sup>, P.P. Allport <sup>20</sup>, A. Aloisio <sup>72a,72b</sup>, F. Alonso <sup>91</sup>, C. Alpigiani <sup>139</sup>, M. Alvarez Estevez <sup>100</sup>, A. Alvarez Fernandez <sup>101</sup>, M. Alves Cardoso <sup>56</sup>, M.G. Alviggi <sup>72a,72b</sup>, M. Aly <sup>102</sup>, Y. Amaral Coutinho <sup>83b</sup>, A. Ambler <sup>105</sup>, C. Amelung <sup>36</sup>, M. Amerl <sup>102</sup>, C.G. Ames <sup>110</sup>, D. Amidei <sup>107</sup>, K.J. Amirie <sup>156</sup>, S.P. Amor Dos Santos <sup>131a</sup>, K.R. Amos <sup>164</sup>, S. An <sup>84</sup>, V. Ananiev <sup>126</sup>, C. Anastopoulos <sup>140</sup>, T. Andeen <sup>11</sup>, J.K. Anders <sup>36</sup>, S.Y. Andrean <sup>47a,47b</sup>, A. Andreazza <sup>71a,71b</sup>, S. Angelidakis <sup>9</sup>, A. Angerami <sup>41,aa</sup>, A.V. Anisenkov <sup>37</sup>, A. Annovi <sup>74a</sup>, C. Antel <sup>56</sup>, M.T. Anthony <sup>140</sup>, E. Antipov <sup>146</sup>, M. Antonelli <sup>53</sup>, F. Anulli <sup>75a</sup>, M. Aoki <sup>84</sup>, T. Aoki <sup>154</sup>, J.A. Aparisi Pozo <sup>164</sup>, M.A. Aparo <sup>147</sup>, L. Aperio Bella <sup>48</sup>, C. Appelt <sup>18</sup>, A. Apyan <sup>26</sup>, S.J. Arbiol Val <sup>87</sup>, C. Arcangeletti <sup>53</sup>, A.T.H. Arce <sup>51</sup>, E. Arena <sup>93</sup>, J-F. Arguin <sup>109</sup>, S. Argyropoulos <sup>54</sup>, J.-H. Arling <sup>48</sup>, O. Arnaez <sup>4</sup>, H. Arnold <sup>115</sup>, G. Artoni <sup>75a,75b</sup>, H. Asada <sup>112</sup>, K. Asai <sup>119</sup>, S. Asai <sup>154</sup>, N.A. Asbah <sup>36</sup>, K. Assamagan <sup>29</sup>, R. Astalos <sup>28a</sup>, K.S.V. Astrand <sup>99</sup>, S. Atashi <sup>160</sup>, R.J. Atkin <sup>33a</sup>, M. Atkinson <sup>163</sup>, H. Atmani <sup>35f</sup>, P.A. Atlasiddha <sup>129</sup>, K. Augsten <sup>133</sup>, S. Auricchio <sup>72a,72b</sup>, A.D. Auriol <sup>20</sup>, V.A. Austrup <sup>102</sup>, G. Avolio <sup>36</sup>, K. Axiotis <sup>56</sup>, G. Azuelos <sup>109,ae</sup>, D. Babal <sup>28b</sup>, H. Bachacou <sup>136</sup>, K. Bachas <sup>153,p</sup>, A. Bachiu <sup>34</sup>, F. Backman <sup>47a,47b</sup>, A. Badea <sup>39</sup>, T.M. Baer <sup>107</sup>, P. Bagnaia <sup>75a,75b</sup>, M. Bahmani <sup>18</sup>, D. Bahner <sup>54</sup>, K. Bai <sup>124</sup>, J.T. Baines <sup>135</sup>, L. Baines <sup>95</sup>, O.K. Baker <sup>173</sup>, E. Bakos <sup>15</sup>, D. Bakshi Gupta <sup>8</sup>, V. Balakrishnan <sup>121</sup>, R. Balasubramanian <sup>115</sup>, E.M. Baldin <sup>37</sup>, P. Balek <sup>86a</sup>, E. Ballabene <sup>23b,23a</sup>, F. Balli <sup>136</sup>, L.M. Baltos <sup>63a</sup>, W.K. Balunas <sup>32</sup>, J. Balz <sup>101</sup>, E. Banas <sup>87</sup>, M. Bandieramonte <sup>130</sup>, A. Bandyopadhyay <sup>24</sup>, S. Bansal <sup>24</sup>, L. Barak <sup>152</sup>, M. Barakat <sup>48</sup>, E.L. Barberio <sup>106</sup>, D. Barberis <sup>57b,57a</sup>, M. Barbero <sup>103</sup>, M.Z. Barel <sup>115</sup>, K.N. Barends <sup>33a</sup>, T. Barillari <sup>111</sup>, M-S. Barisits <sup>36</sup>, T. Barklow <sup>144</sup>, P. Baron <sup>123</sup>, D.A. Baron Moreno <sup>102</sup>, A. Baroncelli <sup>62a</sup>, G. Barone <sup>29</sup>, A.J. Barr <sup>127</sup>, J.D. Barr <sup>97</sup>, F. Barreiro <sup>100</sup>, J. Barreiro Guimarães da Costa <sup>14a</sup>, U. Barron <sup>152</sup>, M.G. Barros Teixeira <sup>131a</sup>, S. Barsov <sup>37</sup>, F. Bartels <sup>63a</sup>, R. Bartoldus <sup>144</sup>, A.E. Barton <sup>92</sup>, P. Bartos <sup>28a</sup>, A. Basan <sup>101</sup>, M. Baselga <sup>49</sup>, A. Bassalat <sup>66,b</sup>, M.J. Basso <sup>157a</sup>, R. Bate <sup>165</sup>, R.L. Bates <sup>59</sup>, S. Batlamous <sup>35e</sup>, B. Batool <sup>142</sup>, M. Battaglia <sup>137</sup>, D. Battulga <sup>18</sup>, M. Baucé <sup>75a,75b</sup>, M. Bauer <sup>36</sup>, P. Bauer <sup>24</sup>, L.T. Bazzano Hurrell <sup>30</sup>, J.B. Beacham <sup>51</sup>, T. Beau <sup>128</sup>, J.Y. Beaucamp <sup>91</sup>, P.H. Beauchemin <sup>159</sup>, P. Bechtel <sup>24</sup>, H.P. Beck <sup>19,o</sup>, K. Becker <sup>168</sup>, A.J. Beddall <sup>82</sup>, V.A. Bednyakov <sup>38</sup>, C.P. Bee <sup>146</sup>, L.J. Beemster <sup>15</sup>, T.A. Beermann <sup>36</sup>, M. Begalli <sup>83d</sup>, M. Begel <sup>29</sup>, A. Behera <sup>146</sup>, J.K. Behr <sup>48</sup>, J.F. Beirer <sup>36</sup>, F. Beisiegel <sup>24</sup>, M. Belfkir <sup>117b</sup>, G. Bella <sup>152</sup>, L. Bellagamba <sup>23b</sup>, A. Bellerive <sup>34</sup>, P. Bellos <sup>20</sup>, K. Beloborodov <sup>37</sup>, D. Bencheekroun <sup>35a</sup>, F. Bendebba <sup>35a</sup>, Y. Benhammou <sup>152</sup>,

K.C. Benkendorfer <sup>61</sup>, L. Beresford <sup>48</sup>, M. Beretta <sup>53</sup>, E. Bergeaas Kuutmann <sup>162</sup>, N. Berger <sup>4</sup>,  
 B. Bergmann <sup>133</sup>, J. Beringer <sup>17a</sup>, G. Bernardi <sup>5</sup>, C. Bernius <sup>144</sup>, F.U. Bernlochner <sup>24</sup>,  
 F. Bernon <sup>36,103</sup>, A. Berrocal Guardia <sup>13</sup>, T. Berry <sup>96</sup>, P. Berta <sup>134</sup>, A. Berthold <sup>50</sup>, S. Bethke <sup>111</sup>,  
 A. Betti <sup>75a,75b</sup>, A.J. Bevan <sup>95</sup>, N.K. Bhalla <sup>54</sup>, M. Bhamjee <sup>33c</sup>, S. Bhatta <sup>146</sup>,  
 D.S. Bhattacharya <sup>167</sup>, P. Bhattarai <sup>144</sup>, K.D. Bhide <sup>54</sup>, V.S. Bhopatkar <sup>122</sup>, R.M. Bianchi <sup>130</sup>,  
 G. Bianco <sup>23b,23a</sup>, O. Biebel <sup>110</sup>, R. Bielski <sup>124</sup>, M. Biglietti <sup>77a</sup>, C.S. Billingsley <sup>44</sup>, M. Bindi <sup>55</sup>,  
 A. Bingul <sup>21b</sup>, C. Bini <sup>75a,75b</sup>, A. Biondini <sup>93</sup>, C.J. Birch-sykes <sup>102</sup>, G.A. Bird <sup>32</sup>, M. Birman <sup>170</sup>,  
 M. Biros <sup>134</sup>, S. Biryukov <sup>147</sup>, T. Bisanz <sup>49</sup>, E. Bisceglie <sup>43b,43a</sup>, J.P. Biswal <sup>135</sup>, D. Biswas <sup>142</sup>,  
 K. Bjørke <sup>126</sup>, I. Bloch <sup>48</sup>, A. Blue <sup>59</sup>, U. Blumenschein <sup>95</sup>, J. Blumenthal <sup>101</sup>,  
 V.S. Bobrovnikov <sup>37</sup>, M. Boehler <sup>54</sup>, B. Boehm <sup>167</sup>, D. Bogovac <sup>36</sup>, A.G. Bogdanchikov <sup>37</sup>,  
 C. Bohm <sup>47a</sup>, V. Boisvert <sup>96</sup>, P. Bokan <sup>36</sup>, T. Bold <sup>86a</sup>, M. Bomben <sup>5</sup>, M. Bona <sup>95</sup>,  
 M. Boonekamp <sup>136</sup>, C.D. Booth <sup>96</sup>, A.G. Borbély <sup>59</sup>, I.S. Bordulev <sup>37</sup>, H.M. Borecka-Bielska <sup>109</sup>,  
 G. Borissov <sup>92</sup>, D. Bortoletto <sup>127</sup>, D. Boscherini <sup>23b</sup>, M. Bosman <sup>13</sup>, J.D. Bossio Sola <sup>36</sup>,  
 K. Bouaouda <sup>35a</sup>, N. Bouchhar <sup>164</sup>, J. Boudreau <sup>130</sup>, E.V. Bouhova-Thacker <sup>92</sup>, D. Boumediene <sup>40</sup>,  
 R. Bouquet <sup>57b,57a</sup>, A. Boveia <sup>120</sup>, J. Boyd <sup>36</sup>, D. Boye <sup>29</sup>, I.R. Boyko <sup>38</sup>, J. Bracinik <sup>20</sup>,  
 N. Brahimy <sup>4</sup>, G. Brandt <sup>172</sup>, O. Brandt <sup>32</sup>, F. Braren <sup>48</sup>, B. Brau <sup>104</sup>, J.E. Brau <sup>124</sup>,  
 R. Brenner <sup>170</sup>, L. Brenner <sup>115</sup>, R. Brenner <sup>162</sup>, S. Bressler <sup>170</sup>, D. Britton <sup>59</sup>, D. Britzger <sup>111</sup>,  
 I. Brock <sup>24</sup>, G. Brooijmans <sup>41</sup>, E. Brost <sup>29</sup>, L.M. Brown <sup>166</sup>, L.E. Bruce <sup>61</sup>, T.L. Bruckler <sup>127</sup>,  
 P.A. Bruckman de Renstrom <sup>87</sup>, B. Brüers <sup>48</sup>, A. Bruni <sup>23b</sup>, G. Bruni <sup>23b</sup>, M. Bruschi <sup>23b</sup>,  
 N. Brusino <sup>75a,75b</sup>, T. Buanes <sup>16</sup>, Q. Buat <sup>139</sup>, D. Buchin <sup>111</sup>, A.G. Buckley <sup>59</sup>, O. Bulekov <sup>37</sup>,  
 B.A. Bullard <sup>144</sup>, S. Burdin <sup>93</sup>, C.D. Burgard <sup>49</sup>, A.M. Burger <sup>36</sup>, B. Burghgrave <sup>8</sup>,  
 O. Burlayenko <sup>54</sup>, J.T.P. Burr <sup>32</sup>, C.D. Burton <sup>11</sup>, J.C. Burzynski <sup>143</sup>, E.L. Busch <sup>41</sup>,  
 V. Büscher <sup>101</sup>, P.J. Bussey <sup>59</sup>, J.M. Butler <sup>25</sup>, C.M. Buttar <sup>59</sup>, J.M. Butterworth <sup>97</sup>,  
 W. Buttinger <sup>135</sup>, C.J. Buxo Vazquez <sup>108</sup>, A.R. Buzykaev <sup>37</sup>, S. Cabrera Urbán <sup>164</sup>,  
 L. Cadamuro <sup>66</sup>, D. Caforio <sup>58</sup>, H. Cai <sup>130</sup>, Y. Cai <sup>14a,14e</sup>, Y. Cai <sup>14c</sup>, V.M.M. Cairo <sup>36</sup>,  
 O. Cakir <sup>3a</sup>, N. Calace <sup>36</sup>, P. Calafiura <sup>17a</sup>, G. Calderini <sup>128</sup>, P. Calfayan <sup>68</sup>, G. Callea <sup>59</sup>,  
 L.P. Caloba <sup>83b</sup>, D. Calvet <sup>40</sup>, S. Calvet <sup>40</sup>, M. Calvetti <sup>74a,74b</sup>, R. Camacho Toro <sup>128</sup>,  
 S. Camarda <sup>36</sup>, D. Camarero Munoz <sup>26</sup>, P. Camarri <sup>76a,76b</sup>, M.T. Camerlingo <sup>72a,72b</sup>,  
 D. Cameron <sup>36</sup>, C. Camincher <sup>166</sup>, M. Campanelli <sup>97</sup>, A. Camplani <sup>42</sup>, V. Canale <sup>72a,72b</sup>,  
 A.C. Canbay <sup>3a</sup>, E. Canonero <sup>96</sup>, J. Cantero <sup>164</sup>, Y. Cao <sup>163</sup>, F. Capocasa <sup>26</sup>, M. Capua <sup>43b,43a</sup>,  
 A. Carbone <sup>71a,71b</sup>, R. Cardarelli <sup>76a</sup>, J.C.J. Cardenas <sup>8</sup>, F. Cardillo <sup>164</sup>, G. Carducci <sup>43b,43a</sup>,  
 T. Carli <sup>36</sup>, G. Carlino <sup>72a</sup>, J.I. Carlotto <sup>13</sup>, B.T. Carlson <sup>130,q</sup>, E.M. Carlson <sup>166,157a</sup>,  
 L. Carminati <sup>71a,71b</sup>, A. Carnelli <sup>136</sup>, M. Carnesale <sup>75a,75b</sup>, S. Caron <sup>114</sup>, E. Carquin <sup>138f</sup>,  
 S. Carrá <sup>71a</sup>, G. Carratta <sup>23b,23a</sup>, A.M. Carroll <sup>124</sup>, T.M. Carter <sup>52</sup>, M.P. Casado <sup>13,i</sup>,  
 M. Caspar <sup>48</sup>, F.L. Castillo <sup>4</sup>, L. Castillo Garcia <sup>13</sup>, V. Castillo Gimenez <sup>164</sup>, N.F. Castro <sup>131a,131e</sup>,  
 A. Catinaccio <sup>36</sup>, J.R. Catmore <sup>126</sup>, T. Cavaliere <sup>4</sup>, V. Cavaliere <sup>29</sup>, N. Cavalli <sup>23b,23a</sup>,  
 Y.C. Cekmecelioglu <sup>48</sup>, E. Celebi <sup>21a</sup>, S. Cella <sup>36</sup>, F. Celli <sup>127</sup>, M.S. Centonze <sup>70a,70b</sup>,  
 V. Cepaitis <sup>56</sup>, K. Cerny <sup>123</sup>, A.S. Cerqueira <sup>83a</sup>, A. Cerri <sup>147</sup>, L. Cerrito <sup>76a,76b</sup>, F. Cerutti <sup>17a</sup>,  
 B. Cervato <sup>142</sup>, A. Cervelli <sup>23b</sup>, G. Cesarini <sup>53</sup>, S.A. Cetin <sup>82</sup>, D. Chakraborty <sup>116</sup>, J. Chan <sup>17a</sup>,  
 W.Y. Chan <sup>154</sup>, J.D. Chapman <sup>32</sup>, E. Chapon <sup>136</sup>, B. Chargeishvili <sup>150b</sup>, D.G. Charlton <sup>20</sup>,  
 M. Chatterjee <sup>19</sup>, C. Chauhan <sup>134</sup>, Y. Che <sup>14c</sup>, S. Chekanov <sup>6</sup>, S.V. Chekulaev <sup>157a</sup>,  
 G.A. Chelkov <sup>38,a</sup>, A. Chen <sup>107</sup>, B. Chen <sup>152</sup>, B. Chen <sup>166</sup>, H. Chen <sup>14c</sup>, H. Chen <sup>29</sup>,  
 J. Chen <sup>62c</sup>, J. Chen <sup>143</sup>, M. Chen <sup>127</sup>, S. Chen <sup>154</sup>, S.J. Chen <sup>14c</sup>, X. Chen <sup>62c,136</sup>,  
 X. Chen <sup>14b,ad</sup>, Y. Chen <sup>62a</sup>, C.L. Cheng <sup>171</sup>, H.C. Cheng <sup>64a</sup>, S. Cheong <sup>144</sup>, A. Cheplakov <sup>38</sup>,  
 E. Cheremushkina <sup>48</sup>, E. Cherepanova <sup>115</sup>, R. Cherkaoui El Moursli <sup>35e</sup>, E. Cheu <sup>7</sup>, K. Cheung <sup>65</sup>,  
 L. Chevalier <sup>136</sup>, V. Chiarella <sup>53</sup>, G. Chiarelli <sup>74a</sup>, N. Chiedde <sup>103</sup>, G. Chiodini <sup>70a</sup>,  
 A.S. Chisholm <sup>20</sup>, A. Chitan <sup>27b</sup>, M. Chitishvili <sup>164</sup>, M.V. Chizhov <sup>38</sup>, K. Choi <sup>11</sup>, Y. Chou <sup>139</sup>,

E.Y.S. Chow <sup>id114</sup>, K.L. Chu <sup>id170</sup>, M.C. Chu <sup>id64a</sup>, X. Chu <sup>id14a,14e</sup>, J. Chudoba <sup>id132</sup>,  
 J.J. Chwastowski <sup>id87</sup>, D. Cieri <sup>id111</sup>, K.M. Ciesla <sup>id86a</sup>, V. Cindro <sup>id94</sup>, A. Ciocio <sup>id17a</sup>, F. Ciroto <sup>id72a,72b</sup>,  
 Z.H. Citron <sup>id170</sup>, M. Citterio <sup>id71a</sup>, D.A. Ciubotaru <sup>id27b</sup>, A. Clark <sup>id56</sup>, P.J. Clark <sup>id52</sup>, C. Clarry <sup>id156</sup>,  
 J.M. Clavijo Columbie <sup>id48</sup>, S.E. Clawson <sup>id48</sup>, C. Clement <sup>id47a,47b</sup>, J. Clercx <sup>id48</sup>, Y. Coadou <sup>id103</sup>,  
 M. Cobal <sup>id69a,69c</sup>, A. Coccaro <sup>id57b</sup>, R.F. Coelho Barrue <sup>id131a</sup>, R. Coelho Lopes De Sa <sup>id104</sup>,  
 S. Coelli <sup>id71a</sup>, B. Cole <sup>id41</sup>, J. Collot <sup>id60</sup>, P. Conde Muiño <sup>id131a,131g</sup>, M.P. Connell <sup>id33c</sup>,  
 S.H. Connell <sup>id33c</sup>, E.I. Conroy <sup>id127</sup>, F. Conventi <sup>id72a,af</sup>, H.G. Cooke <sup>id20</sup>, A.M. Cooper-Sarkar <sup>id127</sup>,  
 F.A. Corchia <sup>id23b,23a</sup>, A. Cordeiro Oudot Choi <sup>id128</sup>, L.D. Corpe <sup>id40</sup>, M. Corradi <sup>id75a,75b</sup>,  
 F. Corriveau <sup>id105,w</sup>, A. Cortes-Gonzalez <sup>id18</sup>, M.J. Costa <sup>id164</sup>, F. Costanza <sup>id4</sup>, D. Costanzo <sup>id140</sup>,  
 B.M. Cote <sup>id120</sup>, G. Cowan <sup>id96</sup>, K. Cranmer <sup>id171</sup>, D. Cremonini <sup>id23b,23a</sup>, S. Crépe-Renaudin <sup>id60</sup>,  
 F. Crescioli <sup>id128</sup>, M. Cristinziani <sup>id142</sup>, M. Cristoforetti <sup>id78a,78b</sup>, V. Croft <sup>id115</sup>, J.E. Crosby <sup>id122</sup>,  
 G. Crosetti <sup>id43b,43a</sup>, A. Cueto <sup>id100</sup>, H. Cui <sup>id14a,14e</sup>, Z. Cui <sup>id7</sup>, W.R. Cunningham <sup>id59</sup>, F. Curcio <sup>id164</sup>,  
 J.R. Curran <sup>id52</sup>, P. Czodrowski <sup>id36</sup>, M.M. Czurylo <sup>id36</sup>, M.J. Da Cunha Sargedas De Sousa <sup>id57b,57a</sup>,  
 J.V. Da Fonseca Pinto <sup>id83b</sup>, C. Da Via <sup>id102</sup>, W. Dabrowski <sup>id86a</sup>, T. Dado <sup>id49</sup>, S. Dahbi <sup>id149</sup>,  
 T. Dai <sup>id107</sup>, D. Dal Santo <sup>id19</sup>, C. Dallapiccola <sup>id104</sup>, M. Dam <sup>id42</sup>, G. D'amen <sup>id29</sup>, V. D'Amico <sup>id110</sup>,  
 J. Damp <sup>id101</sup>, J.R. Dandoy <sup>id34</sup>, M. Danninger <sup>id143</sup>, V. Dao <sup>id36</sup>, G. Darbo <sup>id57b</sup>, S.J. Das <sup>id29,ag</sup>,  
 F. Dattola <sup>id48</sup>, S. D'Auria <sup>id71a,71b</sup>, A. D'avanzo <sup>id72a,72b</sup>, C. David <sup>id33a</sup>, T. Davidek <sup>id134</sup>,  
 B. Davis-Purcell <sup>id34</sup>, I. Dawson <sup>id95</sup>, H.A. Day-hall <sup>id133</sup>, K. De <sup>id8</sup>, R. De Asmundis <sup>id72a</sup>,  
 N. De Biase <sup>id48</sup>, S. De Castro <sup>id23b,23a</sup>, N. De Groot <sup>id114</sup>, P. de Jong <sup>id115</sup>, H. De la Torre <sup>id116</sup>,  
 A. De Maria <sup>id14c</sup>, A. De Salvo <sup>id75a</sup>, U. De Sanctis <sup>id76a,76b</sup>, F. De Santis <sup>id70a,70b</sup>, A. De Santo <sup>id147</sup>,  
 J.B. De Vivie De Regie <sup>id60</sup>, D.V. Dedovich <sup>id38</sup>, J. Degens <sup>id93</sup>, A.M. Deiana <sup>id44</sup>, F. Del Corso <sup>id23b,23a</sup>,  
 J. Del Peso <sup>id100</sup>, F. Del Rio <sup>id63a</sup>, L. Delagrangé <sup>id128</sup>, F. Deliot <sup>id136</sup>, C.M. Delitzsch <sup>id49</sup>,  
 M. Della Pietra <sup>id72a,72b</sup>, D. Della Volpe <sup>id56</sup>, A. Dell'Acqua <sup>id36</sup>, L. Dell'Asta <sup>id71a,71b</sup>, M. Delmastro <sup>id4</sup>,  
 P.A. Delsart <sup>id60</sup>, S. Demers <sup>id173</sup>, M. Demichev <sup>id38</sup>, S.P. Denisov <sup>id37</sup>, L. D'Eramo <sup>id40</sup>,  
 D. Derendarz <sup>id87</sup>, F. Derue <sup>id128</sup>, P. Dervan <sup>id93</sup>, K. Desch <sup>id24</sup>, C. Deutsch <sup>id24</sup>, F.A. Di Bello <sup>id57b,57a</sup>,  
 A. Di Ciaccio <sup>id76a,76b</sup>, L. Di Ciaccio <sup>id4</sup>, A. Di Domenico <sup>id75a,75b</sup>, C. Di Donato <sup>id72a,72b</sup>,  
 A. Di Girolamo <sup>id36</sup>, G. Di Gregorio <sup>id36</sup>, A. Di Luca <sup>id78a,78b</sup>, B. Di Micco <sup>id77a,77b</sup>, R. Di Nardo <sup>id77a,77b</sup>,  
 M. Diamantopoulou <sup>id34</sup>, F.A. Dias <sup>id115</sup>, T. Dias Do Vale <sup>id143</sup>, M.A. Diaz <sup>id138a,138b</sup>,  
 F.G. Diaz Capriles <sup>id24</sup>, M. Didenko <sup>id164</sup>, E.B. Diehl <sup>id107</sup>, S. Díez Cornell <sup>id48</sup>, C. Diez Pardo <sup>id142</sup>,  
 C. Dimitriadi <sup>id162,24</sup>, A. Dimitrievska <sup>id20</sup>, J. Dingfelder <sup>id24</sup>, I-M. Dinu <sup>id27b</sup>, S.J. Dittmeier <sup>id63b</sup>,  
 F. Dittus <sup>id36</sup>, M. Divisek <sup>id134</sup>, F. Djama <sup>id103</sup>, T. Djobava <sup>id150b</sup>, C. Doglioni <sup>id102,99</sup>, A. Dohnalova <sup>id28a</sup>,  
 J. Dolejsi <sup>id134</sup>, Z. Dolezal <sup>id134</sup>, K.M. Dona <sup>id39</sup>, M. Donadelli <sup>id83c</sup>, B. Dong <sup>id108</sup>, J. Donini <sup>id40</sup>,  
 A. D'Onofrio <sup>id72a,72b</sup>, M. D'Onofrio <sup>id93</sup>, J. Dopke <sup>id135</sup>, A. Doria <sup>id72a</sup>, N. Dos Santos Fernandes <sup>id131a</sup>,  
 P. Dougan <sup>id102</sup>, M.T. Dova <sup>id91</sup>, A.T. Doyle <sup>id59</sup>, M.A. Draguet <sup>id127</sup>, E. Dreyer <sup>id170</sup>,  
 I. Drivas-koulouris <sup>id10</sup>, M. Drnevich <sup>id118</sup>, M. Drozdova <sup>id56</sup>, D. Du <sup>id62a</sup>, T.A. du Pree <sup>id115</sup>,  
 F. Dubinin <sup>id37</sup>, M. Dubovsky <sup>id28a</sup>, E. Duchovni <sup>id170</sup>, G. Duckeck <sup>id110</sup>, O.A. Ducu <sup>id27b</sup>, D. Duda <sup>id52</sup>,  
 A. Dudarev <sup>id36</sup>, E.R. Duden <sup>id26</sup>, M. D'uffizi <sup>id102</sup>, L. Duflost <sup>id66</sup>, M. Dührssen <sup>id36</sup>, I. Duminica <sup>id27g</sup>,  
 A.E. Dumitriu <sup>id27b</sup>, M. Dunford <sup>id63a</sup>, S. Dungs <sup>id49</sup>, K. Dunne <sup>id47a,47b</sup>, A. Duperrin <sup>id103</sup>,  
 H. Duran Yildiz <sup>id3a</sup>, M. Düren <sup>id58</sup>, A. Durglishvili <sup>id150b</sup>, B.L. Dwyer <sup>id116</sup>, G.I. Dyckes <sup>id17a</sup>,  
 M. Dyndal <sup>id86a</sup>, B.S. Dziedzic <sup>id87</sup>, Z.O. Earnshaw <sup>id147</sup>, G.H. Eberwein <sup>id127</sup>, B. Eckerova <sup>id28a</sup>,  
 S. Eggebrecht <sup>id55</sup>, E. Egidio Purcino De Souza <sup>id128</sup>, L.F. Ehrke <sup>id56</sup>, G. Eigen <sup>id16</sup>, K. Einsweiler <sup>id17a</sup>,  
 T. Ekelof <sup>id162</sup>, P.A. Ekman <sup>id99</sup>, S. El Farkh <sup>id35b</sup>, Y. El Ghazali <sup>id35b</sup>, H. El Jarrari <sup>id36</sup>,  
 A. El Moussaouy <sup>id109</sup>, V. Ellajosyula <sup>id162</sup>, M. Ellert <sup>id162</sup>, F. Ellinghaus <sup>id172</sup>, N. Ellis <sup>id36</sup>,  
 J. Elmsheuser <sup>id29</sup>, M. Elsayy <sup>id117a</sup>, M. Elsing <sup>id36</sup>, D. Emelianov <sup>id135</sup>, Y. Enari <sup>id154</sup>, I. Ene <sup>id17a</sup>,  
 S. Epari <sup>id13</sup>, P.A. Erland <sup>id87</sup>, M. Errenst <sup>id172</sup>, M. Escalier <sup>id66</sup>, C. Escobar <sup>id164</sup>, E. Etzion <sup>id152</sup>,  
 G. Evans <sup>id131a</sup>, H. Evans <sup>id68</sup>, L.S. Evans <sup>id96</sup>, A. Ezhilov <sup>id37</sup>, S. Ezzarqtouni <sup>id35a</sup>, F. Fabbri <sup>id23b,23a</sup>,  
 L. Fabbri <sup>id23b,23a</sup>, G. Facini <sup>id97</sup>, V. Fadeyev <sup>id137</sup>, R.M. Fakhruddinov <sup>id37</sup>, D. Fakoudis <sup>id101</sup>,

S. Falciano <sup>75a</sup>, L.F. Falda Ulhoa Coelho <sup>36</sup>, P.J. Falke <sup>24</sup>, F. Fallavollita <sup>111</sup>, J. Faltova <sup>134</sup>,  
 C. Fan <sup>163</sup>, Y. Fan <sup>14a</sup>, Y. Fang <sup>14a,14e</sup>, M. Fanti <sup>71a,71b</sup>, M. Faraj <sup>69a,69b</sup>, Z. Farazpay <sup>98</sup>,  
 A. Farbin <sup>8</sup>, A. Farilla <sup>77a</sup>, T. Farooque <sup>108</sup>, S.M. Farrington <sup>52</sup>, F. Fassi <sup>35e</sup>, D. Fassouliotis <sup>9</sup>,  
 M. Faucci Giannelli <sup>76a,76b</sup>, W.J. Fawcett <sup>32</sup>, L. Fayard <sup>66</sup>, P. Federic <sup>134</sup>, P. Federicova <sup>132</sup>,  
 O.L. Fedin <sup>37,a</sup>, M. Feickert <sup>171</sup>, L. Feligioni <sup>103</sup>, D.E. Fellers <sup>124</sup>, C. Feng <sup>62b</sup>, M. Feng <sup>14b</sup>,  
 Z. Feng <sup>115</sup>, M.J. Fenton <sup>160</sup>, L. Ferencz <sup>48</sup>, R.A.M. Ferguson <sup>92</sup>, S.I. Fernandez Luengo <sup>138f</sup>,  
 P. Fernandez Martinez <sup>13</sup>, M.J.V. Fernoux <sup>103</sup>, J. Ferrando <sup>92</sup>, A. Ferrari <sup>162</sup>, P. Ferrari <sup>115,114</sup>,  
 R. Ferrari <sup>73a</sup>, D. Ferrere <sup>56</sup>, C. Ferretti <sup>107</sup>, F. Fiedler <sup>101</sup>, P. Fiedler <sup>133</sup>, A. Filipčič <sup>94</sup>,  
 E.K. Filmer <sup>1</sup>, F. Filthaut <sup>114</sup>, M.C.N. Fiolhais <sup>131a,131c,c</sup>, L. Fiorini <sup>164</sup>, W.C. Fisher <sup>108</sup>,  
 T. Fitschen <sup>102</sup>, P.M. Fitzhugh <sup>136</sup>, I. Fleck <sup>142</sup>, P. Fleischmann <sup>107</sup>, T. Flick <sup>172</sup>, M. Flores <sup>33d,ab</sup>,  
 L.R. Flores Castillo <sup>64a</sup>, L. Flores Sanz De Acedo <sup>36</sup>, F.M. Follega <sup>78a,78b</sup>, N. Fomin <sup>16</sup>,  
 J.H. Foo <sup>156</sup>, A. Formica <sup>136</sup>, A.C. Forti <sup>102</sup>, E. Fortin <sup>36</sup>, A.W. Fortman <sup>17a</sup>, M.G. Foti <sup>17a</sup>,  
 L. Fountas <sup>9j</sup>, D. Fournier <sup>66</sup>, H. Fox <sup>92</sup>, P. Francavilla <sup>74a,74b</sup>, S. Francescato <sup>61</sup>,  
 S. Franchellucci <sup>56</sup>, M. Franchini <sup>23b,23a</sup>, S. Franchino <sup>63a</sup>, D. Francis <sup>36</sup>, L. Franco <sup>114</sup>,  
 V. Franco Lima <sup>36</sup>, L. Franconi <sup>48</sup>, M. Franklin <sup>61</sup>, G. Frattari <sup>26</sup>, W.S. Freund <sup>83b</sup>, Y.Y. Frid <sup>152</sup>,  
 J. Friend <sup>59</sup>, N. Fritzsche <sup>50</sup>, A. Froch <sup>54</sup>, D. Froidevaux <sup>36</sup>, J.A. Frost <sup>127</sup>, Y. Fu <sup>62a</sup>,  
 S. Fuenzalida Garrido <sup>138f</sup>, M. Fujimoto <sup>103</sup>, K.Y. Fung <sup>64a</sup>, E. Furtado De Simas Filho <sup>83e</sup>,  
 M. Furukawa <sup>154</sup>, J. Fuster <sup>164</sup>, A. Gabrielli <sup>23b,23a</sup>, A. Gabrielli <sup>156</sup>, P. Gadow <sup>36</sup>,  
 G. Gagliardi <sup>57b,57a</sup>, L.G. Gagnon <sup>17a</sup>, S. Gaid <sup>161</sup>, S. Galantzan <sup>152</sup>, E.J. Gallas <sup>127</sup>,  
 B.J. Gallop <sup>135</sup>, K.K. Gan <sup>120</sup>, S. Ganguly <sup>154</sup>, Y. Gao <sup>52</sup>, F.M. Garay Walls <sup>138a,138b</sup>, B. Garcia <sup>29</sup>,  
 C. García <sup>164</sup>, A. Garcia Alonso <sup>115</sup>, A.G. Garcia Caffaro <sup>173</sup>, J.E. García Navarro <sup>164</sup>,  
 M. Garcia-Sciveres <sup>17a</sup>, G.L. Gardner <sup>129</sup>, R.W. Gardner <sup>39</sup>, N. Garelli <sup>159</sup>, D. Garg <sup>80</sup>,  
 R.B. Garg <sup>144,m</sup>, J.M. Gargan <sup>52</sup>, C.A. Garner <sup>156</sup>, C.M. Garvey <sup>33a</sup>, P. Gaspar <sup>83b</sup>, V.K. Gassmann <sup>159</sup>,  
 G. Gaudio <sup>73a</sup>, V. Gautam <sup>13</sup>, P. Gauzzi <sup>75a,75b</sup>, I.L. Gavrilenko <sup>37</sup>, A. Gavrilyuk <sup>37</sup>, C. Gay <sup>165</sup>,  
 G. Gaycken <sup>48</sup>, E.N. Gazis <sup>10</sup>, A.A. Geanta <sup>27b</sup>, C.M. Gee <sup>137</sup>, A. Gekow <sup>120</sup>, C. Gemme <sup>57b</sup>,  
 M.H. Genest <sup>60</sup>, A.D. Gentry <sup>113</sup>, S. George <sup>96</sup>, W.F. George <sup>20</sup>, T. Geralis <sup>46</sup>,  
 P. Gessinger-Befurt <sup>36</sup>, M.E. Geyik <sup>172</sup>, M. Ghani <sup>168</sup>, K. Ghorbanian <sup>95</sup>, A. Ghosal <sup>142</sup>,  
 A. Ghosh <sup>160</sup>, A. Ghosh <sup>7</sup>, B. Giacobbe <sup>23b</sup>, S. Giagu <sup>75a,75b</sup>, T. Giani <sup>115</sup>, P. Giannetti <sup>74a</sup>,  
 A. Giannini <sup>62a</sup>, S.M. Gibson <sup>96</sup>, M. Gignac <sup>137</sup>, D.T. Gil <sup>86b</sup>, A.K. Gilbert <sup>86a</sup>, B.J. Gilbert <sup>41</sup>,  
 D. Gillberg <sup>34</sup>, G. Gilles <sup>115</sup>, L. Ginabat <sup>128</sup>, D.M. Gingrich <sup>2,ae</sup>, M.P. Giordani <sup>69a,69c</sup>,  
 P.F. Giraud <sup>136</sup>, G. Giugliarelli <sup>69a,69c</sup>, D. Giugni <sup>71a</sup>, F. Giuli <sup>36</sup>, I. Gkialas <sup>9j</sup>, L.K. Gladilin <sup>37</sup>,  
 C. Glasman <sup>100</sup>, G.R. Gledhill <sup>124</sup>, G. Glemža <sup>48</sup>, M. Glisic <sup>124</sup>, I. Gnesi <sup>43b,f</sup>, Y. Go <sup>29</sup>,  
 M. Goblirsch-Kolb <sup>36</sup>, B. Gocke <sup>49</sup>, D. Godin <sup>109</sup>, B. Gokturk <sup>21a</sup>, S. Goldfarb <sup>106</sup>, T. Golling <sup>56</sup>,  
 M.G.D. Gololo <sup>33g</sup>, D. Golubkov <sup>37</sup>, J.P. Gombas <sup>108</sup>, A. Gomes <sup>131a,131b</sup>, G. Gomes Da Silva <sup>142</sup>,  
 A.J. Gomez Delegido <sup>164</sup>, R. Gonçalo <sup>131a,131c</sup>, L. Gonella <sup>20</sup>, A. Gongadze <sup>150c</sup>, F. Gonnella <sup>20</sup>,  
 J.L. Gonski <sup>144</sup>, R.Y. González Andana <sup>52</sup>, S. González de la Hoz <sup>164</sup>, R. Gonzalez Lopez <sup>93</sup>,  
 C. Gonzalez Renteria <sup>17a</sup>, M.V. Gonzalez Rodrigues <sup>48</sup>, R. Gonzalez Suarez <sup>162</sup>,  
 S. Gonzalez-Sevilla <sup>56</sup>, L. Goossens <sup>36</sup>, B. Gorini <sup>36</sup>, E. Gorini <sup>70a,70b</sup>, A. Gorišek <sup>94</sup>,  
 T.C. Gosart <sup>129</sup>, A.T. Goshaw <sup>51</sup>, M.I. Gostkin <sup>38</sup>, S. Goswami <sup>122</sup>, C.A. Gottardo <sup>36</sup>,  
 S.A. Gotz <sup>110</sup>, M. Gouighri <sup>35b</sup>, V. Goumarre <sup>48</sup>, A.G. Goussiou <sup>139</sup>, N. Govender <sup>33c</sup>,  
 I. Grabowska-Bold <sup>86a</sup>, K. Graham <sup>34</sup>, E. Gramstad <sup>126</sup>, S. Grancagnolo <sup>70a,70b</sup>, C.M. Grant <sup>1,136</sup>,  
 P.M. Gravila <sup>27f</sup>, F.G. Gravili <sup>70a,70b</sup>, H.M. Gray <sup>17a</sup>, M. Greco <sup>70a,70b</sup>, C. Grefe <sup>24</sup>,  
 I.M. Gregor <sup>48</sup>, K.T. Greif <sup>160</sup>, P. Grenier <sup>144</sup>, S.G. Grewe <sup>111</sup>, A.A. Grillo <sup>137</sup>, K. Grimm <sup>31</sup>,  
 S. Grinstein <sup>13,s</sup>, J.-F. Grivaz <sup>66</sup>, E. Gross <sup>170</sup>, J. Grosse-Knetter <sup>55</sup>, J.C. Grundy <sup>127</sup>,  
 L. Guan <sup>107</sup>, C. Gubbels <sup>165</sup>, J.G.R. Guerrero Rojas <sup>164</sup>, G. Guerrieri <sup>69a,69c</sup>, F. Guescini <sup>111</sup>,  
 R. Gugel <sup>101</sup>, J.A.M. Guhit <sup>107</sup>, A. Guida <sup>18</sup>, E. Guilloton <sup>168</sup>, S. Guindon <sup>36</sup>, F. Guo <sup>14a,14e</sup>,  
 J. Guo <sup>62c</sup>, L. Guo <sup>48</sup>, Y. Guo <sup>107</sup>, R. Gupta <sup>48</sup>, R. Gupta <sup>130</sup>, S. Gurbuz <sup>24</sup>, S.S. Gurdasani <sup>54</sup>,



G. Gustavino [id<sup>36</sup>](#), M. Guth [id<sup>56</sup>](#), P. Gutierrez [id<sup>121</sup>](#), L.F. Gutierrez Zagazeta [id<sup>129</sup>](#), M. Gutsche [id<sup>50</sup>](#), C. Gutschow [id<sup>97</sup>](#), C. Gwenlan [id<sup>127</sup>](#), C.B. Gwilliam [id<sup>93</sup>](#), E.S. Haaland [id<sup>126</sup>](#), A. Haas [id<sup>118</sup>](#), M. Habedank [id<sup>48</sup>](#), C. Haber [id<sup>17a</sup>](#), H.K. Hadavand [id<sup>8</sup>](#), A. Hadeif [id<sup>50</sup>](#), S. Hadzic [id<sup>111</sup>](#), A.I. Hagan [id<sup>92</sup>](#), J.J. Hahn [id<sup>142</sup>](#), E.H. Haines [id<sup>97</sup>](#), M. Haleem [id<sup>167</sup>](#), J. Haley [id<sup>122</sup>](#), J.J. Hall [id<sup>140</sup>](#), G.D. Hallewell [id<sup>103</sup>](#), L. Halser [id<sup>19</sup>](#), K. Hamano [id<sup>166</sup>](#), M. Hamer [id<sup>24</sup>](#), G.N. Hamity [id<sup>52</sup>](#), E.J. Hampshire [id<sup>96</sup>](#), J. Han [id<sup>62b</sup>](#), K. Han [id<sup>62a</sup>](#), L. Han [id<sup>14c</sup>](#), L. Han [id<sup>62a</sup>](#), S. Han [id<sup>17a</sup>](#), Y.F. Han [id<sup>156</sup>](#), K. Hanagaki [id<sup>84</sup>](#), M. Hance [id<sup>137</sup>](#), D.A. Hangal [id<sup>41</sup>](#), H. Hanif [id<sup>143</sup>](#), M.D. Hank [id<sup>129</sup>](#), J.B. Hansen [id<sup>42</sup>](#), P.H. Hansen [id<sup>42</sup>](#), K. Hara [id<sup>158</sup>](#), D. Harada [id<sup>56</sup>](#), T. Harenberg [id<sup>172</sup>](#), S. Harkusha [id<sup>37</sup>](#), M.L. Harris [id<sup>104</sup>](#), Y.T. Harris [id<sup>127</sup>](#), J. Harrison [id<sup>13</sup>](#), N.M. Harrison [id<sup>120</sup>](#), P.F. Harrison [id<sup>168</sup>](#), N.M. Hartman [id<sup>111</sup>](#), N.M. Hartmann [id<sup>110</sup>](#), Y. Hasegawa [id<sup>141</sup>](#), S. Hassan [id<sup>16</sup>](#), R. Hauser [id<sup>108</sup>](#), C.M. Hawkes [id<sup>20</sup>](#), R.J. Hawkins [id<sup>36</sup>](#), Y. Hayashi [id<sup>154</sup>](#), S. Hayashida [id<sup>112</sup>](#), D. Hayden [id<sup>108</sup>](#), C. Hayes [id<sup>107</sup>](#), R.L. Hayes [id<sup>115</sup>](#), C.P. Hays [id<sup>127</sup>](#), J.M. Hays [id<sup>95</sup>](#), H.S. Hayward [id<sup>93</sup>](#), F. He [id<sup>62a</sup>](#), M. He [id<sup>14a,14e</sup>](#), Y. He [id<sup>155</sup>](#), Y. He [id<sup>48</sup>](#), Y. He [id<sup>97</sup>](#), N.B. Heatley [id<sup>95</sup>](#), V. Hedberg [id<sup>99</sup>](#), A.L. Heggelund [id<sup>126</sup>](#), N.D. Hehir [id<sup>95,\\*</sup>](#), C. Heidegger [id<sup>54</sup>](#), K.K. Heidegger [id<sup>54</sup>](#), W.D. Heidorn [id<sup>81</sup>](#), J. Heilman [id<sup>34</sup>](#), S. Heim [id<sup>48</sup>](#), T. Heim [id<sup>17a</sup>](#), J.G. Heinlein [id<sup>129</sup>](#), J.J. Heinrich [id<sup>124</sup>](#), L. Heinrich [id<sup>111,ac</sup>](#), J. Hejbal [id<sup>132</sup>](#), A. Held [id<sup>171</sup>](#), S. Hellesund [id<sup>16</sup>](#), C.M. Helling [id<sup>165</sup>](#), S. Hellman [id<sup>47a,47b</sup>](#), R.C.W. Henderson [id<sup>92</sup>](#), L. Henkelmann [id<sup>32</sup>](#), A.M. Henriques Correia [id<sup>36</sup>](#), H. Herde [id<sup>99</sup>](#), Y. Hernández Jiménez [id<sup>146</sup>](#), L.M. Herrmann [id<sup>24</sup>](#), T. Herrmann [id<sup>50</sup>](#), G. Herten [id<sup>54</sup>](#), R. Hertenberger [id<sup>110</sup>](#), L. Hervas [id<sup>36</sup>](#), M.E. Hesping [id<sup>101</sup>](#), N.P. Hessey [id<sup>157a</sup>](#), E. Hill [id<sup>156</sup>](#), S.J. Hillier [id<sup>20</sup>](#), J.R. Hinds [id<sup>108</sup>](#), F. Hinterkeuser [id<sup>24</sup>](#), M. Hirose [id<sup>125</sup>](#), S. Hirose [id<sup>158</sup>](#), D. Hirschbuehl [id<sup>172</sup>](#), T.G. Hitchings [id<sup>102</sup>](#), B. Hiti [id<sup>94</sup>](#), J. Hobbs [id<sup>146</sup>](#), R. Hobincu [id<sup>27e</sup>](#), N. Hod [id<sup>170</sup>](#), M.C. Hodgkinson [id<sup>140</sup>](#), B.H. Hodgkinson [id<sup>127</sup>](#), A. Hoecker [id<sup>36</sup>](#), D.D. Hofer [id<sup>107</sup>](#), J. Hofer [id<sup>48</sup>](#), T. Holm [id<sup>24</sup>](#), M. Holzbock [id<sup>111</sup>](#), L.B.A.H. Hommels [id<sup>32</sup>](#), B.P. Honan [id<sup>102</sup>](#), J. Hong [id<sup>62c</sup>](#), T.M. Hong [id<sup>130</sup>](#), B.H. Hooberman [id<sup>163</sup>](#), W.H. Hopkins [id<sup>6</sup>](#), Y. Horii [id<sup>112</sup>](#), S. Hou [id<sup>149</sup>](#), A.S. Howard [id<sup>94</sup>](#), J. Howarth [id<sup>59</sup>](#), J. Hoya [id<sup>6</sup>](#), M. Hrabovsky [id<sup>123</sup>](#), A. Hrynevich [id<sup>48</sup>](#), T. Hryn'ova [id<sup>4</sup>](#), P.J. Hsu [id<sup>65</sup>](#), S.-C. Hsu [id<sup>139</sup>](#), T. Hsu [id<sup>66</sup>](#), M. Hu [id<sup>17a</sup>](#), Q. Hu [id<sup>62a</sup>](#), S. Huang [id<sup>64b</sup>](#), X. Huang [id<sup>14a,14e</sup>](#), Y. Huang [id<sup>140</sup>](#), Y. Huang [id<sup>101</sup>](#), Y. Huang [id<sup>14a</sup>](#), Z. Huang [id<sup>102</sup>](#), Z. Hubacek [id<sup>133</sup>](#), M. Huebner [id<sup>24</sup>](#), F. Hugging [id<sup>24</sup>](#), T.B. Huffman [id<sup>127</sup>](#), C.A. Hugli [id<sup>48</sup>](#), M. Huhtinen [id<sup>36</sup>](#), S.K. Huiberts [id<sup>16</sup>](#), R. Hulsken [id<sup>105</sup>](#), N. Huseynov [id<sup>12</sup>](#), J. Huston [id<sup>108</sup>](#), J. Huth [id<sup>61</sup>](#), R. Hyneman [id<sup>144</sup>](#), G. Iacobucci [id<sup>56</sup>](#), G. Iakovidis [id<sup>29</sup>](#), I. Ibragimov [id<sup>142</sup>](#), L. Iconomidou-Fayard [id<sup>66</sup>](#), J.P. Iddon [id<sup>36</sup>](#), P. Iengo [id<sup>72a,72b</sup>](#), R. Iguchi [id<sup>154</sup>](#), T. Iizawa [id<sup>127</sup>](#), Y. Ikegami [id<sup>84</sup>](#), N. Ilic [id<sup>156</sup>](#), H. Imam [id<sup>35a</sup>](#), M. Ince Lezki [id<sup>56</sup>](#), T. Ingebretsen Carlson [id<sup>47a,47b</sup>](#), G. Introzzi [id<sup>73a,73b</sup>](#), M. Iodice [id<sup>77a</sup>](#), V. Ippolito [id<sup>75a,75b</sup>](#), R.K. Irwin [id<sup>93</sup>](#), M. Ishino [id<sup>154</sup>](#), W. Islam [id<sup>171</sup>](#), C. Issever [id<sup>18,48</sup>](#), S. Istin [id<sup>21a,ai</sup>](#), H. Ito [id<sup>169</sup>](#), R. Iuppa [id<sup>78a,78b</sup>](#), A. Ivina [id<sup>170</sup>](#), J.M. Izen [id<sup>45</sup>](#), V. Izzo [id<sup>72a</sup>](#), P. Jacka [id<sup>132,133</sup>](#), P. Jackson [id<sup>1</sup>](#), B.P. Jaeger [id<sup>143</sup>](#), C.S. Jagfeld [id<sup>110</sup>](#), G. Jain [id<sup>157a</sup>](#), P. Jain [id<sup>54</sup>](#), K. Jakobs [id<sup>54</sup>](#), T. Jakoubek [id<sup>170</sup>](#), J. Jamieson [id<sup>59</sup>](#), K.W. Janas [id<sup>86a</sup>](#), M. Javurkova [id<sup>104</sup>](#), L. Jeanty [id<sup>124</sup>](#), J. Jejelava [id<sup>150a,z</sup>](#), P. Jenni [id<sup>54,g</sup>](#), C.E. Jessiman [id<sup>34</sup>](#), C. Jia [id<sup>62b</sup>](#), J. Jia [id<sup>146</sup>](#), X. Jia [id<sup>61</sup>](#), X. Jia [id<sup>14a,14e</sup>](#), Z. Jia [id<sup>14c</sup>](#), C. Jiang [id<sup>52</sup>](#), S. Jiggins [id<sup>48</sup>](#), J. Jimenez Pena [id<sup>13</sup>](#), S. Jin [id<sup>14c</sup>](#), A. Jinaru [id<sup>27b</sup>](#), O. Jinnouchi [id<sup>155</sup>](#), P. Johansson [id<sup>140</sup>](#), K.A. Johns [id<sup>7</sup>](#), J.W. Johnson [id<sup>137</sup>](#), D.M. Jones [id<sup>147</sup>](#), E. Jones [id<sup>48</sup>](#), P. Jones [id<sup>32</sup>](#), R.W.L. Jones [id<sup>92</sup>](#), T.J. Jones [id<sup>93</sup>](#), H.L. Joos [id<sup>55,36</sup>](#), R. Joshi [id<sup>120</sup>](#), J. Jovicevic [id<sup>15</sup>](#), X. Ju [id<sup>17a</sup>](#), J.J. Junggeburth [id<sup>104</sup>](#), T. Junkermann [id<sup>63a</sup>](#), A. Juste Rozas [id<sup>13,s</sup>](#), M.K. Juzek [id<sup>87</sup>](#), S. Kabana [id<sup>138e</sup>](#), A. Kaczmarska [id<sup>87</sup>](#), M. Kado [id<sup>111</sup>](#), H. Kagan [id<sup>120</sup>](#), M. Kagan [id<sup>144</sup>](#), A. Kahn [id<sup>41</sup>](#), A. Kahn [id<sup>129</sup>](#), C. Kahra [id<sup>101</sup>](#), T. Kaji [id<sup>154</sup>](#), E. Kajomovitz [id<sup>151</sup>](#), N. Kakati [id<sup>170</sup>](#), I. Kalaitzidou [id<sup>54</sup>](#), C.W. Calderon [id<sup>29</sup>](#), N.J. Kang [id<sup>137</sup>](#), D. Kar [id<sup>33g</sup>](#), K. Karava [id<sup>127</sup>](#), M.J. Kareem [id<sup>157b</sup>](#), E. Karentzos [id<sup>54</sup>](#), I. Karkanas [id<sup>153</sup>](#), O. Karkout [id<sup>115</sup>](#), S.N. Karpov [id<sup>38</sup>](#), Z.M. Karpova [id<sup>38</sup>](#), V. Kartvelishvili [id<sup>92</sup>](#), A.N. Karyukhin [id<sup>37</sup>](#), E. Kasimi [id<sup>153</sup>](#), J. Katzy [id<sup>48</sup>](#), S. Kaur [id<sup>34</sup>](#), K. Kawade [id<sup>141</sup>](#), M.P. Kawale [id<sup>121</sup>](#), C. Kawamoto [id<sup>88</sup>](#), T. Kawamoto [id<sup>62a</sup>](#), E.F. Kay [id<sup>36</sup>](#), F.I. Kaya [id<sup>159</sup>](#), S. Kazakos [id<sup>108</sup>](#), V.F. Kazanin [id<sup>37</sup>](#), Y. Ke [id<sup>146</sup>](#), J.M. Keaveney [id<sup>33a</sup>](#), R. Keeler [id<sup>166</sup>](#), G.V. Kehris [id<sup>61</sup>](#), J.S. Keller [id<sup>34</sup>](#), A.S. Kelly [id<sup>97</sup>](#), J.J. Kempster [id<sup>147</sup>](#), P.D. Kennedy [id<sup>101</sup>](#), O. Kepka [id<sup>132</sup>](#), B.P. Kerridge [id<sup>135</sup>](#), S. Kersten [id<sup>172</sup>](#),

B.P. Kerševan <sup>94</sup>, L. Keszeghova <sup>28a</sup>, S. Ketabchi Haghghat <sup>156</sup>, R.A. Khan <sup>130</sup>, A. Khanov <sup>122</sup>,  
 A.G. Kharlamov <sup>37</sup>, T. Kharlamova <sup>37</sup>, E.E. Khoda <sup>139</sup>, M. Kholodenko <sup>37</sup>, T.J. Khoo <sup>18</sup>,  
 G. Khoriauli <sup>167</sup>, J. Khubua <sup>150b,\*</sup>, Y.A.R. Khwaira <sup>66</sup>, B. Kibirige<sup>33g</sup>, A. Kilgallon <sup>124</sup>,  
 D.W. Kim <sup>47a,47b</sup>, Y.K. Kim <sup>39</sup>, N. Kimura <sup>97</sup>, M.K. Kingston <sup>55</sup>, A. Kirchoff <sup>55</sup>, C. Kirfel <sup>24</sup>,  
 F. Kirfel <sup>24</sup>, J. Kirk <sup>135</sup>, A.E. Kiryunin <sup>111</sup>, C. Kitsaki <sup>10</sup>, O. Kivernyk <sup>24</sup>, M. Klassen <sup>159</sup>,  
 C. Klein <sup>34</sup>, L. Klein <sup>167</sup>, M.H. Klein <sup>44</sup>, S.B. Klein <sup>56</sup>, U. Klein <sup>93</sup>, P. Klimek <sup>36</sup>,  
 A. Klimentov <sup>29</sup>, T. Klioutchnikova <sup>36</sup>, P. Kluit <sup>115</sup>, S. Kluth <sup>111</sup>, E. Kneringer <sup>79</sup>,  
 T.M. Knight <sup>156</sup>, A. Knue <sup>49</sup>, R. Kobayashi <sup>88</sup>, D. Kobylanski <sup>170</sup>, S.F. Koch <sup>127</sup>,  
 M. Kocian <sup>144</sup>, P. Kodyš <sup>134</sup>, D.M. Koeck <sup>124</sup>, P.T. Koenig <sup>24</sup>, T. Koffas <sup>34</sup>, O. Kolay <sup>50</sup>,  
 I. Koletsou <sup>4</sup>, T. Komarek <sup>123</sup>, K. Köneke <sup>54</sup>, A.X.Y. Kong <sup>1</sup>, T. Kono <sup>119</sup>, N. Konstantinidis <sup>97</sup>,  
 P. Kontaxakis <sup>56</sup>, B. Konya <sup>99</sup>, R. Kopeliansky <sup>41</sup>, S. Koperny <sup>86a</sup>, K. Korcyl <sup>87</sup>, K. Kordas <sup>153,e</sup>,  
 A. Korn <sup>97</sup>, S. Korn <sup>55</sup>, I. Korolkov <sup>13</sup>, N. Korotkova <sup>37</sup>, B. Kortman <sup>115</sup>, O. Kortner <sup>111</sup>,  
 S. Kortner <sup>111</sup>, W.H. Kostecka <sup>116</sup>, V.V. Kostyukhin <sup>142</sup>, A. Kotsokechagia <sup>136</sup>, A. Kotwal <sup>51</sup>,  
 A. Koulouris <sup>36</sup>, A. Kourkoumeli-Charalampidi <sup>73a,73b</sup>, C. Kourkoumelis <sup>9</sup>, E. Kourlitis <sup>111,ac</sup>,  
 O. Kovanda <sup>124</sup>, R. Kowalewski <sup>166</sup>, W. Kozanecki <sup>136</sup>, A.S. Kozhin <sup>37</sup>, V.A. Kramarenko <sup>37</sup>,  
 G. Kramberger <sup>94</sup>, P. Kramer <sup>101</sup>, M.W. Krasny <sup>128</sup>, A. Krasznahorkay <sup>36</sup>, J.W. Kraus <sup>172</sup>,  
 J.A. Kremer <sup>48</sup>, T. Kresse <sup>50</sup>, J. Kretschmar <sup>93</sup>, K. Kreul <sup>18</sup>, P. Krieger <sup>156</sup>,  
 S. Krishnamurthy <sup>104</sup>, M. Krivos <sup>134</sup>, K. Krizka <sup>20</sup>, K. Kroeninger <sup>49</sup>, H. Kroha <sup>111</sup>, J. Kroll <sup>132</sup>,  
 J. Kroll <sup>129</sup>, K.S. Krowpman <sup>108</sup>, U. Kruchonak <sup>38</sup>, H. Krüger <sup>24</sup>, N. Krumnack<sup>81</sup>, M.C. Kruse <sup>51</sup>,  
 O. Kuchinskaia <sup>37</sup>, S. Kuday <sup>3a</sup>, S. Kuehn <sup>36</sup>, R. Kuesters <sup>54</sup>, T. Kuhl <sup>48</sup>, V. Kukhtin <sup>38</sup>,  
 Y. Kulchitsky <sup>37,a</sup>, S. Kuleshov <sup>138d,138b</sup>, M. Kumar <sup>33g</sup>, N. Kumari <sup>48</sup>, P. Kumari <sup>157b</sup>,  
 A. Kupco <sup>132</sup>, T. Kupfer<sup>49</sup>, A. Kupich <sup>37</sup>, O. Kuprash <sup>54</sup>, H. Kurashige <sup>85</sup>, L.L. Kurchaninov <sup>157a</sup>,  
 O. Kurdysh <sup>66</sup>, Y.A. Kurochkin <sup>37</sup>, A. Kurova <sup>37</sup>, M. Kuze <sup>155</sup>, A.K. Kvam <sup>104</sup>, J. Kvita <sup>123</sup>,  
 T. Kwan <sup>105</sup>, N.G. Kyriacou <sup>107</sup>, L.A.O. Laatu <sup>103</sup>, C. Lacasta <sup>164</sup>, F. Lacava <sup>75a,75b</sup>,  
 H. Lacker <sup>18</sup>, D. Lacour <sup>128</sup>, N.N. Lad <sup>97</sup>, E. Ladygin <sup>38</sup>, A. Lafarge <sup>40</sup>, B. Laforge <sup>128</sup>,  
 T. Lagouri <sup>173</sup>, F.Z. Lahbabi <sup>35a</sup>, S. Lai <sup>55</sup>, I.K. Lakomic <sup>86a</sup>, J.E. Lambert <sup>166</sup>, S. Lammers <sup>68</sup>,  
 W. Lampl <sup>7</sup>, C. Lampoudis <sup>153,e</sup>, G. Lamprinoudis<sup>101</sup>, A.N. Lancaster <sup>116</sup>, E. Lançon <sup>29</sup>,  
 U. Landgraf <sup>54</sup>, M.P.J. Landon <sup>95</sup>, V.S. Lang <sup>54</sup>, O.K.B. Langrekken <sup>126</sup>, A.J. Lankford <sup>160</sup>,  
 F. Lanni <sup>36</sup>, K. Lantzsch <sup>24</sup>, A. Lanza <sup>73a</sup>, A. Lapertosa <sup>57b,57a</sup>, J.F. Laporte <sup>136</sup>, T. Lari <sup>71a</sup>,  
 F. Lasagni Manghi <sup>23b</sup>, M. Lassnig <sup>36</sup>, V. Latonova <sup>132</sup>, A. Laudrain <sup>101</sup>, A. Laurier <sup>151</sup>,  
 S.D. Lawlor <sup>140</sup>, Z. Lawrence <sup>102</sup>, R. Lazaridou<sup>168</sup>, M. Lazzaroni <sup>71a,71b</sup>, B. Le<sup>102</sup>,  
 E.M. Le Boulicaut <sup>51</sup>, L.T. Le Pottier <sup>17a</sup>, B. Leban <sup>23b,23a</sup>, A. Lebedev <sup>81</sup>, M. LeBlanc <sup>102</sup>,  
 F. Ledroit-Guillon <sup>60</sup>, A.C.A. Lee<sup>97</sup>, S.C. Lee <sup>149</sup>, S. Lee <sup>47a,47b</sup>, T.F. Lee <sup>93</sup>, L.L. Leeuw <sup>33c</sup>,  
 H.P. Lefebvre <sup>96</sup>, M. Lefebvre <sup>166</sup>, C. Leggett <sup>17a</sup>, G. Lehmann Miotto <sup>36</sup>, M. Leigh <sup>56</sup>,  
 W.A. Leight <sup>104</sup>, W. Leinonen <sup>114</sup>, A. Leisos <sup>153,r</sup>, M.A.L. Leite <sup>83c</sup>, C.E. Leitgeb <sup>18</sup>,  
 R. Leitner <sup>134</sup>, K.J.C. Leney <sup>44</sup>, T. Lenz <sup>24</sup>, S. Leone <sup>74a</sup>, C. Leonidopoulos <sup>52</sup>, A. Leopold <sup>145</sup>,  
 C. Leroy <sup>109</sup>, R. Les <sup>108</sup>, C.G. Lester <sup>32</sup>, M. Levchenko <sup>37</sup>, J. Levêque <sup>4</sup>, L.J. Levinson <sup>170</sup>,  
 G. Levrini <sup>23b,23a</sup>, M.P. Lewicki <sup>87</sup>, C. Lewis <sup>139</sup>, D.J. Lewis <sup>4</sup>, A. Li <sup>5</sup>, B. Li <sup>62b</sup>, C. Li<sup>62a</sup>,  
 C-Q. Li <sup>111</sup>, H. Li <sup>62a</sup>, H. Li <sup>62b</sup>, H. Li <sup>14c</sup>, H. Li <sup>14b</sup>, H. Li <sup>62b</sup>, J. Li <sup>62c</sup>, K. Li <sup>139</sup>,  
 L. Li <sup>62c</sup>, M. Li <sup>14a,14e</sup>, Q.Y. Li <sup>62a</sup>, S. Li <sup>14a,14e</sup>, S. Li <sup>62d,62c,d</sup>, T. Li <sup>5</sup>, X. Li <sup>105</sup>, Z. Li <sup>127</sup>,  
 Z. Li <sup>154</sup>, Z. Li <sup>14a,14e</sup>, S. Liang<sup>14a,14e</sup>, Z. Liang <sup>14a</sup>, M. Liberatore <sup>136</sup>, B. Liberti <sup>76a</sup>, K. Lie <sup>64c</sup>,  
 J. Lieber Marin <sup>83e</sup>, H. Lien <sup>68</sup>, K. Lin <sup>108</sup>, R.E. Lindley <sup>7</sup>, J.H. Lindon <sup>2</sup>, E. Lipeles <sup>129</sup>,  
 A. Lipniacka <sup>16</sup>, A. Lister <sup>165</sup>, J.D. Little <sup>4</sup>, B. Liu <sup>14a</sup>, B.X. Liu <sup>143</sup>, D. Liu <sup>62d,62c</sup>,  
 E.H.L. Liu <sup>20</sup>, J.B. Liu <sup>62a</sup>, J.K.K. Liu <sup>32</sup>, K. Liu <sup>62d</sup>, K. Liu <sup>62d,62c</sup>, M. Liu <sup>62a</sup>, M.Y. Liu <sup>62a</sup>,  
 P. Liu <sup>14a</sup>, Q. Liu <sup>62d,139,62c</sup>, X. Liu <sup>62a</sup>, X. Liu <sup>62b</sup>, Y. Liu <sup>14d,14e</sup>, Y.L. Liu <sup>62b</sup>, Y.W. Liu <sup>62a</sup>,  
 J. Llorente Merino <sup>143</sup>, S.L. Lloyd <sup>95</sup>, E.M. Lobodzinska <sup>48</sup>, P. Loch <sup>7</sup>, T. Lohse <sup>18</sup>,  
 K. Lohwasser <sup>140</sup>, E. Loiacono <sup>48</sup>, M. Lokajicek <sup>132,\*</sup>, J.D. Lomas <sup>20</sup>, J.D. Long <sup>163</sup>,












































































































































I. Longarini <sup>160</sup>, L. Longo <sup>70a,70b</sup>, R. Longo <sup>163</sup>, I. Lopez Paz <sup>67</sup>, A. Lopez Solis <sup>48</sup>,  
 N. Lorenzo Martinez <sup>4</sup>, A.M. Lory <sup>110</sup>, G. Löschcke Centeno <sup>147</sup>, O. Loseva <sup>37</sup>, X. Lou <sup>47a,47b</sup>,  
 X. Lou <sup>14a,14e</sup>, A. Lounis <sup>66</sup>, P.A. Love <sup>92</sup>, G. Lu <sup>14a,14e</sup>, M. Lu <sup>66</sup>, S. Lu <sup>129</sup>, Y.J. Lu <sup>65</sup>,  
 H.J. Lubatti <sup>139</sup>, C. Luci <sup>75a,75b</sup>, F.L. Lucio Alves <sup>14c</sup>, F. Luehring <sup>68</sup>, I. Luise <sup>146</sup>,  
 O. Lukianchuk <sup>66</sup>, O. Lundberg <sup>145</sup>, B. Lund-Jensen <sup>145,\*</sup>, N.A. Luongo <sup>6</sup>, M.S. Lutz <sup>36</sup>,  
 A.B. Lux <sup>25</sup>, D. Lynn <sup>29</sup>, R. Lysak <sup>132</sup>, E. Lytken <sup>99</sup>, V. Lyubushkin <sup>38</sup>, T. Lyubushkina <sup>38</sup>,  
 M.M. Lyukova <sup>146</sup>, M.Firdaus M. Soberi <sup>52</sup>, H. Ma <sup>29</sup>, K. Ma <sup>62a</sup>, L.L. Ma <sup>62b</sup>, W. Ma <sup>62a</sup>,  
 Y. Ma <sup>122</sup>, D.M. Mac Donnell <sup>166</sup>, G. Maccarrone <sup>53</sup>, J.C. MacDonald <sup>101</sup>,  
 P.C. Machado De Abreu Farias <sup>83e</sup>, R. Madar <sup>40</sup>, T. Madula <sup>97</sup>, J. Maeda <sup>85</sup>, T. Maeno <sup>29</sup>,  
 H. Maguire <sup>140</sup>, V. Maiboroda <sup>136</sup>, A. Maio <sup>131a,131b,131d</sup>, K. Maj <sup>86a</sup>, O. Majersky <sup>48</sup>,  
 S. Majewski <sup>124</sup>, N. Makovec <sup>66</sup>, V. Maksimovic <sup>15</sup>, B. Malaescu <sup>128</sup>, Pa. Malecki <sup>87</sup>,  
 V.P. Maleev <sup>37</sup>, F. Malek <sup>60,n</sup>, M. Mali <sup>94</sup>, D. Malito <sup>96</sup>, U. Mallik <sup>80</sup>, S. Maltezos <sup>10</sup>,  
 S. Malyukov <sup>38</sup>, J. Mamuzic <sup>13</sup>, G. Mancini <sup>53</sup>, M.N. Mancini <sup>26</sup>, G. Manco <sup>73a,73b</sup>,  
 J.P. Mandalia <sup>95</sup>, I. Mandić <sup>94</sup>, L. Manhaes de Andrade Filho <sup>83a</sup>, I.M. Maniatis <sup>170</sup>,  
 J. Manjarres Ramos <sup>90</sup>, D.C. Mankad <sup>170</sup>, A. Mann <sup>110</sup>, S. Manzoni <sup>36</sup>, L. Mao <sup>62c</sup>,  
 X. Mapekula <sup>33c</sup>, A. Marantis <sup>153,r</sup>, G. Marchiori <sup>5</sup>, M. Marcisovsky <sup>132</sup>, C. Marcon <sup>71a</sup>,  
 M. Marinescu <sup>20</sup>, S. Marium <sup>48</sup>, M. Marjanovic <sup>121</sup>, A. Markhoos <sup>54</sup>, M. Markovitch <sup>66</sup>,  
 E.J. Marshall <sup>92</sup>, Z. Marshall <sup>17a</sup>, S. Marti-Garcia <sup>164</sup>, T.A. Martin <sup>168</sup>, V.J. Martin <sup>52</sup>,  
 B. Martin dit Latour <sup>16</sup>, L. Martinelli <sup>75a,75b</sup>, M. Martinez <sup>13,s</sup>, P. Martinez Agullo <sup>164</sup>,  
 V.I. Martinez Outschoorn <sup>104</sup>, P. Martinez Suarez <sup>13</sup>, S. Martin-Haugh <sup>135</sup>, G. Martinovicova <sup>134</sup>,  
 V.S. Martoiu <sup>27b</sup>, A.C. Martyniuk <sup>97</sup>, A. Marzin <sup>36</sup>, D. Mascione <sup>78a,78b</sup>, L. Masetti <sup>101</sup>,  
 T. Mashimo <sup>154</sup>, J. Masik <sup>102</sup>, A.L. Maslennikov <sup>37</sup>, P. Massarotti <sup>72a,72b</sup>, P. Mastrandrea <sup>74a,74b</sup>,  
 A. Mastroberardino <sup>43b,43a</sup>, T. Masubuchi <sup>154</sup>, T. Mathisen <sup>162</sup>, J. Matousek <sup>134</sup>, N. Matsuzawa <sup>154</sup>,  
 J. Maurer <sup>27b</sup>, A.J. Maury <sup>66</sup>, B. Maček <sup>94</sup>, D.A. Maximov <sup>37</sup>, A.E. May <sup>102</sup>, R. Mazini <sup>149</sup>,  
 I. Maznas <sup>116</sup>, M. Mazza <sup>108</sup>, S.M. Mazza <sup>137</sup>, E. Mazzeo <sup>71a,71b</sup>, C. Mc Ginn <sup>29</sup>,  
 J.P. Mc Gowan <sup>166</sup>, S.P. Mc Kee <sup>107</sup>, C.C. McCracken <sup>165</sup>, E.F. McDonald <sup>106</sup>,  
 A.E. McDougall <sup>115</sup>, J.A. Mcfayden <sup>147</sup>, R.P. McGovern <sup>129</sup>, G. Mchedlidze <sup>150b</sup>,  
 R.P. Mckenzie <sup>33g</sup>, T.C. Mclachlan <sup>48</sup>, D.J. Mclaughlin <sup>97</sup>, S.J. McMahon <sup>135</sup>,  
 C.M. Mccpartland <sup>93</sup>, R.A. McPherson <sup>166,w</sup>, S. Mehlhase <sup>110</sup>, A. Mehta <sup>93</sup>, D. Melini <sup>164</sup>,  
 B.R. Mellado Garcia <sup>33g</sup>, A.H. Melo <sup>55</sup>, F. Meloni <sup>48</sup>, A.M. Mendes Jacques Da Costa <sup>102</sup>,  
 H.Y. Meng <sup>156</sup>, L. Meng <sup>92</sup>, S. Menke <sup>111</sup>, M. Mentink <sup>36</sup>, E. Meoni <sup>43b,43a</sup>, G. Mercado <sup>116</sup>,  
 C. Merlassino <sup>69a,69c</sup>, L. Merola <sup>72a,72b</sup>, C. Meroni <sup>71a,71b</sup>, J. Metcalfe <sup>6</sup>, A.S. Mete <sup>6</sup>,  
 C. Meyer <sup>68</sup>, J-P. Meyer <sup>136</sup>, R.P. Middleton <sup>135</sup>, L. Mijović <sup>52</sup>, G. Mikenberg <sup>170</sup>,  
 M. Mikestikova <sup>132</sup>, M. Mikuž <sup>94</sup>, H. Mildner <sup>101</sup>, A. Milic <sup>36</sup>, D.W. Miller <sup>39</sup>, E.H. Miller <sup>144</sup>,  
 L.S. Miller <sup>34</sup>, A. Milov <sup>170</sup>, D.A. Milstead <sup>47a,47b</sup>, T. Min <sup>14c</sup>, A.A. Minaenko <sup>37</sup>,  
 I.A. Minashvili <sup>150b</sup>, L. Mince <sup>59</sup>, A.I. Mincer <sup>118</sup>, B. Mindur <sup>86a</sup>, M. Mineev <sup>38</sup>, Y. Mino <sup>88</sup>,  
 L.M. Mir <sup>13</sup>, M. Miralles Lopez <sup>59</sup>, M. Mironova <sup>17a</sup>, A. Mishima <sup>154</sup>, M.C. Missio <sup>114</sup>,  
 A. Mitra <sup>168</sup>, V.A. Mitsou <sup>164</sup>, Y. Mitsumori <sup>112</sup>, O. Miu <sup>156</sup>, P.S. Miyagawa <sup>95</sup>,  
 T. Mkrtchyan <sup>63a</sup>, M. Mlinarevic <sup>97</sup>, T. Mlinarevic <sup>97</sup>, M. Mlynarikova <sup>36</sup>, S. Mobius <sup>19</sup>,  
 P. Mogg <sup>110</sup>, M.H. Mohamed Farook <sup>113</sup>, A.F. Mohammed <sup>14a,14e</sup>, S. Mohapatra <sup>41</sup>,  
 G. Mokgatitswane <sup>33g</sup>, L. Moleri <sup>170</sup>, B. Mondal <sup>142</sup>, S. Mondal <sup>133</sup>, K. Mönig <sup>48</sup>,  
 E. Monnier <sup>103</sup>, L. Monsonis Romero <sup>164</sup>, J. Montejo Berlingen <sup>13</sup>, M. Montella <sup>120</sup>,  
 F. Montekali <sup>77a,77b</sup>, F. Monticelli <sup>91</sup>, S. Monzani <sup>69a,69c</sup>, N. Morange <sup>66</sup>,  
 A.L. Moreira De Carvalho <sup>48</sup>, M. Moreno Llácer <sup>164</sup>, C. Moreno Martinez <sup>56</sup>, P. Morettini <sup>57b</sup>,  
 S. Morgenstern <sup>36</sup>, M. Morii <sup>61</sup>, M. Morinaga <sup>154</sup>, F. Morodei <sup>75a,75b</sup>, L. Morvaj <sup>36</sup>,  
 P. Moschovakos <sup>36</sup>, B. Moser <sup>36</sup>, M. Mosidze <sup>150b</sup>, T. Moskalets <sup>54</sup>, P. Moskvitina <sup>114</sup>,  
 J. Moss <sup>31,k</sup>, A. Moussa <sup>35d</sup>, E.J.W. Moyse <sup>104</sup>, O. Mtintsilana <sup>33g</sup>, S. Muanza <sup>103</sup>,

J. Mueller <sup>130</sup>, D. Muenstermann <sup>92</sup>, R. Müller <sup>19</sup>, G.A. Mullier <sup>162</sup>, A.J. Mullin<sup>32</sup>, J.J. Mullin<sup>129</sup>, D.P. Mungo <sup>156</sup>, D. Munoz Perez <sup>164</sup>, F.J. Munoz Sanchez <sup>102</sup>, M. Murin <sup>102</sup>, W.J. Murray <sup>168,135</sup>, M. Muškinja <sup>94</sup>, C. Mwewa <sup>29</sup>, A.G. Myagkov <sup>37,a</sup>, A.J. Myers <sup>8</sup>, G. Myers <sup>107</sup>, M. Myska <sup>133</sup>, B.P. Nachman <sup>17a</sup>, O. Nackenhorst <sup>49</sup>, K. Nagai <sup>127</sup>, K. Nagano <sup>84</sup>, J.L. Nagle <sup>29,ag</sup>, E. Nagy <sup>103</sup>, A.M. Nairz <sup>36</sup>, Y. Nakahama <sup>84</sup>, K. Nakamura <sup>84</sup>, K. Nakkalil <sup>5</sup>, H. Nanjo <sup>125</sup>, R. Narayan <sup>44</sup>, E.A. Narayanan <sup>113</sup>, I. Naryshkin <sup>37</sup>, M. Naseri <sup>34</sup>, S. Nasri <sup>117b</sup>, C. Nass <sup>24</sup>, G. Navarro <sup>22a</sup>, J. Navarro-Gonzalez <sup>164</sup>, R. Nayak <sup>152</sup>, A. Nayaz <sup>18</sup>, P.Y. Nechaeva <sup>37</sup>, S. Nechaeva <sup>23b,23a</sup>, F. Nechansky <sup>48</sup>, L. Nedic <sup>127</sup>, T.J. Neep <sup>20</sup>, A. Negri <sup>73a,73b</sup>, M. Negrini <sup>23b</sup>, C. Nellist <sup>115</sup>, C. Nelson <sup>105</sup>, K. Nelson <sup>107</sup>, S. Nemecek <sup>132</sup>, M. Nessi <sup>36,h</sup>, M.S. Neubauer <sup>163</sup>, F. Neuhaus <sup>101</sup>, J. Neundorf <sup>48</sup>, R. Newhouse <sup>165</sup>, P.R. Newman <sup>20</sup>, C.W. Ng <sup>130</sup>, Y.W.Y. Ng <sup>48</sup>, B. Ngair <sup>117a</sup>, H.D.N. Nguyen <sup>109</sup>, R.B. Nickerson <sup>127</sup>, R. Nicolaidou <sup>136</sup>, J. Nielsen <sup>137</sup>, M. Niemeyer <sup>55</sup>, J. Niermann <sup>55</sup>, N. Nikiforou <sup>36</sup>, V. Nikolaenko <sup>37,a</sup>, I. Nikolic-Audit <sup>128</sup>, K. Nikolopoulos <sup>20</sup>, P. Nilsson <sup>29</sup>, I. Ninca <sup>48</sup>, H.R. Nindhito <sup>56</sup>, G. Ninio <sup>152</sup>, A. Nisati <sup>75a</sup>, N. Nishu <sup>2</sup>, R. Nisius <sup>111</sup>, J-E. Nitschke <sup>50</sup>, E.K. Nkadimeng <sup>33g</sup>, T. Nobe <sup>154</sup>, D.L. Noel <sup>32</sup>, T. Nommensen <sup>148</sup>, M.B. Norfolk <sup>140</sup>, R.R.B. Norisam <sup>97</sup>, B.J. Norman <sup>34</sup>, M. Noury <sup>35a</sup>, J. Novak <sup>94</sup>, T. Novak <sup>48</sup>, L. Novotny <sup>133</sup>, R. Novotny <sup>113</sup>, L. Nozka <sup>123</sup>, K. Ntekas <sup>160</sup>, N.M.J. Nunes De Moura Junior <sup>83b</sup>, J. Ocariz <sup>128</sup>, A. Ochi <sup>85</sup>, I. Ochoa <sup>131a</sup>, S. Oerdek <sup>48,t</sup>, J.T. Offermann <sup>39</sup>, A. Ogrodnik <sup>134</sup>, A. Oh <sup>102</sup>, C.C. Ohm <sup>145</sup>, H. Oide <sup>84</sup>, R. Oishi <sup>154</sup>, M.L. Ojeda <sup>48</sup>, Y. Okumura <sup>154</sup>, L.F. Oleiro Seabra <sup>131a</sup>, S.A. Olivares Pino <sup>138d</sup>, G. Oliveira Correa <sup>13</sup>, D. Oliveira Damazio <sup>29</sup>, D. Oliveira Goncalves <sup>83a</sup>, J.L. Oliver <sup>160</sup>, Ö.O. Öncel <sup>54</sup>, A.P. O'Neill <sup>19</sup>, A. Onofre <sup>131a,131e</sup>, P.U.E. Onyisi <sup>11</sup>, M.J. Oreglia <sup>39</sup>, G.E. Orellana <sup>91</sup>, D. Orestano <sup>77a,77b</sup>, N. Orlando <sup>13</sup>, R.S. Orr <sup>156</sup>, V. O'Shea <sup>59</sup>, L.M. Osojnak <sup>129</sup>, R. Ospanov <sup>62a</sup>, G. Otero y Garzon <sup>30</sup>, H. Otono <sup>89</sup>, P.S. Ott <sup>63a</sup>, G.J. Ottino <sup>17a</sup>, M. Ouchrif <sup>35d</sup>, F. Ould-Saada <sup>126</sup>, T. Ovsiannikova <sup>139</sup>, M. Owen <sup>59</sup>, R.E. Owen <sup>135</sup>, K.Y. Oyulmaz <sup>21a</sup>, V.E. Ozcan <sup>21a</sup>, F. Ozturk <sup>87</sup>, N. Ozturk <sup>8</sup>, S. Ozturk <sup>82</sup>, H.A. Pacey <sup>127</sup>, A. Pacheco Pages <sup>13</sup>, C. Padilla Aranda <sup>13</sup>, G. Padovano <sup>75a,75b</sup>, S. Pagan Griso <sup>17a</sup>, G. Palacino <sup>68</sup>, A. Palazzo <sup>70a,70b</sup>, J. Pampel <sup>24</sup>, J. Pan <sup>173</sup>, T. Pan <sup>64a</sup>, D.K. Panchal <sup>11</sup>, C.E. Pandini <sup>115</sup>, J.G. Panduro Vazquez <sup>96</sup>, H.D. Pandya <sup>1</sup>, H. Pang <sup>14b</sup>, P. Pani <sup>48</sup>, G. Panizzo <sup>69a,69c</sup>, L. Panwar <sup>128</sup>, L. Paolozzi <sup>56</sup>, S. Parajuli <sup>163</sup>, A. Paramonov <sup>6</sup>, C. Paraskevopoulos <sup>53</sup>, D. Paredes Hernandez <sup>64b</sup>, A. Pareti <sup>73a,73b</sup>, K.R. Park <sup>41</sup>, T.H. Park <sup>156</sup>, M.A. Parker <sup>32</sup>, F. Parodi <sup>57b,57a</sup>, E.W. Parrish <sup>116</sup>, V.A. Parrish <sup>52</sup>, J.A. Parsons <sup>41</sup>, U. Parzefall <sup>54</sup>, B. Pascual Dias <sup>109</sup>, L. Pascual Dominguez <sup>152</sup>, E. Pasqualucci <sup>75a</sup>, S. Passaggio <sup>57b</sup>, F. Pastore <sup>96</sup>, P. Patel <sup>87</sup>, U.M. Patel <sup>51</sup>, J.R. Pater <sup>102</sup>, T. Pauly <sup>36</sup>, C.I. Pazos <sup>159</sup>, J. Pearkes <sup>144</sup>, M. Pedersen <sup>126</sup>, R. Pedro <sup>131a</sup>, S.V. Peleganchuk <sup>37</sup>, O. Penc <sup>36</sup>, E.A. Pender <sup>52</sup>, G.D. Penn <sup>173</sup>, K.E. Pensi <sup>110</sup>, M. Penzin <sup>37</sup>, B.S. Peralva <sup>83d</sup>, A.P. Pereira Peixoto <sup>139</sup>, L. Pereira Sanchez <sup>144</sup>, D.V. Perepelitsa <sup>29,ag</sup>, E. Perez Codina <sup>157a</sup>, M. Perganti <sup>10</sup>, H. Pernegger <sup>36</sup>, O. Perrin <sup>40</sup>, K. Peters <sup>48</sup>, R.F.Y. Peters <sup>102</sup>, B.A. Petersen <sup>36</sup>, T.C. Petersen <sup>42</sup>, E. Petit <sup>103</sup>, V. Petousis <sup>133</sup>, C. Petridou <sup>153,e</sup>, T. Petru <sup>134</sup>, A. Petrukhin <sup>142</sup>, M. Pettee <sup>17a</sup>, N.E. Pettersson <sup>36</sup>, A. Petukhov <sup>37</sup>, K. Petukhova <sup>134</sup>, R. Pezoa <sup>138f</sup>, L. Pezzotti <sup>36</sup>, G. Pezzullo <sup>173</sup>, T.M. Pham <sup>171</sup>, T. Pham <sup>106</sup>, P.W. Phillips <sup>135</sup>, G. Piacquadio <sup>146</sup>, E. Pianori <sup>17a</sup>, F. Piazza <sup>124</sup>, R. Piegai <sup>30</sup>, D. Pietreanu <sup>27b</sup>, A.D. Pilkington <sup>102</sup>, M. Pinamonti <sup>69a,69c</sup>, J.L. Pinfeld <sup>2</sup>, B.C. Pinheiro Pereira <sup>131a</sup>, A.E. Pinto Pinoargote <sup>101,136</sup>, L. Pintucci <sup>69a,69c</sup>, K.M. Piper <sup>147</sup>, A. Pirttikoski <sup>56</sup>, D.A. Pizzi <sup>34</sup>, L. Pizzimento <sup>64b</sup>, A. Pizzini <sup>115</sup>, M.-A. Pleier <sup>29</sup>, V. Plesanovs<sup>54</sup>, V. Pleskot <sup>134</sup>, E. Plotnikova<sup>38</sup>, G. Poddar <sup>95</sup>, R. Poettgen <sup>99</sup>, L. Poggioli <sup>128</sup>, I. Pokharel <sup>55</sup>, S. Polacek <sup>134</sup>, G. Polesello <sup>73a</sup>, A. Poley <sup>143,157a</sup>, A. Polini <sup>23b</sup>, C.S. Pollard <sup>168</sup>, Z.B. Pollock <sup>120</sup>, E. Pompa Pacchi <sup>75a,75b</sup>, D. Ponomarenko <sup>114</sup>, L. Pontecorvo <sup>36</sup>, S. Popa <sup>27a</sup>, G.A. Popeneciu <sup>27d</sup>, A. Poreba <sup>36</sup>, D.M. Portillo Quintero <sup>157a</sup>,

S. Pospisil <sup>133</sup>, M.A. Postill <sup>140</sup>, P. Postolache <sup>27c</sup>, K. Potamianos <sup>168</sup>, P.A. Potepa <sup>86a</sup>,  
 I.N. Potrap <sup>38</sup>, C.J. Potter <sup>32</sup>, H. Potti <sup>1</sup>, J. Poveda <sup>164</sup>, M.E. Pozo Astigarraga <sup>36</sup>,  
 A. Prades Ibanez <sup>164</sup>, J. Pretel <sup>54</sup>, D. Price <sup>102</sup>, M. Primavera <sup>70a</sup>, M.A. Principe Martin <sup>100</sup>,  
 R. Privara <sup>123</sup>, T. Procter <sup>59</sup>, M.L. Proffitt <sup>139</sup>, N. Proklova <sup>129</sup>, K. Prokofiev <sup>64c</sup>, G. Proto <sup>111</sup>,  
 J. Proudfoot <sup>6</sup>, M. Przybycien <sup>86a</sup>, W.W. Przygoda <sup>86b</sup>, A. Psallidas <sup>46</sup>, J.E. Puddefoot <sup>140</sup>,  
 D. Pudzha <sup>37</sup>, D. Pyatiizbyantseva <sup>37</sup>, J. Qian <sup>107</sup>, D. Qichen <sup>102</sup>, Y. Qin <sup>13</sup>, T. Qiu <sup>52</sup>,  
 A. Quadt <sup>55</sup>, M. Queitsch-Maitland <sup>102</sup>, G. Quetant <sup>56</sup>, R.P. Quinn <sup>165</sup>, G. Rabanal Bolanos <sup>61</sup>,  
 D. Rafanoharana <sup>54</sup>, F. Ragusa <sup>71a,71b</sup>, J.L. Rainbolt <sup>39</sup>, J.A. Raine <sup>56</sup>, S. Rajagopalan <sup>29</sup>,  
 E. Ramakoti <sup>37</sup>, I.A. Ramirez-Berend <sup>34</sup>, K. Ran <sup>48,14e</sup>, N.P. Rapheeha <sup>33g</sup>, H. Rasheed <sup>27b</sup>,  
 V. Raskina <sup>128</sup>, D.F. Rassloff <sup>63a</sup>, A. Rastogi <sup>17a</sup>, S. Rave <sup>101</sup>, B. Ravina <sup>55</sup>, I. Ravinovich <sup>170</sup>,  
 M. Raymond <sup>36</sup>, A.L. Read <sup>126</sup>, N.P. Readioff <sup>140</sup>, D.M. Rebutzi <sup>73a,73b</sup>, G. Redlinger <sup>29</sup>,  
 A.S. Reed <sup>111</sup>, K. Reeves <sup>26</sup>, J.A. Reidelsturz <sup>172</sup>, D. Reikher <sup>152</sup>, A. Rej <sup>49</sup>, C. Rembser <sup>36</sup>,  
 M. Renda <sup>27b</sup>, M.B. Rendel <sup>111</sup>, F. Renner <sup>48</sup>, A.G. Rennie <sup>160</sup>, A.L. Rescia <sup>48</sup>, S. Resconi <sup>71a</sup>,  
 M. Ressegotti <sup>57b,57a</sup>, S. Rettie <sup>36</sup>, J.G. Reyes Rivera <sup>108</sup>, E. Reynolds <sup>17a</sup>, O.L. Rezanova <sup>37</sup>,  
 P. Reznicek <sup>134</sup>, H. Riani <sup>35d</sup>, N. Ribaric <sup>92</sup>, E. Ricci <sup>78a,78b</sup>, R. Richter <sup>111</sup>, S. Richter <sup>47a,47b</sup>,  
 E. Richter-Was <sup>86b</sup>, M. Ridel <sup>128</sup>, S. Ridouani <sup>35d</sup>, P. Rieck <sup>118</sup>, P. Riedler <sup>36</sup>, E.M. Riefel <sup>47a,47b</sup>,  
 J.O. Rieger <sup>115</sup>, M. Rijssenbeek <sup>146</sup>, M. Rimoldi <sup>36</sup>, L. Rinaldi <sup>23b,23a</sup>, T.T. Rinn <sup>29</sup>,  
 M.P. Rinnagel <sup>110</sup>, G. Ripellino <sup>162</sup>, I. Riu <sup>13</sup>, J.C. Rivera Vergara <sup>166</sup>, F. Rizatdinova <sup>122</sup>,  
 E. Rizvi <sup>95</sup>, B.R. Roberts <sup>17a</sup>, S.H. Robertson <sup>105,w</sup>, D. Robinson <sup>32</sup>, C.M. Robles Gajardo <sup>138f</sup>,  
 M. Robles Manzano <sup>101</sup>, A. Robson <sup>59</sup>, A. Rocchi <sup>76a,76b</sup>, C. Roda <sup>74a,74b</sup>, S. Rodriguez Bosca <sup>36</sup>,  
 Y. Rodriguez Garcia <sup>22a</sup>, A. Rodriguez Rodriguez <sup>54</sup>, A.M. Rodríguez Vera <sup>116</sup>, S. Roe <sup>36</sup>,  
 J.T. Roemer <sup>160</sup>, A.R. Roepe-Gier <sup>137</sup>, J. Roggel <sup>172</sup>, O. Røhne <sup>126</sup>, R.A. Rojas <sup>104</sup>,  
 C.P.A. Roland <sup>128</sup>, J. Roloff <sup>29</sup>, A. Romaniouk <sup>37</sup>, E. Romano <sup>73a,73b</sup>, M. Romano <sup>23b</sup>,  
 A.C. Romero Hernandez <sup>163</sup>, N. Rompotis <sup>93</sup>, L. Roos <sup>128</sup>, S. Rosati <sup>75a</sup>, B.J. Rosser <sup>39</sup>,  
 E. Rossi <sup>127</sup>, E. Rossi <sup>72a,72b</sup>, L.P. Rossi <sup>61</sup>, L. Rossini <sup>54</sup>, R. Rosten <sup>120</sup>, M. Rotaru <sup>27b</sup>,  
 B. Rottler <sup>54</sup>, C. Rougier <sup>90</sup>, D. Rousseau <sup>66</sup>, D. Rouso <sup>48</sup>, A. Roy <sup>163</sup>, S. Roy-Garand <sup>156</sup>,  
 A. Rozanov <sup>103</sup>, Z.M.A. Rozario <sup>59</sup>, Y. Rozen <sup>151</sup>, A. Rubio Jimenez <sup>164</sup>, A.J. Ruby <sup>93</sup>,  
 V.H. Ruelas Rivera <sup>18</sup>, T.A. Ruggeri <sup>1</sup>, A. Ruggiero <sup>127</sup>, A. Ruiz-Martinez <sup>164</sup>, A. Rummler <sup>36</sup>,  
 Z. Rurikova <sup>54</sup>, N.A. Rusakovich <sup>38</sup>, H.L. Russell <sup>166</sup>, G. Russo <sup>75a,75b</sup>, J.P. Rutherford <sup>7</sup>,  
 S. Rutherford Colmenares <sup>32</sup>, K. Rybacki <sup>92</sup>, M. Rybar <sup>134</sup>, E.B. Rye <sup>126</sup>, A. Ryzhov <sup>44</sup>,  
 J.A. Sabater Iglesias <sup>56</sup>, P. Sabatini <sup>164</sup>, H.F.W. Sadrozinski <sup>137</sup>, F. Safai Tehrani <sup>75a</sup>,  
 B. Safarzadeh Samani <sup>135</sup>, S. Saha <sup>1</sup>, M. Sahinsoy <sup>111</sup>, A. Saibel <sup>164</sup>, M. Saimpert <sup>136</sup>,  
 M. Saito <sup>154</sup>, T. Saito <sup>154</sup>, A. Sala <sup>71a,71b</sup>, D. Salamani <sup>36</sup>, A. Salnikov <sup>144</sup>, J. Salt <sup>164</sup>,  
 A. Salvador Salas <sup>152</sup>, D. Salvatore <sup>43b,43a</sup>, F. Salvatore <sup>147</sup>, A. Salzburger <sup>36</sup>, D. Sammel <sup>54</sup>,  
 E. Sampson <sup>92</sup>, D. Sampsonidis <sup>153,e</sup>, D. Sampsonidou <sup>124</sup>, J. Sánchez <sup>164</sup>,  
 V. Sanchez Sebastian <sup>164</sup>, H. Sandaker <sup>126</sup>, C.O. Sander <sup>48</sup>, J.A. Sandesara <sup>104</sup>, M. Sandhoff <sup>172</sup>,  
 C. Sandoval <sup>22b</sup>, D.P.C. Sankey <sup>135</sup>, T. Sano <sup>88</sup>, A. Sansoni <sup>53</sup>, L. Santi <sup>75a,75b</sup>, C. Santoni <sup>40</sup>,  
 H. Santos <sup>131a,131b</sup>, A. Santra <sup>170</sup>, K.A. Saoucha <sup>161</sup>, J.G. Saraiva <sup>131a,131d</sup>, J. Sardain <sup>7</sup>,  
 O. Sasaki <sup>84</sup>, K. Sato <sup>158</sup>, C. Sauer <sup>63b</sup>, F. Sauerburger <sup>54</sup>, E. Sauvan <sup>4</sup>, P. Savard <sup>156,ae</sup>,  
 R. Sawada <sup>154</sup>, C. Sawyer <sup>135</sup>, L. Sawyer <sup>98</sup>, I. Sayago Galvan <sup>164</sup>, C. Sbarra <sup>23b</sup>, A. Sbrizzi <sup>23b,23a</sup>,  
 T. Scanlon <sup>97</sup>, J. Schaarschmidt <sup>139</sup>, U. Schäfer <sup>101</sup>, A.C. Schaffer <sup>66,44</sup>, D. Schaile <sup>110</sup>,  
 R.D. Schamberger <sup>146</sup>, C. Scharf <sup>18</sup>, M.M. Schefer <sup>19</sup>, V.A. Schegelsky <sup>37</sup>, D. Scheirich <sup>134</sup>,  
 F. Schenck <sup>18</sup>, M. Schernau <sup>160</sup>, C. Scheulen <sup>55</sup>, C. Schiavi <sup>57b,57a</sup>, M. Schioppa <sup>43b,43a</sup>,  
 B. Schlag <sup>144,m</sup>, K.E. Schleicher <sup>54</sup>, S. Schlenker <sup>36</sup>, J. Schmeing <sup>172</sup>, M.A. Schmidt <sup>172</sup>,  
 K. Schmieden <sup>101</sup>, C. Schmitt <sup>101</sup>, N. Schmitt <sup>101</sup>, S. Schmitt <sup>48</sup>, L. Schoeffel <sup>136</sup>,  
 A. Schoening <sup>63b</sup>, P.G. Scholer <sup>34</sup>, E. Schopf <sup>127</sup>, M. Schott <sup>101</sup>, J. Schovancova <sup>36</sup>,  
 S. Schramm <sup>56</sup>, T. Schroer <sup>56</sup>, H-C. Schultz-Coulon <sup>63a</sup>, M. Schumacher <sup>54</sup>, B.A. Schumm <sup>137</sup>,

Ph. Schune <sup>136</sup>, A.J. Schuy <sup>139</sup>, H.R. Schwartz <sup>137</sup>, A. Schwartzman <sup>144</sup>, T.A. Schwarz <sup>107</sup>,  
 Ph. Schwemling <sup>136</sup>, R. Schwienhorst <sup>108</sup>, A. Sciandra <sup>29</sup>, G. Sciolla <sup>26</sup>, F. Scuri <sup>74a</sup>,  
 C.D. Sebastiani <sup>93</sup>, K. Sedlaczek <sup>116</sup>, P. Seema <sup>18</sup>, S.C. Seidel <sup>113</sup>, A. Seiden <sup>137</sup>,  
 B.D. Seidlitz <sup>41</sup>, C. Seitz <sup>48</sup>, J.M. Seixas <sup>83b</sup>, G. Sekhniaidze <sup>72a</sup>, L. Selem <sup>60</sup>,  
 N. Semprini-Cesari <sup>23b,23a</sup>, D. Sengupta <sup>56</sup>, V. Senthilkumar <sup>164</sup>, L. Serin <sup>66</sup>, L. Serkin <sup>69a,69b</sup>,  
 M. Sessa <sup>76a,76b</sup>, H. Severini <sup>121</sup>, F. Sforza <sup>57b,57a</sup>, A. Sfyrla <sup>56</sup>, Q. Sha <sup>14a</sup>, E. Shabalina <sup>55</sup>,  
 A.H. Shah <sup>32</sup>, R. Shaheen <sup>145</sup>, J.D. Shahinian <sup>129</sup>, D. Shaked Renous <sup>170</sup>, L.Y. Shan <sup>14a</sup>,  
 M. Shapiro <sup>17a</sup>, A. Sharma <sup>36</sup>, A.S. Sharma <sup>165</sup>, P. Sharma <sup>80</sup>, P.B. Shatalov <sup>37</sup>, K. Shaw <sup>147</sup>,  
 S.M. Shaw <sup>102</sup>, A. Shcherbakova <sup>37</sup>, Q. Shen <sup>62c,5</sup>, D.J. Sheppard <sup>143</sup>, P. Sherwood <sup>97</sup>, L. Shi <sup>97</sup>,  
 X. Shi <sup>14a</sup>, C.O. Shimmin <sup>173</sup>, J.D. Shinner <sup>96</sup>, I.P.J. Shipsey <sup>127</sup>, S. Shirabe <sup>89</sup>,  
 M. Shiyakova <sup>38,u</sup>, J. Shlomi <sup>170</sup>, M.J. Shochet <sup>39</sup>, J. Shojaii <sup>106</sup>, D.R. Shope <sup>126</sup>,  
 B. Shrestha <sup>121</sup>, S. Shrestha <sup>120,ah</sup>, E.M. Shrif <sup>33g</sup>, M.J. Shroff <sup>166</sup>, P. Sicho <sup>132</sup>,  
 A.M. Sickles <sup>163</sup>, E. Sideras Haddad <sup>33g</sup>, A.C. Sidley <sup>115</sup>, A. Sidoti <sup>23b</sup>, F. Siegert <sup>50</sup>,  
 Dj. Sijacki <sup>15</sup>, F. Sili <sup>91</sup>, J.M. Silva <sup>52</sup>, M.V. Silva Oliveira <sup>29</sup>, S.B. Silverstein <sup>47a</sup>, S. Simion <sup>66</sup>,  
 R. Simoniello <sup>36</sup>, E.L. Simpson <sup>102</sup>, H. Simpson <sup>147</sup>, L.R. Simpson <sup>107</sup>, N.D. Simpson <sup>99</sup>,  
 S. Simsek <sup>82</sup>, S. Sindhu <sup>55</sup>, P. Sinervo <sup>156</sup>, S. Singh <sup>156</sup>, S. Sinha <sup>48</sup>, S. Sinha <sup>102</sup>,  
 M. Sioli <sup>23b,23a</sup>, I. Siral <sup>36</sup>, E. Sitnikova <sup>48</sup>, J. Sjölin <sup>47a,47b</sup>, A. Skaf <sup>55</sup>, E. Skorda <sup>20</sup>,  
 P. Skubic <sup>121</sup>, M. Slawinska <sup>87</sup>, V. Smakhtin <sup>170</sup>, B.H. Smart <sup>135</sup>, S.Yu. Smirnov <sup>37</sup>, Y. Smirnov <sup>37</sup>,  
 L.N. Smirnova <sup>37,a</sup>, O. Smirnova <sup>99</sup>, A.C. Smith <sup>41</sup>, D.R. Smith <sup>160</sup>, E.A. Smith <sup>39</sup>,  
 H.A. Smith <sup>127</sup>, J.L. Smith <sup>102</sup>, R. Smith <sup>144</sup>, M. Smizanska <sup>92</sup>, K. Smolek <sup>133</sup>, A.A. Snesarev <sup>37</sup>,  
 S.R. Snider <sup>156</sup>, H.L. Snoek <sup>115</sup>, S. Snyder <sup>29</sup>, R. Sobie <sup>166,w</sup>, A. Soffer <sup>152</sup>,  
 C.A. Solans Sanchez <sup>36</sup>, E.Yu. Soldatov <sup>37</sup>, U. Soldevila <sup>164</sup>, A.A. Solodkov <sup>37</sup>, S. Solomon <sup>26</sup>,  
 A. Soloshenko <sup>38</sup>, K. Solovieva <sup>54</sup>, O.V. Solovyanov <sup>40</sup>, P. Sommer <sup>36</sup>, A. Sonay <sup>13</sup>,  
 W.Y. Song <sup>157b</sup>, A. Sopczak <sup>133</sup>, A.L. Sopio <sup>97</sup>, F. Sopkova <sup>28b</sup>, J.D. Sorenson <sup>113</sup>,  
 I.R. Sotarriva Alvarez <sup>155</sup>, V. Sothilingam <sup>63a</sup>, O.J. Soto Sandoval <sup>138c,138b</sup>, S. Sottocornola <sup>68</sup>,  
 R. Soualah <sup>161</sup>, Z. Soumami <sup>35e</sup>, D. South <sup>48</sup>, N. Soybelman <sup>170</sup>, S. Spagnolo <sup>70a,70b</sup>,  
 M. Spalla <sup>111</sup>, D. Sperlich <sup>54</sup>, G. Spigo <sup>36</sup>, S. Spinali <sup>92</sup>, D.P. Spiteri <sup>59</sup>, M. Spousta <sup>134</sup>,  
 E.J. Staats <sup>34</sup>, R. Stamen <sup>63a</sup>, A. Stampekis <sup>20</sup>, M. Standke <sup>24</sup>, E. Stanecka <sup>87</sup>,  
 W. Stanek-Maslouska <sup>48</sup>, M.V. Stange <sup>50</sup>, B. Stanislaus <sup>17a</sup>, M.M. Stanitzki <sup>48</sup>, B. Stapf <sup>48</sup>,  
 E.A. Starchenko <sup>37</sup>, G.H. Stark <sup>137</sup>, J. Stark <sup>90</sup>, P. Staroba <sup>132</sup>, P. Starovoitov <sup>63a</sup>, S. Stärz <sup>105</sup>,  
 R. Staszewski <sup>87</sup>, G. Stavropoulos <sup>46</sup>, J. Steentoft <sup>162</sup>, P. Steinberg <sup>29</sup>, B. Stelzer <sup>143,157a</sup>,  
 H.J. Stelzer <sup>130</sup>, O. Stelzer-Chilton <sup>157a</sup>, H. Stenzel <sup>58</sup>, T.J. Stevenson <sup>147</sup>, G.A. Stewart <sup>36</sup>,  
 J.R. Stewart <sup>122</sup>, M.C. Stockton <sup>36</sup>, G. Stoicea <sup>27b</sup>, M. Stolarski <sup>131a</sup>, S. Stonjek <sup>111</sup>,  
 A. Straessner <sup>50</sup>, J. Strandberg <sup>145</sup>, S. Strandberg <sup>47a,47b</sup>, M. Stratmann <sup>172</sup>, M. Strauss <sup>121</sup>,  
 T. Streblner <sup>103</sup>, P. Strizenec <sup>28b</sup>, R. Ströhmer <sup>167</sup>, D.M. Strom <sup>124</sup>, R. Stroynowski <sup>44</sup>,  
 A. Strubig <sup>47a,47b</sup>, S.A. Stucci <sup>29</sup>, B. Stugu <sup>16</sup>, J. Stupak <sup>121</sup>, N.A. Styles <sup>48</sup>, D. Su <sup>144</sup>,  
 S. Su <sup>62a</sup>, W. Su <sup>62d</sup>, X. Su <sup>62a</sup>, D. Suchy <sup>28a</sup>, K. Sugizaki <sup>154</sup>, V.V. Sulin <sup>37</sup>, M.J. Sullivan <sup>93</sup>,  
 D.M.S. Sultan <sup>127</sup>, L. Sultanaliyeva <sup>37</sup>, S. Sultansoy <sup>3b</sup>, T. Sumida <sup>88</sup>, S. Sun <sup>107</sup>, S. Sun <sup>171</sup>,  
 O. Sunneborn Gudnadottir <sup>162</sup>, N. Sur <sup>103</sup>, M.R. Sutton <sup>147</sup>, H. Suzuki <sup>158</sup>, M. Svatos <sup>132</sup>,  
 M. Swiatlowski <sup>157a</sup>, T. Swirski <sup>167</sup>, I. Sykora <sup>28a</sup>, M. Sykora <sup>134</sup>, T. Sykora <sup>134</sup>, D. Ta <sup>101</sup>,  
 K. Tackmann <sup>48,t</sup>, A. Taffard <sup>160</sup>, R. Tafirout <sup>157a</sup>, J.S. Tafoya Vargas <sup>66</sup>, Y. Takubo <sup>84</sup>,  
 M. Talby <sup>103</sup>, A.A. Talyshv <sup>37</sup>, K.C. Tam <sup>64b</sup>, N.M. Tamir <sup>152</sup>, A. Tanaka <sup>154</sup>, J. Tanaka <sup>154</sup>,  
 R. Tanaka <sup>66</sup>, M. Tanasini <sup>57b,57a</sup>, Z. Tao <sup>165</sup>, S. Tapia Araya <sup>138f</sup>, S. Tapprogge <sup>101</sup>,  
 A. Tarek Abouelfadl Mohamed <sup>108</sup>, S. Tarem <sup>151</sup>, K. Tariq <sup>14a</sup>, G. Tarna <sup>27b</sup>, G.F. Tartarelli <sup>71a</sup>,  
 M.J. Tartarin <sup>90</sup>, P. Tas <sup>134</sup>, M. Tasevsky <sup>132</sup>, E. Tassi <sup>43b,43a</sup>, A.C. Tate <sup>163</sup>, G. Tateno <sup>154</sup>,  
 Y. Tayalati <sup>35e,v</sup>, G.N. Taylor <sup>106</sup>, W. Taylor <sup>157b</sup>, A.S. Tee <sup>171</sup>, R. Teixeira De Lima <sup>144</sup>,  
 P. Teixeira-Dias <sup>96</sup>, J.J. Teoh <sup>156</sup>, K. Terashi <sup>154</sup>, J. Terron <sup>100</sup>, S. Terzo <sup>13</sup>, M. Testa <sup>53</sup>,

R.J. Teuscher <sup>156,w</sup>, A. Thaler <sup>79</sup>, O. Theiner <sup>56</sup>, N. Themistokleous <sup>52</sup>, T. Thevenaux-Pelzer <sup>103</sup>, O. Thielmann <sup>172</sup>, D.W. Thomas <sup>96</sup>, J.P. Thomas <sup>20</sup>, E.A. Thompson <sup>17a</sup>, P.D. Thompson <sup>20</sup>, E. Thomson <sup>129</sup>, R.E. Thornberry <sup>44</sup>, Y. Tian <sup>55</sup>, V. Tikhomirov <sup>37,a</sup>, Yu.A. Tikhonov <sup>37</sup>, S. Timoshenko <sup>37</sup>, D. Timoshyn <sup>134</sup>, E.X.L. Ting <sup>1</sup>, P. Tipton <sup>173</sup>, S.H. Tlou <sup>33g</sup>, K. Todome <sup>155</sup>, S. Todorova-Nova <sup>134</sup>, S. Todt <sup>50</sup>, M. Togawa <sup>84</sup>, J. Tojo <sup>89</sup>, S. Tokár <sup>28a</sup>, K. Tokushuku <sup>84</sup>, O. Toldaiev <sup>68</sup>, R. Tombs <sup>32</sup>, M. Tomoto <sup>84,112</sup>, L. Tompkins <sup>144,m</sup>, K.W. Topolnicki <sup>86b</sup>, E. Torrence <sup>124</sup>, H. Torres <sup>90</sup>, E. Torró Pastor <sup>164</sup>, M. Toscani <sup>30</sup>, C. Toscirì <sup>39</sup>, M. Tost <sup>11</sup>, D.R. Tovey <sup>140</sup>, A. Traeet <sup>16</sup>, I.S. Trandafir <sup>27b</sup>, T. Trefzger <sup>167</sup>, A. Tricoli <sup>29</sup>, I.M. Trigger <sup>157a</sup>, S. Trincaz-Duvoid <sup>128</sup>, D.A. Trischuk <sup>26</sup>, B. Trocmé <sup>60</sup>, L. Truong <sup>33c</sup>, M. Trzebinski <sup>87</sup>, A. Trzupke <sup>87</sup>, F. Tsai <sup>146</sup>, M. Tsai <sup>107</sup>, A. Tsiamis <sup>153,e</sup>, P.V. Tsiareshka <sup>37</sup>, S. Tsigaridas <sup>157a</sup>, A. Tsirogitis <sup>153,r</sup>, V. Tsiskaridze <sup>156</sup>, E.G. Tskhadadze <sup>150a</sup>, M. Tsopoulou <sup>153</sup>, Y. Tsujikawa <sup>88</sup>, I.I. Tsukerman <sup>37</sup>, V. Tsulaia <sup>17a</sup>, S. Tsuno <sup>84</sup>, K. Tsurì <sup>119</sup>, D. Tsybychev <sup>146</sup>, Y. Tu <sup>64b</sup>, A. Tudorache <sup>27b</sup>, V. Tudorache <sup>27b</sup>, A.N. Tuna <sup>61</sup>, S. Turchikhin <sup>57b,57a</sup>, I. Turk Cakir <sup>3a</sup>, R. Turra <sup>71a</sup>, T. Turtuvshin <sup>38,x</sup>, P.M. Tuts <sup>41</sup>, S. Tzamarias <sup>153,e</sup>, E. Tzovara <sup>101</sup>, F. Ukegawa <sup>158</sup>, P.A. Ulloa Poblete <sup>138c,138b</sup>, E.N. Umaka <sup>29</sup>, G. Unal <sup>36</sup>, A. Undrus <sup>29</sup>, G. Unel <sup>160</sup>, J. Urban <sup>28b</sup>, P. Urquijo <sup>106</sup>, P. Urrejola <sup>138a</sup>, G. Usai <sup>8</sup>, R. Ushioda <sup>155</sup>, M. Usman <sup>109</sup>, Z. Uysal <sup>82</sup>, V. Vacek <sup>133</sup>, B. Vachon <sup>105</sup>, K.O.H. Vadla <sup>126</sup>, T. Vafeiadis <sup>36</sup>, A. Vaitkus <sup>97</sup>, C. Valderanis <sup>110</sup>, E. Valdes Santurio <sup>47a,47b</sup>, M. Valente <sup>157a</sup>, S. Valentinetti <sup>23b,23a</sup>, A. Valero <sup>164</sup>, E. Valiente Moreno <sup>164</sup>, A. Vallier <sup>90</sup>, J.A. Valls Ferrer <sup>164</sup>, D.R. Van Arneman <sup>115</sup>, T.R. Van Daalen <sup>139</sup>, A. Van Der Graaf <sup>49</sup>, P. Van Gemmeren <sup>6</sup>, M. Van Rijnbach <sup>126</sup>, S. Van Stroud <sup>97</sup>, I. Van Vulpen <sup>115</sup>, P. Vana <sup>134</sup>, M. Vanadia <sup>76a,76b</sup>, W. Vandelli <sup>36</sup>, E.R. Vandewall <sup>122</sup>, D. Vannicola <sup>152</sup>, L. Vannoli <sup>53</sup>, R. Vari <sup>75a</sup>, E.W. Varnes <sup>7</sup>, C. Varni <sup>17b</sup>, T. Varol <sup>149</sup>, D. Varouchas <sup>66</sup>, L. Varriale <sup>164</sup>, K.E. Varvell <sup>148</sup>, M.E. Vasile <sup>27b</sup>, L. Vaslin <sup>84</sup>, G.A. Vasquez <sup>166</sup>, A. Vasyukov <sup>38</sup>, R. Vavricka <sup>101</sup>, F. Vazeille <sup>40</sup>, T. Vazquez Schroeder <sup>36</sup>, J. Veatch <sup>31</sup>, V. Vecchio <sup>102</sup>, M.J. Veen <sup>104</sup>, I. Veliscek <sup>29</sup>, L.M. Veloce <sup>156</sup>, F. Veloso <sup>131a,131c</sup>, S. Veneziano <sup>75a</sup>, A. Ventura <sup>70a,70b</sup>, S. Ventura Gonzalez <sup>136</sup>, A. Verbytskyi <sup>111</sup>, M. Verducci <sup>74a,74b</sup>, C. Vergis <sup>95</sup>, M. Verissimo De Araujo <sup>83b</sup>, W. Verkerke <sup>115</sup>, J.C. Vermeulen <sup>115</sup>, C. Vernieri <sup>144</sup>, M. Vessella <sup>104</sup>, M.C. Vetterli <sup>143,ae</sup>, A. Vgenopoulos <sup>153,e</sup>, N. Viaux Maira <sup>138f</sup>, T. Vickey <sup>140</sup>, O.E. Vickey Boeriu <sup>140</sup>, G.H.A. Viehhauser <sup>127</sup>, L. Vignani <sup>63b</sup>, M. Villa <sup>23b,23a</sup>, M. Villaplana Perez <sup>164</sup>, E.M. Villhauer <sup>52</sup>, E. Vilucchi <sup>53</sup>, M.G. Vincter <sup>34</sup>, G.S. Virdee <sup>20</sup>, A. Visibile <sup>115</sup>, C. Vittori <sup>36</sup>, I. Vivarelli <sup>23b,23a</sup>, E. Voevodina <sup>111</sup>, F. Vogel <sup>110</sup>, J.C. Voigt <sup>50</sup>, P. Vokac <sup>133</sup>, Yu. Volkotrub <sup>86b</sup>, J. Von Ahnen <sup>48</sup>, E. Von Toerne <sup>24</sup>, B. Vormwald <sup>36</sup>, V. Vorobel <sup>134</sup>, K. Vorobev <sup>37</sup>, M. Vos <sup>164</sup>, K. Voss <sup>142</sup>, M. Vozak <sup>115</sup>, L. Vozdecky <sup>121</sup>, N. Vranjes <sup>15</sup>, M. Vranjes Milosavljevic <sup>15</sup>, M. Vreeswijk <sup>115</sup>, N.K. Vu <sup>62d,62c</sup>, R. Vuillermet <sup>36</sup>, O. Vujinovic <sup>101</sup>, I. Vukotic <sup>39</sup>, S. Wada <sup>158</sup>, C. Wagner <sup>104</sup>, J.M. Wagner <sup>17a</sup>, W. Wagner <sup>172</sup>, S. Wahdan <sup>172</sup>, H. Wahlberg <sup>91</sup>, M. Wakida <sup>112</sup>, J. Walder <sup>135</sup>, R. Walker <sup>110</sup>, W. Walkowiak <sup>142</sup>, A. Wall <sup>129</sup>, E.J. Wallin <sup>99</sup>, T. Wamorkar <sup>6</sup>, A.Z. Wang <sup>137</sup>, C. Wang <sup>101</sup>, C. Wang <sup>11</sup>, H. Wang <sup>17a</sup>, J. Wang <sup>64c</sup>, R.-J. Wang <sup>101</sup>, R. Wang <sup>61</sup>, R. Wang <sup>6</sup>, S.M. Wang <sup>149</sup>, S. Wang <sup>62b</sup>, S. Wang <sup>14a</sup>, T. Wang <sup>62a</sup>, W.T. Wang <sup>80</sup>, W. Wang <sup>14a</sup>, X. Wang <sup>14c</sup>, X. Wang <sup>163</sup>, X. Wang <sup>62c</sup>, Y. Wang <sup>62d</sup>, Y. Wang <sup>14c</sup>, Z. Wang <sup>107</sup>, Z. Wang <sup>62d,51,62c</sup>, Z. Wang <sup>107</sup>, A. Warburton <sup>105</sup>, R.J. Ward <sup>20</sup>, N. Warrack <sup>59</sup>, S. Waterhouse <sup>96</sup>, A.T. Watson <sup>20</sup>, H. Watson <sup>59</sup>, M.F. Watson <sup>20</sup>, E. Watton <sup>59,135</sup>, G. Watts <sup>139</sup>, B.M. Waugh <sup>97</sup>, J.M. Webb <sup>54</sup>, C. Weber <sup>29</sup>, H.A. Weber <sup>18</sup>, M.S. Weber <sup>19</sup>, S.M. Weber <sup>63a</sup>, C. Wei <sup>62a</sup>, Y. Wei <sup>127</sup>, A.R. Weidberg <sup>127</sup>, E.J. Weik <sup>118</sup>, J. Weingarten <sup>49</sup>, M. Weirich <sup>101</sup>, C. Weiser <sup>54</sup>, C.J. Wells <sup>48</sup>, T. Wenaus <sup>29</sup>, B. Wendland <sup>49</sup>, T. Wengler <sup>36</sup>, N.S. Wenke <sup>111</sup>, N. Wermes <sup>24</sup>, M. Wessels <sup>63a</sup>, A.M. Wharton <sup>92</sup>, A.S. White <sup>61</sup>, A. White <sup>8</sup>, M.J. White <sup>1</sup>, D. Whiteson <sup>160</sup>, L. Wickremasinghe <sup>125</sup>, W. Wiedenmann <sup>171</sup>, M. Wielers <sup>135</sup>,

C. Wigglesworth , D.J. Wilbern<sup>121</sup>, H.G. Wilkens , J.J.H. Wilkinson , D.M. Williams , H.H. Williams<sup>129</sup>, S. Williams , S. Willocq , B.J. Wilson , P.J. Windischhofer , F.I. Winkel , F. Winklmeier , B.T. Winter , J.K. Winter , M. Wittgen<sup>144</sup>, M. Wobisch , T. Wojtkowski<sup>60</sup>, Z. Wolffs , J. Wollrath<sup>160</sup>, M.W. Wolter , H. Wolters , M.C. Wong<sup>137</sup>, E.L. Woodward , S.D. Worm , B.K. Wosiek , K.W. Woźniak , S. Wozniowski , K. Wraight , C. Wu , M. Wu , M. Wu , S.L. Wu , X. Wu , Y. Wu , Z. Wu , J. Wuerzinger , T.R. Wyatt , B.M. Wynne , S. Xella , L. Xia , M. Xia , J. Xiang , M. Xie , X. Xie , S. Xin , A. Xiong , J. Xiong , D. Xu , H. Xu , L. Xu , R. Xu , T. Xu , Y. Xu , Z. Xu , Z. Xu , B. Yabsley , S. Yacoob , Y. Yamaguchi , E. Yamashita , H. Yamauchi , T. Yamazaki , Y. Yamazaki , J. Yan , S. Yan , Z. Yan , H.J. Yang , H.T. Yang , S. Yang , T. Yang , X. Yang , X. Yang , Y. Yang , Y. Yang , Z. Yang , W-M. Yao , H. Ye , H. Ye , J. Ye , S. Ye , X. Ye , Y. Yeh , I. Yeletsikh , B.K. Yeo , M.R. Yexley , T.P. Yildirim , P. Yin , K. Yorita , S. Younas , C.J.S. Young , C. Young , C. Yu , Y. Yu , M. Yuan , R. Yuan , L. Yue , M. Zaazoua , B. Zabinski , E. Zaid<sup>52</sup>, Z.K. Zak , T. Zakareishvili , N. Zakharchuk , S. Zambito , J.A. Zamora Saa , J. Zang , D. Zanzi , O. Zaplatilek , C. Zeitnitz , H. Zeng , J.C. Zeng , D.T. Zenger Jr , O. Zenin , T. Ženiš , S. Zenz , S. Zerradi , D. Zerwas , M. Zhai , D.F. Zhang , J. Zhang , J. Zhang , K. Zhang , L. Zhang , L. Zhang , P. Zhang , R. Zhang , S. Zhang , T. Zhang , X. Zhang , X. Zhang , Y. Zhang , Y. Zhang , Y. Zhang , Z. Zhang , Z. Zhang , H. Zhao , T. Zhao , Y. Zhao , Z. Zhao , Z. Zhao , A. Zhemchugov , J. Zheng , K. Zheng , X. Zheng , Z. Zheng , D. Zhong , B. Zhou , H. Zhou , N. Zhou , Y. Zhou , Y. Zhou<sup>7</sup>, C.G. Zhu , J. Zhu , Y. Zhu , Y. Zhu , X. Zhuang , K. Zhukov , N.I. Zimine , J. Zinsser , M. Ziolkowski , L. Živković , A. Zoccoli , K. Zoch , T.G. Zorbas , O. Zormpa , W. Zou , L. Zwalinski .

<sup>1</sup>Department of Physics, University of Adelaide, Adelaide; Australia.

<sup>2</sup>Department of Physics, University of Alberta, Edmonton AB; Canada.

<sup>3</sup>(<sup>a</sup>)Department of Physics, Ankara University, Ankara; (<sup>b</sup>)Division of Physics, TOBB University of Economics and Technology, Ankara; Türkiye.

<sup>4</sup>LAPP, Université Savoie Mont Blanc, CNRS/IN2P3, Annecy; France.

<sup>5</sup>APC, Université Paris Cité, CNRS/IN2P3, Paris; France.

<sup>6</sup>High Energy Physics Division, Argonne National Laboratory, Argonne IL; United States of America.

<sup>7</sup>Department of Physics, University of Arizona, Tucson AZ; United States of America.

<sup>8</sup>Department of Physics, University of Texas at Arlington, Arlington TX; United States of America.

<sup>9</sup>Physics Department, National and Kapodistrian University of Athens, Athens; Greece.

<sup>10</sup>Physics Department, National Technical University of Athens, Zografou; Greece.

<sup>11</sup>Department of Physics, University of Texas at Austin, Austin TX; United States of America.

<sup>12</sup>Institute of Physics, Azerbaijan Academy of Sciences, Baku; Azerbaijan.

<sup>13</sup>Institut de Física d'Altes Energies (IFAE), Barcelona Institute of Science and Technology, Barcelona; Spain.

<sup>14</sup>(<sup>a</sup>)Institute of High Energy Physics, Chinese Academy of Sciences, Beijing; (<sup>b</sup>)Physics Department, Tsinghua University, Beijing; (<sup>c</sup>)Department of Physics, Nanjing University, Nanjing; (<sup>d</sup>)School of Science, Shenzhen Campus of Sun Yat-sen University; (<sup>e</sup>)University of Chinese Academy of Science (UCAS), Beijing; China.



- <sup>15</sup>Institute of Physics, University of Belgrade, Belgrade; Serbia.
- <sup>16</sup>Department for Physics and Technology, University of Bergen, Bergen; Norway.
- <sup>17</sup>(<sup>a</sup>)Physics Division, Lawrence Berkeley National Laboratory, Berkeley CA;(<sup>b</sup>)University of California, Berkeley CA; United States of America.
- <sup>18</sup>Institut für Physik, Humboldt Universität zu Berlin, Berlin; Germany.
- <sup>19</sup>Albert Einstein Center for Fundamental Physics and Laboratory for High Energy Physics, University of Bern, Bern; Switzerland.
- <sup>20</sup>School of Physics and Astronomy, University of Birmingham, Birmingham; United Kingdom.
- <sup>21</sup>(<sup>a</sup>)Department of Physics, Bogazici University, Istanbul;(<sup>b</sup>)Department of Physics Engineering, Gaziantep University, Gaziantep;(<sup>c</sup>)Department of Physics, Istanbul University, Istanbul; Türkiye.
- <sup>22</sup>(<sup>a</sup>)Facultad de Ciencias y Centro de Investigaciones, Universidad Antonio Nariño, Bogotá;(<sup>b</sup>)Departamento de Física, Universidad Nacional de Colombia, Bogotá; Colombia.
- <sup>23</sup>(<sup>a</sup>)Dipartimento di Fisica e Astronomia A. Righi, Università di Bologna, Bologna;(<sup>b</sup>)INFN Sezione di Bologna; Italy.
- <sup>24</sup>Physikalisches Institut, Universität Bonn, Bonn; Germany.
- <sup>25</sup>Department of Physics, Boston University, Boston MA; United States of America.
- <sup>26</sup>Department of Physics, Brandeis University, Waltham MA; United States of America.
- <sup>27</sup>(<sup>a</sup>)Transilvania University of Brasov, Brasov;(<sup>b</sup>)Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest;(<sup>c</sup>)Department of Physics, Alexandru Ioan Cuza University of Iasi, Iasi;(<sup>d</sup>)National Institute for Research and Development of Isotopic and Molecular Technologies, Physics Department, Cluj-Napoca;(<sup>e</sup>)National University of Science and Technology Politehnica, Bucharest;(<sup>f</sup>)West University in Timisoara, Timisoara;(<sup>g</sup>)Faculty of Physics, University of Bucharest, Bucharest; Romania.
- <sup>28</sup>(<sup>a</sup>)Faculty of Mathematics, Physics and Informatics, Comenius University, Bratislava;(<sup>b</sup>)Department of Subnuclear Physics, Institute of Experimental Physics of the Slovak Academy of Sciences, Kosice; Slovak Republic.
- <sup>29</sup>Physics Department, Brookhaven National Laboratory, Upton NY; United States of America.
- <sup>30</sup>Universidad de Buenos Aires, Facultad de Ciencias Exactas y Naturales, Departamento de Física, y CONICET, Instituto de Física de Buenos Aires (IFIBA), Buenos Aires; Argentina.
- <sup>31</sup>California State University, CA; United States of America.
- <sup>32</sup>Cavendish Laboratory, University of Cambridge, Cambridge; United Kingdom.
- <sup>33</sup>(<sup>a</sup>)Department of Physics, University of Cape Town, Cape Town;(<sup>b</sup>)iThemba Labs, Western Cape;(<sup>c</sup>)Department of Mechanical Engineering Science, University of Johannesburg, Johannesburg;(<sup>d</sup>)National Institute of Physics, University of the Philippines Diliman (Philippines);(<sup>e</sup>)University of South Africa, Department of Physics, Pretoria;(<sup>f</sup>)University of Zululand, KwaDlangezwa;(<sup>g</sup>)School of Physics, University of the Witwatersrand, Johannesburg; South Africa.
- <sup>34</sup>Department of Physics, Carleton University, Ottawa ON; Canada.
- <sup>35</sup>(<sup>a</sup>)Faculté des Sciences Ain Chock, Réseau Universitaire de Physique des Hautes Energies - Université Hassan II, Casablanca;(<sup>b</sup>)Faculté des Sciences, Université Ibn-Tofail, Kénitra;(<sup>c</sup>)Faculté des Sciences Semlalia, Université Cadi Ayyad, LPHEA-Marrakech;(<sup>d</sup>)LPMR, Faculté des Sciences, Université Mohamed Premier, Oujda;(<sup>e</sup>)Faculté des sciences, Université Mohammed V, Rabat;(<sup>f</sup>)Institute of Applied Physics, Mohammed VI Polytechnic University, Ben Guerir; Morocco.
- <sup>36</sup>CERN, Geneva; Switzerland.
- <sup>37</sup>Affiliated with an institute covered by a cooperation agreement with CERN.
- <sup>38</sup>Affiliated with an international laboratory covered by a cooperation agreement with CERN.
- <sup>39</sup>Enrico Fermi Institute, University of Chicago, Chicago IL; United States of America.
- <sup>40</sup>LPC, Université Clermont Auvergne, CNRS/IN2P3, Clermont-Ferrand; France.
- <sup>41</sup>Nevis Laboratory, Columbia University, Irvington NY; United States of America.

- <sup>42</sup>Niels Bohr Institute, University of Copenhagen, Copenhagen; Denmark.
- <sup>43</sup>(<sup>a</sup>)Dipartimento di Fisica, Università della Calabria, Rende; (<sup>b</sup>)INFN Gruppo Collegato di Cosenza, Laboratori Nazionali di Frascati; Italy.
- <sup>44</sup>Physics Department, Southern Methodist University, Dallas TX; United States of America.
- <sup>45</sup>Physics Department, University of Texas at Dallas, Richardson TX; United States of America.
- <sup>46</sup>National Centre for Scientific Research "Demokritos", Agia Paraskevi; Greece.
- <sup>47</sup>(<sup>a</sup>)Department of Physics, Stockholm University; (<sup>b</sup>)Oskar Klein Centre, Stockholm; Sweden.
- <sup>48</sup>Deutsches Elektronen-Synchrotron DESY, Hamburg and Zeuthen; Germany.
- <sup>49</sup>Fakultät Physik , Technische Universität Dortmund, Dortmund; Germany.
- <sup>50</sup>Institut für Kern- und Teilchenphysik, Technische Universität Dresden, Dresden; Germany.
- <sup>51</sup>Department of Physics, Duke University, Durham NC; United States of America.
- <sup>52</sup>SUPA - School of Physics and Astronomy, University of Edinburgh, Edinburgh; United Kingdom.
- <sup>53</sup>INFN e Laboratori Nazionali di Frascati, Frascati; Italy.
- <sup>54</sup>Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, Freiburg; Germany.
- <sup>55</sup>II. Physikalisches Institut, Georg-August-Universität Göttingen, Göttingen; Germany.
- <sup>56</sup>Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève; Switzerland.
- <sup>57</sup>(<sup>a</sup>)Dipartimento di Fisica, Università di Genova, Genova; (<sup>b</sup>)INFN Sezione di Genova; Italy.
- <sup>58</sup>II. Physikalisches Institut, Justus-Liebig-Universität Giessen, Giessen; Germany.
- <sup>59</sup>SUPA - School of Physics and Astronomy, University of Glasgow, Glasgow; United Kingdom.
- <sup>60</sup>LPSC, Université Grenoble Alpes, CNRS/IN2P3, Grenoble INP, Grenoble; France.
- <sup>61</sup>Laboratory for Particle Physics and Cosmology, Harvard University, Cambridge MA; United States of America.
- <sup>62</sup>(<sup>a</sup>)Department of Modern Physics and State Key Laboratory of Particle Detection and Electronics, University of Science and Technology of China, Hefei; (<sup>b</sup>)Institute of Frontier and Interdisciplinary Science and Key Laboratory of Particle Physics and Particle Irradiation (MOE), Shandong University, Qingdao; (<sup>c</sup>)School of Physics and Astronomy, Shanghai Jiao Tong University, Key Laboratory for Particle Astrophysics and Cosmology (MOE), SKLPPC, Shanghai; (<sup>d</sup>)Tsung-Dao Lee Institute, Shanghai; (<sup>e</sup>)School of Physics and Microelectronics, Zhengzhou University; China.
- <sup>63</sup>(<sup>a</sup>)Kirchhoff-Institut für Physik, Ruprecht-Karls-Universität Heidelberg, Heidelberg; (<sup>b</sup>)Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg; Germany.
- <sup>64</sup>(<sup>a</sup>)Department of Physics, Chinese University of Hong Kong, Shatin, N.T., Hong Kong; (<sup>b</sup>)Department of Physics, University of Hong Kong, Hong Kong; (<sup>c</sup>)Department of Physics and Institute for Advanced Study, Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong; China.
- <sup>65</sup>Department of Physics, National Tsing Hua University, Hsinchu; Taiwan.
- <sup>66</sup>IJCLab, Université Paris-Saclay, CNRS/IN2P3, 91405, Orsay; France.
- <sup>67</sup>Centro Nacional de Microelectrónica (IMB-CNM-CSIC), Barcelona; Spain.
- <sup>68</sup>Department of Physics, Indiana University, Bloomington IN; United States of America.
- <sup>69</sup>(<sup>a</sup>)INFN Gruppo Collegato di Udine, Sezione di Trieste, Udine; (<sup>b</sup>)ICTP, Trieste; (<sup>c</sup>)Dipartimento Politecnico di Ingegneria e Architettura, Università di Udine, Udine; Italy.
- <sup>70</sup>(<sup>a</sup>)INFN Sezione di Lecce; (<sup>b</sup>)Dipartimento di Matematica e Fisica, Università del Salento, Lecce; Italy.
- <sup>71</sup>(<sup>a</sup>)INFN Sezione di Milano; (<sup>b</sup>)Dipartimento di Fisica, Università di Milano, Milano; Italy.
- <sup>72</sup>(<sup>a</sup>)INFN Sezione di Napoli; (<sup>b</sup>)Dipartimento di Fisica, Università di Napoli, Napoli; Italy.
- <sup>73</sup>(<sup>a</sup>)INFN Sezione di Pavia; (<sup>b</sup>)Dipartimento di Fisica, Università di Pavia, Pavia; Italy.
- <sup>74</sup>(<sup>a</sup>)INFN Sezione di Pisa; (<sup>b</sup>)Dipartimento di Fisica E. Fermi, Università di Pisa, Pisa; Italy.
- <sup>75</sup>(<sup>a</sup>)INFN Sezione di Roma; (<sup>b</sup>)Dipartimento di Fisica, Sapienza Università di Roma, Roma; Italy.
- <sup>76</sup>(<sup>a</sup>)INFN Sezione di Roma Tor Vergata; (<sup>b</sup>)Dipartimento di Fisica, Università di Roma Tor Vergata, Roma; Italy.

- <sup>77(a)</sup>INFN Sezione di Roma Tre; <sup>(b)</sup>Dipartimento di Matematica e Fisica, Università Roma Tre, Roma; Italy.
- <sup>78(a)</sup>INFN-TIFPA; <sup>(b)</sup>Università degli Studi di Trento, Trento; Italy.
- <sup>79</sup>Universität Innsbruck, Department of Astro and Particle Physics, Innsbruck; Austria.
- <sup>80</sup>University of Iowa, Iowa City IA; United States of America.
- <sup>81</sup>Department of Physics and Astronomy, Iowa State University, Ames IA; United States of America.
- <sup>82</sup>Istinye University, Sariyer, Istanbul; Türkiye.
- <sup>83(a)</sup>Departamento de Engenharia Elétrica, Universidade Federal de Juiz de Fora (UFJF), Juiz de Fora; <sup>(b)</sup>Universidade Federal do Rio De Janeiro COPPE/EE/IF, Rio de Janeiro; <sup>(c)</sup>Instituto de Física, Universidade de São Paulo, São Paulo; <sup>(d)</sup>Rio de Janeiro State University, Rio de Janeiro; <sup>(e)</sup>Federal University of Bahia, Bahia; Brazil.
- <sup>84</sup>KEK, High Energy Accelerator Research Organization, Tsukuba; Japan.
- <sup>85</sup>Graduate School of Science, Kobe University, Kobe; Japan.
- <sup>86(a)</sup>AGH University of Krakow, Faculty of Physics and Applied Computer Science, Krakow; <sup>(b)</sup>Marian Smoluchowski Institute of Physics, Jagiellonian University, Krakow; Poland.
- <sup>87</sup>Institute of Nuclear Physics Polish Academy of Sciences, Krakow; Poland.
- <sup>88</sup>Faculty of Science, Kyoto University, Kyoto; Japan.
- <sup>89</sup>Research Center for Advanced Particle Physics and Department of Physics, Kyushu University, Fukuoka ; Japan.
- <sup>90</sup>L2IT, Université de Toulouse, CNRS/IN2P3, UPS, Toulouse; France.
- <sup>91</sup>Instituto de Física La Plata, Universidad Nacional de La Plata and CONICET, La Plata; Argentina.
- <sup>92</sup>Physics Department, Lancaster University, Lancaster; United Kingdom.
- <sup>93</sup>Oliver Lodge Laboratory, University of Liverpool, Liverpool; United Kingdom.
- <sup>94</sup>Department of Experimental Particle Physics, Jožef Stefan Institute and Department of Physics, University of Ljubljana, Ljubljana; Slovenia.
- <sup>95</sup>School of Physics and Astronomy, Queen Mary University of London, London; United Kingdom.
- <sup>96</sup>Department of Physics, Royal Holloway University of London, Egham; United Kingdom.
- <sup>97</sup>Department of Physics and Astronomy, University College London, London; United Kingdom.
- <sup>98</sup>Louisiana Tech University, Ruston LA; United States of America.
- <sup>99</sup>Fysiska institutionen, Lunds universitet, Lund; Sweden.
- <sup>100</sup>Departamento de Física Teórica C-15 and CIAFF, Universidad Autónoma de Madrid, Madrid; Spain.
- <sup>101</sup>Institut für Physik, Universität Mainz, Mainz; Germany.
- <sup>102</sup>School of Physics and Astronomy, University of Manchester, Manchester; United Kingdom.
- <sup>103</sup>CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille; France.
- <sup>104</sup>Department of Physics, University of Massachusetts, Amherst MA; United States of America.
- <sup>105</sup>Department of Physics, McGill University, Montreal QC; Canada.
- <sup>106</sup>School of Physics, University of Melbourne, Victoria; Australia.
- <sup>107</sup>Department of Physics, University of Michigan, Ann Arbor MI; United States of America.
- <sup>108</sup>Department of Physics and Astronomy, Michigan State University, East Lansing MI; United States of America.
- <sup>109</sup>Group of Particle Physics, University of Montreal, Montreal QC; Canada.
- <sup>110</sup>Fakultät für Physik, Ludwig-Maximilians-Universität München, München; Germany.
- <sup>111</sup>Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), München; Germany.
- <sup>112</sup>Graduate School of Science and Kobayashi-Maskawa Institute, Nagoya University, Nagoya; Japan.
- <sup>113</sup>Department of Physics and Astronomy, University of New Mexico, Albuquerque NM; United States of America.
- <sup>114</sup>Institute for Mathematics, Astrophysics and Particle Physics, Radboud University/Nikhef, Nijmegen;

Netherlands.

<sup>115</sup>Nikhef National Institute for Subatomic Physics and University of Amsterdam, Amsterdam; Netherlands.

<sup>116</sup>Department of Physics, Northern Illinois University, DeKalb IL; United States of America.

<sup>117</sup>(<sup>a</sup>)New York University Abu Dhabi, Abu Dhabi;(<sup>b</sup>)United Arab Emirates University, Al Ain; United Arab Emirates.

<sup>118</sup>Department of Physics, New York University, New York NY; United States of America.

<sup>119</sup>Ochanomizu University, Otsuka, Bunkyo-ku, Tokyo; Japan.

<sup>120</sup>Ohio State University, Columbus OH; United States of America.

<sup>121</sup>Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, Norman OK; United States of America.

<sup>122</sup>Department of Physics, Oklahoma State University, Stillwater OK; United States of America.

<sup>123</sup>Palacký University, Joint Laboratory of Optics, Olomouc; Czech Republic.

<sup>124</sup>Institute for Fundamental Science, University of Oregon, Eugene, OR; United States of America.

<sup>125</sup>Graduate School of Science, Osaka University, Osaka; Japan.

<sup>126</sup>Department of Physics, University of Oslo, Oslo; Norway.

<sup>127</sup>Department of Physics, Oxford University, Oxford; United Kingdom.

<sup>128</sup>LPNHE, Sorbonne Université, Université Paris Cité, CNRS/IN2P3, Paris; France.

<sup>129</sup>Department of Physics, University of Pennsylvania, Philadelphia PA; United States of America.

<sup>130</sup>Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh PA; United States of America.

<sup>131</sup>(<sup>a</sup>)Laboratório de Instrumentação e Física Experimental de Partículas - LIP, Lisboa;(<sup>b</sup>)Departamento de Física, Faculdade de Ciências, Universidade de Lisboa, Lisboa;(<sup>c</sup>)Departamento de Física, Universidade de Coimbra, Coimbra;(<sup>d</sup>)Centro de Física Nuclear da Universidade de Lisboa, Lisboa;(<sup>e</sup>)Departamento de Física, Universidade do Minho, Braga;(<sup>f</sup>)Departamento de Física Teórica y del Cosmos, Universidad de Granada, Granada (Spain);(<sup>g</sup>)Departamento de Física, Instituto Superior Técnico, Universidade de Lisboa, Lisboa; Portugal.

<sup>132</sup>Institute of Physics of the Czech Academy of Sciences, Prague; Czech Republic.

<sup>133</sup>Czech Technical University in Prague, Prague; Czech Republic.

<sup>134</sup>Charles University, Faculty of Mathematics and Physics, Prague; Czech Republic.

<sup>135</sup>Particle Physics Department, Rutherford Appleton Laboratory, Didcot; United Kingdom.

<sup>136</sup>IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette; France.

<sup>137</sup>Santa Cruz Institute for Particle Physics, University of California Santa Cruz, Santa Cruz CA; United States of America.

<sup>138</sup>(<sup>a</sup>)Departamento de Física, Pontificia Universidad Católica de Chile, Santiago;(<sup>b</sup>)Millennium Institute for Subatomic physics at high energy frontier (SAPHIR), Santiago;(<sup>c</sup>)Instituto de Investigación Multidisciplinario en Ciencia y Tecnología, y Departamento de Física, Universidad de La Serena;(<sup>d</sup>)Universidad Andres Bello, Department of Physics, Santiago;(<sup>e</sup>)Instituto de Alta Investigación, Universidad de Tarapacá, Arica;(<sup>f</sup>)Departamento de Física, Universidad Técnica Federico Santa María, Valparaíso; Chile.

<sup>139</sup>Department of Physics, University of Washington, Seattle WA; United States of America.

<sup>140</sup>Department of Physics and Astronomy, University of Sheffield, Sheffield; United Kingdom.

<sup>141</sup>Department of Physics, Shinshu University, Nagano; Japan.

<sup>142</sup>Department Physik, Universität Siegen, Siegen; Germany.

<sup>143</sup>Department of Physics, Simon Fraser University, Burnaby BC; Canada.

<sup>144</sup>SLAC National Accelerator Laboratory, Stanford CA; United States of America.

<sup>145</sup>Department of Physics, Royal Institute of Technology, Stockholm; Sweden.

- <sup>146</sup>Departments of Physics and Astronomy, Stony Brook University, Stony Brook NY; United States of America.
- <sup>147</sup>Department of Physics and Astronomy, University of Sussex, Brighton; United Kingdom.
- <sup>148</sup>School of Physics, University of Sydney, Sydney; Australia.
- <sup>149</sup>Institute of Physics, Academia Sinica, Taipei; Taiwan.
- <sup>150</sup><sup>(a)</sup>E. Andronikashvili Institute of Physics, Iv. Javakhishvili Tbilisi State University, Tbilisi;<sup>(b)</sup>High Energy Physics Institute, Tbilisi State University, Tbilisi;<sup>(c)</sup>University of Georgia, Tbilisi; Georgia.
- <sup>151</sup>Department of Physics, Technion, Israel Institute of Technology, Haifa; Israel.
- <sup>152</sup>Raymond and Beverly Sackler School of Physics and Astronomy, Tel Aviv University, Tel Aviv; Israel.
- <sup>153</sup>Department of Physics, Aristotle University of Thessaloniki, Thessaloniki; Greece.
- <sup>154</sup>International Center for Elementary Particle Physics and Department of Physics, University of Tokyo, Tokyo; Japan.
- <sup>155</sup>Department of Physics, Tokyo Institute of Technology, Tokyo; Japan.
- <sup>156</sup>Department of Physics, University of Toronto, Toronto ON; Canada.
- <sup>157</sup><sup>(a)</sup>TRIUMF, Vancouver BC;<sup>(b)</sup>Department of Physics and Astronomy, York University, Toronto ON; Canada.
- <sup>158</sup>Division of Physics and Tomonaga Center for the History of the Universe, Faculty of Pure and Applied Sciences, University of Tsukuba, Tsukuba; Japan.
- <sup>159</sup>Department of Physics and Astronomy, Tufts University, Medford MA; United States of America.
- <sup>160</sup>Department of Physics and Astronomy, University of California Irvine, Irvine CA; United States of America.
- <sup>161</sup>University of Sharjah, Sharjah; United Arab Emirates.
- <sup>162</sup>Department of Physics and Astronomy, University of Uppsala, Uppsala; Sweden.
- <sup>163</sup>Department of Physics, University of Illinois, Urbana IL; United States of America.
- <sup>164</sup>Instituto de Física Corpuscular (IFIC), Centro Mixto Universidad de Valencia - CSIC, Valencia; Spain.
- <sup>165</sup>Department of Physics, University of British Columbia, Vancouver BC; Canada.
- <sup>166</sup>Department of Physics and Astronomy, University of Victoria, Victoria BC; Canada.
- <sup>167</sup>Fakultät für Physik und Astronomie, Julius-Maximilians-Universität Würzburg, Würzburg; Germany.
- <sup>168</sup>Department of Physics, University of Warwick, Coventry; United Kingdom.
- <sup>169</sup>Waseda University, Tokyo; Japan.
- <sup>170</sup>Department of Particle Physics and Astrophysics, Weizmann Institute of Science, Rehovot; Israel.
- <sup>171</sup>Department of Physics, University of Wisconsin, Madison WI; United States of America.
- <sup>172</sup>Fakultät für Mathematik und Naturwissenschaften, Fachgruppe Physik, Bergische Universität Wuppertal, Wuppertal; Germany.
- <sup>173</sup>Department of Physics, Yale University, New Haven CT; United States of America.
- <sup>a</sup> Also Affiliated with an institute covered by a cooperation agreement with CERN.
- <sup>b</sup> Also at An-Najah National University, Nablus; Palestine.
- <sup>c</sup> Also at Borough of Manhattan Community College, City University of New York, New York NY; United States of America.
- <sup>d</sup> Also at Center for High Energy Physics, Peking University; China.
- <sup>e</sup> Also at Center for Interdisciplinary Research and Innovation (CIRI-AUTH), Thessaloniki; Greece.
- <sup>f</sup> Also at Centro Studi e Ricerche Enrico Fermi; Italy.
- <sup>g</sup> Also at CERN, Geneva; Switzerland.
- <sup>h</sup> Also at Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève; Switzerland.
- <sup>i</sup> Also at Departament de Física de la Universitat Autònoma de Barcelona, Barcelona; Spain.
- <sup>j</sup> Also at Department of Financial and Management Engineering, University of the Aegean, Chios; Greece.

- k* Also at Department of Physics, California State University, Sacramento; United States of America.
- l* Also at Department of Physics, King's College London, London; United Kingdom.
- m* Also at Department of Physics, Stanford University, Stanford CA; United States of America.
- n* Also at Department of Physics, Stellenbosch University; South Africa.
- o* Also at Department of Physics, University of Fribourg, Fribourg; Switzerland.
- p* Also at Department of Physics, University of Thessaly; Greece.
- q* Also at Department of Physics, Westmont College, Santa Barbara; United States of America.
- r* Also at Hellenic Open University, Patras; Greece.
- s* Also at Institutio Catalana de Recerca i Estudis Avancats, ICREA, Barcelona; Spain.
- t* Also at Institut für Experimentalphysik, Universität Hamburg, Hamburg; Germany.
- u* Also at Institute for Nuclear Research and Nuclear Energy (INRNE) of the Bulgarian Academy of Sciences, Sofia; Bulgaria.
- v* Also at Institute of Applied Physics, Mohammed VI Polytechnic University, Ben Guerir; Morocco.
- w* Also at Institute of Particle Physics (IPP); Canada.
- x* Also at Institute of Physics and Technology, Mongolian Academy of Sciences, Ulaanbaatar; Mongolia.
- y* Also at Institute of Physics, Azerbaijan Academy of Sciences, Baku; Azerbaijan.
- z* Also at Institute of Theoretical Physics, Ilia State University, Tbilisi; Georgia.
- aa* Also at Lawrence Livermore National Laboratory, Livermore; United States of America.
- ab* Also at National Institute of Physics, University of the Philippines Diliman (Philippines); Philippines.
- ac* Also at Technical University of Munich, Munich; Germany.
- ad* Also at The Collaborative Innovation Center of Quantum Matter (CICQM), Beijing; China.
- ae* Also at TRIUMF, Vancouver BC; Canada.
- af* Also at Università di Napoli Parthenope, Napoli; Italy.
- ag* Also at University of Colorado Boulder, Department of Physics, Colorado; United States of America.
- ah* Also at Washington College, Chestertown, MD; United States of America.
- ai* Also at Yeditepe University, Physics Department, Istanbul; Türkiye.
- \* Deceased