



Letter

Exploring scalar contributions with $K^+ \rightarrow \pi^+ \ell^+ \ell^-$ G. D'Ambrosio^a, A.M. Iyer^b, F. Mahmoudi^{c,d,e}, S. Neshatpour^{c, ,*}^a INFN-Sezione di Napoli, Complesso Universitario di Monte S. Angelo, Via Cintia Edificio 6, 80126 Napoli, Italy^b Department of Physics, Indian Institute of Technology Delhi, Hauz Khas, New Delhi, 110016, India^c Université Claude Bernard Lyon 1, CNRS/IN2P3, Institut de Physique des 2 Infinis de Lyon, UMR 5822, F-69622, Villeurbanne, France^d Theoretical Physics Department, CERN, CH-1211 Geneva 23, Switzerland^e Institut Universitaire de France (IUF), 75005 Paris, France

ARTICLE INFO

Editor: A. Ringwald

ABSTRACT

The rare kaon decay $K^+ \rightarrow \pi^+ \ell^+ \ell^-$ offers insights into Standard Model (SM) physics and beyond. Driven by vector form factor in the SM, it can also probe non-standard contributions. In this letter we study the scalar contribution, f_S . Using differential decay width and Forward-Backward Asymmetry we propose a simultaneous fit to vector and scalar contributions which is necessary for a consistent analysis. Novel bounds on $|f_S|$ are presented for the first time through a reinterpretation of the E865, NA48/2, and NA62 experimental results. The analysis results in the most precise bound $f_S < 7.9 \times 10^{-6}$ at 90% confidence level.

1. Introduction

Kaon physics has been fundamental for our understanding of the structure of weak interactions: discovery of the GIM mechanism, CP violation, P violation, etc. There are several ongoing efforts for a deeper understanding of rare kaon decays: They include processes like $K_L \rightarrow \pi^0 \nu \bar{\nu}$, $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ [1,2], and $K^+ \rightarrow \pi^+ \ell^+ \ell^-$ among others. The interest in semi-leptonic $K^+ \rightarrow \pi^+ \ell^+ \ell^-$ decay is to extract short-distance information: lepton flavour universality violation test [3–8] and other short-distance probes like P and CP violation [9,10]. Chiral Perturbation Theory (ChPT) [10–17] is the appropriate theory framework for describing this decay. Within the Standard Model (SM), this decay at short distances is induced at the loop level and predominantly occurs via single virtual photon exchange. Due to its suppression in the SM it offers a glimpse into different types of short-distance physics.

In this letter, we study the scalar contribution to $K^+ \rightarrow \pi^+ \ell^+ \ell^-$. It also affects processes such as $K_{L,S} \rightarrow \mu^+ \mu^-$, $K_L \rightarrow \pi^0 \ell^+ \ell^-$ in addition to $K^+ \rightarrow \pi^+ \ell^+ \ell^-$. The existing datasets from the electron and the muon channels [18–20] and the ongoing measurements of $K^+ \rightarrow \pi^+ \mu^+ \mu^-$ decay at NA62 [21] make it an exciting probe to test the limits of the scalar operators. We address the fact that the only existing bound on scalar contributions from the E865 experiment from 1999 is derived from the branching ratio. While the Forward-Backward Asymmetry (A_{FB}) in these decays vanishes in the SM [13,22–24], the presence of scalar in-

teractions introduces non-zero A_{FB} , making it a powerful probe of New Physics. Typical scenarios that can give rise to such effects have been studied for example in [25–27]. Although A_{FB} serves as a robust probe, its effectiveness can be influenced by the lepton mass, leading to suppression in the electron mode, where direct measurements are currently lacking. Alternatively, the branching ratio (BR) offers another avenue for investigating scalar interactions. While slightly less sensitive than A_{FB} in the muon channel, the branching ratio serves as a primary probe in the electron channel, providing insights into the presence of scalar contributions [18].

We propose a more thorough investigation into the scalar contribution by conducting a simultaneous fit to both the vector and the scalar contributions using the differential decay width bins, both when the A_{FB} measurement is included or not. Specifically, when conducting a combined investigation with A_{FB} , which only exhibits non-zero values in the presence of scalar contribution, the inclusion of the scalar form factor in the fit is mandatory in order to have a consistent study. We obtain bounds on scalar contributions from the different available experimental datasets.

After discussing the framework in section 2, in section 3 through the examination of the A_{FB} and branching ratio, we find an estimate of the sensitivity to scalar contributions. In section 4 we obtain a bound from our proposed fit to data where in addition to the vector contributions,

* Corresponding author.

E-mail addresses: gdambros@na.infn.it (G. D'Ambrosio), iyerabhishek@physics.iitd.ac.in (A.M. Iyer), nazila@cern.ch (F. Mahmoudi), s.neshatpour@ip2i.in2p3.fr (S. Neshatpour).<https://doi.org/10.1016/j.physletb.2024.138824>

Received 9 June 2024; Accepted 19 June 2024

Available online 26 June 2024

0370-2693/© 2024 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

scalar contributions are taken into account and we summarise our results in section 5.

2. Framework

The amplitude of the $K^+ \rightarrow \pi^+ \ell^+ \ell^-$ decay, when taking into account only the vector and scalar interactions, as denoted by the form factors f_V , and f_S respectively, can be written as [18,28]

$$\mathcal{M} = \frac{\alpha G_F}{4\pi} f_V(z) P^\mu \bar{\ell} \gamma_\mu \ell + G_F M_K f_S \bar{\ell} \ell, \quad (1)$$

where $P = p_K + p_\pi$ and $q = p_K - p_\pi$, with p_K and p_π the momenta of the kaon and the pion, respectively and the dilepton invariant mass squared can be written as $q^2 = z M_K^2$.

Given the above amplitude, the double-differential decay width, in terms of the vector and scalar form factors, is expressed as [23,24]

$$\frac{d^2\Gamma}{dz d\cos\theta} = \frac{G_F^2 M_K^5}{2^8 \pi^3} \beta_\ell \lambda^{1/2}(z) \times \left\{ |f_V|^2 \frac{\alpha^2}{16\pi^2} \lambda(z) (1 - \beta_\ell^2 \cos^2\theta) + |f_S|^2 z \beta_\ell^2 + \text{Re}(f_V^* f_S) \frac{\alpha r_\ell}{\pi} \beta_\ell \lambda^{1/2}(z) \cos\theta \right\}, \quad (2)$$

where θ is the angle between the negatively charged lepton and the kaon in the dilepton rest frame, $r_\ell = m_\ell/M_K$, $r_\pi = m_\pi/M_K$, $\beta_\ell = \sqrt{1 - 4r_\ell^2/z}$, and $\lambda(z) \equiv \lambda(1, z, r_\pi^2)$ is the Källén function.

The familiar z -spectrum is recovered by integrating over $\cos\theta$

$$\frac{d\Gamma}{dz} = \frac{2}{3} \frac{G_F^2 M_K^5}{2^8 \pi^3} \beta_\ell \lambda^{1/2}(z) \times \left\{ |f_V|^2 2 \frac{\alpha^2}{16\pi^2} \lambda(z) \left(1 + 2 \frac{r_\ell^2}{z}\right) |f_S|^2 3 z \beta_\ell^2 \right\}, \quad (3)$$

which upon further integration with respect to z yields the branching ratio.

Another interesting observable is obtained by considering the angular behaviour of the decay, with the Forward-Backward Asymmetry defined as

$$A_{\text{FB}}(z) = \frac{\int_0^1 \left(\frac{d\Gamma}{dz d\cos\theta}\right) d\cos\theta - \int_{-1}^0 \left(\frac{d\Gamma}{dz d\cos\theta}\right) d\cos\theta}{\int_0^1 \left(\frac{d\Gamma}{dz d\cos\theta}\right) d\cos\theta + \int_{-1}^0 \left(\frac{d\Gamma}{dz d\cos\theta}\right) d\cos\theta}. \quad (4)$$

Considering Eq. (2), we have [23,24]

$$A_{\text{FB}}(z) = \frac{\alpha G_F^2 M_K^5}{2^8 \pi^4} r_\ell \beta_\ell^2(z) \lambda(z) \text{Re}(f_V^* f_S) \left/ \left(\frac{d\Gamma(z)}{dz}\right)\right., \quad (5)$$

which is non-zero only in case vector and scalar contributions are simultaneously present.

In the SM, the $K^+ \rightarrow \pi^+ \ell^+ \ell^-$ decay is completely governed by the vector form factor $f_V(z)$ which can be described as a linear contribution in z accompanied by the unitarity loop correction [10],¹ expressed as

$$f_V(z) = a_+ + b_+ z + V^{\pi\pi}(z). \quad (6)$$

Here, $V^{\pi\pi}(z)$ accounts for the pion loop contribution calculated at $\mathcal{O}(p^6)$ in ChPT [10]. The parameters a_+ and b_+ are considered as phenomenological constants, typically extracted from experimental data. Nevertheless, recent advancements have been achieved in theoretical calculations concerning these parameters (e.g., see [17,32,33]). On the other hand, the scalar form factor is highly suppressed and negligible in the SM. Regarding the available data, precise measurements for the differential decay width distributions of $K^\pm \rightarrow \pi^\pm \ell^+ \ell^-$ decay have been conducted

¹ The vector form factor has been described with several formulations, from a simplistic model with only a linear parameterisation, $f_V = f_0(1 + \delta z)$, to more complicated models [10,29–31]. In this study, for f_V we considered the ‘‘Linear + Chiral’’ description from Ref. [10] as given in Eq. (6), where f_V here in terms of $W(z)$ of Ref. [10] is given by $f_V = W(z)/(G_F M_K^2)$.

since the initial observation of $K^+ \rightarrow \pi^+ e^+ e^-$ [34] at CERN. In the electron channel, the most events have been observed by BNL-E865 [18] and NA48/2 [19], while for the muon channel, there are results from NA48/2 [20] and more recently NA62 [21].

To investigate scalar contributions we consider both the branching ratio and the Forward-Backward Asymmetry. The latter is clearly dependent on scalar contributions, where a non-vanishing A_{FB} necessitates non-zero scalar contributions. However, due to its proportionality to the lepton mass ($r_\ell = m_\ell/M_K$), it is highly suppressed in the electron channel. This suppression does not apply to the branching ratio which is obtained by integrating the differential decay width, Eq. (3), over z . Section 3 provides further elaboration on the approach.

A more concrete analysis assuming the presence of scalar contributions in addition to the vector form factors is given in section 4. This choice is strongly motivated by the possible non-zero value of A_{FB} . The measurement of the A_{FB} makes it mandatory to have a three parameter fit in general. While an experimental measurement of the A_{FB} is extremely difficult owing to the electron mass suppression, we can extract a more consistent bound on the scalar contribution by means of a three-parameter fit.

3. Branching ratio vs. forward-backward asymmetry

Historically, the scalar contributions were constrained by the branching ratio [18] and A_{FB} [20]. The muon mode enjoys a model-independent measurement of the branching ratio. In this case, we have two independent sources for estimating the sensitivity on scalar contributions: 1) The measured branching ratio, 2) The measurement of A_{FB} . On the other hand, there is no model-independent measurement of the branching ratio for the electron mode. The existing method for measuring the branching ratio for the electron mode is by assuming a model-specific vector form factor (see footnote 1). There is only the branching fraction which allows for a measure of the sensitivity of the scalar contributions and there are no measurements on A_{FB} .

For the branching ratio, to constrain f_S , we evaluate the permissible contributions from scalar interactions at 90% confidence level (CL), considering the uncertainty in the measured branching ratio of $K^+ \rightarrow \pi^+ \ell^+ \ell^-$. Regarding A_{FB} , we examine the experimental results and derive the corresponding 90% CL upper bound on f_S employing the relation in Eq. (5). Experimental data for A_{FB} is, however, as mentioned before, only available for the muon channel. The bounds on f_S that we derived from both the BR and the A_{FB} measurements are presented in Table 1, considering various experimental measurements. In each panel of Table 1, the second column corresponds to the experimental value, while the last column denotes the upper bound on $|f_S|$ that we obtain.

Notice that the bound from the NA62 measurement of A_{FB} is approximately seven times stronger than the one from the branching ratio as measured by the same experiment. However, it is important to note that this constraint may not apply to scenarios involving lepton flavour universality violating scalar contributions. Regarding the electron mode, current constraints are solely derived from the BR measurements, utilising either the NA48/2 [19] or E865 [18] results. Our result is in agreement with the upper bound given by the E865 experiment [18]. While the NA62 collaboration has yet to measure the electron mode, assuming a similar enhancement as observed in the muon channel compared to NA48/2 (a factor of 3), the upper limit for $|f_S|$ in the electron channel could potentially decrease to $\sim 4 \times 10^{-5}$.

4. Bound on f_S from three-parameter fit

The experimentally determined values of a_+ and b_+ are obtained by analysing $d\Gamma/dz$ data, assuming only vector contributions. It is possible to extend this analysis to include scalar contributions and re-evaluate the fit for a_+ , b_+ , and f_S . For the muon channel, measurements of A_{FB} can also be incorporated into the analysis.

Table 1

Bound on $|f_S|$ at 90% CL, from A_{FB} and the uncertainty of the branching ratio. In each panel, the last column corresponds to the upper bound obtained from the experimental measurement of the column to its left. For the electron channel, there are no measurements available for the Forward-Backward Asymmetry.

$(K^+ \rightarrow \pi^+ \mu^+ \mu^-)$			$(K^+ \rightarrow \pi^+ e^+ e^-)$		
NA48/2	exp	$ f_S <$	E865	exp	$ f_S <$
BR	$(9.62 \pm 0.21) \times 10^{-8}$	1.0×10^{-4}	BR	$(2.988 \pm 0.040) \times 10^{-7}$	6.8×10^{-5}
A_{FB}	$(-2.4 \pm 1.8) \times 10^{-2}$	4.2×10^{-5}	A_{FB}	–	–
<hr/>			<hr/>		
NA62	exp	$ f_S <$	NA48/2	exp	$ f_S <$
BR	$(9.16 \pm 0.06) \times 10^{-8}$	5.6×10^{-5}	BR	$(3.14 \pm 0.04) \times 10^{-7}$	6.8×10^{-5}
A_{FB}	$(0.0 \pm 0.7) \times 10^{-2}$	7.7×10^{-6}	A_{FB}	–	–




Fig. 1. The 90% and 95% CL upper bound on $|f_S|$ obtained by our analysis of different experimental datasets. Left: bound from BR or A_{FB} as given in Table 1. Right: bound from 3-dim. fit to f_V and f_S as given in Table 2. The vertical black line in the left plot corresponds to the only existing experimental upper bound from E865 [18].

Table 2

Upper bound for $|f_S|$ at 90% CL, from the three parameter fit to a , b and f_S using various datasets. For the relevant inputs regarding the theoretical calculations we have considered PDG 2022 [42], and the external parameters α_+ , β_+ are taken from [43], in agreement with NA62 [21].

	$d\Gamma/dz$	$d\Gamma/dz + A_{FB}$
$K \rightarrow \pi ee$	$ f_S <$	$ f_S <$
E865	8.0×10^{-5}	–
NA48/2	4.0×10^{-5}	–
<hr/>		
$K \rightarrow \pi \mu \mu$	$ f_S <$	$ f_S <$
NA48/2	10.0×10^{-5}	4.1×10^{-5}
NA62	9.0×10^{-5}	7.9×10^{-6}

In the previous section, it is practically assumed that the size of $|f_S|$ is small compared to the vector form factor. In this section with a three-parameter fit, we abandon this assumption to obtain a more solid evaluation of the scalar contributions.

In Table 2 we give the 90% CL upper bound on $|f_S|$, obtained from the three-parameter fit to a_+ , b_+ , and f_S with the different datasets [35–38]. The $d\Gamma/dz$ data of the NA48/2 and NA62 measurements are available on the HEPData repository [39–41]. The second (third) column corresponds to the fit considering $d\Gamma/dz$ bins, while excluding (including) A_{FB} . When A_{FB} is not included, the results are similar to what one gets from the branching ratio in the previous section. This is expected as BR effectively encapsulates the information from the differential decay width bins. Nonetheless, it would be interesting to see experiments also explore a three-parameter model, including f_S in addition to a_+ and b_+ when analysing the data. On the other hand, the bound from the fit including both the $d\Gamma/dz$ and A_{FB} (last column in Table 2) is very similar

to the bound obtained from only A_{FB} (as given in Table 1). This is due to the fact that the bound from A_{FB} is far stronger than the decay width and adding the latter does not offer much further information. The consistency between the fit and the constraints from the preceding section on f_S is reassuring and justifies the assumption of the dominance of the vector form factor made in the previous section.

5. Summary

In this letter, we analysed the data on the $K^+ \rightarrow \pi^+ \ell^+ \ell^-$ decay and constrained scalar contributions, f_S . Using BR or A_{FB} we obtain 90% upper bounds on f_S as given in Table 1. The shortcoming of this simple approach is the assumption of the relative smallness of f_S compared to f_V . For a concrete and precise bound on f_S , we proposed a new approach to analyse the experimental data via a 3-dimensional simultaneous fit to f_V and f_S (Table 2). This method is necessary to have a consistent analysis of the data, especially given the direct measurement of A_{FB} . It is particularly crucial if a non-zero A_{FB} is measured, as is the case for the NA48/2 measurement of the muon mode.

A comparison between the two methods of sections 3 and 4 is depicted in the left and right plots of Fig. 1, respectively. The only upper bound available in the literature, which is obtained using the method of section 3 via BR, is indicated by the vertical black line in the left plot as given by [18]. All the bounds represented by the colored bars are derived in this paper through the analysis of various experimental datasets. The slight difference of our upper bound obtained via the BR result of the E865 [18] experiment compared to the value given in Ref. [18] is expected due to updated input parameters.

The most precise limit on f_S that we obtain is 7.9×10^{-6} at 90% CL which is about one order of magnitude stronger than the bound given in Ref. [18] by the E865 experiment. Our upper bound arises from the 3-dimensional fit to NA62 data when A_{FB} is included, highlighting the potential of the latter to probe scalar interactions. It would be interesting

to analyse A_{FB} in smaller bins, as it could enhance our ability to further scrutinise scalar contributions. As the NA62 collaboration is planning to have an analysis of the full dataset of the $K^+ \rightarrow \pi^+ \ell^+ \ell^-$ decay, this consideration will be particularly pertinent to further constrain the size of the scalar contributions.

Declaration of competing interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: S. Neshatpour reports financial support was provided by French National Research Agency. A.M. Iyer reports financial support was provided by SERB India. G. D'Ambrosio reports was provided by INFN research initiative Exploring New Physics (ENP). If there are other authors, they declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Data will be made available on request.

Acknowledgements

We appreciate the valuable input from E. Goudzovski, M. Koval, C. Lazzeroni, and G. Ruggiero of the NA62 collaboration. The stimulating discussions and correspondence have enhanced our comprehension concerning the data and their treatment. This research was funded in part by the National Research Agency (ANR) under project ANR-21-CE31-0002-01. AMI would like to acknowledge the generous support by SERB India through the project SRG/2022/001003. AMI would also like to thank the French Institute- Embassy of France in India for facilitating research trip to IP2I Lyon in December 2022. AMI would also like to thank the hospitality of INFN Sezione di Napoli and IP2I Lyon during the academic visit in December 2023. GD was supported in part by the INFN research initiative Exploring New Physics (ENP).

References

- [1] KOTO collaboration, Study of the $K_L \rightarrow \pi^0 \nu \bar{\nu}$ decay at the J-PARC KOTO experiment, Phys. Rev. Lett. 126 (2021) 121801, arXiv:2012.07571.
- [2] NA62 collaboration, Measurement of the very rare $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ decay, J. High Energy Phys. 06 (2021) 093, arXiv:2103.15389.
- [3] A. Crivellin, G. D'Ambrosio, M. Hoferichter, L.C. Tunstall, Violation of lepton flavor and lepton flavor universality in rare kaon decays, Phys. Rev. D 93 (2016) 074038, arXiv:1601.00970.
- [4] G. D'Ambrosio, A.M. Iyer, F. Mahmoudi, S. Neshatpour, Anatomy of kaon decays and prospects for lepton flavour universality violation, J. High Energy Phys. 09 (2022) 148, arXiv:2206.14748.
- [5] HIKE collaboration, HIKE, high intensity kaon experiments at the CERN SPS: letter of intent, arXiv:2211.16586.
- [6] C. Ahdida, et al., Post-LS3 experimental options in ECN3, arXiv:2310.17726.
- [7] HIKE collaboration, High intensity kaon experiments (HIKE) at the CERN SPS proposal for phases 1 and 2, arXiv:2311.08231.
- [8] G. D'Ambrosio, F. Mahmoudi, S. Neshatpour, Beyond the Standard Model prospects for kaon physics at future experiments, J. High Energy Phys. 02 (2024) 166, arXiv:2311.04878.
- [9] G. D'Ambrosio, G. Isidori, CP violation in kaon decays, Int. J. Mod. Phys. A 13 (1998) 1, arXiv:hep-ph/9611284.
- [10] G. D'Ambrosio, G. Ecker, G. Isidori, J. Portoles, The decays $K \rightarrow \pi \ell^+ \ell^-$ beyond leading order in the chiral expansion, J. High Energy Phys. 08 (1998) 004, arXiv:hep-ph/9808289.
- [11] F.J. Gilman, M.B. Wise, $K \rightarrow \pi e^+ e^-$ in the six quark model, Phys. Rev. D 21 (1980) 3150.
- [12] G. Ecker, A. Pich, E. de Rafael, $K \rightarrow \pi l^+ l^-$ decays in the effective chiral Lagrangian of the standard model, Nucl. Phys. B 291 (1987) 692.
- [13] G. Ecker, A. Pich, E. de Rafael, Radiative kaon decays and CP violation in chiral perturbation theory, Nucl. Phys. B 303 (1988) 665.
- [14] G. D'Ambrosio, G. Ecker, G. Isidori, H. Neufeld, Radiative nonleptonic kaon decays, in: The Second DAΦNE Physics Handbook, 11, 1994, pp. 265–312, arXiv:hep-ph/9411439.
- [15] V. Cirigliano, G. Ecker, H. Neufeld, A. Pich, J. Portoles, Kaon decays in the standard model, Rev. Mod. Phys. 84 (2012) 399, arXiv:1107.6001.
- [16] B. Ananthanarayan, I. Sentitemsu Imsong, The 27-plet contributions to the CP-conserving $K \rightarrow \pi l^+ l^-$ decays, J. Phys. G 39 (2012) 095002, arXiv:1207.0567.
- [17] G. D'Ambrosio, D. Greynat, M. Knecht, On the amplitudes for the CP-conserving $K^\pm(K_S) \rightarrow \pi^\pm(\pi^0) \ell^+ \ell^-$ rare decay modes, J. High Energy Phys. 02 (2019) 049, arXiv:1812.00735.
- [18] E865 collaboration, A new measurement of the properties of the rare decay $K^+ \rightarrow \pi^+ e^+ e^-$, Phys. Rev. Lett. 83 (1999) 4482, arXiv:hep-ex/9907045.
- [19] NA48/2 collaboration, Precise measurement of the $K^\pm \rightarrow \pi^\pm e^+ e^-$ decay, Phys. Lett. B 677 (2009) 246, arXiv:0903.3130.
- [20] NA48/2 collaboration, New measurement of the $K^\pm \rightarrow \pi^\pm \mu^+ \mu^-$ decay, Phys. Lett. B 697 (2011) 107, arXiv:1011.4817.
- [21] NA62 collaboration, A measurement of the $K^+ \rightarrow \pi^+ \mu^+ \mu^-$ decay, J. High Energy Phys. 11 (2022) 011, arXiv:2209.05076.
- [22] G. Ecker, A. Pich, The longitudinal muon polarization in $K_L \rightarrow \mu^+ \mu^-$, Nucl. Phys. B 366 (1991) 189.
- [23] D.-N. Gao, Long distance contribution to the forward backward asymmetry in decays $K^+ \rightarrow \pi^+ \ell^+ \ell^-$, Phys. Rev. D 69 (2004) 094030, arXiv:hep-ph/0311253.
- [24] C.-H. Chen, C.Q. Geng, L.-L. Ho, Forward backward asymmetry in $K^+ \rightarrow \pi^+ \ell^+ \ell^-$, Phys. Rev. D 67 (2003) 074029, arXiv:hep-ph/0302207.
- [25] A. Dedes, The Higgs penguin and its applications: an overview, Mod. Phys. Lett. A 18 (2003) 2627, arXiv:hep-ph/0309233.
- [26] D.A. Demir, Higgs boson couplings to quarks with supersymmetric CP and flavor violations, Phys. Lett. B 571 (2003) 193, arXiv:hep-ph/0303249.
- [27] G. Isidori, F. Mescia, P. Paradisi, C. Smith, S. Trine, Exploring the flavour structure of the MSSM with rare K decays, J. High Energy Phys. 08 (2006) 064, arXiv:hep-ph/0604074.
- [28] D.S. Beder, G.V. Dass, Leptonic coupling types in $K^+ \rightarrow \pi^+ \ell^+ \ell^-$ decays and the question of strangeness-changing semileptonic neutral currents, Phys. Lett. B 59 (1975) 444.
- [29] S. Friot, D. Greynat, E. De Rafael, Rare kaon decays revisited, Phys. Lett. B 595 (2004) 301, arXiv:hep-ph/0404136.
- [30] E. Coluccio Leskow, G. D'Ambrosio, D. Greynat, A. Nath, $K \rightarrow \pi \ell^+ \ell^-$ form factor in the Large- N_c and cut-off regularization method, Phys. Rev. D 93 (2016) 094031, arXiv:1603.09721.
- [31] A.Z. Dubnickova, S. Dubnicka, E. Goudzovski, V.N. Pervushin, M. Secansky, Kaon decay probe of the weak static interaction, Phys. Part. Nucl. Lett. 5 (2008) 76, arXiv:hep-ph/0611175.
- [32] G. D'Ambrosio, D. Greynat, M. Knecht, Matching long and short distances at order $\mathcal{O}(\alpha_s)$ in the form factors for $K \rightarrow \pi \ell^+ \ell^-$, Phys. Lett. B 797 (2019) 134891, arXiv:1906.03046.
- [33] RBC, UKQCD collaboration, Simulating rare kaon decays $K^+ \rightarrow \pi^+ \ell^+ \ell^-$ using domain wall lattice QCD with physical light quark masses, Phys. Rev. D 107 (2023) L011503, arXiv:2202.08795.
- [34] P. Bloch, et al., Observation of the $K^+ \rightarrow \pi^+ e^+ e^-$ decay, Phys. Lett. B 56 (1975) 201.
- [35] F. Mahmoudi, SuperIso: a program for calculating the isospin asymmetry of $B \rightarrow K^* \gamma$ in the MSSM, Comput. Phys. Commun. 178 (2008) 745, arXiv:0710.2067.
- [36] F. Mahmoudi, SuperIso v2.3: a program for calculating flavor physics observables in supersymmetry, Comput. Phys. Commun. 180 (2009) 1579, arXiv:0808.3144.
- [37] F. Mahmoudi, SuperIso v3.0, flavor physics observables calculations: extension to NMSSM, Comput. Phys. Commun. 180 (2009) 1718.
- [38] S. Neshatpour, F. Mahmoudi, Flavour physics phenomenology with SuperIso, PoS CompTools2021 (2022) 010, arXiv:2207.04956.
- [39] NA48/2 Collaboration, <https://doi.org/10.17182/hepdata.35365>, 2015.
- [40] NA48/2 Collaboration, <https://doi.org/10.17182/hepdata.69636>, 2015.
- [41] NA62 Collaboration, <https://doi.org/10.17182/hepdata.135498>, 2022.
- [42] Particle Data Group collaboration, Review of Particle Physics, PTEP 2022 (2022) 083C01.
- [43] G. D'Ambrosio, M. Knecht, S. Neshatpour, Determination of the structure of the $K \rightarrow \pi \pi \pi$ amplitudes from recent data, Phys. Lett. B 835 (2022) 137594, arXiv:2209.02143.