

Observation of double J/ψ meson production in pPb collisions at $\sqrt{s_{NN}} = 8.16$ TeV

The CMS Collaboration*

Abstract

The first observation of the concurrent production of two J/ψ mesons in proton-nucleus collisions is presented. The analysis is based on a proton-lead (pPb) data sample recorded at a nucleon-nucleon center-of-mass energy of 8.16 TeV by the CMS experiment at the CERN LHC and corresponding to an integrated luminosity of 174.6 nb^{-1} . The two J/ψ mesons are reconstructed in their $\mu^+\mu^-$ decay channels with transverse momenta $p_T > 6.5 \text{ GeV}$ and rapidity $|y| < 2.4$. Events where one of the J/ψ mesons is reconstructed in the dielectron channel are also considered in the search. The $\text{pPb} \rightarrow J/\psi J/\psi + X$ process is observed with a significance of 5.3 standard deviations. The measured inclusive fiducial cross section, using the four-muon channel alone, is $\sigma(\text{pPb} \rightarrow J/\psi J/\psi + X) = 22.0 \pm 8.9 (\text{stat}) \pm 1.5 (\text{syst}) \text{ nb}$. A fit of the data to the expected rapidity separation for pairs of J/ψ mesons produced in single (SPS) and double (DPS) parton scatterings yields $\sigma_{\text{SPS}}^{\text{pPb} \rightarrow J/\psi J/\psi + X} = 16.5 \pm 10.8 (\text{stat}) \pm 0.1 (\text{syst}) \text{ nb}$ and $\sigma_{\text{DPS}}^{\text{pPb} \rightarrow J/\psi J/\psi + X} = 5.4 \pm 6.2 (\text{stat}) \pm 0.4 (\text{syst}) \text{ nb}$, respectively. This latter result can be transformed into a lower bound on the effective DPS cross section, closely related to the squared average interparton transverse separation in the collision, of $\sigma_{\text{eff}} > 1.0 \text{ mb}$ at 95% confidence level.

Submitted to Physical Review D

1 Introduction

High-energy collisions of protons and nuclei at the CERN LHC are characterized by multiple interactions of their underlying partonic constituents (quarks and gluons). Such multiple parton interactions can lead to the simultaneous production of several particles with values of transverse momentum and/or mass above a few GeV, much larger than the typical non-perturbative $\mathcal{O}(0.2 \text{ GeV})$ scale of quantum chromodynamics (QCD). Research on multiparton interactions has attracted an increasing interest at hadron colliders as a means to probe the generalized parton densities of the proton, to determine the unknown energy evolution of the partonic transverse profile of the proton, and to study the role of partonic correlations (in position and momentum phase space, as well as in flavor, color, and spin quantum numbers) in the hadronic wave functions [1, 2]. A good understanding of multiple hard scatterings of partons is also crucial to describe the continuum backgrounds in studies of rare standard model resonance decays, as well as in searches for new physics, in final states where various heavy particles are produced [3–5].

Following a purely geometric approach, i.e., ignoring any parton correlations, the probability to produce n high- p_T particles in a given hadron-hadron collision is proportional to the n^{th} -product of the probabilities to independently produce each of them [6]. Thus, the probability to produce two high- p_T particles in a double parton scattering (DPS) would scale with the square of the corresponding single parton scattering (SPS) probabilities. The occurrence of DPS processes is therefore more likely for final states with large SPS cross sections, as is the case for quarkonium mesons compared to rarer heavy particles such as electroweak bosons [7]. Under this approximation, the cross section to produce two charmonium mesons, ψ_1 and ψ_2 , via DPS in a proton-proton (pp) collision can be expressed as the product of the SPS cross sections for the production of each individual meson (calculated using perturbative QCD or directly measured using collision data) normalized by an effective cross section (σ_{eff}) to ensure the proper units of the ratio,

$$\sigma_{\text{DPS}}^{\text{pp} \rightarrow \psi_1 \psi_2 + X} = \left(\frac{m}{2}\right) \frac{\sigma_{\text{SPS}}^{\text{pp} \rightarrow \psi_1 + X} \sigma_{\text{SPS}}^{\text{pp} \rightarrow \psi_2 + X}}{\sigma_{\text{eff}}}. \quad (1)$$

Here, m is a combinatorial factor to avoid double counting the same process, $m = 1$ (2) if $\psi_1 = \psi_2$ ($\psi_1 \neq \psi_2$). In a purely geometric approach, the σ_{eff} value can be determined via

$$\sigma_{\text{eff}} = \left[\int d^2\mathbf{b} T^2(\mathbf{b}) \right]^{-1}, \quad (2)$$

where $T(\mathbf{b})$ is the pp transverse overlap function at the vector impact parameter \mathbf{b} of the collision [6]. In terms of the transverse parton density of the proton, $\rho(\mathbf{b})$, the overlap function is given by $T(\mathbf{b}) = \int \rho(\mathbf{b}_1) \rho(\mathbf{b}_1 - \mathbf{b}) d^2\mathbf{b}_1$, and the σ_{eff} parameter can be interpreted as the squared average interparton distance in the pp collision where the two hard scatterings take place. Thus, a smaller value of σ_{eff} implies larger DPS yields. For the typical proton transverse density profiles implemented in common Monte Carlo (MC) event generators used at collider energies, e.g., PYTHIA [8] or HERWIG++ [9], $\sigma_{\text{eff}} \approx 20\text{--}30 \text{ mb}$. These values are about a factor of two larger than those experimentally derived via Eq. (1) for a variety of different pairs of particles measured at the Tevatron and the LHC [2]. This discrepancy points to the presence of various types of parton correlations in the proton [10] not accounted for in the purely geometrical approach, as represented in Eq. (2). In addition, the experimentally extracted values of σ_{eff} seemingly depend on the particular final state under consideration, with multi-quarkonium production processes yielding lower values, $\sigma_{\text{eff}} \approx 5\text{--}10 \text{ mb}$ from $J/\psi/J/\psi$ [11–15], $J/\psi/\psi$ [16],

$J/\psi Y$ [17], and YY [4, 18] studies, than the $\sigma_{\text{eff}} \approx 15$ mb obtained from final states involving pairs of high- p_T jets and/or electroweak bosons [19–26]. Such process-dependent σ_{eff} determinations appear consistent with flavor-dependent correlations that follow the different quark or gluon densities in the parton distribution functions (PDFs) probed at varying values of the momentum fraction by each scattering [27].

To advance understanding of multiple hard scatterings, it has been suggested to study DPS processes in proton-nucleus (pA) collisions [28–30] in which the large transverse parton density of the nucleus significantly enhances the DPS contributions compared to the pp case. Measurements of DPS processes in pA collisions have also been proposed as a means to shed light on the impact-parameter dependence of nuclear PDFs (nPDFs) [31]. For pPb collisions, the equivalent of Eq. (1) for the inclusive production of two J/ψ mesons is

$$\sigma_{\text{DPS}}^{\text{pPb} \rightarrow J/\psi J/\psi + X} = \left(\frac{1}{2}\right) \frac{\sigma_{\text{SPS}}^{\text{pPb} \rightarrow J/\psi + X} \sigma_{\text{SPS}}^{\text{pPb} \rightarrow J/\psi + X}}{\sigma_{\text{eff,pPb}}}, \quad (3)$$

where the effective pPb cross section in the denominator accounts now for two possible DPS contributions coming from collisions of partons of the “projectile” proton with partons of one or two nucleons of the “target” lead ion (center and right diagrams of Fig. 1, respectively). Under the same assumptions of factorization of hard-scattering probabilities that lead to Eq. (2) in the pp case, the value of $\sigma_{\text{eff,pPb}}$ can be connected to that of σ_{eff} through the relationship [6, 30]

$$\sigma_{\text{eff,pPb}} = \frac{A \sigma_{\text{eff}}}{1 + \sigma_{\text{eff}} F_{\text{pPb}}/A}, \quad (4)$$

where $A = 208$ is the Pb mass number, and the quantity $F_{\text{pPb}} = (A - 1)/A \int d^2\mathbf{b} T^2(\mathbf{b}) \approx A^{1/3}/14\pi \text{ mb}^{-1}$, similarly to Eq. (2), is the integral over impact parameter of the squared pPb transverse profile, which can be determined through a Glauber MC model [32]. In this approximation, the value of $\sigma_{\text{eff,pPb}}$ is expected to be about half the total hadronic pPb cross section that amounts to $\sigma_{\text{pPb}} \approx 2$ b at the LHC [33, 34]. Using Eq. (4), measurements of the DPS cross sections in pPb collisions can therefore provide alternative extractions of the σ_{eff} parameter, independently of pp data. Such studies with pPb collisions extend the physics reach of the heavy ion program at the LHC [35].

A first study of DPS in pPb collisions was carried out by the LHCb experiment through measurements of the simultaneous production of different pairs of charmed hadrons at nucleon-nucleon center-of-mass energy $\sqrt{s_{\text{NN}}} = 8.16$ TeV [36]. This paper reports the measurement of the concurrent production of two J/ψ vector mesons in pPb collisions at the same center-of-mass energy. This process has not been previously observed and is of interest to clarify the SPS production of multiple quarkonium states [7, 37], to improve our understanding of DPS processes [6, 10], as well as to study nPDF properties [31]. The generic types of partonic scatterings contributing to prompt $J/\psi J/\psi + X$ production are schematically shown in Fig. 1. In SPS processes (Fig. 1, left), the two vector mesons are produced by a single pair of interacting partons, whereas two different pairs of partons produce the final $J/\psi J/\psi + X$ state in DPS (Fig. 1, center and right). The relatively high mass ($m_{J/\psi} = 3.1$ GeV), combined with the experimental requirement of $p_T^{J/\psi} > 6.5$ GeV, ensure that both charmonium mesons are produced in hard partonic scatterings with virtualities above $Q \approx \sqrt{m_{J/\psi}^2 + p_T^2} \approx 7$ GeV, which can be well described theoretically with perturbative calculations based, e.g., on the nonrelativistic QCD approach [38].

This analysis studies DPS in double charmonium production in pPb collisions at the LHC, following an approach similar to that used in the triple- J/ψ observation in pp collisions [16].




Figure 1: Representative diagrams of SPS (left) and DPS (center and right) contributions to the production of two J/ψ mesons in pPb collisions.

Exploiting the pPb data sample collected at $\sqrt{s_{\text{NN}}} = 8.16$ TeV in 2016, we measure the fiducial cross section for the rare final state of interest. The $J/\psi J/\psi + X$ cross section is measured in the fiducial volume of the detector, defined in Table 1, correcting for experimental inefficiencies. There is no extrapolation of the cross section for the final state beyond the fiducial acceptance, thus avoiding potential systematic uncertainties coming, e.g., from the polarization of the J/ψ mesons. The tabulated results of this study are provided in a HEPData record [39].

Table 1: Definition of the fiducial phase space for the $p\text{Pb} \rightarrow J/\psi J/\psi + X$ cross section measurement in pPb collisions at $\sqrt{s_{\text{NN}}} = 8.16$ TeV.

Particle	Fiducial requirement
Muons	$p_{\text{T}} > 3.4$ GeV for $ \eta < 0.3$
	$p_{\text{T}} > 3.3$ GeV for $0.3 < \eta < 1.1$
	$p_{\text{T}} > 5.5 - 2.0 \eta $ GeV for $1.1 < \eta < 2.1$
	$p_{\text{T}} > 1.3$ GeV for $2.1 < \eta < 2.4$
J/ψ mesons	$p_{\text{T}} > 6.5$ GeV and $ y < 2.4$

2 The CMS detector and data selection

The CMS apparatus [40] is a multipurpose, nearly hermetic detector, designed to trigger on [41, 42] and identify electrons, muons, photons, and charged and neutral hadrons [43–45]. A global “particle-flow” algorithm [46] aims to reconstruct all individual particles in an event, combining information provided by the all-silicon inner tracker and by the crystal electromagnetic and brass-scintillator hadron calorimeters, operating inside a 3.8 T superconducting solenoid, with data from the gas-ionization muon detectors embedded in the flux-return yoke outside the solenoid. The silicon tracker measures charged particles within the pseudorapidity range $|\eta| < 2.5$. During the LHC running period when the data used in this Letter were recorded, the silicon tracker consisted of 1440 silicon pixel and 15 148 silicon strip detector modules. For nonisolated particles of $1 < p_{\text{T}} < 10$ GeV and $|\eta| < 1.4$, the track resolutions are typically 1.5% in p_{T} and 25–90 (45–150) μm in the transverse (longitudinal) impact parameter [45]. The electromagnetic calorimeter (ECAL) consists of 75 848 lead tungstate crystals, which provide coverage in pseudorapidity $|\eta| < 1.48$ in a barrel region and $1.48 < |\eta| < 3.0$ in two endcap regions. Muons are measured in the pseudorapidity range $|\eta| < 2.4$, with detection planes made using three technologies: drift tubes, cathode strip chambers, and resistive plate chambers. The

forward hadron (HF) calorimeter extends the pseudorapidity coverage provided by the barrel and endcap detectors, and uses steel as an absorber and quartz fibers as the sensitive material. The two halves of the HF are located 11.2 m from the interaction region, one on each end, and together they provide coverage in the range $3.0 < |\eta| < 5.2$. They also serve as luminosity monitors. A more detailed description of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, can be found in Ref. [40].

Events are filtered using a two-tiered trigger system [42]. The first level, composed of custom hardware processors, uses information from the calorimeters and muon detectors [41]. The second level, known as the high-level trigger, consists of a farm of processors running a version of the full event reconstruction software optimized for fast processing, and reduces the event rate to around 1 kHz before data storage [42].

The events are collected with a trigger requiring two muon candidates found in the muon detectors at level-1. In order to maximize the detection efficiency [47], no explicit selection is made on their transverse momentum or pseudorapidity. During the 2016 pPb run, this trigger was operated without any prescale, recording an integrated luminosity of 174.6 nb^{-1} .

For the offline analysis, events have to pass a set of selection criteria designed to reject events from background processes (beam-gas collisions) as described in Ref. [48]. Events are required to have at least one reconstructed interaction vertex, formed by two or more associated tracks, with a distance from the center of the nominal interaction region of less than 25 cm along the beam direction and 0.2 cm in the plane transverse to the beam direction. The primary vertex (PV) is taken to be the vertex corresponding to the hardest scattering in the event, evaluated using tracking information alone, as described in Section 9.4.1 of Ref. [49]. In order to select inelastic hadronic collisions, the pPb events are also required to have at least one tower in each of the HF detectors with energy deposits of more than 3 GeV per tower. Any potential event containing more than one pPb interaction (pileup) is removed by the offline requirement of a common vertex for the four selected muons, as described below.

3 Event reconstruction and efficiency

Muons are reconstructed by combining information from the silicon tracker and the muon system composed of four layers of detectors (muon “stations”) [44]. The matching between tracks reconstructed in each of the subsystems proceeds either outside-in, starting from a track in the muon system, or inside-out, starting from a track provided by the silicon tracker. In the latter case, charged particles in the tracker that match track segments in only one or two muon stations are also considered in the analysis to collect low- p_T muons that may not have sufficient energy to traverse the entire muon system. The muons are selected from the reconstructed track candidates that match with at least one segment in any muon station in both x and y coordinates. Furthermore, the muon candidates are required to be identified as “soft”, as defined in Ref. [44]. Matching muons to tracks measured in the silicon tracker results in a relative transverse momentum resolution, for muons with p_T up to 100 GeV, of 1% in the barrel and 3% in the endcaps [44, 47].

All muons are required to pass a variable minimum p_T value based on their pseudorapidity, as described in Table 1. Muons with opposite-sign charge are combined into pairs with invariant masses $2.6 < m_{\mu\mu} < 3.6 \text{ GeV}$ to form a J/ψ candidate. No pair ambiguities (namely, cases where the two J/ψ mesons can be simultaneously reconstructed with two different combinations of the four muons), nor events with more than four muons satisfying the selection criteria, are found in data. The reconstructed J/ψ candidates are required to have $p_T > 6.5 \text{ GeV}$ and $|y| < 2.4$. The

four muons are fit to a common vertex, and the χ^2 probability of the fit is required to be larger than 0.1% [50]. A total of 16 candidate di- J/ψ events pass all selection criteria (Fig. 2). The requirement of the four muons to share the same vertex suppresses contributions from non-prompt J/ψ mesons originating from the decays of b quark hadrons. All selected events have vertices consistent with the PV within $100 \mu\text{m}$ and, therefore, the nonprompt single J/ψ meson contribution is expected to be, at most, a few percent of the prompt one within the selected kinematical range [51, 52]. This feature further confirms that the J/ψ mesons considered in this analysis are promptly produced.

Simulations based on the PYTHIA 8.303 [53] and EPOS-LHC [54] MC event generators are used to determine the overall efficiency of the analysis including the impact of the dimuon triggers, as well as of the single muon and J/ψ meson reconstruction within the fiducial volume defined in Table 1. The generated prompt J/ψ mesons are produced unpolarized in agreement with LHC data [55–57]. Generated PYTHIA-8 $pp \rightarrow J/\psi + X$ events are embedded into pPb events simulated with EPOS-LHC that replicate the hadronic activity from the underlying event. To improve the agreement between the simulated and measured charged particle multiplicities, a weighting procedure based on the number of tracks per event is applied when combining both MC samples. The generated events are then processed through a detailed simulation of the CMS detector based on the GEANT4 [58] package. The overall reconstruction efficiency is calculated as the ratio of the number of reconstructed events in the MC simulation (after all selection criteria are applied) to the number of generated events (after generator-level filters) within the fiducial phase space of the analysis.

4 Signal extraction

The signal is extracted from all events passing the fiducial requirements listed in Table 1, with a two-dimensional (2D) unbinned extended maximum likelihood fit that exploits the invariant masses of the two J/ψ meson candidates as variables. The signal J/ψ mass distribution is modeled with a Crystal-Ball function [59] and the combinatorial background is described by an exponential function. Both probability density functions (pdfs) are used twice, for the J/ψ_1 and J/ψ_2 states, per event, where the indices 1 and 2 indicate, respectively, the leading (i.e., highest p_T) and subleading J/ψ meson. The parameters describing the Crystal-Ball J/ψ signal model are fixed (for both J/ψ mesons) to values obtained from fits of the simulated distributions, the mean being essentially identical to the world-average J/ψ mass, $m_{J/\psi} = 3.1 \text{ GeV}$ [60], and the relative dimuon mass resolution being close to 1% ($\delta m_{J/\psi} \approx 30 \text{ MeV}$). The parameters of the exponential background are left free in the fit. The per-event 2D map of the two measured dimuon invariant mass distributions is shown in Fig. 2 (left) with the red circle indicating the double- J/ψ signal region within $\pm 5\delta m_{J/\psi}$ around the two J/ψ meson masses.

Table 2: Signal and background yields obtained from the likelihood fit procedure in the four-muon and dimuon-dielectron analyses over the $m_{\mu\mu,ee} = 2.6\text{--}3.6 \text{ GeV}$ mass range.

Yield	Final state	
	$\mu^+\mu^-\mu^+\mu^-$	$\mu^+\mu^-e^+e^-$
$N(J/\psi_1^{\text{sig}}, J/\psi_2^{\text{sig}})$	8.5 ± 3.4	5.7 ± 4.0
$N(J/\psi_1^{\text{bkg}}, J/\psi_2^{\text{sig}})$	0.9 ± 1.6	4.9 ± 3.8
$N(J/\psi_1^{\text{sig}}, J/\psi_2^{\text{bkg}})$	5.2 ± 3.0	4.5 ± 3.4
$N(J/\psi_1^{\text{bkg}}, J/\psi_2^{\text{bkg}})$	1.4 ± 1.9	6.0 ± 3.5

In total, four parameters representing the four combination of yields, $N(J/\psi_1^{\text{sig,bkg}}, J/\psi_2^{\text{sig,bkg}})$ for

each of the two J/ψ meson candidates to be either signal or background, are incorporated into the likelihood fit. The number of signal and background events derived from the fit procedure are listed in Table 2.

The projection of the result of the 2D fit in the two variables, $m_{\mu\mu,1}$ and $m_{\mu\mu,2}$, is shown in Fig. 2 (center and right panels). The data points indicate the measured counts, and the red hashed area shows the fitted signal component. The derived signal yield is $N_{\text{sig}} \equiv N(J/\psi_1^{\text{sig}}, J/\psi_2^{\text{sig}}) = 8.5 \pm 3.4$ events. The difference between the red area and the fit (solid black curve) indicates the size of the backgrounds, listed in Table 2, from events with four uncorrelated muons (mostly filling the area to the left of the J/ψ peaks) and with two uncorrelated muons coming from the $J/\psi_{1,2} + \mu^+\mu^-$ combination (filling the empty area below the peaks).




Figure 2: Dimuon invariant mass distributions of the pPb events passing all selection criteria, shown as a 2D map (left panel, with the numbers indicating the counts per bin and the red circle indicating the double J/ψ meson signal region) and its 1D projections for each pair (center and right panels). In the 1D projections, the measured data are represented by the points (with vertical bars showing their Poisson statistical uncertainties), the solid curve shows the overall signal + background fit to the data, and the red hashed area shows the signal yields for the $J/\psi J/\psi + X$ process.

The signal has a significance of 4.9 standard deviations, obtained from the likelihood ratio of the default signal-plus-background fit and the background-only fit (imposing $N_{\text{sig}} = 0$), using the standard asymptotic formula [61] assuming that the Wilks theorem conditions apply [62]. As discussed below, the systematic uncertainties are small (6.1%, with a fraction of them impacting equally signal and background) and do not alter the statistical significance evaluation. This significance was also validated with MC pseudo-experiments.

To fully confirm the observation of the double- J/ψ process, events are also analyzed where the leading J/ψ meson is reconstructed in its $\mu^+\mu^-$ decay mode and the other decays via $J/\psi \rightarrow e^+e^-$. This channel has a larger background than the 4-muon one, but increases the total size of the data sample and provides a cross-check of the latter. The electron momentum is estimated by combining the ECAL energy measurement (including the energy sum of all bremsstrahlung photons spatially compatible with originating from the electron trajectory) with the momentum measurement in the tracker. The momentum resolution is typically smaller than 5% for electrons in the range $2 < p_T < 10$ GeV. It is generally better in the barrel region than in the endcaps, and also depends on the fraction of the electron energy emitted through bremsstrahlung (f_{brem}) as it traverses the material in front of the ECAL [43, 63]. Both electrons are required to have $f_{\text{brem}} > 0.01$ (to remove any potential charged-pion contamination) and to be of opposite-sign charge.

For the $\mu^+\mu^- + e^+e^-$ channel, the selection of the dimuon candidates remains the same as

for the double- $\mu^+\mu^-$ analysis, and the candidate e^\pm are required to have $p_T > 2.5$ GeV and $|\eta| < 2.5$ and to share the same common vertex. The parameters of the Crystal-Ball J/ψ meson signal are derived from the MC simulation, with the resulting mass resolution of the dielectron decay mode being $\delta m_{J/\psi} \approx 70$ MeV. A total number of 21 events pass the $(\mu^+\mu^-, e^+e^-)$ criteria with both dileptons having invariant masses over $m_{\mu\mu,ee} = 2.6\text{--}3.6$ GeV. No events passing the selection criteria of the dimuon-dielectron channel are present in the double-dimuon sample.

The fit procedure followed is the same as for the four-muon final state. The per-event 2D map of the dimuon-dielectron invariant mass distributions is shown in Fig. 3 (left) with the red circle indicating the double- J/ψ signal region within $\pm 5\delta m_{J/\psi}$ around the masses of the two J/ψ mesons. The projection of the result of the 2D fit in the two variables, $m_{\mu\mu,1}$ and $m_{ee,2}$, is shown in Fig. 3 (center and right panels). The signal yield (red hashed area) is 5.7 ± 4.0 events.




Figure 3: Dilepton invariant mass distributions of the pPb events passing all selection criteria, shown as a 2D map (left panel, with the numbers indicating the counts per bin and the red circle indicating the double J/ψ meson signal region) and its 1D projections for each pair (center and right panels). In the 1D projections, the measured data are represented by the points (vertical bars showing their Poisson statistical uncertainties), the solid curve shows the overall signal + background fit to the data, and the red hashed area shows the signal yields for the $J/\psi J/\psi + X$ process.

The significance of the $(J/\psi \rightarrow e^+e^- + J/\psi \rightarrow \mu^+\mu^-)$ mode, calculated in the same way as for the $(J/\psi \rightarrow \mu^+\mu^- + J/\psi \rightarrow \mu^+\mu^-)$ channel, is 2.3 standard deviations. The combination of the two decay modes, using the Fisher formalism [64] with $p^{\text{value}} = p_1^{\text{value}} p_2^{\text{value}} (1 - \log(p_1^{\text{value}} p_2^{\text{value}}))$, where $p_{1,2}$ are the p -values of each independent analysis, gives a total statistical significance of 5.3 standard deviations.

5 Results

The cross section of the double- J/ψ production reconstructed in the four-muon decay mode is obtained through the expression

$$\sigma(\text{pPb} \rightarrow J/\psi J/\psi + X) = \frac{N_{\text{sig}}}{\epsilon \mathcal{L}_{\text{int}} \mathcal{B}_{J/\psi \rightarrow \mu^+\mu^-}^2}, \quad (5)$$

where $N_{\text{sig}} = 8.5 \pm 3.4$ is the number of signal events reconstructed with an overall efficiency ϵ in the fiducial region defined in Table 1, $\mathcal{L} = 174.6 \pm 6.1 \text{ nb}^{-1}$ is the total integrated luminosity, and $\mathcal{B}_{J/\psi \rightarrow \mu^+\mu^-} = (5.961 \pm 0.033)\%$ is the dimuon decay branching fraction [60]. We do not compute the cross section corresponding to the $(J/\psi \rightarrow e^+e^- + J/\psi \rightarrow \mu^+\mu^-)$ decay mode

because it has much larger backgrounds and is not precise enough to quantitatively improve the overall cross section extraction. To determine the number of signal events corrected for the reconstruction efficiency, we use the expression $N_{\text{sig}}/\epsilon = \sum_i N_{\text{sig}}^i/\epsilon^i$, where N_{sig}^i are statistical per-event weights (having a value between 1 and 0, indicating events being more or less signal-like) obtained using the SPLOT technique [65]. Based on the kinematics of each J/ψ meson in the p_T and rapidity plane, the ϵ^i values are calculated from a 2D efficiency map obtained from the simulated signal. Any mismatches on the reconstruction and trigger efficiency observed between the MC simulation and the measured data are corrected by applying the scale factors described in Ref. [44]. The double- J/ψ efficiency is $\epsilon = 62.1\%$, consistent with the squared single- J/ψ efficiency, as expected, given that it is driven by the kinematics of the two individual J/ψ mesons.

The systematic uncertainties in the cross section determined with Eq. (5) are listed in Table 3. The dependence of the measured cross section on the modeling of the J/ψ meson signal and background subtraction is examined by replacing the Crystal-Ball pdf of the nominal model by a Crystal-Ball-plus-Gaussian function, as well as the exponential background by a first-order polynomial distribution. The differences observed in the signal yields are 4.0 and 2.5%, for the variations of the signal and background pdfs, respectively. Releasing the constraint of the J/ψ signal mean and width parameters from the MC simulation results in a negligible difference (0.1%) and is not considered as a source of systematic uncertainty. A 3.5% uncertainty is added from the integrated luminosity measurement [66, 67]. The uncertainty in the scale factors applied to the MC simulation to correct for possible data/MC differences amounts to 1.3%. Finally, the uncertainty of $\mathcal{B}_{J/\psi \rightarrow \mu^+\mu^-}$ propagates into an uncertainty of 1.1% in the cross section measurement.

Table 3: Relative contributions to the systematic uncertainty in the $\sigma(\text{pPb} \rightarrow J/\psi J/\psi + X)$ measurement. The last row lists the sum in quadrature of all components.

Source of uncertainty	$\sigma(\text{pPb} \rightarrow J/\psi J/\psi + X)$
J/ψ meson signal shape	4.0%
Dimuon continuum background shape	2.5%
Integrated luminosity	3.5%
Efficiency scale factors	1.3%
Branching fraction	1.1%
Total	6.1%

The resulting fiducial cross section is $\sigma(\text{pPb} \rightarrow J/\psi J/\psi) = 22.0 \pm 8.9$ (stat) ± 1.5 (syst) nb. Such a cross section contains in principle contributions from the SPS and DPS production channels shown in Fig. 1. Namely, the result is $\sigma(\text{pPb} \rightarrow J/\psi J/\psi + X) = \sigma_{\text{SPS}}^{\text{pPb} \rightarrow J/\psi J/\psi + X} + \sigma_{\text{DPS}}^{\text{pPb} \rightarrow J/\psi J/\psi + X}$, and the size of each contribution is estimated from theoretical calculations (matched to existing LHC experimental J/ψ data) and the measured double J/ψ meson differential distributions, as explained in the next section.

6 Discussion

In the simplest factorized approach represented by Eq. (1), the DPS double J/ψ meson cross section can be derived from the squared single J/ψ meson cross section divided by σ_{eff} . The single- J/ψ cross sections in pPb and pp collisions are well understood both experimentally and theoretically thanks to multiple measurements and calculations that agree with each other within less than 10% [7]. We use here the perturbative QCD calculation at next-to-leading

order (NLO) accuracy from HELAC-ONIA [68, 69] using the CT14NLO PDF [70] for the proton and the EPPS16 nPDF for the Pb nucleus [71], reweighted as discussed in Refs. [72, 73]. The predicted SPS cross section (Table 4, first row) is normalized so as to reproduce the inclusive J/ψ measurements in pPb collisions at $\sqrt{s_{\text{NN}}} = 8.16$ TeV [74], and thus it does not have any theoretical scale uncertainties assigned but only a $\approx 6\%$ data normalization uncertainty. The proton-nucleon (pN) cross section (obtained by dividing the pPb cross section by $A = 208$) within the fiducial requirements of Table 1 amounts to $\sigma_{\text{SPS}}^{\text{pPb} \rightarrow J/\psi + X} \mathcal{B}(J/\psi \rightarrow \mu^+ \mu^-) / A = 21.7 \pm 1.4$ (nPDF) ± 1.4 (norm) nb, to be compared with the corresponding pp cross section of $\sigma_{\text{SPS}}^{\text{pp} \rightarrow J/\psi + X} \mathcal{B}(J/\psi \rightarrow \mu^+ \mu^-) = 25.0 \pm 1.8$ (PDF) nb, indicating a $\approx 10\%$ reduction due to nPDF shadowing. The theoretical prediction for the prompt double J/ψ meson SPS cross section (Table 4, second row) is obtained using the color singlet model at approximately NLO accuracy [75, 76] computed with HELAC-ONIA using the CT14NLO proton PDF and the reweighted EPPS16 nPDF for the Pb nucleus. The double J/ψ meson SPS cross section in pN collisions within the fiducial phase space of Table 1 amounts to $\sigma_{\text{SPS}}^{\text{pPb} \rightarrow J/\psi J/\psi + X} \mathcal{B}^2(J/\psi \rightarrow \mu^+ \mu^-) / A = 0.097_{-0.063}^{+0.185}$ (scale) ± 0.004 (nPDF) pb, to be compared with a pp cross section of $\sigma_{\text{SPS}}^{\text{pp} \rightarrow J/\psi J/\psi + X} \mathcal{B}^2(J/\psi \rightarrow \mu^+ \mu^-) = 0.105_{-0.068}^{+0.200}$ (scale) ± 0.004 (PDF) pb, indicating about 8% nPDF shadowing corrections.

Table 4: Predictions for single and double J/ψ meson production cross sections in SPS processes in pPb collisions at $\sqrt{s_{\text{NN}}} = 8.16$ TeV within the fiducial phase space of Table 1, obtained with the HELAC-ONIA code using the CT14NLO proton PDF and the reweighted EPPS16 lead nPDF, as explained in the text. The quoted uncertainty of the single (double) J/ψ meson cross section includes nPDF and normalization (nPDF and scale) uncertainties, added in quadrature.

Process	Theoretical cross section
$\sigma_{\text{SPS}}^{\text{pPb} \rightarrow J/\psi + X} \mathcal{B}(J/\psi \rightarrow \mu^+ \mu^-)$	$4.51 \pm 0.42 \mu\text{b}$
$\sigma_{\text{SPS}}^{\text{pPb} \rightarrow J/\psi J/\psi + X} \mathcal{B}^2(J/\psi \rightarrow \mu^+ \mu^-)$	$20.2_{-13.1}^{+38.5} \text{pb}$

The theoretical SPS single J/ψ meson cross section has an overall uncertainty of $\pm 9\%$, and thus so will have the corresponding DPS cross section (for a given σ_{eff}). However, the SPS double J/ψ meson cross section is poorly constrained, and can vary within a $+200\%$, -65% range [76]. Therefore, we use our own data in order to constrain the SPS and DPS contributions to the measured double- J/ψ cross section. The most discriminating variables to distinguish SPS from DPS processes are the separation in azimuth $\Delta\phi$ and rapidity Δy between the two produced particles [2]. The J/ψ mesons produced by DPS processes are expected to be completely decorrelated in their production and, thus, should feature flatter distributions in their relative azimuthal and rapidity separations (blue histograms in Fig. 4). On the other hand, the SPS mechanism leads to a correlated production of both J/ψ mesons that prefer to be produced closer in rapidity and with a small or a maximum separation in azimuthal angle. Theoretical expectations based on contributions from multiple SPS channels, computed up to $\mathcal{O}(\alpha_s^6)$ perturbative accuracy and cross-checked with existing data [76], indicate that the distributions of the two J/ψ mesons from SPS processes tend to have a rapidity separation peaked at $\Delta y = 0$ and maxima at $\Delta\phi \approx 0, \pi$ (Fig. 4).

Due to the limited size of our signal data sample, it is not possible to perform a 2D template fit of the measured $\Delta\phi$ - Δy distributions with shapes dictated by the theoretical expectations to extract the absolute normalization of the poorly known SPS double J/ψ meson yields. Instead, we identify a phase space region where the SPS contribution is expected to be minimal, and constrain there the expected size of the DPS yield. For this purpose, we use the predicted

shape of the Δy distribution, which shows that the $\Delta y \gtrsim 2$ region is free from SPS contributions (red histogram in Fig. 4, left) and, therefore, any signal count measured there is dominated by DPS production.

To extract the number of DPS events, we perform a 1D fit to the Δy variable. Using a DPS template built by combining J/ψ mesons from different events, we fit the weighted data in the DPS dominated region of $\Delta y > 2$. The fit results in a yield of 2.1 DPS events and $\sigma_{\text{eff}} = 4.0 \text{ mb}$, as explained in more detail below, shown as a blue histogram in Fig. 4. The left plot shows the fit result of the Δy distribution compared with the data (markers). We also overlay the combined SPS + DPS distribution shown with the red histogram (and hashed uncertainty bands in red). Figure 4 (right) shows the corresponding $\Delta\phi$ distribution, with the DPS and SPS + DPS distributions having the same normalization as in the left plot.




Figure 4: Distribution of rapidity separation (left) between the two J/ψ mesons in data (black markers), fitted template of the expected DPS contribution (blue histogram) normalized to the data outside of the SPS-dominated region (as described in the text), and fitted SPS + DPS distribution (red histogram). Distribution of azimuthal separation (right) between the two J/ψ mesons measured in data (black markers) and expected DPS contribution (blue histogram), and SPS + DPS sum (red histogram), with the relative normalization derived from the Δy distribution (left panel), as explained in the text.

This fit procedure allows the extraction of the number of SPS and DPS signal counts: $N_{\text{SPS}} = 6.4 \pm 4.2$ and $N_{\text{DPS}} = 2.1 \pm 2.4$, respectively. The corresponding SPS and DPS fiducial cross sections, derived using Eq. (5), amount to $\sigma_{\text{SPS}}^{\text{pPb} \rightarrow J/\psi J/\psi + X} = 16.5 \pm 10.8 \text{ (stat)} \pm 0.1 \text{ (syst)} \text{ nb}$ and $\sigma_{\text{DPS}}^{\text{pPb} \rightarrow J/\psi J/\psi + X} = 5.4 \pm 6.2 \text{ (stat)} \pm 0.4 \text{ (syst)} \text{ nb}$. From this latter value, we can extract the effective $\sigma_{\text{eff,pPb}}$ cross section via Eq. (3). Using the theoretical predictions (constrained by existing data) for $\sigma_{\text{SPS}}^{\text{pPb} \rightarrow J/\psi + X}$ from Table 4, we obtain $\sigma_{\text{eff,pPb}} = 0.53_{-0.2}^{+\infty} \text{ b}$. The extracted $\sigma_{\text{eff,pPb}}$ value has an arbitrarily large upper uncertainty that indicates that the SPS yield alone would be compatible with the data.

By neglecting parton correlations, and assuming that the double PDF of the nucleons can be factorized in longitudinal and transverse components, the effective DPS cross section for pPb collisions can be converted into a pp-equivalent σ_{eff} one, through Eq. (4). Using $A = 208$ for a Pb nucleus and a factor $F_{\text{pPb}} = 29.5 \text{ mb}^{-1}$ derived from the pPb thickness function with a

Glauber MC model, a value $\sigma_{\text{eff}} = 4.0^{+\infty}_{-1.5}$ mb is obtained, where the unbound upper uncertainty indicates that the double J/ψ meson yields could in principle be produced without DPS contributions. Therefore, the final result is better expressed as a limit on the maximum amount of DPS allowed by the data, which translates into a lower limit on the effective DPS cross section of $\sigma_{\text{eff}} > 1.0$ mb at 95% confidence level (CL).

In Fig. 5, the lower σ_{eff} bound determined in this work is compared with other extractions of the effective DPS cross section derived from double- and triple-quarkonium production measurements [12, 14–17, 37, 77–80] (blue circles), as well as from processes with jets, photons, and W bosons [19, 26, 81–84] (black squares). A few of the quarkonium σ_{eff} values plotted (indicated with an asterisk in the legend) are those derived by more recent phenomenological studies [37, 77, 79, 80] of the experimental data, with an improved evaluation of the SPS contributions. Overall, the effective cross sections obtained from multi-quarkonium production favor a smaller value of $\sigma_{\text{eff}} \approx 3\text{--}10$ mb compared with the $\sigma_{\text{eff}} \approx 10\text{--}20$ mb result of the other types of multiple hard scattering processes. Our measurement of $\sigma_{\text{eff}} > 1.0$ mb at 95% CL calls for larger pPb data sets to confirm, or not, its consistency with the σ_{eff} values derived from pp data and, eventually, better constrain the average effective interparton distance squared (σ_{eff}) involved in the production of a pair of quarkonium mesons in hadronic collisions.




Figure 5: Comparison of the effective DPS cross sections σ_{eff} extracted in this work (red arrow lower limit) with the same parameter derived in pp measurements of double- and triple-quarkonium production [12, 14–16, 37, 77, 79, 80] (blue circles), as well as in final states with jets [81, 82, 85], γ + jets [83, 84, 86], W+ jets [19, 20], and double W bosons [26] (black squares). The asterisk shown in a few legend entries indicates that the result has been obtained by more recent phenomenological analyses of the experimental data.

7 Summary

The first observation of the concurrent production of two J/ψ mesons in proton-lead (pPb) collisions has been reported. Events with two J/ψ mesons, each decaying into two muons, have been reconstructed in pPb collisions at a nucleon-nucleon center-of-mass-energy of $\sqrt{s_{\text{NN}}} = 8.16$ TeV. The data sample corresponds to an integrated luminosity of 174.6 nb^{-1} collected by the CMS experiment at the CERN LHC. After all selection requirements, 8.5 ± 3.4 events are found that confirm the production of the $J/\psi J/\psi + X$ final state. Events where the least energetic J/ψ meson is reconstructed in the dielectron channel are also considered in the search, leading to a signal yield of 5.7 ± 4.0 events. The statistical significance of the signal, relative to the background-only expectation, corresponds to 5.3 standard deviations (4.9 standard deviations in the four-muon channel alone). The fiducial cross section for $J/\psi J/\psi + X$ production is $\sigma(\text{pPb} \rightarrow J/\psi J/\psi + X) = 22.0 \pm 8.9 (\text{stat}) \pm 1.5 (\text{syst}) \text{ nb}$. This result is compared with the theoretical expectations for $J/\psi J/\psi + X$ production via the sum of contributions from single- and double-parton scatterings (SPS and DPS, respectively). Under the simplest assumption of factorization of multiple hard scattering probabilities in terms of SPS cross sections, the measured $J/\psi J/\psi + X$ cross section is consistent with the production of SPS + DPS processes with fiducial cross sections $\sigma_{\text{SPS}}^{\text{pPb} \rightarrow J/\psi J/\psi + X} = 16.5 \pm 10.8 (\text{stat}) \pm 0.1 (\text{syst}) \text{ nb}$ and $\sigma_{\text{DPS}}^{\text{pPb} \rightarrow J/\psi J/\psi + X} = 5.4 \pm 6.2 (\text{stat}) \pm 0.4 (\text{syst}) \text{ nb}$. The derived DPS cross section can be transformed into a lower bound on the effective DPS cross section parameter (σ_{eff}) that is closely related to the squared average interparton transverse separation in the collision. A limit of $\sigma_{\text{eff}} > 1.0 \text{ mb}$ at 95% confidence level is set, which is consistent with σ_{eff} values obtained from double and triple quarkonium measurements in pp collisions. The present analysis supports the interest of exploiting the production of multiple heavy and/or high- p_{T} particles in pPb collisions at the LHC as a novel means to study the dynamics of multiple independent hard scatterings.

Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid and other centers for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: SC (Armenia), BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); ERC PRG, RVTT3 and MoER TK202 (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); SRNSF (Georgia); BMBF, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LMTLT (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES and NSC (Poland); FCT (Portugal); MESTD (Serbia); MCIN/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI and NSTDA (Thailand); TUBITAK and TENMAK (Turkey); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (USA).

Individuals have received support from the Marie-Curie program and the European Research

Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, 101115353, 101002207, and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Science Committee, project no. 22rl-037 (Armenia); the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the F.R.S.-FNRS and FWO (Belgium) under the "Excellence of Science – EOS" – be.h project n. 30820817; the Beijing Municipal Science & Technology Commission, No. Z191100007219010 and Fundamental Research Funds for the Central Universities (China); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Shota Rustaveli National Science Foundation, grant FR-22-985 (Georgia); the Deutsche Forschungsgemeinschaft (DFG), under Germany's Excellence Strategy – EXC 2121 "Quantum Universe" – 390833306, and under project number 400140256 - GRK2497; the Hellenic Foundation for Research and Innovation (HFRI), Project Number 2288 (Greece); the Hungarian Academy of Sciences, the New National Excellence Program - ÚNKP, the NKFIH research grants K 131991, K 133046, K 138136, K 143460, K 143477, K 146913, K 146914, K 147048, 2020-2.2.1-ED-2021-00181, and TKP2021-NKTA-64 (Hungary); the Council of Science and Industrial Research, India; ICSC – National Research Center for High Performance Computing, Big Data and Quantum Computing and FAIR – Future Artificial Intelligence Research, funded by the NextGenerationEU program (Italy); the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369 and 2021/43/B/ST2/01552 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; MCIN/AEI/10.13039/501100011033, ERDF "a way of making Europe", and the Programa Estatal de Fomento de la Investigación Científica y Técnica de Excelencia María de Maeztu, grant MDM-2017-0765 and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, and the National Science, Research and Innovation Fund via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, grant B37G660013 (Thailand); the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (USA).

References

- [1] M. Diehl and J. R. Gaunt, "Double parton scattering theory overview", *Adv. Ser. Direct. High Energy Phys.* **29** (2018) 7, doi:10.1142/9789813227767_0002, arXiv:1710.04408.
- [2] P. Bartalini and J. R. Gaunt, eds., "Multiple parton interactions at the LHC", volume 29. World Scientific, 2019. doi:10.1142/10646, ISBN 978-981-322-775-0, 978-981-322-777-4.
- [3] CMS Collaboration, "Search for Higgs and Z boson decays to J/ψ or Y pairs in the four-muon final state in proton-proton collisions at $\sqrt{s} = 13$ TeV", *Phys. Lett. B* **797** (2019) 134811, doi:10.1016/j.physletb.2019.134811, arXiv:1905.10408.
- [4] CMS Collaboration, "Measurement of the $Y(1S)$ pair production cross section and search for resonances decaying to $Y(1S)\mu^+\mu^-$ in proton-proton collisions at $\sqrt{s} = 13$ TeV", *Phys. Lett. B* **808** (2020) 135578, doi:10.1016/j.physletb.2020.135578, arXiv:2002.06393.

-
- [5] CMS Collaboration, “New structures in the $J/\psi/\psi$ mass spectrum in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *Phys. Rev. Lett.* **132** (2024) 111901, doi:10.1103/PhysRevLett.132.111901, arXiv:2306.07164.
- [6] D. d’Enterria and A. Snigirev, “Double, triple, and n -parton scatterings in high-energy proton and nuclear collisions”, *Adv. Ser. Direct. High Energy Phys.* **29** (2018) 159, doi:10.1142/9789813227767_0009, arXiv:1708.07519.
- [7] E. Chapon et al., “Prospects for quarkonium studies at the high-luminosity LHC”, *Prog. Part. Nucl. Phys.* **122** (2022) 103906, doi:10.1016/j.pnpnp.2021.103906, arXiv:2012.14161.
- [8] T. Sjöstrand, “The development of MPI modeling in PYTHIA”, *Adv. Ser. Direct. High Energy Phys.* **29** (2019) 191, doi:10.1142/9789813227767_0010, arXiv:1706.02166.
- [9] M. H. Seymour and A. Siodmok, “Constraining MPI models using σ_{eff} and recent Tevatron and LHC underlying event data”, *JHEP* **10** (2013) 113, doi:10.1007/JHEP10(2013)113, arXiv:1307.5015.
- [10] B. Blok and M. Strikman, “Multiparton pp and pA collisions: From geometry to parton-parton correlations”, *Adv. Ser. Direct. High Energy Phys.* **29** (2018) 63, doi:10.1142/9789813227767_0005, arXiv:1709.00334.
- [11] LHCb Collaboration, “Observation of J/ψ pair production in pp collisions at $\sqrt{s} = 7$ TeV”, *Phys. Lett. B* **707** (2012) 52, doi:10.1016/j.physletb.2011.12.015, arXiv:1109.0963.
- [12] D0 Collaboration, “Observation and studies of double J/ψ production at the Tevatron”, *Phys. Rev. D* **90** (2014) 111101, doi:10.1103/PhysRevD.90.111101, arXiv:1406.2380.
- [13] CMS Collaboration, “Measurement of prompt J/ψ pair production in pp collisions at $\sqrt{s} = 7$ TeV”, *JHEP* **09** (2014) 094, doi:10.1007/JHEP09(2014)094, arXiv:1406.0484.
- [14] ATLAS Collaboration, “Measurement of the prompt J/ψ pair production cross-section in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **77** (2017) 76, doi:10.1140/epjc/s10052-017-4644-9, arXiv:1612.02950.
- [15] LHCb Collaboration, “Measurement of the J/ψ pair production cross-section in pp collisions at $\sqrt{s} = 13$ TeV”, *JHEP* **06** (2017) 047, doi:10.1007/JHEP06(2017)047, arXiv:1612.07451. [Erratum: doi:10.1007/JHEP10(2017)068].
- [16] CMS Collaboration, “Observation of triple J/ψ meson production in proton-proton collisions”, *Nature Phys.* **19** (2023) 338, doi:10.1038/s41567-022-01838-y, arXiv:2111.05370. [Erratum: doi:10.1038/s41567-023-01992-x].
- [17] D0 Collaboration, “Evidence for simultaneous production of J/ψ and Y mesons”, *Phys. Rev. Lett.* **116** (2016) 082002, doi:10.1103/PhysRevLett.116.082002, arXiv:1511.02428.
- [18] CMS Collaboration, “Observation of $Y(1S)$ pair production in proton-proton collisions at $\sqrt{s} = 8$ TeV”, *JHEP* **05** (2017) 013, doi:10.1007/JHEP05(2017)013, arXiv:1610.07095.

- [19] ATLAS Collaboration, “Measurement of hard double-parton interactions in $W(\rightarrow \ell\nu)+2$ jet events at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *New J. Phys.* **15** (2013) 033038, doi:10.1088/1367-2630/15/3/033038, arXiv:1301.6872.
- [20] CMS Collaboration, “Study of double parton scattering using $W + 2$ -jet events in proton-proton collisions at $\sqrt{s} = 7$ TeV”, *JHEP* **03** (2014) 032, doi:10.1007/JHEP03(2014)032, arXiv:1312.5729.
- [21] ATLAS Collaboration, “Study of hard double-parton scattering in four-jet events in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS experiment”, *JHEP* **11** (2016) 110, doi:10.1007/JHEP11(2016)110, arXiv:1608.01857.
- [22] CMS Collaboration, “Constraints on the double-parton scattering cross section from same-sign W boson pair production in proton-proton collisions at $\sqrt{s} = 8$ TeV”, *JHEP* **02** (2018) 032, doi:10.1007/JHEP02(2018)032, arXiv:1712.02280.
- [23] ATLAS Collaboration, “Study of the hard double-parton scattering contribution to inclusive four-lepton production in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Phys. Lett. B* **790** (2019) 595, doi:10.1016/j.physletb.2019.01.062, arXiv:1811.11094.
- [24] CMS Collaboration, “Evidence for WW production from double-parton interactions in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *Eur. Phys. J. C* **80** (2020) 41, doi:10.1140/epjc/s10052-019-7541-6, arXiv:1909.06265.
- [25] CMS Collaboration, “Measurement of double-parton scattering in inclusive production of four jets with low transverse momentum in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JHEP* **01** (2022) 177, doi:10.1007/JHEP01(2022)177, arXiv:2109.13822.
- [26] CMS Collaboration, “Observation of same-sign WW production from double parton scattering in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *Phys. Rev. Lett.* **131** (2023) 091803, doi:10.1103/PhysRevLett.131.091803, arXiv:2206.02681.
- [27] E. Huayra, J. V. C. Lovato, and E. G. de Oliveira, “Valence and sea parton correlations in double parton scattering from data”, *JHEP* **09** (2023) 177, doi:10.1007/JHEP09(2023)177, arXiv:2305.11106.
- [28] M. Strikman and D. Treleani, “Measuring double parton distributions in nucleons at proton nucleus colliders”, *Phys. Rev. Lett.* **88** (2002) 031801, doi:10.1103/PhysRevLett.88.031801, arXiv:hep-ph/0111468.
- [29] E. Cattaruzza, A. Del Fabbro, and D. Treleani, “Heavy-quark production in proton-nucleus collision at the CERN LHC”, *Phys. Rev. D* **70** (2004) 034022, doi:10.1103/PhysRevD.70.034022, arXiv:hep-ph/0404177.
- [30] D. d’Enterria and A. M. Snigirev, “Same-sign WW production in proton-nucleus collisions at the LHC as a signal for double parton scattering”, *Phys. Lett. B* **718** (2013) 1395, doi:10.1016/j.physletb.2012.12.032, arXiv:1211.0197.
- [31] H.-S. Shao, “Probing impact-parameter dependent nuclear parton densities from double parton scatterings in heavy-ion collisions”, *Phys. Rev. D* **101** (2020) 054036, doi:10.1103/PhysRevD.101.054036, arXiv:2001.04256.

-
- [32] D. d’Enterria and C. Loizides, “Progress in the Glauber model at collider energies”, *Ann. Rev. Nucl. Part. Sci.* **71** (2021) 315, doi:10.1146/annurev-nucl-102419-060007, arXiv:2011.14909.
- [33] CMS Collaboration, “Measurement of the inelastic cross section in proton-lead collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV”, *Phys. Lett. B* **759** (2016) 641, doi:10.1016/j.physletb.2016.06.027, arXiv:1509.03893.
- [34] C. Loizides, J. Kamin, and D. d’Enterria, “Improved Monte Carlo Glauber predictions at present and future nuclear colliders”, *Phys. Rev. C* **97** (2018) 054910, doi:10.1103/PhysRevC.97.054910, arXiv:1710.07098. [Erratum: doi:10.1103/PhysRevC.99.019901].
- [35] CMS Collaboration, “Overview of high-density QCD studies with the CMS experiment at the LHC”, 2024. arXiv:2405.10785. Submitted to *Phys. Rep.*
- [36] LHCb Collaboration, “Observation of enhanced double parton scattering in proton-lead collisions at $\sqrt{s_{\text{NN}}} = 8.16$ TeV”, *Phys. Rev. Lett.* **125** (2020) 212001, doi:10.1103/PhysRevLett.125.212001, arXiv:2007.06945.
- [37] J.-P. Lansberg, “New observables in inclusive production of quarkonia”, *Phys. Rept.* **889** (2020) 1, doi:10.1016/j.physrep.2020.08.007, arXiv:1903.09185.
- [38] G. T. Bodwin, E. Braaten, and G. P. Lepage, “Rigorous QCD analysis of inclusive annihilation and production of heavy quarkonium”, *Phys. Rev. D* **51** (1995) 1125, doi:10.1103/PhysRevD.55.5853, arXiv:hep-ph/9407339. [Erratum: doi:10.1103/PhysRevD.55.5853].
- [39] HEPData record for this analysis, 2024. doi:10.17182/hepdata.152618.
- [40] CMS Collaboration, “The CMS experiment at the CERN LHC”, *JINST* **3** (2008) S08004, doi:10.1088/1748-0221/3/08/S08004.
- [41] CMS Collaboration, “Performance of the CMS level-1 trigger in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JINST* **15** (2020) P10017, doi:10.1088/1748-0221/15/10/P10017, arXiv:2006.10165.
- [42] CMS Collaboration, “The CMS trigger system”, *JINST* **12** (2017) P01020, doi:10.1088/1748-0221/12/01/P01020, arXiv:1609.02366.
- [43] CMS Collaboration, “Electron and photon reconstruction and identification with the CMS experiment at the CERN LHC”, *JINST* **16** (2021) P05014, doi:10.1088/1748-0221/16/05/P05014, arXiv:2012.06888.
- [44] CMS Collaboration, “Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JINST* **13** (2018) P06015, doi:10.1088/1748-0221/13/06/P06015, arXiv:1804.04528.
- [45] CMS Collaboration, “Description and performance of track and primary-vertex reconstruction with the CMS tracker”, *JINST* **9** (2014) P10009, doi:10.1088/1748-0221/9/10/P10009, arXiv:1405.6569.
- [46] CMS Collaboration, “Particle-flow reconstruction and global event description with the CMS detector”, *JINST* **12** (2017) P10003, doi:10.1088/1748-0221/12/10/P10003, arXiv:1706.04965.

- [47] CMS Collaboration, “Performance of CMS muon reconstruction from proton-proton to heavy ion collisions”, 2024. arXiv:2404.17377. Submitted to *JINST*.
- [48] CMS Collaboration, “Charged-particle nuclear modification factors in PbPb and pPb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV”, *JHEP* **04** (2017) 039, doi:10.1007/JHEP04(2017)039, arXiv:1611.01664.
- [49] CMS Collaboration, “Technical proposal for the Phase-II upgrade of the Compact Muon Solenoid”, CMS Technical Proposal CERN-LHCC-2015-010, CMS-TDR-15-02, 2015.
- [50] R. Frühwirth, “Application of Kalman filtering to track and vertex fitting”, *Nucl. Instrum. Meth. A* **262** (1987) 444, doi:10.1016/0168-9002(87)90887-4.
- [51] CMS Collaboration, “Measurement of quarkonium production cross sections in pp collisions at $\sqrt{s} = 13$ TeV”, *Phys. Lett. B* **780** (2018) 251, doi:10.1016/j.physletb.2018.02.033, arXiv:1710.11002.
- [52] ATLAS Collaboration, “Measurement of the differential cross-sections of inclusive, prompt and non-prompt J/ψ production in proton-proton collisions at $\sqrt{s} = 7$ TeV”, *Nucl. Phys. B* **850** (2011) 387, doi:10.1016/j.nuclphysb.2011.05.015, arXiv:1104.3038.
- [53] T. Sjöstrand et al., “An introduction to PYTHIA 8.2”, *Comput. Phys. Commun.* **191** (2015) 159, doi:10.1016/j.cpc.2015.01.024, arXiv:1410.3012.
- [54] T. Pierog et al., “EPOS LHC: Test of collective hadronization with data measured at the CERN Large Hadron Collider”, *Phys. Rev. C* **92** (2015) 034906, doi:10.1103/PhysRevC.92.034906, arXiv:1306.0121.
- [55] CMS Collaboration, “Measurement of the prompt J/ψ and $\psi(2S)$ polarizations in pp collisions at $\sqrt{s} = 7$ TeV”, *Phys. Lett. B* **727** (2013) 381, doi:10.1016/j.physletb.2013.10.055, arXiv:1307.6070.
- [56] LHCb Collaboration, “Measurement of J/ψ polarization in pp collisions at $\sqrt{s} = 7$ TeV”, *Eur. Phys. J. C* **73** (2013) 2631, doi:10.1140/epjc/s10052-013-2631-3, arXiv:1307.6379.
- [57] ALICE Collaboration, “First measurement of quarkonium polarization in nuclear collisions at the LHC”, *Phys. Lett. B* **815** (2021) 136146, doi:10.1016/j.physletb.2021.136146, arXiv:2005.11128.
- [58] GEANT4 Collaboration, “GEANT4 — a simulation toolkit”, *Nucl. Instrum. Meth. A* **506** (2003) 250, doi:10.1016/S0168-9002(03)01368-8.
- [59] M. J. Oreglia, “A study of the reactions $\psi' \rightarrow \gamma\gamma\psi$ ”. PhD thesis, Stanford University, 1980. SLAC Report SLAC-R-236.
- [60] Particle Data Group Collaboration, “Review of particle physics”, *PTEP* **2022** (2022) 083C01, doi:10.1093/ptep/ptac097.
- [61] G. Cowan, K. Cranmer, E. Gross, and O. Vitells, “Asymptotic formulae for likelihood-based tests of new physics”, *Eur. Phys. J. C* **71** (2011) 1554, doi:10.1140/epjc/s10052-011-1554-0, arXiv:1007.1727. [Erratum: doi:10.1140/epjc/s10052-013-2501-z].

-
- [62] S. S. Wilks, "The large-sample distribution of the likelihood ratio for testing composite hypotheses", *Annals Math. Statist.* **9** (1938) 60, doi:10.1214/aoms/1177732360.
- [63] CMS Collaboration, "ECAL 2016 refined calibration and Run2 summary plots", CMS Detector Performance Note CMS-DP-2020-021, 2020.
- [64] F. Mosteller and R. A. Fisher, "Questions and answers", *Amer. Statist.* **2** (1948) 30, doi:10.2307/2681650.
- [65] M. Pivk and F. R. Le Diberder, "*s*Plot: A statistical tool to unfold data distributions", *Nucl. Instrum. Meth. A* **555** (2005) 356, doi:10.1016/j.nima.2005.08.106, arXiv:physics/0402083.
- [66] CMS Collaboration, "Precision luminosity measurement in proton-proton collisions at $\sqrt{s} = 13$ TeV in 2015 and 2016 at CMS", *Eur. Phys. J. C* **81** (2021) 800, doi:10.1140/epjc/s10052-021-09538-2, arXiv:2104.01927.
- [67] CMS Collaboration, "CMS luminosity measurement for the 2018 data-taking period at $\sqrt{s} = 13$ TeV", CMS Physics Analysis Summary CMS-PAS-LUM-18-002, 2019.
- [68] H.-S. Shao, "HELAC-Onia: An automatic matrix element generator for heavy quarkonium physics", *Comput. Phys. Commun.* **184** (2013) 2562, doi:10.1016/j.cpc.2013.05.023, arXiv:1212.5293.
- [69] H.-S. Shao, "HELAC-Onia 2.0: an upgraded matrix-element and event generator for heavy quarkonium physics", *Comput. Phys. Commun.* **198** (2016) 238, doi:10.1016/j.cpc.2015.09.011, arXiv:1507.03435.
- [70] S. Dulat et al., "New parton distribution functions from a global analysis of quantum chromodynamics", *Phys. Rev. D* **93** (2016) 033006, doi:10.1103/PhysRevD.93.033006, arXiv:1506.07443.
- [71] K. J. Eskola, P. Paakkinen, H. Paukkunen, and C. A. Salgado, "EPPS16: Nuclear parton distributions with LHC data", *Eur. Phys. J. C* **77** (2017) 163, doi:10.1140/epjc/s10052-017-4725-9, arXiv:1612.05741.
- [72] A. Kusina, J.-P. Lansberg, I. Schienbein, and H.-S. Shao, "Gluon shadowing in heavy-flavor production at the LHC", *Phys. Rev. Lett.* **121** (2018) 052004, doi:10.1103/PhysRevLett.121.052004, arXiv:1712.07024.
- [73] A. Kusina, J.-P. Lansberg, I. Schienbein, and H.-S. Shao, "Reweighted nuclear PDFs using heavy-flavor production data at the LHC", *Phys. Rev. D* **104** (2021) 014010, doi:10.1103/PhysRevD.104.014010, arXiv:2012.11462.
- [74] J.-P. Lansberg and H.-S. Shao, "Towards an automated tool to evaluate the impact of the nuclear modification of the gluon density on quarkonium, D and B meson production in proton-nucleus collisions", *Eur. Phys. J. C* **77** (2017) 1, doi:10.1140/epjc/s10052-016-4575-x, arXiv:1610.05382.
- [75] J.-P. Lansberg and H.-S. Shao, "Production of $J/\psi + \eta_c$ versus $J/\psi + J/\psi$ at the LHC: Importance of real α_s^5 corrections", *Phys. Rev. Lett.* **111** (2013) 122001, doi:10.1103/PhysRevLett.111.122001, arXiv:1308.0474.

- [76] J.-P. Lansberg and H.-S. Shao, "J/ ψ -pair production at large momenta: Indications for double parton scatterings and large α_s^5 contributions", *Phys. Lett. B* **751** (2015) 479, doi:10.1016/j.physletb.2015.10.083, arXiv:1410.8822.
- [77] J.-P. Lansberg, H.-S. Shao, and N. Yamanaka, "Indication for double parton scatterings in W^+ prompt J/ ψ production at the LHC", *Phys. Lett. B* **781** (2018) 485, doi:10.1016/j.physletb.2018.04.020, arXiv:1707.04350.
- [78] ATLAS Collaboration, "Observation and measurements of the production of prompt and non-prompt J/ ψ mesons in association with a Z boson in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector", *Eur. Phys. J. C* **75** (2015) 229, doi:10.1140/epjc/s10052-015-3406-9, arXiv:1412.6428.
- [79] H.-S. Shao and Y.-J. Zhang, "Complete study of hadroproduction of a Y meson associated with a prompt J/ ψ ", *Phys. Rev. Lett.* **117** (2016) 062001, doi:10.1103/PhysRevLett.117.062001, arXiv:1605.03061.
- [80] J.-P. Lansberg and H.-S. Shao, "Phenomenological analysis of associated production of $Z^0 + b$ in the $b \rightarrow J/\psi X$ decay channel at the LHC", *Nucl. Phys. B* **916** (2017) 132, doi:10.1016/j.nuclphysb.2017.01.002, arXiv:1611.09303.
- [81] UA2 Collaboration, "A study of multi-jet events at the CERN $\bar{p}p$ collider and a search for double parton scattering", *Phys. Lett. B* **268** (1991) 145, doi:10.1016/0370-2693(91)90937-L.
- [82] CDF Collaboration, "Study of four jet events and evidence for double parton interactions in $\bar{p}p$ collisions at $\sqrt{s} = 1.8$ TeV", *Phys. Rev. D* **47** (1993) 4857, doi:10.1103/PhysRevD.47.4857.
- [83] CDF Collaboration, "Double parton scattering in $\bar{p}p$ collisions at $\sqrt{s} = 1.8$ TeV", *Phys. Rev. D* **56** (1997) 3811, doi:10.1103/PhysRevD.56.3811.
- [84] D0 Collaboration, "Double parton interactions in $\gamma+3$ jet events in $\bar{p}p$ collisions $\sqrt{s} = 1.96$ TeV", *Phys. Rev. D* **81** (2010) 052012, doi:10.1103/PhysRevD.81.052012, arXiv:0912.5104.
- [85] D0 Collaboration, "Study of double parton interactions in diphoton + dijet events in $\bar{p}p$ collisions at $\sqrt{s} = 1.96$ TeV", *Phys. Rev. D* **93** (2016) 052008, doi:10.1103/PhysRevD.93.052008, arXiv:1512.05291.
- [86] D0 Collaboration, "Double Parton Interactions in $\gamma + 3$ jet and $\gamma + b/c$ jet + 2 jet events in $\bar{p}p$ collisions at $\sqrt{s} = 1.96$ TeV", *Phys. Rev. D* **89** (2014) 072006, doi:10.1103/PhysRevD.89.072006, arXiv:1402.1550.

A The CMS Collaboration

Yerevan Physics Institute, Yerevan, Armenia

A. Hayrapetyan, A. Tumasyan¹ 













Institut für Hochenergiephysik, Vienna, Austria

W. Adam , J.W. Andrejkovic, T. Bergauer , S. Chatterjee , K. Damanakis , M. Dragicevic , P.S. Hussain , M. Jeitler² , N. Krammer , A. Li , D. Liko , I. Mikulec , J. Schieck² , R. Schöfbeck , D. Schwarz , M. Sonawane , S. Templ , W. Waltenberger , C.-E. Wulz² 

















Universiteit Antwerpen, Antwerpen, Belgium

M.R. Darwish³ , T. Janssen , T. Van Laer, P. Van Mechelen 













Vrije Universiteit Brussel, Brussel, Belgium

N. Breugelmans, J. D'Hondt , S. Dansana , A. De Moor , M. Delcourt , F. Heyen, S. Lowette , I. Makarenko , D. Müller , S. Tavernier , M. Tytgat⁴ , G.P. Van Onsem , S. Van Putte , D. Vannerom 















Université Libre de Bruxelles, Bruxelles, Belgium

B. Bilin , B. Clerbaux , A.K. Das, G. De Lentdecker , H. Evard , L. Favart , P. Giannelos , J. Jaramillo , A. Khalilzadeh, F.A. Khan , K. Lee , M. Mahdavihorrani , A. Malara , S. Paredes , M.A. Shahzad, L. Thomas , M. Vanden Bemden , C. Vander Velde , P. Vanlaer 










Ghent University, Ghent, Belgium

M. De Coen , D. Dobur , G. Gokbulut , Y. Hong , J. Knolle , L. Lambrecht , D. Marckx , K. Mota Amarilo , A. Samalan, K. Skovpen , N. Van Den Bossche , J. van der Linden , L. Wezenbeek 


















Université Catholique de Louvain, Louvain-la-Neuve, Belgium

A. Benecke , A. Bethani , G. Bruno , C. Caputo , J. De Favereau De Jeneret , C. Delaere , I.S. Donertas , A. Giammanco , A.O. Guzel , Sa. Jain , V. Lemaitre, J. Lidrych , P. Mastrapasqua , T.T. Tran , S. Wertz 










Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil

G.A. Alves , M. Alves Gallo Pereira , E. Coelho , G. Correia Silva , C. Hensel , T. Menezes De Oliveira , A. Moraes , P. Rebello Teles , M. Soeiro, A. Vilela Pereira⁵ 

Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W.L. Aldá Júnior , M. Barroso Ferreira Filho , H. Brandao Malbouisson , W. Carvalho , J. Chinellato⁶, E.M. Da Costa , G.G. Da Silveira⁷ , D. De Jesus Damiao , S. Fonseca De Souza , R. Gomes De Souza, M. Macedo , J. Martins⁸ , C. Mora Herrera , L. Mundim , H. Nogima , J.P. Pinheiro , A. Santoro , A. Sznajder , M. Thiel 

Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

C.A. Bernardes⁷ , L. Calligaris , T.R. Fernandez Perez Tomei , E.M. Gregores , I. Maietto Silverio , P.G. Mercadante , S.F. Novaes , B. Orzari , Sandra S. Padula 

Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Aleksandrov , G. Antchev , R. Hadjiiska , P. Iaydjiev , M. Misheva , M. Shopova , G. Sultanov 




University of Sofia, Sofia, Bulgaria

A. Dimitrov , L. Litov , B. Pavlov , P. Petkov , A. Petrov , E. Shumka 



Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile

S. Keshri , S. Thakur 













Beihang University, Beijing, China

T. Cheng , T. Javaid , L. Yuan 

Department of Physics, Tsinghua University, Beijing, China

Z. Hu , Z. Liang, J. Liu, K. Yi^{9,10} 

Institute of High Energy Physics, Beijing, China

G.M. Chen¹¹ , H.S. Chen¹¹ , M. Chen¹¹ , F. Iemmi , C.H. Jiang, A. Kapoor¹² , H. Liao , Z.-A. Liu¹³ , R. Sharma¹⁴ , J.N. Song¹³, J. Tao , C. Wang¹¹, J. Wang , Z. Wang¹¹, H. Zhang , J. Zhao 


State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

A. Agapitos , Y. Ban , S. Deng , B. Guo, C. Jiang , A. Levin , C. Li , Q. Li , Y. Mao, S. Qian, S.J. Qian , X. Qin, X. Sun , D. Wang , H. Yang, L. Zhang , Y. Zhao, C. Zhou 

Guangdong Provincial Key Laboratory of Nuclear Science and Guangdong-Hong Kong Joint Laboratory of Quantum Matter, South China Normal University, Guangzhou, China

S. Yang 

Sun Yat-Sen University, Guangzhou, China

Z. You 

University of Science and Technology of China, Hefei, China

K. Jaffel , N. Lu 

Nanjing Normal University, Nanjing, China

G. Bauer¹⁵, B. Li, J. Zhang 

Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) - Fudan University, Shanghai, China

X. Gao¹⁶ 

Zhejiang University, Hangzhou, Zhejiang, China

Z. Lin , C. Lu , M. Xiao 





Universidad de Los Andes, Bogota, Colombia

C. Avila , D.A. Barbosa Trujillo, A. Cabrera , C. Florez , J. Fraga , J.A. Reyes Vega

Universidad de Antioquia, Medellin, Colombia

F. Ramirez , C. Rendón, M. Rodriguez , A.A. Ruales Barbosa , J.D. Ruiz Alvarez 

University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia

D. Giljanovic , N. Godinovic , D. Lelas , A. Sculac 


University of Split, Faculty of Science, Split, Croatia

M. Kovac , A. Petkovic, T. Sculac 




Institute Rudjer Boskovic, Zagreb, Croatia

P. Bargassa , V. Brigljevic , B.K. Chitroda , D. Ferencek , K. Jakovic, S. Mishra , A. Starodumov¹⁷ , T. Susa 

University of Cyprus, Nicosia, Cyprus

A. Attikis , K. Christoforou , A. Hadjiagapiou, C. Leonidou, J. Mousa , C. Nicolaou, L. Paizanos, F. Ptochos , P.A. Razis , H. Rykaczewski, H. Saka , A. Stepennov 



Charles University, Prague, Czech Republic

M. Finger , M. Finger Jr. , A. Kveton 

Universidad San Francisco de Quito, Quito, Ecuador

E. Carrera Jarrin 

Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt

B. El-mahdy, S. Khalil¹⁸ , E. Salama^{19,20} 

Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt

A. Lotfy , M.A. Mahmoud 
















National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

K. Ehataht , M. Kadastik, T. Lange , S. Nandan , C. Nielsen , J. Pata , M. Raidal , L. Tani , C. Veelken 

Department of Physics, University of Helsinki, Helsinki, Finland

H. Kirschenmann , K. Osterberg , M. Voutilainen 



















Helsinki Institute of Physics, Helsinki, Finland

S. Bharthuar , N. Bin Norjoharuddeen , E. Brücken , F. Garcia , P. Inkaew , K.T.S. Kallonen , T. Lampén , K. Lassila-Perini , S. Lehti , T. Lindén , L. Martikainen , M. Myllymäki , M.m. Rantanen , H. Siikonen , J. Tuominiemi 




























Lappeenranta-Lahti University of Technology, Lappeenranta, Finland

P. Luukka , H. Petrow 
















IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France

M. Besancon , F. Couderc , M. Dejardin , D. Denegri, J.L. Faure, F. Ferri , S. Ganjour , P. Gras , G. Hamel de Monchenault , M. Kumar , V. Lohezic , J. Malcles , F. Orlandi , L. Portales , A. Rosowsky , M.Ö. Sahin , A. Savoy-Navarro²¹ , P. Simkina , M. Titov , M. Tornago 

Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

F. Beaudette , G. Boldrini , P. Busson , A. Cappati , C. Charlot , M. Chiusi , F. Damas , O. Davignon , A. De Wit , I.T. Ehle , B.A. Fontana Santos Alves , S. Ghosh , A. Gilbert , R. Granier de Cassagnac , A. Hakimi , B. Harikrishnan , L. Kalipoliti , G. Liu , M. Nguyen , C. Ochando , R. Salerno , J.B. Sauvan , Y. Sirois , L. Urda Gómez , E. Vernazza , A. Zabi , A. Zghiche 






Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France













J.-L. Agram²² , J. Andrea , D. Apparú , D. Bloch , J.-M. Brom , E.C. Chabert , C. Collard , S. Falke , U. Goerlach , R. Haeberle , A.-C. Le Bihan , M. Meena , O. Poncet , G. Saha , M.A. Sessini , P. Van Hove , P. Vaucelle 

Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France



A. Di Florio 

Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France

D. Amram, S. Beauceron , B. Blancon , G. Boudoul , N. Chanon , D. Contardo 

P. Depasse , C. Dozen²³ , H. El Mamouni, J. Fay , S. Gascon , M. Gouzevitch , C. Greenberg, G. Grenier , B. Ille , E. Jourd'huy, I.B. Laktineh, M. Lethuillier , L. Mirabito, S. Perries, A. Purohit , M. Vander Donckt , P. Verdier , J. Xiao 






























Georgian Technical University, Tbilisi, Georgia

D. Lomidze , I. Lomidze , Z. Tsamalaidze¹⁷ 









RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany

V. Botta , S. Consuegra Rodríguez , L. Feld , K. Klein , M. Lipinski , D. Meuser , A. Pauls , D. Pérez Adán , N. Röwert , M. Teroerde 

RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

S. Diekmann , A. Dodonova , N. Eich , D. Eliseev , F. Engelke , J. Erdmann , M. Erdmann , P. Fackeldey , B. Fischer , T. Hebbeker , K. Hoepfner , F. Ivone , A. Jung , M.y. Lee , F. Mausolf , M. Merschmeyer , A. Meyer , S. Mukherjee , D. Noll , F. Nowotny, A. Pozdnyakov , Y. Rath, W. Redjeb , F. Rehm, H. Reithler , V. Sarkisovi , A. Schmidt , A. Sharma , J.L. Spah , A. Stein , F. Torres Da Silva De Araujo²⁴ , S. Wiedenbeck , S. Zaleski







































RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany

C. Dziwok , G. Flügge , T. Kress , A. Nowack , O. Pooth , A. Stahl , T. Ziemons , A. Zotz 
















Deutsches Elektronen-Synchrotron, Hamburg, Germany























H. Aarup Petersen , M. Aldaya Martin , J. Alimena , S. Amoroso, Y. An , J. Bach , S. Baxter , M. Bayatmakou , H. Becerril Gonzalez , O. Behnke , A. Belvedere , S. Bhattacharya , F. Blekman²⁵ , K. Borrás²⁶ , A. Campbell , A. Cardini , C. Cheng, F. Colombina , M. De Silva , G. Eckerlin, D. Eckstein , L.I. Estevez Banos , O. Filatov , E. Gallo²⁵ , A. Geiser , V. Guglielmi , M. Guthoff , A. Hinzmann , L. Jeppe , B. Kaech , M. Kasemann , C. Kleinwort , R. Kogler , M. Komm , D. Krücker , W. Lange, D. Leyva Pernia , K. Lipka²⁷ , W. Lohmann²⁸ , F. Lorkowski , R. Mankel , I.-A. Melzer-Pellmann , M. Mendizabal Morentin , A.B. Meyer , G. Milella , K. Moral Figueroa , A. Mussgiller , L.P. Nair , J. Niedziela , A. Nürnberg , Y. Otariid, J. Park , E. Ranken , A. Raspereza , D. Rastorguev , J. Rübenach, L. Rygaard, A. Saggio , M. Scham^{29,26} , S. Schnake²⁶ , P. Schütze , C. Schwanenberger²⁵ , D. Selivanova , K. Sharko , M. Shchedrolosiev , D. Stafford, F. Vazzoler , A. Ventura Barroso , R. Walsh , D. Wang , Q. Wang , Y. Wen , K. Wichmann, L. Wiens²⁶ , C. Wissing , Y. Yang , A. Zimmermann Castro Santos 

University of Hamburg, Hamburg, Germany

A. Albrecht , S. Albrecht , M. Antonello , S. Bein , L. Benato , S. Bollweg, M. Bonanomi , P. Connor , K. El Morabit , Y. Fischer , E. Garutti , A. Grohsjean , J. Haller , H.R. Jabusch , G. Kasieczka , P. Keicher, R. Klanner , W. Korcari , T. Kramer , C.c. Kuo, V. Kutzner , F. Labe , J. Lange , A. Lobanov , C. Matthies , L. Moureaux , M. Mrowietz, A. Nigamova , Y. Nissan, A. Paasch , K.J. Pena Rodriguez , T. Quadfasel , B. Raciti , M. Rieger , D. Savoio , J. Schindler , P. Schleper , M. Schröder , J. Schwandt , M. Sommerhalder , H. Stadie , G. Steinbrück , A. Tews, M. Wolf 

Karlsruher Institut fuer Technologie, Karlsruhe, Germany






S. Brommer , M. Burkart, E. Butz , T. Chwalek , A. Dierlamm , A. Droll, N. Faltermann , M. Giffels , A. Gottmann , F. Hartmann³⁰ , R. Hofsaess , M. Horzela , U. Husemann , J. Kieseler , M. Klute , R. Koppenhöfer , J.M. Lawhorn , M. Link,

A. Lintuluoto , B. Maier , S. Maier , S. Mitra , M. Mornile , Th. Müller , M. Neukum, M. Oh , E. Pfeffer , M. Presilla , G. Quast , K. Rabbertz , B. Regnery , N. Shadskiy , I. Shvetsov , H.J. Simonis , L. Sowa, L. Stockmeier, K. Tauqeer, M. Toms , N. Trevisani , R.F. Von Cube , M. Wassmer , S. Wieland , F. Wittig, R. Wolf , X. Zuo 

Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

G. Anagnostou, G. Daskalakis , A. Kyriakis, A. Papadopoulos³⁰, A. Stakia 

National and Kapodistrian University of Athens, Athens, Greece

P. Kontaxakis , G. Melachroinos, Z. Painesis , I. Papavergou , I. Paraskevas , N. Saoulidou , K. Theofilatos , E. Tziaferi , K. Vellidis , I. Zisopoulos 






National Technical University of Athens, Athens, Greece

G. Bakas , T. Chatzistavrou, G. Karapostoli , K. Kousouris , I. Papakrivopoulos , E. Siamarkou, G. Tsiopolitis , A. Zacharopoulou

University of Ioánnina, Ioánnina, Greece

K. Adamidis, I. Bestintzanos, I. Evangelou , C. Foudas, C. Kamtsikis, P. Katsoulis, P. Kokkas , P.G. Kosmoglou Kioseoglou , N. Manthos , I. Papadopoulos , J. Strologas 

HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

C. Hajdu , D. Horvath^{31,32} , K. Márton, A.J. Rádl³³ , F. Sikler , V. Veszpremi 

MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

M. Csanád , K. Farkas , A. Fehérkuti³⁴ , M.M.A. Gadallah³⁵ , Á. Kadlecik , P. Major , G. Pásztor , G.I. Veres 

Faculty of Informatics, University of Debrecen, Debrecen, Hungary

B. Ujvari , G. Zilizi 















Institute of Nuclear Research ATOMKI, Debrecen, Hungary

G. Bencze, S. Czellar, J. Molnar, Z. Szillasi

Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary

F. Nemes³⁴ , T. Novak 

Panjab University, Chandigarh, India

J. Babbar , S. Bansal , S.B. Beri, V. Bhatnagar , G. Chaudhary , S. Chauhan , N. Dhingra³⁶ , A. Kaur , A. Kaur , H. Kaur , M. Kaur , S. Kumar , K. Sandeep , T. Sheokand, J.B. Singh , A. Singla 
















University of Delhi, Delhi, India

A. Ahmed , A. Bhardwaj , A. Chhetri , B.C. Choudhary , A. Kumar , A. Kumar , M. Naimuddin , K. Ranjan , M.K. Saini, S. Saumya 

Saha Institute of Nuclear Physics, HBNI, Kolkata, India

S. Baradia , S. Barman³⁷ , S. Bhattacharya , S. Das Gupta, S. Dutta , S. Dutta, S. Sarkar










Indian Institute of Technology Madras, Madras, India

M.M. Ameen , P.K. Behera , S.C. Behera , S. Chatterjee , G. Dash , P. Jana , P. Kalbhor , S. Kamble , J.R. Komaragiri³⁸ , D. Kumar³⁸ , P.R. Pujahari , N.R. Saha , A. Sharma , A.K. Sikdar , R.K. Singh, P. Verma, S. Verma , A. Vijay












Tata Institute of Fundamental Research-A, Mumbai, India

S. Dugad, G.B. Mohanty , B. Parida , M. Shelake, P. Suryadevara

Tata Institute of Fundamental Research-B, Mumbai, India

A. Bala , S. Banerjee , R.M. Chatterjee, M. Guchait , Sh. Jain , A. Jaiswal, S. Kumar , G. Majumder , K. Mazumdar , S. Parolia , A. Thachayath 

National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India

S. Bahinipati³⁹ , C. Kar , D. Maity⁴⁰ , P. Mal , T. Mishra , V.K. Muraleedharan Nair Bindhu⁴⁰ , K. Naskar⁴⁰ , A. Nayak⁴⁰ , S. Nayak, K. Pal, P. Sadangi, S.K. Swain , S. Varghese⁴⁰ , D. Vats⁴⁰ 

Indian Institute of Science Education and Research (IISER), Pune, India

S. Acharya⁴¹ , A. Alpana , S. Dube , B. Gomber⁴¹ , P. Hazarika , B. Kansal , A. Laha , B. Sahu⁴¹ , S. Sharma , K.Y. Vaish 

Isfahan University of Technology, Isfahan, Iran

H. Bakhshiansohi⁴² , A. Jafari⁴³ , M. Zeinali⁴⁴ 































Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

S. Bashiri, S. Chenarani⁴⁵ , S.M. Etesami , Y. Hosseini , M. Khakzad , E. Khazaie⁴⁶ , M. Mohammadi Najafabadi , S. Tizchang⁴⁷ 






























University College Dublin, Dublin, Ireland

M. Felcini , M. Grunewald 






INFN Sezione di Bari^a, Università di Bari^b, Politecnico di Bari^c, Bari, Italy

M. Abbrescia^{a,b} , A. Colaleo^{a,b} , D. Creanza^{a,c} , B. D'Anzi^{a,b} , N. De Filippis^{a,c} , M. De Palma^{a,b} , W. Elmetenawee^{a,b,48} , L. Fiore^a , G. Iaselli^{a,c} , L. Longo^a , M. Louka^{a,b}, G. Maggi^{a,c} , M. Maggi^a , I. Margjeka^a , V. Mastrapasqua^{a,b} , S. My^{a,b} , S. Nuzzo^{a,b} , A. Pellecchia^{a,b} , A. Pompili^{a,b} , G. Pugliese^{a,c} , R. Radogna^{a,b} , D. Ramos^a , A. Ranieri^a , L. Silvestris^a , F.M. Simone^{a,c} , Ü. Sözbilir^a , A. Stamerra^{a,b} , D. Troiano^{a,b} , R. Venditti^{a,b} , P. Verwilligen^a , A. Zaza^{a,b} 


















INFN Sezione di Bologna^a, Università di Bologna^b, Bologna, Italy

G. Abbiendi^a , C. Battilana^{a,b} , D. Bonacorsi^{a,b} , P. Capiluppi^{a,b} , A. Castro^{†a,b} , F.R. Cavallo^a , M. Cuffiani^{a,b} , G.M. Dallavalle^a , T. Diotallevi^{a,b} , F. Fabbri^a , A. Fanfani^{a,b} , D. Fasanella^a , P. Giacomelli^a , L. Giommi^{a,b} , C. Grandi^a , L. Guiducci^{a,b} , S. Lo Meo^{a,49} , M. Lorusso^{a,b} , L. Lunerti^a , S. Marcellini^a , G. Masetti^a , F.L. Navarria^{a,b} , G. Paggi^{a,b} , A. Perrotta^a , F. Primavera^{a,b} , A.M. Rossi^{a,b} , S. Rossi Tisbeni^{a,b} , T. Rovelli^{a,b} , G.P. Siroli^{a,b} 

INFN Sezione di Catania^a, Università di Catania^b, Catania, Italy





















































S. Costa^{a,b,50} , A. Di Mattia^a , A. Lapertosa^a , R. Potenza^{a,b}, A. Tricomi^{a,b,50} , C. Tuve^{a,b} 

























INFN Sezione di Firenze^a, Università di Firenze^b, Firenze, Italy

P. Assiouras^a , G. Barbagli^a , G. Bardelli^{a,b} , B. Camaiani^{a,b} , A. Cassese^a , R. Ceccarelli^a , V. Ciulli^{a,b} , C. Civinini^a , R. D'Alessandro^{a,b} , E. Focardi^{a,b} , T. Kello^a, G. Latino^{a,b} , P. Lenzi^{a,b} , M. Lizzo^a , M. Meschini^a , S. Paoletti^a , A. Papanastassiou^{a,b}, G. Sguazzoni^a , L. Viliani^a 







INFN Laboratori Nazionali di Frascati, Frascati, Italy

L. Benussi , S. Bianco , S. Meola⁵¹ , D. Piccolo 

INFN Sezione di Genova^a, Università di Genova^b, Genova, ItalyP. Chatagnon^a , F. Ferro^a , E. Robutti^a , S. Tosi^{a,b} **INFN Sezione di Milano-Bicocca^a, Università di Milano-Bicocca^b, Milano, Italy**A. Benaglia^a , F. Brivio^a , F. Cetorelli^{a,b} , F. De Guio^{a,b} , M.E. Dinardo^{a,b} , P. Dini^a , S. Gennai^a , R. Gerosa^{a,b} , A. Ghezzi^{a,b} , P. Govoni^{a,b} , L. Guzzi^a , M.T. Lucchini^{a,b} , M. Malberti^a , S. Malvezzi^a , A. Massironi^a , D. Menasce^a , L. Moroni^a , M. Paganoni^{a,b} , S. Palluotto^{a,b} , D. Pedrini^a , A. Perego^{a,b} , B.S. Pinolini^a, G. Pizzati^{a,b}, S. Ragazzi^{a,b} , T. Tabarelli de Fatis^{a,b} **INFN Sezione di Napoli^a, Università di Napoli 'Federico II'^b, Napoli, Italy; Università della Basilicata^c, Potenza, Italy; Scuola Superiore Meridionale (SSM)^d, Napoli, Italy**S. Buontempo^a , A. Cagnotta^{a,b} , F. Carnevali^{a,b}, N. Cavallo^{a,c} , F. Fabozzi^{a,c} , A.O.M. Iorio^{a,b} , L. Lista^{a,b,52} , P. Paolucci^{a,30} , B. Rossi^a **INFN Sezione di Padova^a, Università di Padova^b, Padova, Italy; Università di Trento^c, Trento, Italy**R. Ardino^a , P. Azzi^a , N. Bacchetta^{a,53} , D. Bisello^{a,b} , P. Bortignon^a , G. Bortolato^{a,b}, A. Bragagnolo^{a,b} , A.C.M. Bulla^a , R. Carlin^{a,b} , P. Checchia^a , T. Dorigo^a , F. Gasparini^{a,b} , U. Gasparini^{a,b} , M. Gulmini^{a,54} , E. Lusiani^a , M. Margoni^{a,b} , A.T. Meneguzzo^{a,b} , M. Migliorini^{a,b} , J. Pazzini^{a,b} , P. Ronchese^{a,b} , R. Rossin^{a,b} , F. Simonetto^{a,b} , M. Tosi^{a,b} , A. Triossi^{a,b} , S. Ventura^a , M. Zanetti^{a,b} , P. Zotto^{a,b} , A. Zucchetta^{a,b} **INFN Sezione di Pavia^a, Università di Pavia^b, Pavia, Italy**C. Aimè^a , A. Braghieri^a , S. Calzaferri^a , D. Fiorina^a , P. Montagna^{a,b} , V. Re^a , C. Riccardi^{a,b} , P. Salvini^a , I. Vai^{a,b} , P. Vitulo^{a,b} **INFN Sezione di Perugia^a, Università di Perugia^b, Perugia, Italy**S. Ajmal^{a,b} , M.E. Ascioti^{a,b}, G.M. Bilei^a , C. Carrivale^{a,b}, D. Ciangottini^{a,b} , L. Fanò^{a,b} , M. Magherini^{a,b} , V. Mariani^{a,b} , M. Menichelli^a , F. Moscatelli^{a,55} , A. Rossi^{a,b} , A. Santocchia^{a,b} , D. Spiga^a , T. Tedeschi^{a,b} **INFN Sezione di Pisa^a, Università di Pisa^b, Scuola Normale Superiore di Pisa^c, Pisa, Italy; Università di Siena^d, Siena, Italy**C.A. Alexe^{a,c} , P. Asenov^{a,b} , P. Azzurri^a , G. Bagliesi^a , R. Bhattacharya^a , L. Bianchini^{a,b} , T. Boccali^a , E. Bossini^a , D. Bruschini^{a,c} , R. Castaldi^a , M.A. Ciocci^{a,b} , M. Cipriani^{a,b} , V. D'Amante^{a,d} , R. Dell'Orso^a , S. Donato^a , A. Giassi^a , F. Ligabue^{a,c} , A.C. Marini^a , D. Matos Figueiredo^a , A. Messineo^{a,b} , M. Musich^{a,b} , F. Palla^a , A. Rizzi^{a,b} , G. Rolandi^{a,c} , S. Roy Chowdhury^a , T. Sarkar^a , A. Scribano^a , P. Spagnolo^a , R. Tenchini^a , G. Tonelli^{a,b} , N. Turini^{a,d} , F. Vaselli^{a,c} , A. Venturi^a , P.G. Verdini^a **INFN Sezione di Roma^a, Sapienza Università di Roma^b, Roma, Italy**C. Baldenegro Barrera^{a,b} , P. Barria^a , C. Basile^{a,b} , M. Campana^{a,b} , F. Cavallari^a , L. Cunqueiro Mendez^{a,b} , D. Del Re^{a,b} , E. Di Marco^{a,b} , M. Diemoz^a , F. Errico^{a,b} , E. Longo^{a,b} , J. Mijuskovic^{a,b} , G. Organtini^{a,b} , F. Pandolfi^a , R. Paramatti^{a,b} , C. Quaranta^{a,b} , S. Rahatlou^{a,b} , C. Rovelli^a , F. Santanastasio^{a,b} , L. Soffi^a **INFN Sezione di Torino^a, Università di Torino^b, Torino, Italy; Università del Piemonte Orientale^c, Novara, Italy**N. Amapane^{a,b} , R. Arcidiacono^{a,c} , S. Argiro^{a,b} , M. Arneodo^{a,c} , N. Bartosik^a , R. Bellan^{a,b} , A. Bellora^{a,b} , C. Biino^a , C. Borca^{a,b} , N. Cartiglia^a , M. Costa^{a,b}

R. Covarelli^{a,b} , N. Demaria^a , L. Finco^a , M. Grippo^{a,b} , B. Kiani^{a,b} , F. Legger^a , F. Luongo^{a,b} , C. Mariotti^a , L. Markovic^{a,b} , S. Maselli^a , A. Mecca^{a,b} , L. Menzio^{a,b}, P. Meridiani^a , E. Migliore^{a,b} , M. Monteno^a , R. Mulargia^a , M.M. Obertino^{a,b} , G. Ortona^a , L. Pacher^{a,b} , N. Pastrone^a , M. Pelliccioni^a , M. Ruspa^{a,c} , F. Siviero^{a,b} , V. Sola^{a,b} , A. Solano^{a,b} , A. Staiano^a , C. Tarricone^{a,b} , D. Trocino^a , G. Umoret^{a,b} , R. White^{a,b}

INFN Sezione di Trieste^a, Università di Trieste^b, Trieste, Italy

S. Belforte^a , V. Candelise^{a,b} , M. Casarsa^a , F. Cossutti^a , K. De Leo^a , G. Della Ricca^{a,b} 


Kyungpook National University, Daegu, Korea

S. Dogra , J. Hong , C. Huh , B. Kim , J. Kim, D. Lee, H. Lee, S.W. Lee , C.S. Moon , Y.D. Oh , M.S. Ryu , S. Sekmen , B. Tae, Y.C. Yang 

Department of Mathematics and Physics - GWNNU, Gangneung, Korea

M.S. Kim 

Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

G. Bak , P. Gwak , H. Kim , D.H. Moon 

Hanyang University, Seoul, Korea

E. Asilar , J. Choi , D. Kim , T.J. Kim , J.A. Merlin, Y. Ryou

Korea University, Seoul, Korea

S. Choi , S. Han, B. Hong , K. Lee, K.S. Lee , S. Lee , J. Yoo 

Kyung Hee University, Department of Physics, Seoul, Korea

J. Goh , S. Yang 








Sejong University, Seoul, Korea

H. S. Kim , Y. Kim, S. Lee



Seoul National University, Seoul, Korea

J. Almond, J.H. Bhyun, J. Choi , J. Choi, W. Jun , J. Kim , S. Ko , H. Kwon , H. Lee , J. Lee , J. Lee , B.H. Oh , S.B. Oh , H. Seo , U.K. Yang, I. Yoon 

University of Seoul, Seoul, Korea

W. Jang , D.Y. Kang, Y. Kang , S. Kim , B. Ko, J.S.H. Lee , Y. Lee , I.C. Park , Y. Roh, I.J. Watson 

Yonsei University, Department of Physics, Seoul, Korea

S. Ha , H.D. Yoo 


Sungkyunkwan University, Suwon, Korea

M. Choi , M.R. Kim , H. Lee, Y. Lee , I. Yu 

College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait

T. Beyrouthy, Y. Gharbia


Kuwait University - College of Science - Department of Physics, Safat, Kuwait

F. Alazemi 

Riga Technical University, Riga, Latvia

K. Dreimanis , A. Gaile , G. Pikurs, A. Potrebko , M. Seidel , D. Sidiropoulos Kontos

University of Latvia (LU), Riga, Latvia

N.R. Strautnieks 







Vilnius University, Vilnius, Lithuania

M. Ambrozas , A. Juodagalvis , A. Rinkevicius , G. Tamulaitis 








National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia

I. Yusuff⁵⁶ , Z. Zolkapli



Universidad de Sonora (UNISON), Hermosillo, Mexico

J.F. Benitez , A. Castaneda Hernandez , H.A. Encinas Acosta, L.G. Gallegos Maríñez, M. León Coello , J.A. Murillo Quijada , A. Sehrawat , L. Valencia Palomo 

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

G. Ayala , H. Castilla-Valdez , H. Crotte Ledesma, E. De La Cruz-Burelo , I. Heredia-De La Cruz⁵⁷ , R. Lopez-Fernandez , J. Mejia Guisao , C.A. Mondragon Herrera, A. Sánchez Hernández 

Universidad Iberoamericana, Mexico City, Mexico

C. Oropeza Barrera , D.L. Ramirez Guadarrama, M. Ramírez García 

Benemerita Universidad Autonoma de Puebla, Puebla, Mexico

I. Bautista , I. Pedraza , H.A. Salazar Ibarquen , C. Uribe Estrada 

University of Montenegro, Podgorica, Montenegro

I. Bubanja , N. Raicevic 

University of Canterbury, Christchurch, New Zealand

P.H. Butler 

National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan

A. Ahmad , M.I. Asghar, A. Awais , M.I.M. Awan, H.R. Hoorani , W.A. Khan 







AGH University of Krakow, Faculty of Computer Science, Electronics and Telecommunications, Krakow, Poland

V. Avati, L. Grzanka , M. Malawski 

National Centre for Nuclear Research, Swierk, Poland

H. Bialkowska , M. Bluj , M. Górski , M. Kazana , M. Szleper , P. Zalewski 

















Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland

K. Bunkowski , K. Doroba , A. Kalinowski , M. Konecki , J. Krolikowski , A. Muhammad 



Warsaw University of Technology, Warsaw, Poland

K. Pozniak , W. Zabolotny 

Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal

M. Araujo , D. Bastos , C. Beirão Da Cruz E Silva , A. Boletti , M. Bozzo , T. Camporesi , G. Da Molin , P. Faccioli , M. Gallinaro , J. Hollar , N. Leonardo , G.B. Marozzo, T. Niknejad , A. Petrilli , M. Pisano , J. Seixas , J. Varela , J.W. Wulff





























Faculty of Physics, University of Belgrade, Belgrade, Serbia

P. Adzic , P. Milenovic 

VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia

M. Dordevic , J. Milosevic , L. Nadderd , V. Rekovic















Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain

J. Alcaraz Maestre , Cristina F. Bedoya , Oliver M. Carretero , M. Cepeda , M. Cerrada , N. Colino , B. De La Cruz , A. Delgado Peris , A. Escalante Del Valle , D. Fernández Del Val , J.P. Fernández Ramos , J. Flix , M.C. Fouz , O. Gonzalez Lopez , S. Goy Lopez , J.M. Hernandez , M.I. Josa , E. Martin Viscasillas , D. Moran , C. M. Morcillo Perez , Á. Navarro Tobar , C. Perez Dengra , A. Pérez-Calero Yzquierdo , J. Puerta Pelayo , I. Redondo , S. Sánchez Navas , J. Sastre , J. Vazquez Escobar 




















Universidad Autónoma de Madrid, Madrid, Spain

J.F. de Trocóniz 



Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain

B. Alvarez Gonzalez , J. Cuevas , J. Fernandez Menendez , S. Folgueras , I. Gonzalez Caballero , J.R. González Fernández , P. Leguina , E. Palencia Cortezon , J. Prado Pico, C. Ramón Álvarez , V. Rodríguez Bouza , A. Soto Rodríguez , A. Trapote , C. Vico Villalba , P. Vischia 

Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

S. Bhowmik , S. Blanco Fernández , J.A. Brochero Cifuentes , I.J. Cabrillo , A. Calderon , J. Duarte Campderros , M. Fernandez , G. Gomez , C. Lasasa García , R. Lopez Ruiz , C. Martinez Rivero , P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas , E. Navarrete Ramos , J. Piedra Gomez , L. Scodellaro , I. Vila , J.M. Vizan Garcia 

University of Colombo, Colombo, Sri Lanka

B. Kailasapathy⁵⁸ , D.D.C. Wickramarathna 













University of Ruhuna, Department of Physics, Matara, Sri Lanka

W.G.D. Dharmaratna⁵⁹ , K. Liyanage , N. Perera 






































CERN, European Organization for Nuclear Research, Geneva, Switzerland

D. Abbaneo , C. Amendola , E. Auffray , G. Auzinger , J. Baechler, D. Barney , A. Bermúdez Martínez , M. Bianco , A.A. Bin Anuar , A. Bocci , L. Boronovi , C. Botta , E. Brondolin , C. Caillol , G. Cerminara , N. Chernyavskaya , D. d'Enterria , A. Dabrowski , A. David , A. De Roeck , M.M. Defranchis , M. Deile , M. Dobson , G. Franzoni , W. Funk , S. Giani, D. Gigi, K. Gill , F. Glege , J. Hegeman , J.K. Heikkilä , B. Huber, V. Innocente , T. James , P. Janot , O. Kaluzinska , O. Karacheban²⁸ , S. Laurila , P. Lecoq , E. Leutgeb , C. Lourenço , L. Malgeri , M. Mannelli , M. Matthewman, A. Mehta , F. Meijers , S. Mersi , E. Meschi , V. Milosevic , F. Monti , F. Moortgat , M. Mulders , I. Neutelings , S. Orfanelli, F. Pantaleo , G. Petrucciani , A. Pfeiffer , M. Pierini , H. Qu , D. Rabadý , B. Ribeiro Lopes , M. Rovere , H. Sakulin , S. Sanchez Cruz , S. Scarfi , C. Schwick, M. Selvaggi , A. Sharma , K. Shchelina , P. Silva , P. Sphicas⁶⁰ , A.G. Stahl Leitner , A. Steen , S. Summers , D. Treille , P. Tropea , D. Walter , J. Wanczyk⁶¹ , J. Wang, S. Wuchterl , P. Zehetner , P. Zejdl , W.D. Zeuner




















Paul Scherrer Institut, Villigen, Switzerland

T. Bevilacqua⁶² , L. Caminada⁶² , A. Ebrahimi , W. Erdmann , R. Horisberger , Q. Ingram , H.C. Kaestli , D. Kotlinski , C. Lange , M. Missiroli⁶² , L. Noehte⁶² , T. Rohe 




ETH Zurich - Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

T.K. Aarrestad , K. Androsov⁶¹ , M. Backhaus , G. Bonomelli, A. Calandri , C. Cazaniga , K. Datta , P. De Bryas Dexmiers D'archiac⁶¹ , A. De Cosa , G. Dissertori , M. Dittmar, M. Donegà , F. Eble , M. Galli , K. Gedia , F. Glessgen , C. Grab , N. Härringer , T.G. Harte, D. Hits , W. Luster mann , A.-M. Lyon , R.A. Manzoni , M. Marchegiani , L. Marchese , C. Martin Perez , A. Mascellani⁶¹ , F. Nessi-Tedaldi , F. Pauss , V. Perovic , S. Pigazzini , C. Reissel , T. Reitenspiess , B. Ristic , F. Riti , R. Seidita , J. Steggemann⁶¹ , A. Tarabini , D. Valsecchi , R. Wallny 












Universität Zürich, Zurich, Switzerland

C. Amsler⁶³ , P. Bärtschi , M.F. Canelli , K. Cormier , M. Huwiler , W. Jin , A. Jofrehei , B. Kilminster , S. Leontsinis , S.P. Liehti , A. Macchiolo , P. Meiring , F. Meng , U. Molinatti , J. Motta , A. Reimers , P. Robmann, M. Senger , E. Shokr, F. Stäger , R. Tramontano 




National Central University, Chung-Li, Taiwan

C. Adloff⁶⁴, D. Bhowmik, C.M. Kuo, W. Lin, P.K. Rout , P.C. Tiwari³⁸ , S.S. Yu 



















National Taiwan University (NTU), Taipei, Taiwan

L. Ceard, K.F. Chen , P.s. Chen, Z.g. Chen, A. De Iorio , W.-S. Hou , T.h. Hsu, Y.w. Kao, S. Karmakar , G. Kole , Y.y. Li , R.-S. Lu , E. Paganis , X.f. Su , J. Thomas-Wilsker , L.s. Tsai, H.y. Wu, E. Yazgan 


High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

C. Asawatangtrakuldee , N. Srimanobhas , V. Wachirapusanand 

Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

D. Agyel , F. Boran , F. Dolek , I. Dumanoglu⁶⁵ , E. Eskut , Y. Guler⁶⁶ , E. Gurpınar Guler⁶⁶ , C. Isik , O. Kara, A. Kayis Topaksu , U. Kiminsu , G. Onengut , K. Ozdemir⁶⁷ , A. Polatoz , B. Tali⁶⁸ , U.G. Tok , S. Turkcapar , E. Uslan , I.S. Zorbakir 






Middle East Technical University, Physics Department, Ankara, Turkey

G. Sokmen, M. Yalvac⁶⁹ 








Bogazici University, Istanbul, Turkey

B. Akgun , I.O. Atakisi , E. Gülmez , M. Kaya⁷⁰ , O. Kaya⁷¹ , S. Tekten⁷² 





Istanbul Technical University, Istanbul, Turkey

A. Cakir , K. Cankocak^{65,73} , G.G. Dincer⁶⁵ , Y. Komurcu , S. Sen⁷⁴ 

Istanbul University, Istanbul, Turkey

O. Aydilek⁷⁵ , B. Hacisahinoglu , I. Hos⁷⁶ , B. Kaynak , S. Ozkorucuklu , O. Potok , H. Sert , C. Simsek , C. Zorbilmez 


Yildiz Technical University, Istanbul, Turkey

S. Cerci⁶⁸ , B. Isildak⁷⁷ , D. Sunar Cerci , T. Yetkin 















Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine

A. Boyaryntsev , B. Grynyov 




















National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine

L. Levchuk 




































University of Bristol, Bristol, United Kingdom

D. Anthony , J.J. Brooke , A. Bundock , F. Bury , E. Clement , D. Cussans , H. Flacher , M. Glowacki, J. Goldstein , H.F. Heath , M.-L. Holmberg , L. Kreczko , S. Paramesvaran , L. Robertshaw, S. Seif El Nasr-Storey, V.J. Smith , N. Stylianou⁷⁸ , K. Walkingshaw Pass




Rutherford Appleton Laboratory, Didcot, United Kingdom

A.H. Ball, K.W. Bell , A. Belyaev⁷⁹ , C. Brew , R.M. Brown , D.J.A. Cockerill , C. Cooke , A. Elliot , K.V. Ellis, K. Harder , S. Harper , J. Linacre , K. Manolopoulos, D.M. Newbold , E. Olaiya, D. Petyt , T. Reis , A.R. Sahasransu , G. Salvi , T. Schuh, C.H. Shepherd-Themistocleous , I.R. Tomalin , K.C. Whalen , T. Williams 

Imperial College, London, United Kingdom

I. Andreou , R. Bainbridge , P. Bloch , C.E. Brown , O. Buchmuller, V. Cacchio, C.A. Carrillo Montoya , G.S. Chahal⁸⁰ , D. Colling , J.S. Dancu, I. Das , P. Dauncey , G. Davies , J. Davies, M. Della Negra , S. Fayer, G. Fedi , G. Hall , M.H. Hassanshahi , A. Howard, G. Iles , M. Knight , J. Langford , J. León Holgado , L. Lyons , A.-M. Magnan , S. Mallios, M. Mieskolainen , J. Nash⁸¹ , M. Pesaresi , P.B. Pradeep, B.C. Radburn-Smith , A. Richards, A. Rose , K. Savva , C. Seez , R. Shukla , A. Tapper , K. Uchida , G.P. Uttley , L.H. Vage, T. Virdee³⁰ , M. Vojinovic , N. Wardle , D. Winterbottom 






Brunel University, Uxbridge, United Kingdom

K. Coldham, J.E. Cole , A. Khan, P. Kyberd , I.D. Reid 

Baylor University, Waco, Texas, USA

S. Abdullin , A. Brinkerhoff , E. Collins , J. Dittmann , K. Hatakeyama , J. Hiltbrand , B. McMaster , J. Samudio , S. Sawant , C. Sutantawibul , J. Wilson 



















Catholic University of America, Washington, DC, USA

R. Bartek , A. Dominguez , C. Huerta Escamilla, A.E. Simsek , R. Uniyal , A.M. Vargas Hernandez 



















The University of Alabama, Tuscaloosa, Alabama, USA

B. Bam , A. Buchot Perraguin , R. Chudasama , S.I. Cooper , C. Crovella , S.V. Gleyzer , E. Pearson, C.U. Perez , P. Rumerio⁸² , E. Usai , R. Yi 




















Boston University, Boston, Massachusetts, USA

A. Akpinar , C. Cosby , G. De Castro, Z. Demiragli , C. Erice , C. Fangmeier , C. Fernandez Madrazo , E. Fontanesi , D. Gastler , F. Golf , S. Jeon , J. O'cain, I. Reed , J. Rohlf , K. Salyer , D. Sperka , D. Spitzbart , I. Suarez , A. Tsatsos , A.G. Zecchinelli 




Brown University, Providence, Rhode Island, USA

G. Benelli , D. Cutts , L. Gouskos , M. Hadley , U. Heintz , J.M. Hogan⁸³ , T. Kwon , G. Landsberg , K.T. Lau , D. Li , J. Luo , S. Mondal , M. Narain[†] , N. Pervan , T. Russell, S. Sagir⁸⁴ , F. Simpson , M. Stamenkovic , N. Venkatasubramanian, X. Yan 

University of California, Davis, Davis, California, USA

S. Abbott , C. Brainerd , R. Breedon , H. Cai , M. Calderon De La Barca Sanchez , M. Chertok , M. Citron , J. Conway , P.T. Cox , R. Erbacher , F. Jensen , O. Kukral , G. Mocellin , M. Mulhearn , S. Ostrom , W. Wei , Y. Yao , S. Yoo , F. Zhang 

























University of California, Los Angeles, California, USA

M. Bachtis , R. Cousins , A. Datta , G. Flores Avila , J. Hauser , M. Ignatenko , M.A. Iqbal , T. Lam , E. Manca , A. Nunez Del Prado, D. Saltzberg , V. Valuev 

University of California, Riverside, Riverside, California, USA

R. Clare , J.W. Gary , M. Gordon, G. Hanson , W. Si 

University of California, San Diego, La Jolla, California, USA

A. Aportela, A. Arora , J.G. Branson , S. Cittolin , S. Cooperstein , D. Diaz , J. Duarte , L. Giannini , Y. Gu, J. Guiang , R. Kansal , V. Krutelyov , R. Lee , J. Letts , M. Masciovecchio , F. Mokhtar , S. Mukherjee , M. Pieri , M. Quinnan , B.V. Sathia Narayanan , V. Sharma , M. Tadel , E. Vourliotis , F. Würthwein , Y. Xiang , A. Yagil 
















University of California, Santa Barbara - Department of Physics, Santa Barbara, California, USA

A. Barzdukas , L. Brennan , C. Campagnari , K. Downham , C. Grieco , J. Incandela , J. Kim , A.J. Li , P. Masterson , H. Mei , J. Richman , S.N. Santpur , U. Sarica , R. Schmitz , F. Setti , J. Sheplock , D. Stuart , T.Á. Vámi , S. Wang , D. Zhang















California Institute of Technology, Pasadena, California, USA

A. Bornheim , O. Cerri, A. Latorre, J. Mao , H.B. Newman , G. Reales Gutiérrez, M. Spiropulu , J.R. Vlimant , C. Wang , S. Xie , R.Y. Zhu 
















Carnegie Mellon University, Pittsburgh, Pennsylvania, USA

J. Alison , S. An , P. Bryant , M. Cremonesi, V. Dutta , T. Ferguson , T.A. Gómez Espinosa , A. Harilal , A. Kallil Tharayil, C. Liu , T. Mudholkar , S. Murthy , P. Palit , K. Park, M. Paulini , A. Roberts , A. Sanchez , W. Terrill 


































































University of Colorado Boulder, Boulder, Colorado, USA

J.P. Cumalad , W.T. Ford , A. Hart , A. Hassani , G. Karathanasis , N. Manganelli , A. Perloff , C. Savard , N. Schonbeck , K. Stenson , K.A. Ulmer , S.R. Wagner , N. Zipper , D. Zuolo 



















Cornell University, Ithaca, New York, USA

J. Alexander , S. Bright-Thonney , X. Chen , D.J. Cranshaw , J. Fan , X. Fan , S. Hogan , P. Kotamnives, J. Monroy , M. Oshiro , J.R. Patterson , M. Reid , A. Ryd , J. Thom , P. Wittich , R. Zou 














Fermi National Accelerator Laboratory, Batavia, Illinois, USA

M. Albrow , M. Alyari , O. Amram , G. Apollinari , A. Apresyan , L.A.T. Bauerdick , D. Berry , J. Berryhill , P.C. Bhat , K. Burkett , J.N. Butler , A. Canepa , G.B. Cerati , H.W.K. Cheung , F. Chlebana , G. Cummings , J. Dickinson , I. Dutta , V.D. Elvira , Y. Feng , J. Freeman , A. Gandrakota , Z. Gecse , L. Gray , D. Green, A. Grummer , S. Grünendahl , D. Guerrero , O. Gutsche , R.M. Harris , R. Heller , T.C. Herwig , J. Hirschauer , B. Jayatilaka , S. Jindariani , M. Johnson , U. Joshi , T. Klijnsma , B. Klima , K.H.M. Kwok , S. Lammel , D. Lincoln , R. Lipton , T. Liu , C. Madrid , K. Maeshima , C. Mantilla , D. Mason , P. McBride , P. Merkel , S. Mrenna , S. Nahn , J. Ngadiuba , D. Noonan , S. Norberg, V. Papadimitriou , N. Pastika , K. Pedro , C. Pena⁸⁵ , F. Ravera , A. Reinsvold Hall⁸⁶ , L. Ristori , M. Safdari , E. Sexton-Kennedy , N. Smith , A. Soha , L. Spiegel , S. Stoynev , J. Strait , L. Taylor , S. Tkaczyk , N.V. Tran , L. Uplegger , E.W. Vaandering , I. Zoi 






University of Florida, Gainesville, Florida, USA

C. Aruta , P. Avery , D. Bourilkov , P. Chang , V. Cherepanov , R.D. Field, E. Koenig , M. Kolosova , J. Konigsberg , A. Korytov , K. Matchev , N. Menendez , G. Mitselmakher , K. Mohrman , A. Muthirakalayil Madhu , N. Rawal , S. Rosenzweig , Y. Takahashi , J. Wang 
























Florida State University, Tallahassee, Florida, USA

T. Adams , A. Al Kadhim , A. Askew , S. Bower , R. Habibullah , V. Hagopian , R. Hashmi , R.S. Kim , S. Kim , T. Kolberg , G. Martinez, H. Prosper , P.R. Prova, M. Wulansatiti , R. Yohay , J. Zhang











Florida Institute of Technology, Melbourne, Florida, USA

B. Alsufyani, M.M. Baarmand , S. Butalla , S. Das , T. Elkafrawy²⁰ , M. Hohlmann , M. Rahmani, E. Yanes












University of Illinois Chicago, Chicago, USA, Chicago, USA

M.R. Adams , A. Baty , C. Bennett, R. Cavanaugh , R. Escobar Franco , O. Evdokimov , C.E. Gerber , M. Hawksworth, A. Hingrajiya, D.J. Hofman , J.h. Lee , D. S. Lemos , A.H. Merrit , C. Mills , S. Nanda , G. Oh , B. Ozek , D. Pilipovic , R. Pradhan , E. Prifti, T. Roy , S. Rudrabhatla , M.B. Tonjes , N. Varelas , M.A. Wadud , Z. Ye , J. Yoo 
























The University of Iowa, Iowa City, Iowa, USA

M. Alhousseini , D. Blend, K. Dilsiz⁸⁷ , L. Emediato , G. Karaman , O.K. Köseyan , J.-P. Merlo, A. Mestvirishvili⁸⁸ , O. Neogi, H. Ogul⁸⁹ , Y. Onel , A. Penzo , C. Snyder, E. Tiras⁹⁰ 









Johns Hopkins University, Baltimore, Maryland, USA

B. Blumenfeld , L. Corcodilos , J. Davis , A.V. Gritsan , L. Kang , S. Kyriacou , P. Maksimovic , M. Roguljic , J. Roskes , S. Sekhar , M. Swartz 
















The University of Kansas, Lawrence, Kansas, USA

A. Abreu , L.F. Alcerro Alcerro , J. Anguiano , S. Arteaga Escatel , P. Baringer , A. Bean , Z. Flowers , D. Grove , J. King , G. Krintiras , M. Lazarovits , C. Le Mahieu , J. Marquez , M. Murray , M. Nickel , M. Pitt , S. Popescu⁹¹ , C. Rogan , C. Royon , R. Salvatico , S. Sanders , C. Smith , G. Wilson 






















Kansas State University, Manhattan, Kansas, USA

B. Allmond , R. Gujju Gurunadha , A. Ivanov , K. Kaadze , Y. Maravin , J. Natoli , D. Roy , G. Sorrentino 

University of Maryland, College Park, Maryland, USA

A. Baden , A. Belloni , J. Bistany-riebman, Y.M. Chen , S.C. Eno , N.J. Hadley , S. Jabeen , R.G. Kellogg , T. Koeth , B. Kronheim, Y. Lai , S. Lascio , A.C. Mignerey , S. Nabili , C. Palmer , C. Papageorgakis , M.M. Paranjpe, L. Wang 

Massachusetts Institute of Technology, Cambridge, Massachusetts, USA













J. Bendavid , I.A. Cali , P.c. Chou , M. D'Alfonso , J. Eysermans , C. Freer , G. Gomez-Ceballos , M. Goncharov, G. Grosso, P. Harris, D. Hoang, D. Kovalskyi , J. Krupa , L. Lavezzo , Y.-J. Lee , K. Long , C. Mcginn, A. Novak , C. Paus , C. Roland , G. Roland , S. Rothman , G.S.F. Stephans , Z. Wang , B. Wyslouch , T. J. Yang 

University of Minnesota, Minneapolis, Minnesota, USA











B. Crossman , B.M. Joshi , C. Kapsiak , M. Krohn , D. Mahon , J. Mans 

B. Marzocchi , M. Revering , R. Rusack , R. Saradhy , N. Strobbe 

















University of Nebraska-Lincoln, Lincoln, Nebraska, USA

K. Bloom , D.R. Claes , G. Haza , J. Hossain , C. Joo , I. Kravchenko , J.E. Siado , W. Tabb , A. Vagnerini , A. Wightman , F. Yan , D. Yu 







State University of New York at Buffalo, Buffalo, New York, USA

H. Bandyopadhyay , L. Hay , H.w. Hsia, I. Iashvili , A. Kalogeropoulos , A. Kharchilava , M. Morris , D. Nguyen , S. Rappoccio , H. Rejeb Sfar, A. Williams , P. Young 



























Northeastern University, Boston, Massachusetts, USA

G. Alverson , E. Barberis , J. Bonilla , J. Dervan, Y. Haddad , Y. Han , A. Krishna , J. Li , M. Lu , G. Madigan , R. Mccarthy , D.M. Morse , V. Nguyen , T. Orimoto , A. Parker , L. Skinnari , D. Wood 









Northwestern University, Evanston, Illinois, USA

J. Bueghly, S. Dittmer , K.A. Hahn , Y. Liu , Y. Miao , D.G. Monk , M.H. Schmitt , A. Taliercio , M. Velasco


















University of Notre Dame, Notre Dame, Indiana, USA

G. Agarwal , R. Band , R. Bucci, S. Castells , A. Das , R. Goldouzian , M. Hildreth , K.W. Ho , K. Hurtado Anampa , T. Ivanov , C. Jessop , K. Lannon , J. Lawrence , N. Loukas , L. Lutton , J. Mariano, N. Marinelli, I. Mcalister, T. McCauley , C. Mcgrady , C. Moore , Y. Musienko¹⁷ , H. Nelson , M. Osherson , A. Piccinelli , R. Ruchti , A. Townsend , Y. Wan, M. Wayne , H. Yockey, M. Zarucki , L. Zygalá 

The Ohio State University, Columbus, Ohio, USA

A. Basnet , B. Bylsma, M. Carrigan , L.S. Durkin , C. Hill , M. Joyce , M. Nunez Ornelas , K. Wei, B.L. Winer , B. R. Yates 


















Princeton University, Princeton, New Jersey, USA

H. Bouchamaoui , P. Das , G. Dezoort , P. Elmer , A. Frankenthal , B. Greenberg , N. Haubrich , K. Kennedy, G. Kopp , S. Kwan , D. Lange , A. Loeliger , D. Marlow , I. Ojalvo , J. Olsen , A. Shevelev , D. Stickland , C. Tully 




University of Puerto Rico, Mayaguez, Puerto Rico, USA

S. Malik 

Purdue University, West Lafayette, Indiana, USA

A.S. Bakshi , S. Chandra , R. Chawla , A. Gu , L. Gutay, M. Jones , A.W. Jung , A.M. Koshy, M. Liu , G. Negro , N. Neumeister , G. Paspalaki , S. Piperov , V. Scheurer, J.F. Schulte , M. Stojanovic , J. Thieman , A. K. Viridi , F. Wang , W. Xie 

Purdue University Northwest, Hammond, Indiana, USA

J. Dolen , N. Parashar , A. Pathak 







Rice University, Houston, Texas, USA

D. Acosta , T. Carnahan , K.M. Ecklund , P.J. Fernández Manteca , S. Freed, P. Gardner, E.J.M. Geurts , W. Li , J. Lin , O. Miguel Colin , B.P. Padley , R. Redjimi, J. Rotter , E. Yigitbasi , Y. Zhang 








University of Rochester, Rochester, New York, USA

A. Bodek , P. de Barbaro , R. Demina , J.L. Dulemba , A. Garcia-Bellido , O. Hindrichs , A. Khukhunaishvili , N. Parmar, P. Parygin⁹² , E. Popova⁹² , R. Taus 








Rutgers, The State University of New Jersey, Piscataway, New Jersey, USA

B. Chiarito, J.P. Chou , S.V. Clark , D. Gadkari , Y. Gershtein , E. Halkiadakis , M. Heindl , C. Houghton , D. Jaroslowski , S. Konstantinou , I. Laflotte , A. Lath , R. Montalvo, K. Nash, J. Reichert , H. Routray , P. Saha , S. Salur , S. Schnetzer, S. Somalwar , R. Stone , S.A. Thayil , S. Thomas, J. Vora , H. Wang

University of Tennessee, Knoxville, Tennessee, USA

D. Ally , A.G. Delannoy , S. Fiorendi , S. Higginbotham , T. Holmes , A.R. Kanuganti , N. Karunarathna , L. Lee , E. Nibigira , S. Spanier








Texas A&M University, College Station, Texas, USA

D. Aebi , M. Ahmad , T. Akhter , O. Bouhali⁹³ , R. Eusebi , J. Gilmore , T. Huang , T. Kamon⁹⁴ , H. Kim , S. Luo , R. Mueller , D. Overton , D. Rathjens , A. Safonov







Texas Tech University, Lubbock, Texas, USA

N. Akchurin , J. Damgov , N. Gogate , V. Hegde , A. Hussain , Y. Kazhykarim, K. Lamichhane , S.W. Lee , A. Mankel , T. Peltola , I. Volobouev

Vanderbilt University, Nashville, Tennessee, USA

E. Appelt , Y. Chen , S. Greene, A. Gurrola , W. Johns , R. Kunnawalkam Elayavalli , A. Melo , F. Romeo , P. Sheldon , S. Tuo , J. Velkovska , J. Viinikainen







University of Virginia, Charlottesville, Virginia, USA

B. Cardwell , B. Cox , J. Hakala , R. Hirosky , A. Ledovskoy , C. Neu 

Wayne State University, Detroit, Michigan, USA

S. Bhattacharya , P.E. Karchin 


University of Wisconsin - Madison, Madison, Wisconsin, USA

A. Aravind, S. Banerjee , K. Black , T. Bose , S. Dasu , I. De Bruyn , P. Everaerts , C. Galloni, H. He , M. Herndon , A. Herve , C.K. Koraka , A. Lanaro, R. Loveless , J. Madhusudanan Sreekala , A. Mallampalli , A. Mohammadi , S. Mondal, G. Parida , L. Pétré , D. Pinna, A. Savin, V. Shang , V. Sharma , W.H. Smith , D. Teague, H.F. Tsoi , W. Vetens , A. Warden

Authors affiliated with an institute or an international laboratory covered by a cooperation agreement with CERN

S. Afanasiev , V. Alexakhin , D. Budkouski , I. Golutvin[†] , I. Gorbunov , V. Karjavine , V. Korenkov , A. Lanev , A. Malakhov , V. Matveev⁹⁵ , V. Palichik , V. Perelygin , M. Savina , V. Shalaev , S. Shmatov , S. Shulha , V. Smirnov , O. Teryaev , N. Voytishin , B.S. Yuldashev⁹⁶, A. Zarubin , I. Zhizhin , G. Gavrilo , V. Golovtsov , Y. Ivanov , V. Kim⁹⁵ , P. Levchenko⁹⁷ , V. Murzin , V. Oreshkin , D. Sosnov , V. Sulimov , L. Uvarov , A. Vorobyev[†], Yu. Andreev , A. Dermenev , S. Gninenko , N. Golubev , A. Karneyeu , D. Kirpichnikov , M. Kirsanov , N. Krasnikov , I. Tlisova , A. Toropin , T. Aushev , V. Gavrilo , N. Lychkovskaya , A. Nikitenko^{98,99} , V. Popov , A. Zhokin , M. Chadeeva⁹⁵ , R. Chistov⁹⁵ , S. Polikarpov⁹⁵ , V. Andreev , M. Azarkin , M. Kirakosyan, A. Terkulov , E. Boos , A. Ershov , A. Gribushin , A. Kaminskiy , L. Khein, O. Kodolova⁹⁹ , V. Korotkikh, S. Obraztsov , S. Petrushanko , V. Savrin , A. Snigirev , I. Vardanyan , V. Blinov⁹⁵, T. Dimova⁹⁵ , A. Kozyrev⁹⁵ , O. Radchenko⁹⁵ , Y. Skovpen⁹⁵ , V. Kachanov , D. Konstantinov , S. Slabospitskii , A. Uzunian , A. Babaev , V. Borshch , D. Druzhkin¹⁰⁰

Authors affiliated with an institute formerly covered by a cooperation agreement with CERN

V. Chekhovsky, V. Makarenko 

†: Deceased

¹Also at Yerevan State University, Yerevan, Armenia

²Also at TU Wien, Vienna, Austria

³Also at Institute of Basic and Applied Sciences, Faculty of Engineering, Arab Academy for Science, Technology and Maritime Transport, Alexandria, Egypt

⁴Also at Ghent University, Ghent, Belgium

⁵Also at Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

⁶Also at Universidade Estadual de Campinas, Campinas, Brazil

⁷Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil

⁸Also at UFMS, Nova Andradina, Brazil

⁹Also at Nanjing Normal University, Nanjing, China

¹⁰Now at The University of Iowa, Iowa City, Iowa, USA

¹¹Also at University of Chinese Academy of Sciences, Beijing, China

¹²Also at China Center of Advanced Science and Technology, Beijing, China

¹³Also at University of Chinese Academy of Sciences, Beijing, China

¹⁴Also at China Spallation Neutron Source, Guangdong, China

¹⁵Now at Henan Normal University, Xinxiang, China

¹⁶Also at Université Libre de Bruxelles, Bruxelles, Belgium

¹⁷Also at an institute or an international laboratory covered by a cooperation agreement with CERN

¹⁸Also at Zewail City of Science and Technology, Zewail, Egypt

¹⁹Also at British University in Egypt, Cairo, Egypt

²⁰Now at Ain Shams University, Cairo, Egypt

²¹Also at Purdue University, West Lafayette, Indiana, USA

²²Also at Université de Haute Alsace, Mulhouse, France

²³Also at Istinye University, Istanbul, Turkey

²⁴Also at The University of the State of Amazonas, Manaus, Brazil

²⁵Also at University of Hamburg, Hamburg, Germany

²⁶Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

²⁷Also at Bergische University Wuppertal (BUW), Wuppertal, Germany

²⁸Also at Brandenburg University of Technology, Cottbus, Germany

²⁹Also at Forschungszentrum Jülich, Juelich, Germany

³⁰Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland

³¹Also at Institute of Nuclear Research ATOMKI, Debrecen, Hungary

³²Now at Universitatea Babeş-Bolyai - Facultatea de Fizica, Cluj-Napoca, Romania

³³Also at MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

³⁴Also at HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

³⁵Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt

³⁶Also at Punjab Agricultural University, Ludhiana, India

³⁷Also at University of Visva-Bharati, Santiniketan, India

³⁸Also at Indian Institute of Science (IISc), Bangalore, India

³⁹Also at IIT Bhubaneswar, Bhubaneswar, India

⁴⁰Also at Institute of Physics, Bhubaneswar, India

⁴¹Also at University of Hyderabad, Hyderabad, India

⁴²Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany

⁴³Also at Isfahan University of Technology, Isfahan, Iran

⁴⁴Also at Sharif University of Technology, Tehran, Iran

⁴⁵Also at Department of Physics, University of Science and Technology of Mazandaran,

Behshahr, Iran

⁴⁶Also at Department of Physics, Isfahan University of Technology, Isfahan, Iran

⁴⁷Also at Department of Physics, Faculty of Science, Arak University, ARAK, Iran

⁴⁸Also at Helwan University, Cairo, Egypt

⁴⁹Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy

⁵⁰Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy

⁵¹Also at Università degli Studi Guglielmo Marconi, Roma, Italy

⁵²Also at Scuola Superiore Meridionale, Università di Napoli 'Federico II', Napoli, Italy

⁵³Also at Fermi National Accelerator Laboratory, Batavia, Illinois, USA

⁵⁴Also at Laboratori Nazionali di Legnaro dell'INFN, Legnaro, Italy

⁵⁵Also at Consiglio Nazionale delle Ricerche - Istituto Officina dei Materiali, Perugia, Italy

⁵⁶Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia

⁵⁷Also at Consejo Nacional de Ciencia y Tecnología, Mexico City, Mexico

⁵⁸Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka

⁵⁹Also at Saegis Campus, Nugegoda, Sri Lanka

⁶⁰Also at National and Kapodistrian University of Athens, Athens, Greece

⁶¹Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland

⁶²Also at Universität Zürich, Zurich, Switzerland

⁶³Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria

⁶⁴Also at Laboratoire d'Annecy-le-Vieux de Physique des Particules, IN2P3-CNRS, Annecy-le-Vieux, France

⁶⁵Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey

⁶⁶Also at Konya Technical University, Konya, Turkey

⁶⁷Also at Izmir Bakircay University, Izmir, Turkey

⁶⁸Also at Adiyaman University, Adiyaman, Turkey

⁶⁹Also at Bozok Universitetesi Rektörlüğü, Yozgat, Turkey

⁷⁰Also at Marmara University, Istanbul, Turkey

⁷¹Also at Milli Savunma University, Istanbul, Turkey

⁷²Also at Kafkas University, Kars, Turkey

⁷³Now at Istanbul Okan University, Istanbul, Turkey

⁷⁴Also at Hacettepe University, Ankara, Turkey

⁷⁵Also at Erzincan Binali Yildirim University, Erzincan, Turkey

⁷⁶Also at Istanbul University - Cerrahpasa, Faculty of Engineering, Istanbul, Turkey

⁷⁷Also at Yildiz Technical University, Istanbul, Turkey

⁷⁸Also at Vrije Universiteit Brussel, Brussel, Belgium

⁷⁹Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom

⁸⁰Also at IPPP Durham University, Durham, United Kingdom

⁸¹Also at Monash University, Faculty of Science, Clayton, Australia

⁸²Also at Università di Torino, Torino, Italy

⁸³Also at Bethel University, St. Paul, Minnesota, USA

⁸⁴Also at Karamanoğlu Mehmetbey University, Karaman, Turkey

⁸⁵Also at California Institute of Technology, Pasadena, California, USA

⁸⁶Also at United States Naval Academy, Annapolis, Maryland, USA

⁸⁷Also at Bingol University, Bingol, Turkey

⁸⁸Also at Georgian Technical University, Tbilisi, Georgia

⁸⁹Also at Sinop University, Sinop, Turkey

⁹⁰Also at Erciyes University, Kayseri, Turkey

⁹¹Also at Horia Hulubei National Institute of Physics and Nuclear Engineering (IFIN-HH), Bucharest, Romania

⁹²Now at another institute or international laboratory covered by a cooperation agreement with CERN

⁹³Also at Texas A&M University at Qatar, Doha, Qatar

⁹⁴Also at Kyungpook National University, Daegu, Korea

⁹⁵Also at another institute or international laboratory covered by a cooperation agreement with CERN

⁹⁶Also at Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan

⁹⁷Also at Northeastern University, Boston, Massachusetts, USA

⁹⁸Also at Imperial College, London, United Kingdom

⁹⁹Now at Yerevan Physics Institute, Yerevan, Armenia

¹⁰⁰Also at Universiteit Antwerpen, Antwerpen, Belgium