






RECEIVED: August 21, 2024

REVISED: August 28, 2024

ACCEPTED: September 3, 2024

PUBLISHED: October 1, 2024

# An FONLL prescription with coexisting flavor number PDFs

Andrea Barontini <sup>a</sup>, Alessandro Candido <sup>a,b</sup>, Felix Hekhorn <sup>a,c,d</sup>, Giacomo Magni <sup>e,f</sup>  
and Roy Stegeman <sup>g</sup>

<sup>a</sup>*Tif Lab, Dipartimento di Fisica, Università di Milano and INFN, Sezione di Milano,  
Via Celoria 16, I-20133 Milano, Italy*

<sup>b</sup>*CERN, Theoretical Physics Department,  
CH-1211 Geneva 23, Switzerland*

<sup>c</sup>*Department of Physics, University of Jyväskylä,  
P.O. Box 35, FI-40014, University of Jyväskylä, Finland*

<sup>d</sup>*Helsinki Institute of Physics, University of Helsinki,  
P.O. Box 64, FI-00014, University of Helsinki, Finland*

<sup>e</sup>*Department of Physics and Astronomy,  
Vrije Universiteit, NL-1081 HV Amsterdam, The Netherlands*

<sup>f</sup>*Nikhef Theory Group,  
Science Park 105, 1098 XG Amsterdam, The Netherlands*

<sup>g</sup>*The Higgs Centre for Theoretical Physics, University of Edinburgh,  
JCMB, KB, Mayfield Rd, Edinburgh EH9 3JZ, U.K.*

*E-mail:* [andrea.barontini@unimi.it](mailto:andrea.barontini@unimi.it), [a.candido@cern.ch](mailto:a.candido@cern.ch),  
[felix.a.hekhorn@jyu.fi](mailto:felix.a.hekhorn@jyu.fi), [gmagni@nikhef.nl](mailto:gmagni@nikhef.nl), [r.stegeman@ed.ac.uk](mailto:r.stegeman@ed.ac.uk)

**ABSTRACT:** We present a new prescription to account for heavy quark mass effects in the determination of parton distribution functions (PDFs) based on the FONLL scheme. Our prescription makes explicit use of the freedom to choose the number of active flavors at a given scale and, thus, use coexisting PDFs with different active flavor number. This new prescription is perturbatively equivalent to the former but improves the implementation in two ways. First, it can be naturally generalized to account simultaneously for multiple heavy quark effects, such as charm and bottom effects, which can both be relevant at the same scale due to the small mass difference. Second, it can be trivially generalized to use at any fixed-order or collinear resummed accuracy, while previous prescriptions required ad-hoc expansions of the DGLAP evolution kernels for each coefficient. We supplement the paper with codes for the computation of deep inelastic scattering observables in this new prescription.

**KEYWORDS:** Parton Distributions, Deep Inelastic Scattering or Small-x Physics, Quark Masses

ARXIV EPRINT: [2408.07383](https://arxiv.org/abs/2408.07383)

---

**Contents**

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Constructing flavor number schemes</b>	<b>2</b>
<b>3</b>	<b>FONLL with coexisting flavor number PDFs</b>	<b>6</b>
3.1	Single mass case	6
3.2	Two masses case	9
3.3	A combined prescription for all scales	11
<b>4</b>	<b>Generalizations</b>	<b>12</b>
<b>5</b>	<b>Conclusion</b>	<b>14</b>
<b>A</b>	<b>Simultaneous two mass case</b>	<b>15</b>
<b>B</b>	<b>Implementation</b>	<b>15</b>

---

**1 Introduction**

The high energy physics community has seen significant progress in increasing both the theoretical and experimental precision in the last few years [1]. In the context of hadronic scattering, one of the major source of uncertainties in the prediction of accurate observables comes from the determination of parton distribution functions (PDFs). Recently, several possible lines of improvement have been investigated, from the inclusion of photon evolution effects [2, 3], to the extension to next-to-next-to-next-to-leading order (N<sup>3</sup>LO) in the perturbative strong coupling [4–9] and the inclusion of theoretical uncertainties [10, 11]. In this work, we are interested in the description of heavy quark mass effects, especially in deep inelastic scattering (DIS), which can become significant if the relevant process scale is of the order of any heavy quark mass.

For instance, in the case of the HERA data these corrections turned out to be up to 20% of the observed cross-section [12, 13] and this feature is also expected at the planned Electron-Ion Collider in the US [14] and China [15], as both are designed as low scale colliders. However, also for experiments at the LHC, these corrections can be of great relevance if the scale of the process of interest is low, e.g., B-meson production [16, 17].

As there is no unique prescription on how to include heavy quark mass effects into the theoretical calculations, several approaches have been suggested [18–24]. The general idea of these schemes, known as general mass variable flavor number schemes (GM-VFNSs), is to combine fixed order calculations, which retain all heavy quark mass effects, and collinear resummed calculations. The usage of GM-VFNSs is necessary due to the finite perturbative knowledge and as such all schemes attempt to mimic the exact analytic behavior. Here, we focus on the FONLL approach which was originally suggested for heavy flavor hadroproduction [25], later applied to DIS structure functions at NLO [23] and extended to intrinsic charm [26].

We propose a new prescription of the FONLL scheme which is perturbatively equivalent to the former, but is capable of dealing with an arbitrary accuracy in either the fixed-order calculations or the collinear resummed calculations and also allows a direct application to the case of any number of parton distributions.

Indeed, a complication with the original prescription [23] is the way in which the final coefficient functions are constructed: the procedure rewrites all expression using a single PDF (as is the case for all heavy quark mass schemes so far), which results in a prescription that is non-trivial to follow in practice. This problem becomes apparent when dealing with high perturbative orders (i.e. N<sup>3</sup>LO) or with hadronic collision, e.g. at the LHC, with multiple PDFs involved. The assumptions of using only a single PDF is however not required in FONLL and our new approach primarily relies on lifting this assumption.

Furthermore, the former FONLL prescription only considers a single mass problem without giving a clear recipe on how to deal with the multi-mass case. In practice this is, however, a relevant question as the charm quark mass and the bottom quark mass are of similar order of magnitude. We address this issue specifically and show how our new approach can resolve the issue in a natural way.

By using the same initial PDF defined in different flavour number schemes, instead of rewriting the expressions in terms of a single PDF, we achieve a clear separation between evolution and partonic matrix elements. The `EK0` package [27] for solving the DGLAP evolution equations, supports the computation of such coexisting flavor number PDFs for a given factorization scale, while the `yadism` [28] library allows for the computation of DIS structure functions. Together these codes allow for the evaluation of DIS structure functions in this new FONLL prescription. Coexisting PDFs computed with `EK0` could be used to compute observables for other processes in the FONLL scheme, but we do not provide such codes.

The rest of the paper proceeds as follows. In section 2 we establish the notation used in this paper and recall the necessary ingredients to perform PDF evolution. In section 3.1 we explicitly construct the FONLL scheme for DIS structure functions for the case of a single heavy quark following the procedures of ref. [23] and the new way proposed in this paper. In section 3.2 we examine the case of multiple heavy quarks and in section 3.3 we combine the two cases to a prescription valid at all scales. In section 4 we discuss the generalization to arbitrary accuracy of the perturbative calculation or of the resummed calculation, to intrinsic heavy quark treatment [29] and to arbitrary observables. In section 5 we summarize our results and, finally, we provide two appendices with further details. In appendix A we comment on a generalized treatment of the two mass case and in appendix B we briefly comment on the practical implementation of our new prescription in the `pineline` framework [30].

## 2 Constructing flavor number schemes

High energy hadronic scattering can be described using the collinear factorization theorem [31]. In doing so, the physical observable is computed as a convolution between the short-distance (high scale) contribution, given by the partonic matrix elements, and a long-distance (low scale) contribution, given by the universal PDFs. The former is computed at a given accuracy using perturbative QCD, while the latter is intrinsically related to how the partons are distributed inside the colliding hadron. The dependence of PDFs  $f(x, \mu^2)$  on the factorization

Scheme name	up/down/strange	charm	bottom	top
FFNS3	light	decoupled	decoupled	decoupled
FFNS3c	light	massive	decoupled	decoupled
FFNS3cb	light	massive	massive	decoupled
FFNS4	light	light	decoupled	decoupled
FFNS4b	light	light	massive	decoupled
FFNS5	light	light	light	decoupled

**Table 1.** Mapping of several example configuration of a FFNS to the respective quark masses. Quarks can be either light ( $m = 0$ ), massive ( $m$  finite), or decoupled ( $m = \infty$ ).

scale  $\mu^2$ , is determined by the associated renormalization group equation (RGE), commonly referred to as DGLAP equations [32–34], whose kernels admit a perturbative QCD expansion.

When solving the DGLAP equations or computing the partonic matrix elements, we need to consider the necessary Feynman diagrams and also make an assumption on which flavors may, or may not, participate in each line of the diagram. The value of the active number of flavors is typically associated to the relevant scale  $Q^2$  of the process, but there is no unique prescription on how to treat the heavy quark flavors. A specific resolution of this ambiguity is referred to as a flavor number scheme (FNS). In particular, to define a FNS it is sufficient consider the masses of the quarks which can take three different states: light ( $m = 0$ ), heavy ( $m$  finite), and decoupled ( $m = \infty$ ). The option of using a configuration in which the number of contributing flavors is constant, i.e. it does not depend on any scale, is called the fixed flavor number scheme (FFNS). We denote a fixed flavor number scheme with just  $n_f$  light flavors by FFNS $n_f$ . If a FFNS also accounts for finite mass contributions of the heavy quarks we add the corresponding symbols as suffixes (FFNS $n_f h$ ). Our notation differs from that found in previous literature where the heavy quark contributions are usually not explicitly denoted in the symbol for the FNS, as such we give some examples of our notation in table 1.

Let us now analyze, in more detail, how the strong coupling and PDFs are evolved in different FNSs. In the following, when using FFNS $n_f$ , we add an explicit superscript ( $n_f$ ) to all ingredients, which are computed with this configuration. First, we consider the RGE of the strong coupling  $a_s(\mu^2)$ , the  $\beta$  function,

$$\mu^2 \frac{da_s^{(n_f)}(\mu^2)}{d\mu^2} = \beta^{(n_f)}(a_s^{(n_f)}(\mu^2)) = - \sum_{k=0}^{\infty} (a_s^{(n_f)}(\mu^2))^{2+k} \beta^{(k),(n_f)}, \quad (2.1)$$

where the coefficients of the beta function  $\beta^{(k),(n_f)}$  are known up to 5-loop [35–38]. For any arbitrary final scale  $\mu^2$ , we can solve eq. (2.1) with a given boundary condition  $a_s^{(n_f)}(\mu_0^2)$ , at the initial scale  $\mu_0^2$ , in terms of a couplings operator  $T$

$$a_s^{(n_f)}(\mu^2) = T^{(n_f)}(\mu^2 \leftarrow \mu_0^2) a_s^{(n_f)}(\mu_0^2). \quad (2.2)$$

In an analogous way the DGLAP equations, are given by

$$\mu^2 \frac{df_i^{(n_f)}(\mu^2)}{d\mu^2} = -\gamma_{ij}^{(n_f)}(a_s^{(n_f)}(\mu^2)) f_j^{(n_f)}(\mu^2) = - \sum_{k=0}^{\infty} (a_s^{(n_f)}(\mu^2))^{1+k} \gamma_{ij}^{(k),(n_f)} f_j^{(n_f)}(\mu^2), \quad (2.3)$$

where  $i, j$  are the flavor indices, here and in the rest of the paper, the standard convolution in momentum space is not written explicitly. The coefficients of the anomalous dimension  $\gamma_{ij}^{(k),(n_f)}$  are known up to approximate 4-loop order [5, 7, 39–47]. Again, we can solve the RGEs with a given boundary value  $f_i^{(n_f)}(\mu_0^2)$  with an evolution kernel operator (EKO)  $E$  via

$$f_i^{(n_f)}(\mu^2) = E_{ij}^{(n_f)}(\mu^2 \leftarrow \mu_0^2) f_j^{(n_f)}(\mu_0^2). \quad (2.4)$$

eqs. (2.1) and (2.3) arise from the cancellation of ultra-violet (UV) or collinear poles associated to the  $n_f$  light quarks, respectively. This leads to the logarithmic nature of the RGEs. Indeed, both RGEs correspond to a resummation: due to the logarithmic differential on the l.h.s., when solving the equations, we collect logarithms of the form  $\log(\mu^2/\mu_0^2)$ . Other than these logarithms there is no dependence on the scale, i.e. both the coupling operator  $T$  and the EKO  $E$  are only functions of  $\log(\mu^2/\mu_0^2)$ . We stress that to obtain a FFNS $n_f$  result, we need to use a consistent configuration throughout, i.e. the boundary conditions are given in FFNS $n_f$  and the operators use FFNS $n_f$  ingredients.


Next, we move to the computation of observables: to simplify the discussion in the beginning we consider an observable  $F$  which depends linearly on a PDF  $f$ . In section 4 we explicitly remove this assumption and show how to construct the new prescription for any observable. In an FFNS setup, the observable  $F$  can be computed as a convolution between the respective PDF  $f^{(n_f)}$  and a coefficient function  $C^{(n_f)}$ ,

$$F^{(n_f)} = f^{(n_f)} C^{(n_f)} \quad \text{with} \quad C^{(n_f)}(x) = \sum_k \left(a_s^{(n_f)}\right)^k C_k^{(n_f)}(x), \quad (2.5)$$

where we expanded the coefficient function into a power series of  $a_s^{(n_f)}$ . Note that we suppressed any flavor dependence on the PDF or the coefficient function for the sake of clarity and that the lower bound of the series expansion depends on the observable  $F$ . When computing the higher order contributions to the coefficient functions  $C_k^{(n_f)}$  one always encounters collinear configuration which have to be reabsorbed into the PDF definition and using the  $\overline{\text{MS}}$  scheme this can leave logarithms of  $\ln(1-x)$  and  $\ln(x)$  in the coefficient functions due to over-subtractions [48].

The simplest way to include heavy quark mass effects is then to move to a FFNS $n_f h$  scheme and to thus add an explicit mass dependency to the coefficient functions. The coefficient functions contain again collinear configuration, which, however, do not manifest as poles when they involve heavy quarks but instead yield collinear logarithms of the form  $\ln(Q^2/m_h^2)$  inside the coefficient functions  $C_k^{(n_f h)}$ . Eventually, due to the finite perturbative expansion in  $a_s$ , we only get a finite number of such logarithms which turn the calculation unstable in the region of large- $Q^2$ . By allowing one light flavor more in the RGEs, in the region where  $Q^2 \gg m_h^2$ , these logarithms are properly resummed. A variable flavor number scheme (VFNS) is a FNS where the number of active flavours participating in the evolution equations is dynamic.

It is thus important to use a FNS which consistently combines several FFNSs at a given matching scale. A common choice of matching scales are the heavy quark masses such that logarithms of the form  $\ln(Q^2/m_h^2)$  are included in the region where  $Q > \lambda m_h$  for some constant  $\lambda$ . When transitioning from one set of active flavors to the next, we need



**Figure 1.** Schematic representation of a VFNS. Each horizontal line indicates a FFNS scheme with  $n_l$  light flavor and  $n_h$  massive quarks. Vertical lines can coexist for any scale  $Q$  denoted on the abscissa axis; the solid black line corresponds to a VFNS, which changes FNS at a scale equal to the quark threshold, see eq. (2.7): for scales below  $Q \leq m_c$  it accounts for massive charm corrections, while otherwise treats all the quarks a massless, with 4, 5 or 6 active flavors.

to apply a matching operator which can be computed perturbatively. For matching the strong coupling  $a_s$  at  $\lambda_\alpha m_h^2$  we find

$$\begin{aligned}
 a_s^{(n_f+1)}(\lambda_\alpha m_h^2) &= \zeta^{(n_f+1)}\left(a_s^{(n_f+1)}(\lambda_\alpha m_h^2), \ln(\lambda_\alpha)\right) a_s^{(n_f)}(\lambda_\alpha m_h^2) \\
 &= \sum_{k=0}^{\infty} \left(a_s^{(n_f+1)}(\lambda_\alpha m_h^2)\right)^k \sum_{l=0}^k \zeta^{kl, (n_f+1)} \ln^k(\lambda_\alpha) a_s^{(n_f)}(\lambda_\alpha m_h^2),
 \end{aligned}
 \tag{2.6}$$

where the decoupling constants  $\zeta^{kl, (n_f)}$  are known up to 4-loop [49, 50], while for matching the PDFs at  $\lambda_f m_h^2$  we find

$$\begin{aligned}
 f_i^{(n_f+1)}(\lambda_f m_h^2) &= A_{ij}^{(n_f+1)}\left(a_s^{(n_f+1)}(\lambda_f m_h^2), \ln(\lambda_f)\right) f_i^{(n_f)}(\lambda_f m_h^2) \\
 &= \sum_{k=0}^{\infty} \left(a_s^{(n_f+1)}(\lambda_f m_h^2)\right)^k \sum_{l=0}^k A_{ij}^{kl, (n_f+1)} \ln^l(\lambda_f) f_j^{(n_f)}(\lambda_f m_h^2)
 \end{aligned}
 \tag{2.7}$$

where the coefficients of the matching matrix  $A_{ij}^{kl, (n_f)}$  are known up to 4-loop order [51–65]. In the following we assume that the matching is performed at the mass,  $\lambda_\alpha = 1 = \lambda_f$ , but we comment on the general case, which can be easily recovered in our new prescription, in section 4.

The central observation of this paper is that, at any scale  $Q$ , we can theoretically use any FFNS and, moreover, we could even use them *simultaneously*. To visualize this concept, we provide figure 1 where we indicate the different FFNSs by the horizontal lines. The RGEs, eqs. (2.2) and (2.4), allow to navigate in the scale  $Q$ , i.e. along the horizontal axis, while the matching equations, eqs. (2.6) and (2.7), allow to navigate in the number of flavors  $n_f$ , i.e. along the vertical axis. So far most evolution codes do not consider this coexistence of FNS

and hence provide no native interface to access them, but our recently developed DGLAP solver EKO [27] explicitly allows for this possibility.

Before moving to the explicit FONLL prescription, we list another VFNS example, the simple but common choice of the zero mass-variable flavor number scheme (ZM-VFNS). In the ZM-VFNS quarks are treated as decoupled below their respective matching scale and light above. We can thus write for an arbitrary observable  $\sigma(Q^2)$  at a given scale  $Q$

$$\sigma^{\text{ZM-VFNS}}(Q^2) = \begin{cases} \sigma^{(3)}(Q^2) & \text{if } Q^2 < m_c^2 \\ \sigma^{(4)}(Q^2) & \text{if } m_c^2 \leq Q^2 < m_b^2 \\ \sigma^{(5)}(Q^2) & \text{if } m_b^2 \leq Q^2 < m_t^2 \\ \sigma^{(6)}(Q^2) & \text{if } Q^2 \geq m_t^2 \end{cases} \quad (2.8)$$

This scheme usually generates unphysical discontinuities where the scale  $Q$  equals any of the heavy quark masses. It does not account for mass effects of heavy quarks in coefficient functions, which simplifies the calculation of the observable, however these effects can be significant around  $Q^2 \sim m_h^2$ .

### 3 FONLL with coexisting flavor number PDFs

Here we discuss the FONLL prescription applied to DIS structure functions. First, in section 3.1 we review the prescription accounting for the massive contributions of a single heavy quark as introduced in refs. [23, 26]. Then, in section 3.2 we discuss the case of accounting for the massive contributions of two heavy quarks. As in the previous section, we assume that the matching scale is equal to the quark masses,  $\lambda_f = 1$ , though the general case is easily recovered. While the FONLL prescription may be applied to other observables, as discussed in section 4, in this section we focus the discussion on DIS.

#### 3.1 Single mass case

In the following we consider, without loss of generality, the case of the charm quark and assume that the boundary conditions for the strong coupling  $a_s$  and the PDF  $f$  are given in the three flavor scheme at the charm mass, i.e.  $a_s^{(3)}(m_c)$  and  $f^{(3)}(m_c)$ .

The main idea of the FONLL prescription is to enhance the fixed order calculation by accounting for the resummation of collinear logarithms, which can become arbitrarily large. In practice we sum the observables defined in FFNS3c and FFNS4, while taking care of the double counting and write

$$F^{\text{FONLL}}(Q^2, m_c^2) = F^{(3c)}(Q^2, m_c^2) + F^{(4)}(Q^2) - F^{(3c \cap 4)}(Q^2, m_c^2), \quad (3.1)$$

where we suppress the  $x$  dependence and borrow the intersection operation  $\cap$  from set theory language to indicate the subset of terms present in both schemes.<sup>1</sup> Each ingredient obeys factorization and is thus given by a convolution between a PDF  $f$  and a coefficient function  $C$ .

In FFNS3c, the observable is given by

$$F^{(3c)}(Q^2, m_c^2) = C^{(3c)}(Q^2/m_c^2) f^{(3c)}(Q^2) = C^{(3c)}(Q^2/m_c^2) E^{(3c)}(\log(Q^2/m_c^2)) f^{(3)}(m_c^2) \quad (3.2)$$

<sup>1</sup>In the notation of refs. [23, 26] the term  $F^{(3c \cap 4)}$  is denoted by  $F^{(3,0)}$ .



with the coefficient function  $C^{(3c)}$ , which can depend both power-like and logarithmically on the ratio  $Q^2/m_c^2$ , and the EKO  $E^{(3c)}$ , which contributes only logarithmically. Note that both, the EKO  $E^{(3c)}$  and the coefficient function  $C^{(3c)}$  also depend on the strong coupling  $a_s^{(3c)}$  defined in FFNS3c, but for the sake for readability we suppress this dependency.

Analogously, in FFNS4 the observable is given by

$$F^{(4)}(Q^2) = C^{(4)} f^{(4)}(Q^2) = C^{(4)} E^{(4)}(\log(Q^2/m_c^2)) A^{(4)}(\log(\mu_c^2/m_c^2) = 0) f^{(3)}(m_c^2) \quad (3.3)$$

with the coefficient function  $C^{(4)}$ , which is independent of the charm mass as charm is considered light, and the EKO  $E^{(4)}$ , which encodes the logarithmic dependency including that on the charm mass. We abbreviate the dependency of the matching condition  $A^{(4)}$  to the logarithmic ratio between the matching scale  $\mu_c^2 = \lambda_f m_c^2$  and the heavy quark mass  $m_c^2$ . Remember, that here we assumed the matching scale to be at the corresponding quark mass, thus  $\lambda_f = 1$ , and  $A^{(4)}(0)$  indicates the non-trivial scale independent part of the matching between FFNS4 and FFNS3c. Again, we suppress the dependency of all elements on the strong coupling  $a_s^{(4)}$  for readability.

Finally, we can determine the overlap,  $F^{(3c\cap 4)}$ , between the two expressions, eqs. (3.2) and (3.3), and we find

$$\begin{aligned} F^{(3c\cap 4)}(Q^2, m_c^2) &= C^{(3c\cap 4)}(\log(Q^2/m_c^2)) f^{(3c)}(Q^2) \\ &= C^{(3c\cap 4)}(\log(Q^2/m_c^2)) E^{(3c)}(\log(Q^2/m_c^2)) f^{(3)}(m_c^2), \end{aligned} \quad (3.4)$$

where the coefficient function  $C^{(3c\cap 4)}$  depends only logarithmically on the ratio  $Q^2/m_c^2$ , since eq. (3.3) has only a logarithmic dependency. Moreover, we can give an explicit expression for  $C^{(3c\cap 4)}$ :

$$C^{(3c\cap 4)}(\log(Q^2/m_c^2)) = C^{(4)} A^{(4)}(\log(Q^2/m_c^2)) \quad (3.5)$$

which can be obtained from eq. (3.3) by displacing the matching from  $\mu_c^2 = m_c^2$  (thus  $A^{(4)}(0)$ ) to  $\mu_c^2 = Q^2$  (thus  $A^{(4)}(\log(Q^2/m_c^2))$ ). Displacing the matching reshuffles collinear logarithms from the resummation in  $E^{(4)}$  to the matching operator  $A^{(4)}$ . eq. (3.5) gives a definition of  $C^{(3c\cap 4)}$  in terms of FFNS4 ingredients, i.e. specifically of  $a_s^{(4)}$ , which would naively make eq. (3.5) use multiple schemes simultaneously (comparing l.h.s. and r.h.s.). However, we can also give a definition based on FFNS3c by observing that  $C^{(3c\cap 4)}$  may only depend logarithmically on  $Q^2/m_c^2$ . Thus we can extract  $C^{(3c\cap 4)}$  from  $C^{(3c)}$  by only retaining the collinear logarithms  $\log(Q^2/m_c^2)$  and neglecting the power-like contributions, i.e. we can write:

$$C^{(3c)}(Q^2/m_c^2) = C^{(3c\cap 4)}(\log(Q^2/m_c^2)) + \tilde{C}^{(3c)}(m_c^2/Q^2), \quad (3.6)$$


where  $\tilde{C}^{(3c)}(m_c^2/Q^2)$  contains only terms proportional to mass power corrections. Note that the constant term  $\mathcal{O}(1)$  is considered part of  $C^{(3c\cap 4)}$ .

The prescription of refs. [23, 26] now proceeds in rewriting eq. (3.1) in terms of a single PDF  $f^{(4)}(Q^2)$ , i.e. by transforming the coefficient function into FFNS4:

$$F^{\text{FONLL}}(Q^2, m_c^2) = C^{\text{FONLL}}(Q^2, m_c^2) f^{(4)}(Q^2) \quad (3.7)$$

$$= \left( C^{(4)} + B^{[3c]}(Q^2, m_c^2) - B^{[3c\cap 4]}(\log(Q^2/m_c^2)) \right) f^{(4)}(Q^2), \quad (3.8)$$





**Figure 2.** Relative (left) and absolute (right) difference of the charm-tagged structure function  $F_{2c}$  evaluated with the method presented in this work and the prescription of refs. [23, 26].


where  $B^{[3c]}$  and  $B^{[3c\cap 4]}$  are related to the FFNS3c contribution, eq. (3.2), and the overlap contribution, eq. (3.4), respectively, written as a function of FFNS4 quantities. Both can be computed by explicitly reverting the PDF scheme change,  $(A^{(4)}(Q^2/m_c^2))^{-1}$ , and by transforming the strong coupling  $a_s$  into the required FFNS4 scheme:

$$B^{[3c]}(Q^2, m_c^2) = C^{(3c)}(Q^2, m_c^2) \left( A^{(4)}(Q^2/m_c^2) \right)^{-1}. \quad (3.9)$$


However, using EKO we can compute  $f^{(3c)}(Q^2)$  and  $f^{(4)}(Q^2)$  at any scale  $Q^2$ , which allows us to simply use eqs. (3.2)–(3.4) as they are without re-expressing any terms as a function of FFNS4 quantities. This, which is the central argument of the paper, means we are dealing with *coexisting flavor number PDFs*, as we have explicitly decoupled the scale and the FNS here. Differences between this prescription, and that of refs. [23, 26] are higher order with respect to the required perturbative accuracy.

While in the prescription of refs. [23, 26] the necessary cancellations in the low or high scale region respectively happen analytically at the level of coefficient functions, i.e. *before* multiplying with the PDF  $f^{(4)}(Q^2)$  in eq. (3.8), in our new prescription this happens numerically at the level of structure functions, i.e. on the r.h.s. of eq. (3.1).

In figure 2 we show the difference on the FONLL charm tagged structure function  $F_{2c}$ , as a function of  $x$  and  $Q^2$ , evaluated with the new prescription described above and the one presented in refs. [23, 26]. We indicate both, the relative difference (left panel) and absolute difference (right panel) in the considered kinematic range of  $x \in [10^{-5}, 1]$  and  $Q^2 \in [25, 100]$  GeV<sup>2</sup>. The calculations are performed at NNLO accuracy and use as boundary condition NNPDF40\_nnlo\_as\_01180 at  $Q^2 = 1.65$  GeV (and thus  $n_f = 4$ ). [66]. The same comparison for the total structure function  $F_2$  is displayed in figure 3. While we observe a good overall agreement between the two prescriptions, in the small- $x$  region differences can be significant. This difference can be directly associated to the collinear



**Figure 3.** Same as figure 2, but now for the total structure function  $F_2$ .



**Figure 4.** The ratio  $\mathcal{R}_{2c}$  of eq. (3.10) representing the relative impact of the heavy charm contribution to  $F_{2c}^{\text{FONLL}}$  at scales above the bottom quark mass.

resummation: while the prescription of refs. [23, 26] relies on the finite order expansion of the resummation, entering through the inverse scheme change in eq. (3.9), the prescription of this paper retains the full resummation, by using the full EKO in eqs. (3.2)–(3.4). These resummation terms are precisely the higher order difference between the two prescriptions. Performing the comparison at NLO accuracy or with input PDFs containing a fitted charm component [29] yields similar conclusions.

### 3.2 Two masses case

For the case of multiple masses it is possible to apply the prescription of ref. [23] consecutively for each quark mass. However, this only yields faithful results if the quark masses are strongly ordered (e.g.  $m_c \ll m_b$ ), which is not realized in nature. In particular, the charm mass

corrections remains quite large for scales above the bottom mass. This is shown explicitly in figure 4 in which we plot the ratio

$$\mathcal{R}_{2c} \equiv \frac{F_{2c}^{(4)}(Q^2, m_c^2)}{F_{2c}^{\text{FONLL}}(Q^2, m_c^2)}, \quad (3.10)$$

where  $F_{2c}^{\text{FONLL}}$  is defined as in eq. (3.1) for the structure function  $F_{2c}$ . In the calculation we use a charm quark mass  $m_c = 1.51$  GeV and a bottom quark mass of  $m_b = 4.92$  GeV, meaning that the plotted kinematic domain corresponds to scales  $Q^2 > m_b^2$ . It is clear that even in this region, contributions due to the heavy charm mass effects are up to 30% and thus can not be neglected.

To produce accurate predictions, we therefore need to include simultaneously the massive effects of multiple heavy quarks. To account for both the charm and bottom masses, the approach of [23] would require rewriting both FFNS3c and FFNS4b contributions in terms of FFNS5 (see eq. (3.9)), while in our new method the inclusion is significantly simpler.

In practice we need to combine FFNS3c, where power-like terms for charm are present, FFNS4b, where power-like terms for bottom are present, and FFNS5, where collinear resummation for both quarks are fully accounted for. Using again set theory language we write

$$\begin{aligned} F^{\text{FONLL}}(Q^2, m_c^2, m_b^2) &= F^{(3c)}(Q^2, m_c^2) + F^{(4b)}(Q^2, m_c^2, m_b^2) + F^{(5)}(Q^2, m_c^2, m_b^2) \\ &\quad - F^{(3c \cap 4b)}(Q^2, m_c^2) - F^{(4b \cap 5)}(Q^2, m_c^2, m_b^2) - F^{(3c \cap 5)}(Q^2, m_c^2, m_b^2) \\ &\quad + F^{(3c \cap 4b \cap 5)}(Q^2, m_c^2, m_b^2) \end{aligned} \quad (3.11)$$

where the first line corresponds to the plain combination we want to combine, the second line removes the double-counting between any two out of the three schemes, and the third line adds back the triple-overlap between all schemes. Note that here we are considering a consecutive approach, where we only consider one massive quark at a time, however, we comment on the case of simultaneous heavy quark effects in appendix A.

The expression for FFNS3c is given in eq. (3.2), while for FFNS4b we have

$$F^{(4b)}(Q^2, m_c^2, m_b^2) = C^{(4b)}(Q^2/m_b^2) f^{(4b)}(Q^2) \quad (3.12)$$

$$= C^{(4b)}(Q^2/m_b^2) E^{(4b)}(\log(Q^2/m_c^2)) A^{(4b)}(0) f^{(3)}(m_c^2), \quad (3.13)$$

where  $C^{(4b)}(Q^2/m_b^2)$  contains both logarithm and powers of  $Q^2/m_b^2$ , but has no dependence the charm mass. For FFNS5 we have, similarly to eq. (3.3),

$$F^{(5)}(Q^2, m_c^2, m_b^2) = C^{(5)} f^{(5)}(Q^2) \quad (3.14)$$

$$= C^{(5)} E^{(4)}(\log(Q^2/m_b^2)) A^{(5)}(0) E^{(4)}(\log(m_b^2/m_c^2)) A^{(4)}(0) f^{(3)}(m_c^2) \quad (3.15)$$

where  $C^{(5)}$  is a massless coefficient function and we suppress again all dependency on the strong coupling  $a_s$ .

In analogy to the single mass case, the neighboring overlaps  $F^{(3c \cap 4b)}$  and  $F^{(4b \cap 5)}$  are obtained by

$$F^{(3c \cap 4b)}(Q^2, m_c^2) = C^{(3c \cap 4b)}(\log(Q^2/m_c^2)) f^{(3c)}(Q^2) \quad (3.16)$$

$$= C^{(3c \cap 4b)}(\log(Q^2/m_c^2)) E^{(3c)}(\log(Q^2/m_c^2)) f^{(3)}(m_c^2), \quad (3.17)$$

and

$$F^{(4b\cap 5)}(Q^2, m_b^2) = C^{(4b\cap 5)}(\log(Q^2/m_b^2))f^{(4b)}(Q^2) \quad (3.18)$$

$$= C^{(4b\cap 5)}(\log(Q^2/m_b^2))E^{(4b)}(\log(Q^2/m_c^2))A^{(4b)}(0)f^{(3)}(m_c^2), \quad (3.19)$$

where the corresponding coefficient functions are given by

$$C^{(3c\cap 4b)}(\log(Q^2/m_c^2)) = C^{(4)}A^{(4)}(\log(Q^2/m_c^2)) = C^{(3c\cap 4)}(\log(Q^2/m_c^2)), \quad (3.20)$$

$$C^{(4b\cap 5)}(\log(Q^2/m_b^2)) = C^{(5)}A^{(5)}(\log(Q^2/m_b^2)). \quad (3.21)$$

Before discussing the remaining double overlap  $F^{(3c\cap 5)}$  in eq. (3.11) we turn to the triple overlap  $F^{(3c\cap 4b\cap 5)}$  for which we observe the following identity:

$$F^{(3c\cap 4b\cap 5)} = F^{(3c\cap 4b\cap 4b\cap 5)} = F^{((3c\cap 4)\cap (4b\cap 5))} \quad (3.22)$$

where for the first equality we used the fact that the intersection of a set with itself is an identity operation in set theory and for the second equality we used the commutativity of the intersection operation and the fact that  $F^{(3c\cap 4b)} = F^{(3c\cap 4)}$ , as FFNS3c does not contain any bottom mass corrections as the bottom is decoupled.

From eq. (3.22) we recover the well known fact that the matching procedure is a sequential procedure, i.e. we first add the charm contributions and then the bottom contributions. This immediately implies that  $F^{(3c\cap 5)} = F^{(3c\cap 4b\cap 5)}$  as the former would need to go through the charm matching in any case. This reduces the number of contributions in eq. (3.11) in practice to just the first five terms.

Nevertheless, using similar arguments as before we can give an explicit expression for the remaining double overlap  $F^{(3c\cap 5)}$  in eq. (3.11) and write

$$F^{(3c\cap 5)}(Q^2, m_c^2, m_b^2) = C^{(3c\cap 5)}(\log(Q^2/m_c^2), \log(Q^2/m_b^2))f^{(3c)}(Q^2) \quad (3.23)$$

$$= C^{(3c\cap 5)}(\log(Q^2/m_c^2), \log(Q^2/m_b^2))E^{(3c)}(\log(Q^2/m_c^2))f^{(3)}(m_c^2), \quad (3.24)$$

where the coefficient function

$$C^{(3c\cap 5)}(\log(Q^2/m_c^2), \log(Q^2/m_b^2)) = C^{(5)}A^{(5)}(\log(Q^2/m_b^2))A^{(4)}(\log(Q^2/m_c^2)), \quad (3.25)$$

can again be found by shifting both the charm and bottom matching scale to  $Q^2$ .

Generalizing these expressions to the case of more than two masses is straightforward but tedious. One would just follow the same procedure together with the corresponding set theory combination. Moreover, the three mass case is less phenomenologically relevant than the two mass case: the strange quark has a low enough mass to be always considered massless, and the top quark mass is much larger than the bottom quark mass.

### 3.3 A combined prescription for all scales

We now address the issue of combining the above discussion into a combined prescription which can be applied at all scales  $Q^2$ .

While one possible choice would be to use the prescription of section 3.2 at all scales, this turns out to be phenomenologically not well-behaved. Indeed, when  $Q^2$  is smaller than

one of the relevant quark masses, the resummation of the corresponding collinear logarithms is power suppressed and the fixed order calculation is reliable. In fact, also the prescription of ref. [23] is meant to be used only with  $Q^2 > m_h^2$ . In our case this means that we use the single mass case described in section 3.1 after crossing the first quark threshold and the two mass case described in section 3.2 after crossing the second quark threshold. For scales higher than the top mass, one can, as mentioned before, apply an extension of the FONLL prescription for a simultaneous treatment of charm, bottom, and top mass effects; though because  $m_c, m_b \ll m_t$ , the effects of  $m_c$  and  $m_b$  are small at such high scales and for simplicity, one may use a prescription neglecting charm and bottom mass effects. Thus we have

$$F(Q^2, m_c^2, m_b^2) = \begin{cases} F^{(3cb)}(Q^2, m_c^2, m_b^2) & \text{if } Q^2 < m_c^2 \\ F^{(3c)}(Q^2, m_c^2) + F^{(4b)}(Q^2, m_c^2, m_b^2) - F^{(3c \cap 4)}(Q^2, m_c^2) & \text{if } m_c^2 \leq Q^2 < m_b^2 \\ \text{eq. (3.11)} & \text{if } m_b^2 \leq Q^2 < m_t^2 \\ F^{(6)}(Q^2) & \text{if } Q^2 \geq m_t^2 \end{cases}, \quad (3.26)$$

where for  $Q^2 < m_c^2$  charm mass corrections are dealt in a FFNS setup (i.e.  $F^{(3cb)}$ ) and above  $m_c$  the FONLL prescription applies. The same holds for the bottom mass effects: below  $m_b$  bottom mass corrections are dealt in a FFNS setup by either  $F^{(3cb)}$  or  $F^{(4b)}$  and above  $m_b$  the FONLL prescription applies.

At low perturbative order it was found [23] that in the single mass case the cancellation between the FFNS4 structure functions, eq. (3.3), and the overlap, eq. (3.4), is incomplete near the heavy quark threshold. To make the definition of the structure function continuous across the mass thresholds, ref. [23] introduces a damping function

$$\chi(Q^2, m^2) = \left(1 - \frac{m^2}{Q^2}\right)^2. \quad (3.27)$$

For example, we can then modify eq. (3.11) to be

$$F^{\text{FONLL}}(Q^2, m_c^2, m_b^2) = F^{(3c)}(Q^2, m_c^2) + \chi(Q^2, m_c^2) \left[ F^{(4b)}(Q^2, m_c^2, m_b^2) - F^{(3c \cap 4b)}(Q^2, m_c^2) \right. \\ \left. + \chi(Q^2, m_b^2) \left[ F^{(5)}(Q^2, m_c^2, m_b^2) - F^{(4b \cap 5)}(Q^2, m_c^2, m_b^2) \right] \right], \quad (3.28)$$

to ensure a smooth transition at the bottom threshold in eq. (3.26).

We briefly comment on the practical implementation of our new prescription in appendix B.

## 4 Generalizations

In the following we discuss a series of generalizations which demonstrate the improved applicability of our new prescription in comparison to ref. [23]. We stress that all of these considerations are also possible in their prescription, but we would like to point out the advantages gained on both theoretical grounds and practical grounds when implementing the FONLL scheme using our new prescription.

**Perturbative and logarithmic accuracy.** It is the limited perturbative knowledge of the various ingredients  $(C, \beta, \gamma, A, d)$  which make a GM-VFNS prescription necessary. In fact, at all-order accuracy eqs. (3.3) and (3.4) would be identical and we would just be left with eq. (3.2). However, in practice we have to work at a finite order and thus a practical limitation of the prescription of ref. [23] is the required perturbative expansion of eq. (3.9), which becomes technically challenging with increasing perturbative and/or logarithmic accuracy.

To give an example: the recently presented extraction of PDFs at aN<sup>3</sup>LO [7] requires a correct treatment of DIS structure functions at the quoted accuracy. While it is possible [67] to use the prescription of ref. [23], the resulting formulas are lengthy and error prone. Instead, in our new prescription no additional transformations are needed, as we obtained a clear separation between evolution kernels and coefficient functions. The new prescription does not use eq. (3.9) but only relies on the correct determination of the overlap coefficient function eq. (3.5), which is needed in either prescription.

**Mixing accuracies.** As was pointed out in ref. [23] the naive first non-vanishing order for FFNS3c and FFNS4 differ, and this is directly related to the different FONLL schemes (FONLL-A, FONLL-B and FONLL-C) described in ref. [23], since they correspond to different accuracies of eqs. (3.2) and (3.3) respectively. Especially the case of FONLL-B requires additional attention as it relies on the observation that the first non-vanishing order in the strong coupling is different for heavy quark production in massive and massless schemes when neglecting intrinsic heavy quark contributions.

For example, in DIS, the former emerges by photon-gluon fusion which opens at  $O(a_s)$ , the latter is just ordinary photon-quark fusion that first contributes at  $O(a_s^0)$ . Extending the computation to higher perturbative orders in our new prescription is straightforward. Given the coefficients in FFNS3c, eq. (3.2), and FFNS4, eq. (3.3), at a certain order, one just needs to identify the collinear logarithmically order in FFNS3c such that only those also present in FFNS4 are retained in the overlap eq. (3.4).

**Changing the matching point.** In the above discussion we always assumed that the transition scale  $\mu_h$  between the FFNS in which the heavy quark  $h$  is considered massive and the FFNS in which the same quark is considered massless coincides with its mass  $\mu_h = m_h$  — this is a natural choice and indeed the most common convention. However, this is still a choice and one may vary this scale to investigate its dependence. In our new prescription this freedom is naturally embedded by the requirement to compute ingredients which all follow a strict FFNS definition and thus include by definition this matching procedure. Indeed, the full dependency is contained in the evolution operators  $E$  as they naturally define the transition between schemes.

**Treatment of intrinsic heavy quarks.** While ref. [23] only considers the case of perturbative generation of heavy quarks its prescription was later extended to the presence of intrinsic heavy quark components in ref. [26]. In fact, the authors realized that this generalization simplifies the prescription of ref. [23] because several intermediate terms now cancel amongst each other, as the first non-vanishing order is readjusted.

In our new prescription the inclusion of intrinsic heavy quark contributions fall naturally into place as they simply contribute to the FFNS3c structure function, eq. (3.2), and hence also to the overlap, eq. (3.4).

**Observables beyond structure functions and with more collinear distributions.**

Due to the clear separation between the partonic coefficient functions and the contributions from collinear evolution our new prescription can be applied easily to any observable which may also contain more than one collinear distribution (such as e.g. observables at the LHC [25]). We can give a straight generalization of eq. (3.1) for any observable  $\sigma$ :

$$\sigma^{\text{FONLL}}(Q^2, m_c^2) = \sigma^{(3c)}(Q^2, m_c^2) + \sigma^{(4)}(Q^2, m_c^2) - \sigma^{(3c\cap 4)}(Q^2, m_c^2) \quad (4.1)$$

Just as before,  $\sigma^{(3c)}$  and  $\sigma^{(4)}$  have to be computed in a strict FFNS3c and FFNS4 setup respectively, and  $\sigma^{(3c\cap 4)}$  can be deduced from  $\sigma^{(3c)}$  by identifying the collinear logarithms which are fully resummed in the FFNS4 setup. If the observable  $\sigma$  is not sensitive to heavy quark mass effects,  $\sigma^{(3c)}$  and  $\sigma^{(3c\cap 4)}$  cancel each other exactly, as the former does not develop any collinear logarithms.

**Longitudinally polarized structure functions.** Finally, a straight-forward generalization can be obtained for the case of longitudinally polarized structure functions [68] as, in fact, all of the discussion also holds for a given spin state.

## 5 Conclusion

We have presented a new prescription of the FONLL scheme, which is perturbatively equivalent to the former prescription [23] but can be implemented more efficiently. The new prescription relies on the existence of coexisting flavor number PDFs, i.e. an explicit decoupling of the number of active quarks and the factorization scale.

These PDFs can in practice be accessed using the EKO program in an efficient way, while the coefficient functions can be computed using `yadism` [28]. We have implemented two ways how these numerical tools can be used to calculate predictions such as the ones presented in the figures of this paper. In particular, one can store the computed coefficients and DGLAP kernels in PDF-independent FastKernel (FK) tables [69] such that the computation can efficiently be performed for many different PDFs without the need to repeat the often costly computation.

The new prescription features a clear separation between partonic matrix elements and evolution ingredients which make it an ideal tool to investigate higher order corrections. Moreover, our new prescription features a clear extension to the multi-mass case which is relevant, e.g., for mid- and low-scale structure functions in DIS, such as the ones used in all modern PDF determinations.

The new prescription has been adopted by the NNPDF collaboration where it is used to compute predictions for DIS observables [3, 7, 11, 68, 70, 71]. Our new prescription does not imply further restrictions on the kinematically applicable domain and, indeed, eq. (3.26) gives a prescription which can be evaluated at all perturbative scales. The actual kinematic cuts used in NNPDF fits are discussed in the respective references and may differ for unpolarized and longitudinally polarized PDF extractions.



## Acknowledgments

We thank the members of the NNPDF collaboration for useful and insightful discussions at the various stages of this project. We are grateful to Richard Ball and Stefano Forte for comments on the manuscript.

F. H. is supported by the Academy of Finland project 358090 and is funded as a part of the Center of Excellence in Quark Matter of the Academy of Finland, project 346326. G. M. is supported by NWO, the Dutch Research Council. R. S. is supported by the U.K. Science and Technology Facility Council (STFC) consolidated grants ST/T000600/1 and ST/X000494/1.

## A Simultaneous two mass case

In section 3.2 we consider the case of a consecutive decoupling for charm and bottom, i.e. we add heavy quark effects one at a time while still accounting for the other. However, the FONLL scheme and, specifically, the new prescription we propose in this paper can also be applied to the case where one considers simultaneous two mass effects [72, 73]. In this case the massive corrections for either charm or bottom are always contributed by the FFNS3cb structure function, while the collinear resummed parts do not carry any mass dependency.

Eventually, we can give an analogue expression to eq. (3.26):

$$F(Q^2, m_c^2, m_b^2) = \begin{cases} F^{(3cb)}(Q^2, m_c^2, m_b^2) & \text{if } Q^2 < m_c^2 \\ F^{(3cb)}(Q^2, m_c^2, m_b^2) + F^{(4)}(Q^2) - F^{(3cb \cap 4)}(Q^2, m_c^2, m_b^2) & \text{if } m_c^2 \leq Q^2 < m_b^2 \\ F^{(3cb)}(Q^2, m_c^2, m_b^2) + F^{(5)}(Q^2) - F^{(3cb \cap 5)}(Q^2, m_c^2, m_b^2) & \text{if } m_b^2 \leq Q^2 < m_t^2 \\ F^{(6)}(Q^2) & \text{if } Q^2 \geq m_t^2 \end{cases} \quad (\text{A.1})$$

The overlap subtraction terms  $F^{(3cb \cap 4)}$  and  $F^{(3cb \cap 5)}$  follow from the same guiding principle that they can only consist of collinear contributions of either just the charm quark (in the first case) or both charm and bottom quarks (in the second case). Again, recall that the matching is done in a consecutive manner and, thus,  $F^{(5)}$  resums both charm and bottom collinear logarithms.

The implementation in the `yadism` library discussed in appendix B follows the prescription of section 3.2.

## B Implementation

The actual implementation of our new prescription requires the possibility to compute coexisting flavor number PDFs, which is provided by the `EK0` library [27], and the computation of the partonic matrix elements using different FFNS settings. In practice we apply our new prescription only to DIS structure functions and we use the `yadism` library [28] to provide the respective coefficient functions.

An explicit example of a DIS observable calculation using a fixed input PDF is available in the `yadism` documentation at

<https://yadism.readthedocs.io/en/latest/overview/tutorials/fonll.html>.

Instead, if one may wish to compute the same set of observables for many different PDF sets, renormalization or factorization scales, or values of  $\alpha_s$ , as is done in PDF determinations or systematic parameter studies, it is recommended to create interpolation grids in the PineAPPL format [74] that store the combination of DIS coefficients and DGLAP evolution in PDF independent grids called fast-kernel (FK) tables [69] which can be used to compute observables:

$$\sigma = \sum_i \sum_a f_a(x_i, \mu_0^2) \text{FK}_a(x_i, \mu_0^2). \quad (\text{B.1})$$

In this case the implementation is a multi-step process which is driven by the `pineko` code, as illustrated at

<https://pineko.readthedocs.io/en/latest/theory/fonll.html>.

**Open Access.** This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

## References

- [1] S. Amoroso et al., *Snowmass 2021 Whitepaper: Proton Structure at the Precision Frontier*, *Acta Phys. Polon. B* **53** (2022) 12 [[arXiv:2203.13923](https://arxiv.org/abs/2203.13923)] [[INSPIRE](#)].
- [2] T. Cridge, L.A. Harland-Lang, A.D. Martin and R.S. Thorne, *QED parton distribution functions in the MSHT20 fit*, *Eur. Phys. J. C* **82** (2022) 90 [[arXiv:2111.05357](https://arxiv.org/abs/2111.05357)] [[INSPIRE](#)].
- [3] NNPDF collaboration, *Photons in the proton: implications for the LHC*, *Eur. Phys. J. C* **84** (2024) 540 [[arXiv:2401.08749](https://arxiv.org/abs/2401.08749)] [[INSPIRE](#)].
- [4] F. Caola et al., *The path forward to  $N^3$ LO*, in the proceedings of the *Snowmass 2021*, Seattle, U.S.A., July 17–26 (2022) [[arXiv:2203.06730](https://arxiv.org/abs/2203.06730)] [[INSPIRE](#)].
- [5] J. McGowan, T. Cridge, L.A. Harland-Lang and R.S. Thorne, *Approximate  $N^3$ LO parton distribution functions with theoretical uncertainties: MSHT20a $N^3$ LO PDFs*, *Eur. Phys. J. C* **83** (2023) 185 [*Erratum ibid.* **83** (2023) 302] [[arXiv:2207.04739](https://arxiv.org/abs/2207.04739)] [[INSPIRE](#)].
- [6] T. Cridge, L.A. Harland-Lang and R.S. Thorne, *Combining QED and Approximate  $N^3$ LO QCD Corrections in a Global PDF Fit: MSHT20qed\_an3lo PDFs*, *SciPost Phys.* **17** (2024) 026 [[arXiv:2312.07665](https://arxiv.org/abs/2312.07665)] [[INSPIRE](#)].
- [7] NNPDF collaboration, *The path to  $N^3$ LO parton distributions*, *Eur. Phys. J. C* **84** (2024) 659 [[arXiv:2402.18635](https://arxiv.org/abs/2402.18635)] [[INSPIRE](#)].
- [8] A. Barontini, N. Laurenti and J. Rojo, *NNPDF4.0 a $N^3$ LO PDFs with QED corrections*, in the proceedings of the *31st International Workshop on Deep-Inelastic Scattering and Related Subjects*, Grenoble, France, April 08–12 (2024) [[arXiv:2406.01779](https://arxiv.org/abs/2406.01779)] [[INSPIRE](#)].
- [9] A. Cooper-Sarkar et al., *A Benchmarking of QCD Evolution at Approximate  $N^3$ LO*, [arXiv:2406.16188](https://arxiv.org/abs/2406.16188) [[INSPIRE](#)].
- [10] NNPDF collaboration, *Parton Distributions with Theory Uncertainties: General Formalism and First Phenomenological Studies*, *Eur. Phys. J. C* **79** (2019) 931 [[arXiv:1906.10698](https://arxiv.org/abs/1906.10698)] [[INSPIRE](#)].
- [11] NNPDF collaboration, *Determination of the theory uncertainties from missing higher orders on NNLO parton distributions with percent accuracy*, *Eur. Phys. J. C* **84** (2024) 517 [[arXiv:2401.10319](https://arxiv.org/abs/2401.10319)] [[INSPIRE](#)].

- [12] H1 and ZEUS collaborations, *Combination of measurements of inclusive deep inelastic  $e^\pm p$  scattering cross sections and QCD analysis of HERA data*, *Eur. Phys. J. C* **75** (2015) 580 [[arXiv:1506.06042](#)] [[INSPIRE](#)].
- [13] H1 and ZEUS collaborations, *Combination and QCD analysis of charm and beauty production cross-section measurements in deep inelastic ep scattering at HERA*, *Eur. Phys. J. C* **78** (2018) 473 [[arXiv:1804.01019](#)] [[INSPIRE](#)].
- [14] R. Abdul Khalek et al., *Science Requirements and Detector Concepts for the Electron-Ion Collider: EIC Yellow Report*, *Nucl. Phys. A* **1026** (2022) 122447 [[arXiv:2103.05419](#)] [[INSPIRE](#)].
- [15] D.P. Anderle et al., *Electron-ion collider in China*, *Front. Phys. (Beijing)* **16** (2021) 64701 [[arXiv:2102.09222](#)] [[INSPIRE](#)].
- [16] I. Helenius and H. Paukkunen, *B-meson hadroproduction in the SACOT- $m_T$  scheme*, *JHEP* **07** (2023) 054 [[arXiv:2303.17864](#)] [[INSPIRE](#)].
- [17] M. Cacciari et al., *Theoretical predictions for charm and bottom production at the LHC*, *JHEP* **10** (2012) 137 [[arXiv:1205.6344](#)] [[INSPIRE](#)].
- [18] M.A.G. Aivazis, J.C. Collins, F.I. Olness and W.-K. Tung, *Leptoproduction of heavy quarks. II. A unified QCD formulation of charged and neutral current processes from fixed target to collider energies*, *Phys. Rev. D* **50** (1994) 3102 [[hep-ph/9312319](#)] [[INSPIRE](#)].
- [19] R.S. Thorne and R.G. Roberts, *A practical procedure for evolving heavy flavor structure functions*, *Phys. Lett. B* **421** (1998) 303 [[hep-ph/9711223](#)] [[INSPIRE](#)].
- [20] M. Krämer, F.I. Olness and D.E. Soper, *Treatment of heavy quarks in deeply inelastic scattering*, *Phys. Rev. D* **62** (2000) 096007 [[hep-ph/0003035](#)] [[INSPIRE](#)].
- [21] W.-K. Tung, S. Kretzer and C. Schmidt, *Open heavy flavor production in QCD: Conceptual framework and implementation issues*, *J. Phys. G* **28** (2002) 983 [[hep-ph/0110247](#)] [[INSPIRE](#)].
- [22] P.M. Nadolsky and W.-K. Tung, *Improved Formulation of Global QCD Analysis with Zero-mass Matrix Elements*, *Phys. Rev. D* **79** (2009) 113014 [[arXiv:0903.2667](#)] [[INSPIRE](#)].
- [23] S. Forte, E. Laenen, P. Nason and J. Rojo, *Heavy quarks in deep-inelastic scattering*, *Nucl. Phys. B* **834** (2010) 116 [[arXiv:1001.2312](#)] [[INSPIRE](#)].
- [24] M. Guzzi, P.M. Nadolsky, H.-L. Lai and C.-P. Yuan, *General-Mass Treatment for Deep Inelastic Scattering at Two-Loop Accuracy*, *Phys. Rev. D* **86** (2012) 053005 [[arXiv:1108.5112](#)] [[INSPIRE](#)].
- [25] M. Cacciari, M. Greco and P. Nason, *The  $p_T$  spectrum in heavy-flavour hadroproduction*, *JHEP* **05** (1998) 007 [[hep-ph/9803400](#)] [[INSPIRE](#)].
- [26] R.D. Ball, M. Bonvini and L. Rottoli, *Charm in Deep-Inelastic Scattering*, *JHEP* **11** (2015) 122 [[arXiv:1510.02491](#)] [[INSPIRE](#)].
- [27] A. Candido, F. Hekhorn and G. Magni, *EKO: evolution kernel operators*, *Eur. Phys. J. C* **82** (2022) 976 [[arXiv:2202.02338](#)] [[INSPIRE](#)].
- [28] A. Candido et al., *Yadism: yet another deep-inelastic scattering module*, *Eur. Phys. J. C* **84** (2024) 698 [[arXiv:2401.15187](#)] [[INSPIRE](#)].
- [29] NNPDF collaboration, *Evidence for intrinsic charm quarks in the proton*, *Nature* **608** (2022) 483 [[arXiv:2208.08372](#)] [[INSPIRE](#)].
- [30] A. Barontini et al., *Pineline: Industrialization of high-energy theory predictions*, *Comput. Phys. Commun.* **297** (2024) 109061 [[arXiv:2302.12124](#)] [[INSPIRE](#)].

- [31] J.C. Collins, D.E. Soper and G.F. Sterman, *Factorization of Hard Processes in QCD*, *Adv. Ser. Direct. High Energy Phys.* **5** (1989) 1 [[hep-ph/0409313](#)] [[INSPIRE](#)].
- [32] G. Altarelli and G. Parisi, *Asymptotic Freedom in Parton Language*, *Nucl. Phys. B* **126** (1977) 298 [[INSPIRE](#)].
- [33] Y.L. Dokshitzer, *Calculation of the Structure Functions for Deep Inelastic Scattering and  $e^+e^-$  Annihilation by Perturbation Theory in Quantum Chromodynamics* (in Russian), *Sov. Phys. JETP* **46** (1977) 641 [[INSPIRE](#)].
- [34] V.N. Gribov and L.N. Lipatov, *Deep inelastic  $e p$  scattering in perturbation theory*, *Sov. J. Nucl. Phys.* **15** (1972) 438 [[INSPIRE](#)].
- [35] P.A. Baikov, K.G. Chetyrkin and J.H. Kühn, *Five-Loop Running of the QCD coupling constant*, *Phys. Rev. Lett.* **118** (2017) 082002 [[arXiv:1606.08659](#)] [[INSPIRE](#)].
- [36] K.G. Chetyrkin, G. Falcioni, F. Herzog and J.A.M. Vermaseren, *Five-loop renormalisation of QCD in covariant gauges*, *JHEP* **10** (2017) 179 [*Addendum ibid.* **12** (2017) 006] [[arXiv:1709.08541](#)] [[INSPIRE](#)].
- [37] F. Herzog et al., *The five-loop beta function of Yang-Mills theory with fermions*, *JHEP* **02** (2017) 090 [[arXiv:1701.01404](#)] [[INSPIRE](#)].
- [38] T. Luthe, A. Maier, P. Marquard and Y. Schroder, *The five-loop Beta function for a general gauge group and anomalous dimensions beyond Feynman gauge*, *JHEP* **10** (2017) 166 [[arXiv:1709.07718](#)] [[INSPIRE](#)].
- [39] J. Davies et al., *Large- $n_f$  contributions to the four-loop splitting functions in QCD*, *Nucl. Phys. B* **915** (2017) 335 [[arXiv:1610.07477](#)] [[INSPIRE](#)].
- [40] S. Moch et al., *Four-Loop Non-Singlet Splitting Functions in the Planar Limit and Beyond*, *JHEP* **10** (2017) 041 [[arXiv:1707.08315](#)] [[INSPIRE](#)].
- [41] J. Davies, C.-H. Kom, S. Moch and A. Vogt, *Resummation of small- $x$  double logarithms in QCD: inclusive deep-inelastic scattering*, *JHEP* **08** (2022) 135 [[arXiv:2202.10362](#)] [[INSPIRE](#)].
- [42] J.M. Henn, G.P. Korchemsky and B. Mistlberger, *The full four-loop cusp anomalous dimension in  $\mathcal{N} = 4$  super Yang-Mills and QCD*, *JHEP* **04** (2020) 018 [[arXiv:1911.10174](#)] [[INSPIRE](#)].
- [43] M. Bonvini and S. Marzani, *Four-loop splitting functions at small  $x$* , *JHEP* **06** (2018) 145 [[arXiv:1805.06460](#)] [[INSPIRE](#)].
- [44] S. Moch et al., *Low moments of the four-loop splitting functions in QCD*, *Phys. Lett. B* **825** (2022) 136853 [[arXiv:2111.15561](#)] [[INSPIRE](#)].
- [45] G. Soar, S. Moch, J.A.M. Vermaseren and A. Vogt, *On Higgs-exchange DIS, physical evolution kernels and fourth-order splitting functions at large  $x$* , *Nucl. Phys. B* **832** (2010) 152 [[arXiv:0912.0369](#)] [[INSPIRE](#)].
- [46] G. Falcioni, F. Herzog, S. Moch and A. Vogt, *Four-loop splitting functions in QCD — The quark-quark case*, *Phys. Lett. B* **842** (2023) 137944 [[arXiv:2302.07593](#)] [[INSPIRE](#)].
- [47] F. Hekhorn and G. Magni, *DGLAP evolution of parton distributions at approximate  $N^3$ LO*, [[arXiv:2306.15294](#)] [[INSPIRE](#)].
- [48] A. Candido, S. Forte and F. Hekhorn, *Can  $\overline{MS}$  parton distributions be negative?*, *JHEP* **11** (2020) 129 [[arXiv:2006.07377](#)] [[INSPIRE](#)].
- [49] K.G. Chetyrkin, J.H. Kühn and C. Sturm, *QCD decoupling at four loops*, *Nucl. Phys. B* **744** (2006) 121 [[hep-ph/0512060](#)] [[INSPIRE](#)].

- [50] Y. Schroder and M. Steinhauser, *Four-loop decoupling relations for the strong coupling*, *JHEP* **01** (2006) 051 [[hep-ph/0512058](#)] [[INSPIRE](#)].
- [51] M. Buza, Y. Matiounine, J. Smith and W.L. van Neerven, *Charm electroproduction viewed in the variable flavor number scheme versus fixed order perturbation theory*, *Eur. Phys. J. C* **1** (1998) 301 [[hep-ph/9612398](#)] [[INSPIRE](#)].
- [52] I. Bierenbaum, J. Blümlein, S. Klein and C. Schneider, *Two-Loop Massive Operator Matrix Elements for Unpolarized Heavy Flavor Production to  $O(\epsilon)$* , *Nucl. Phys. B* **803** (2008) 1 [[arXiv:0803.0273](#)] [[INSPIRE](#)].
- [53] I. Bierenbaum, J. Blümlein and S. Klein, *The Gluonic Operator Matrix Elements at  $O(\alpha_s^{**2})$  for DIS Heavy Flavor Production*, *Phys. Lett. B* **672** (2009) 401 [[arXiv:0901.0669](#)] [[INSPIRE](#)].
- [54] I. Bierenbaum, J. Blümlein and S. Klein, *Mellin Moments of the  $O(\alpha_s^3)$  Heavy Flavor Contributions to unpolarized Deep-Inelastic Scattering at  $Q^2 \gg m^2$  and Anomalous Dimensions*, *Nucl. Phys. B* **820** (2009) 417 [[arXiv:0904.3563](#)] [[INSPIRE](#)].
- [55] J. Ablinger et al., *The  $O(\alpha_s^3)$  Massive Operator Matrix Elements of  $O(n_f)$  for the Structure Function  $F_2(x, Q^2)$  and Transversity*, *Nucl. Phys. B* **844** (2011) 26 [[arXiv:1008.3347](#)] [[INSPIRE](#)].
- [56] J. Ablinger et al., *The 3-Loop Non-Singlet Heavy Flavor Contributions and Anomalous Dimensions for the Structure Function  $F_2(x, Q^2)$  and Transversity*, *Nucl. Phys. B* **886** (2014) 733 [[arXiv:1406.4654](#)] [[INSPIRE](#)].
- [57] J. Ablinger et al., *The  $O(\alpha_s^3 T_F^2)$  Contributions to the Gluonic Operator Matrix Element*, *Nucl. Phys. B* **885** (2014) 280 [[arXiv:1405.4259](#)] [[INSPIRE](#)].
- [58] A. Behring et al., *The logarithmic contributions to the  $O(\alpha_s^3)$  asymptotic massive Wilson coefficients and operator matrix elements in deeply inelastic scattering*, *Eur. Phys. J. C* **74** (2014) 3033 [[arXiv:1403.6356](#)] [[INSPIRE](#)].
- [59] J. Ablinger et al., *The transition matrix element  $A_{gq}(N)$  of the variable flavor number scheme at  $O(\alpha_s^3)$* , *Nucl. Phys. B* **882** (2014) 263.
- [60] J. Ablinger et al., *The 3-loop pure singlet heavy flavor contributions to the structure function  $F_2(x, Q^2)$  and the anomalous dimension*, *Nucl. Phys. B* **890** (2014) 48 [[arXiv:1409.1135](#)] [[INSPIRE](#)].
- [61] J. Ablinger et al., *The 3-loop pure singlet heavy flavor contributions to the structure function  $F_2(x, Q^2)$  and the anomalous dimension*, *Nucl. Phys. B* **890** (2014) 48 [[arXiv:1409.1135](#)] [[INSPIRE](#)].
- [62] J. Blümlein et al., *Heavy Flavor Wilson Coefficients in Deep-Inelastic Scattering: Recent Results*, *PoS QCDEV2017* (2017) 031 [[arXiv:1711.07957](#)] [[INSPIRE](#)].
- [63] J. Ablinger et al., *The unpolarized and polarized single-mass three-loop heavy flavor operator matrix elements  $A_{gg,Q}$  and  $\Delta A_{gg,Q}$* , *JHEP* **12** (2022) 134 [[arXiv:2211.05462](#)] [[INSPIRE](#)].
- [64] J. Ablinger et al., *The first-order factorizable contributions to the three-loop massive operator matrix elements  $A_{Qg}^{(3)}$  and  $\Delta A_{Qg}^{(3)}$* , *Nucl. Phys. B* **999** (2024) 116427 [[arXiv:2311.00644](#)] [[INSPIRE](#)].
- [65] J. Ablinger et al., *The non-first-order-factorizable contributions to the three-loop single-mass operator matrix elements  $A_{Qg}^{(3)}$  and  $\Delta A_{Qg}^{(3)}$* , *Phys. Lett. B* **854** (2024) 138713 [[arXiv:2403.00513](#)] [[INSPIRE](#)].

- [66] NNPDF collaboration, *The path to proton structure at 1% accuracy*, *Eur. Phys. J. C* **82** (2022) 428 [[arXiv:2109.02653](#)] [[INSPIRE](#)].
- [67] S. Zanolì, *Higher-order matching for heavy quarks in perturbative qcd*, MSc thesis, Università degli Studi di Milano, Milano, Italy (2020).
- [68] F. Hekhorn et al., *Heavy quarks in polarised deep-inelastic scattering at the electron-ion collider*, *Eur. Phys. J. C* **84** (2024) 189 [[arXiv:2401.10127](#)] [[INSPIRE](#)].
- [69] NNPDF collaboration, *A determination of parton distributions with faithful uncertainty estimation*, *Nucl. Phys. B* **809** (2009) 1 [Erratum *ibid.* **816** (2009) 293] [[arXiv:0808.1231](#)] [[INSPIRE](#)].
- [70] F. Hekhorn, *Towards NNPDFpol2.0*, in the proceedings of the *31st International Workshop on Deep-Inelastic Scattering and Related Subjects*, Grenoble, France, April 08–12 (2024) [[arXiv:2406.06083](#)] [[INSPIRE](#)].
- [71] NNPDF collaboration, *NNPDFpol2.0: a first global determination of polarised parton distributions at NNLO accuracy with theory uncertainties*, in preparation.
- [72] J. Ablinger et al., *Three Loop Massive Operator Matrix Elements and Asymptotic Wilson Coefficients with Two Different Masses*, *Nucl. Phys. B* **921** (2017) 585 [[arXiv:1705.07030](#)] [[INSPIRE](#)].
- [73] J. Blümlein, A. De Freitas, C. Schneider and K. Schönwald, *The Variable Flavor Number Scheme at Next-to-Leading Order*, *Phys. Lett. B* **782** (2018) 362 [[arXiv:1804.03129](#)] [[INSPIRE](#)].
- [74] S. Carrazza, E.R. Nocera, C. Schwan and M. Zaro, *PineAPPL: combining EW and QCD corrections for fast evaluation of LHC processes*, *JHEP* **12** (2020) 108 [[arXiv:2008.12789](#)] [[INSPIRE](#)].