

CLNS 95/1343

CLEO 95-9

CALTECH UC - SAN DIEGO UC - SANTA BARBARA CARLETON COLORADO  
CORNELL FLORIDA HARVARD ILLINOIS ITHACA KANSAS MCGILL  
MINNESOTA OHIO STATE OKLAHOMA PURDUE ROCHESTER  
SOUTHERN - METHODIST SYRACUSE SUNY - ALBANY VANDERBILT VIRGINIA TECH

CERN LIBRARIES, GENEVA



SCAN-9511139

509547

Observation of the Cabibbo Suppressed  
Charmed Baryon Decay  $\Lambda_c^+ \rightarrow p\phi$

**Observation of the Cabibbo Suppressed  
Charmed Baryon Decay  $\Lambda_c^+ \rightarrow p\phi$**

J.P. Alexander,<sup>1</sup> C. Bebek,<sup>1</sup> B.E. Berger,<sup>1</sup> K. Berkelman,<sup>1</sup> K. Bloom,<sup>1</sup> T.E. Browder,<sup>1\*</sup>  
 D.G. Cassel,<sup>1</sup> H.A. Cho,<sup>1</sup> D.M. Coffman,<sup>1</sup> D.S. Crowcroft,<sup>1</sup> M. Dickson,<sup>1</sup> P.S. Drell,<sup>1</sup>  
 D.J. Dumas,<sup>1</sup> R. Ehrlich,<sup>1</sup> R. Elia,<sup>1</sup> P. Gaidarev,<sup>1</sup> M. Garcia-Sciveres,<sup>1</sup> B. Gittelman,<sup>1</sup>  
 S.W. Gray,<sup>1</sup> D.L. Hartill,<sup>1</sup> B.K. Heltsley,<sup>1</sup> S. Henderson,<sup>1</sup> C.D. Jones,<sup>1</sup> S.L. Jones,<sup>1</sup>  
 J. Kandaswamy,<sup>1</sup> N. Katayama,<sup>1</sup> P.C. Kim,<sup>1</sup> D.L. Kreinick,<sup>1</sup> T. Lee,<sup>1</sup> Y. Liu,<sup>1</sup> G.S. Ludwig,<sup>1</sup>  
 J. Masui,<sup>1</sup> J. Mevissen,<sup>1</sup> N.B. Mistry,<sup>1</sup> C.R. Ng,<sup>1</sup> E. Nordberg,<sup>1</sup> J.R. Patterson,<sup>1</sup> D. Peterson,<sup>1</sup>  
 D. Riley,<sup>1</sup> A. Soffer,<sup>1</sup> P. Avery,<sup>2</sup> A. Freyberger,<sup>2</sup> K. Lingel,<sup>2</sup> C. Prescott,<sup>2</sup> J. Rodriguez,<sup>2</sup>  
 S. Yang,<sup>2</sup> J. Yelton,<sup>2</sup> G. Brandenburg,<sup>3</sup> D. Cinabro,<sup>3</sup> T. Liu,<sup>3</sup> M. Saulnier,<sup>3</sup> R. Wilson,<sup>3</sup>  
 H. Yamamoto,<sup>3</sup> T. Bergfeld,<sup>4</sup> B.I. Eisenstein,<sup>4</sup> J. Ernst,<sup>4</sup> G.E. Gladding,<sup>4</sup> G.D. Gollin,<sup>4</sup>  
 M. Palmer,<sup>4</sup> M. Selen,<sup>4</sup> J.J. Thaler,<sup>4</sup> K.W. Edwards,<sup>5</sup> K.W. McLean,<sup>5</sup> M. Ogg,<sup>5</sup> A. Bellerive,<sup>6</sup>  
 D.I. Britton,<sup>6</sup> E.R.F. Hyatt,<sup>6</sup> R. Janicek,<sup>6</sup> D.B. MacFarlane,<sup>6</sup> P.M. Patel,<sup>6</sup> B. Spaan,<sup>6</sup>  
 A.J. Sadoff,<sup>7</sup> R. Ammar,<sup>8</sup> P. Baringer,<sup>8</sup> A. Bean,<sup>8</sup> D. Besson,<sup>8</sup> D. Coppage,<sup>8</sup> N. Coptly,<sup>8</sup>  
 R. Davis,<sup>8</sup> N. Hancock,<sup>8</sup> S. Kotov,<sup>8</sup> I. Kravchenko,<sup>8</sup> N. Kwak,<sup>8</sup> Y. Kubota,<sup>9</sup> M. Lattery,<sup>9</sup>  
 M. Momayezi,<sup>9</sup> J.K. Nelson,<sup>9</sup> S. Patton,<sup>9</sup> R. Poling,<sup>9</sup> V. Savinov,<sup>9</sup> S. Schrenk,<sup>9</sup> R. Wang,<sup>9</sup>  
 M.S. Alam,<sup>10</sup> I.J. Kim,<sup>10</sup> Z. Ling,<sup>10</sup> A.H. Mahmood,<sup>10</sup> J.J. O'Neill,<sup>10</sup> H. Severini,<sup>10</sup> C.R. Sun,<sup>10</sup>  
 F. Wappler,<sup>10</sup> G. Crawford,<sup>11</sup> R. Fulton,<sup>11</sup> D. Fujino,<sup>11</sup> K.K. Gan,<sup>11</sup> K. Honscheid,<sup>11</sup> H. Kagan,<sup>11</sup>  
 R. Kass,<sup>11</sup> J. Lee,<sup>11</sup> M. Sung,<sup>11</sup> C. White,<sup>11</sup> A. Wolf,<sup>11</sup> M.M. Zoeller,<sup>11</sup> X. Fu,<sup>12</sup> B. Nemati,<sup>12</sup>  
 W.R. Ross,<sup>12</sup> P. Skubic,<sup>12</sup> M. Wood,<sup>12</sup> M. Bishai,<sup>13</sup> J. Fast,<sup>13</sup> E. Gerndt,<sup>13</sup> J.W. Hinson,<sup>13</sup>  
 T. Miao,<sup>13</sup> D.H. Miller,<sup>13</sup> M. Modesitt,<sup>13</sup> E.I. Shibata,<sup>13</sup> I.P.J. Shipsey,<sup>13</sup> P.N. Wang,<sup>13</sup>  
 L. Gibbons,<sup>14</sup> S.D. Johnson,<sup>14</sup> Y. Kwon,<sup>14</sup> S. Roberts,<sup>14</sup> E.H. Thorndike,<sup>14</sup> T.E. Coan,<sup>15</sup>  
 J. Dominick,<sup>15</sup> V. Fadayev,<sup>15</sup> I. Korolkov,<sup>15</sup> M. Lambrecht,<sup>15</sup> S. Sanghera,<sup>15</sup> V. Shelkov,<sup>15</sup>  
 T. Skwarnicki,<sup>15</sup> R. Stroynowski,<sup>15</sup> I. Volobouev,<sup>15</sup> G. Wei,<sup>15</sup> M. Artuso,<sup>16</sup> M. Gao,<sup>16</sup>  
 M. Goldberg,<sup>16</sup> D. He,<sup>16</sup> N. Horwitz,<sup>16</sup> S. Kopp,<sup>16</sup> G.C. Moneti,<sup>16</sup> R. Mountain,<sup>16</sup> F. Muheim,<sup>16</sup>  
 Y. Mukhin,<sup>16</sup> S. Playfer,<sup>16</sup> S. Stone,<sup>16</sup> X. Xing,<sup>16</sup> J. Bartelt,<sup>17</sup> S.E. Csorna,<sup>17</sup> V. Jain,<sup>17</sup>  
 S. Marka,<sup>17</sup> D. Gibaut,<sup>18</sup> K. Kinoshita,<sup>18</sup> P. Pomianowski,<sup>18</sup> B. Barish,<sup>19</sup> M. Chadha,<sup>19</sup>  
 S. Chan,<sup>19</sup> D.F. Cowen,<sup>19</sup> G. Eigen,<sup>19</sup> J.S. Miller,<sup>19</sup> C. O'Grady,<sup>19</sup> J. Urheim,<sup>19</sup> A.J. Weinstein,<sup>19</sup>  
 F. Würthwein,<sup>19</sup> D.M. Asner,<sup>20</sup> M. Athanas,<sup>20</sup> D.W. Bliss,<sup>20</sup> W.S. Brower,<sup>20</sup> G. Masek,<sup>20</sup>  
 H.P. Paar,<sup>20</sup> J. Gronberg,<sup>21</sup> C.M. Korte,<sup>21</sup> R. Kutschke,<sup>21</sup> S. Menary,<sup>21</sup> R.J. Morrison,<sup>21</sup>  
 S. Nakanishi,<sup>21</sup> H.N. Nelson,<sup>21</sup> T.K. Nelson,<sup>21</sup> C. Qiao,<sup>21</sup> J.D. Richman,<sup>21</sup> D. Roberts,<sup>21</sup>  
 A. Ryd,<sup>21</sup> H. Tajima,<sup>21</sup> M.S. Witherell,<sup>21</sup> R. Balest,<sup>22</sup> K. Cho,<sup>22</sup> W.T. Ford,<sup>22</sup> M. Lohner,<sup>22</sup>  
 H. Park,<sup>22</sup> P. Rankin,<sup>22</sup> and J.G. Smith<sup>22</sup>

<sup>1</sup> *Cornell University, Ithaca, New York 14853*

<sup>2</sup> *University of Florida, Gainesville, Florida 32611*

<sup>3</sup> *Harvard University, Cambridge, Massachusetts 02138*

<sup>4</sup> *University of Illinois, Champaign-Urbana, Illinois, 61801*

<sup>5</sup> *Carleton University, Ottawa, Ontario K1S 5B6 and the Institute of Particle Physics, Canada*

<sup>6</sup> *McGill University, Montréal, Québec H3A 2T8 and the Institute of Particle Physics, Canada*

<sup>7</sup> *Ithaca College, Ithaca, New York 14850*

<sup>8</sup> *University of Kansas, Lawrence, Kansas 66045*

- <sup>9</sup> *University of Minnesota, Minneapolis, Minnesota 55455*  
<sup>10</sup> *State University of New York at Albany, Albany, New York 12222*  
<sup>11</sup> *Ohio State University, Columbus, Ohio, 43210*  
<sup>12</sup> *University of Oklahoma, Norman, Oklahoma 73019*  
<sup>13</sup> *Purdue University, West Lafayette, Indiana 47907*  
<sup>14</sup> *University of Rochester, Rochester, New York 14627*  
<sup>15</sup> *Southern Methodist University, Dallas, Texas 75275*  
<sup>16</sup> *Syracuse University, Syracuse, New York 13244*  
<sup>17</sup> *Vanderbilt University, Nashville, Tennessee 37235*  
<sup>18</sup> *Virginia Polytechnic Institute and State University, Blacksburg, Virginia, 24061*  
<sup>19</sup> *California Institute of Technology, Pasadena, California 91125*  
<sup>20</sup> *University of California, San Diego, La Jolla, California 92093*  
<sup>21</sup> *University of California, Santa Barbara, California 93106*  
<sup>22</sup> *University of Colorado, Boulder, Colorado 80309-0390*

### Abstract

We report the observation of the Cabibbo-suppressed decays  $\Lambda_c^+ \rightarrow pK^-K^+$  and  $\Lambda_c^+ \rightarrow p\phi$  using data collected with the CLEO II detector at CESR. The latter mode, observed for the first time with significant statistics, is of interest as a test of color-suppression in charm decays. We have determined the branching ratios for these modes relative to  $\Lambda_c^+ \rightarrow pK^-\pi^+$  and compared our results with theory.

The strength of color-suppression in internal W-emission charmed meson decays has long been in question. For example,  $B(D_s^+ \rightarrow \bar{K}^{*0}K^+)/B(D_s^+ \rightarrow \phi\pi^+) \simeq 1$  [1,2], while the expectation from color-matching requirements is that this ratio should be about 1/9. Reasonable overall agreement with the experimental data in the charm sector has been obtained using factorization and taking the large  $N_c$  limit in a  $1/N_c$  expansion approach, where  $N_c$  is the number of quark colors [3,4]. The Cabibbo-suppressed charmed baryon decay  $\Lambda_c^+ \rightarrow p\phi$ , shown in Figure 1, is also naively expected to be color-suppressed. However, using factorization and taking the limit  $N_c \rightarrow \infty$  leads to a prediction of no color-suppression [5]. Since the  $\Lambda_c^+ \rightarrow p\phi$  decay receives contributions only from factorizable diagrams, a reliable calculation should be obtained using factorization. Observation of the  $\Lambda_c^+ \rightarrow p\phi$  decay was first reported by the ACCMOR collaboration with  $2.8 \pm 1.9$  events [9]. Last year the E687 collaboration published results on the first observation of the Cabibbo-suppressed

---

\*Permanent address: University of Hawaii at Manoa

charmed baryon decay  $\Lambda_c^+ \rightarrow pK^-K^+$ , along with an upper limit on the resonant substructure  $\Lambda_c^+ \rightarrow p\phi$  [10]. Herein we present new CLEO results on the observation of  $\Lambda_c^+ \rightarrow pK^-K^+$  and  $\Lambda_c^+ \rightarrow p\phi$  decays and discuss the implications of the results.

We use a data sample recorded with the CLEO II detector operating at the Cornell Electron Storage Ring (CESR). The sample consists of  $e^+e^-$  annihilations taken at and slightly below the  $\Upsilon(4S)$  resonance, for a total integrated luminosity of  $3.46 \text{ fb}^{-1}$ . The main detector components which are important for this analysis are the tracking system and the barrel Time-of-Flight (TOF) particle identification system. Additional particle ID is provided by specific ionization ( $dE/dx$ ) information from the tracking system's main drift chamber. A more detailed description of the CLEO II detector has been provided elsewhere [11].

To search for the  $\Lambda_c^+$  signals, we study  $pK^-K^+$  track combinations found by the tracking system. The  $p$  and  $K^\pm$  candidates are identified by combining information from the TOF and  $dE/dx$  systems to form a combined  $\chi^2$  probability  $\mathcal{P}_i$  for each mass hypothesis  $i = \pi, K, p$ . Using these probabilities  $\mathcal{P}_i$ , a normalized probability ratio  $L_i$  is evaluated for each track according to the formula:  $L_i \equiv \mathcal{P}_i / (\mathcal{P}_\pi + \mathcal{P}_K + \mathcal{P}_p)$ . Well-identified protons form a sharp peak near  $L_p = 1$ , while tracks identified as *not* being protons form a peak near  $L_p = 0$ . The remainder of the candidates fall in the region between 0 and 1. For the proton involved in each decay mode under study, we require  $L_p > 0.9$ , which constitutes a strong cut. For the kaons, we apply a loose cut of  $L_K > 0.1$ . In addition, all protons and kaons must pass a minimum requirement of  $\mathcal{P}_p > 0.001$  and  $\mathcal{P}_K > 0.001$ , respectively. In order to reduce the large combinatoric background, the candidate  $\Lambda_c^+$  scaled momentum  $x_p = P_{\Lambda_c} / \sqrt{E_{beam}^2 - m_{\Lambda_c}^2}$  is limited to  $x_p > 0.5$ .

The  $pK^-K^+$  invariant mass is shown in Figure 2. The broad enhancement in the mass region above  $2.37 \text{ GeV}/c^2$  is a reflection from the decay mode  $\Lambda_c^+ \rightarrow pK^-\pi^+$ , where the pion has been misidentified as a kaon. The spectrum is fitted to a Gaussian for the signal with width fixed to  $\sigma = 4.9 \text{ MeV}/c^2$  determined from Monte Carlo simulation [12], and a 2<sup>nd</sup> order Chebychev polynomial for the smooth background. This fit yields  $214 \pm 50$  events for the inclusive  $\Lambda_c^+ \rightarrow pK^-K^+$  signal with a mean mass of  $2285.5 \pm 1.2 \text{ MeV}/c^2$  [13].

To find a  $\Lambda_c^+ \rightarrow p\phi$  signal, we reconstruct  $\phi$  candidates through their decays  $\phi \rightarrow K^-K^+$ .

Because the width of the  $\phi$  is comparable to the detector mass resolution, the  $\phi$  signal shape is best described by a convolution of a Gaussian and a Breit-Wigner of width  $\Gamma = 4.43 \text{ MeV}/c^2$  [1]. The background is parameterized by a function of the form  $b(m) = N(m - m_0)^\alpha e^{\beta(m - m_0)}$ . The measured Gaussian resolution from the fit is  $\sigma = 1.6 \pm 0.2 \text{ MeV}/c^2$ . In order to perform background subtractions,  $1.0121 < m_{KK} < 1.0273 \text{ GeV}/c^2$  is designated as the  $\phi$  “signal” region, while  $0.990 < m_{KK} < 1.005 \text{ GeV}/c^2$  and  $1.035 < m_{KK} < 1.050 \text{ GeV}/c^2$  are designated as the “sideband” regions. Integrating the background function over the sideband and signal regions gives a signal-to-sideband scale factor  $R_\phi = 0.560 \pm 0.016$ , which is used in the  $\phi$  background subtraction below.

In order to obtain the  $\Lambda_c^+ \rightarrow p\phi$  signal, the  $pK^-K^+$  mass plot is made both for  $m_{K^-K^+}$  in the  $\phi$  signal region and the  $\phi$  sideband regions. Figure 3 shows the results. The spectra are fitted to a Gaussian for the signal with width fixed to  $\sigma = 4.9 \text{ MeV}/c^2$  from Monte Carlo, and a 2nd order Chebychev polynomial for the smooth background. The fit to the  $pK^-K^+$  mass spectrum corresponding to the  $\phi$  signal region yields  $54 \pm 12$  events with a confidence level of 97%. The mean mass for the signal is measured to be  $2288.2 \pm 1.3 \text{ MeV}/c^2$ . In fitting the  $pK^-K^+$  mass corresponding to the  $\phi$  sideband region, the mean  $\Lambda_c^+$  mass is fixed to that obtained from the  $\phi$  signal region and the  $\sigma$  is fixed to the Monte Carlo value as before. This gives  $-16.4 \pm 9.6$  events for the  $\phi$ -sideband  $\Lambda_c^+$  yield. Since the true contribution must be positive-definite, we set the central value to zero and use  $0 \pm 9.6$  as the best estimate of the  $\Lambda_c^+ \rightarrow pK^-K^+$  contribution. After scaling this by  $R_\phi$  and subtracting, we find that the net  $\Lambda_c^+ \rightarrow p\phi$  yield is  $54 \pm 13$  events.

As a check of the non-resonant contribution to the  $\Lambda_c^+ \rightarrow p\phi$  signal, we fit the  $K^-K^+$  mass spectra corresponding to the  $\Lambda_c^+$  signal and sideband regions as determined from the inclusive  $pK^-K^+$  mass spectrum. The  $\phi$  yield obtained from the  $\Lambda_c^+$  sideband regions,  $2.246 < m_{pKK} < 2.266$  and  $2.306 < m_{pKK} < 2.326 \text{ GeV}/c^2$ , is subtracted from that for the  $\Lambda_c^+$  signal region,  $2.276 < m_{pKK} < 2.296 \text{ GeV}/c^2$ . Figure 4 shows the fits to the  $K^-K^+$  spectra from the  $\Lambda_c^+$  signal and sideband regions, which yield  $\phi$  signals of  $92.2 \pm 17.0$  events and  $36.5 \pm 13.5$  events, respectively. The  $\Lambda_c^+$  sideband  $K^-K^+$  mass spectrum in the Figure 4 has been scaled by the  $\Lambda_c^+$  signal-to-sideband scale factor of  $R_{\Lambda_c^+} = 0.502 \pm 0.013$ , obtained by integrating the background

function in Figure 2 over the  $\Lambda_c^+$  signal and sideband regions. This gives  $56 \pm 22$  events for the  $\Lambda_c^+ \rightarrow p\phi$  signal, which is in agreement with the first method.

A check is also made for a possible reflection from  $D_s^+ \rightarrow \phi\pi^+$ , where the pion is misidentified as a proton. It is found that the reflection is a broad enhancement in the mass region above the signal. The effect of this background is minimized by the tight particle-ID requirement on the proton. Consequently, the overall fake rate is less than 1%, causing negligible effect on the  $\Lambda_c^+ \rightarrow p\phi$  signal yield from the fit.

The decay  $\Lambda_c^+ \rightarrow pK^-\pi^+$  is used as the normalization mode for the  $\Lambda_c^+ \rightarrow p\phi$  relative branching ratio. In finding the  $\Lambda_c^+ \rightarrow pK^-\pi^+$  yield, the same cuts are applied as in the  $\Lambda_c^+ \rightarrow pK^-K^+$  analysis to minimize systematic errors, except that the particle-ID for the  $\pi^+$  is loosened to a consistency requirement:  $\mathcal{P}_\pi > 0.001$ . The  $\Lambda_c^+ \rightarrow pK^-\pi^+$  mass spectrum is shown in Figure 5. The parameterization of the fit is the same as the  $\Lambda_c^+ \rightarrow p\phi$  mass fit in Figure 3, except that the width of the Gaussian is allowed to vary. The fit yields  $5683 \pm 138$  observed signal events with a mean of  $2286.8 \pm 0.2$  MeV/ $c^2$  and a width of  $6.4 \pm 0.2$  MeV/ $c^2$ . If the width of the Gaussian is fixed to the Monte Carlo prediction of 5.8 MeV/ $c^2$ , the yield changes by 4%. This dependence is included in the systematic error.

Monte Carlo simulation is used to determine all aspects of the detection efficiency except particle-ID. The particle-ID efficiency for protons is obtained using a sample of 33000  $\Lambda \rightarrow p\pi^-$  decays with a signal-to-background ratio of 50:1 [14]. For protons thus identified, the momentum spectrum after the particle-ID cuts ( $L_p > 0.9$ ,  $\mathcal{P}_p > 0.001$ ) is divided by the momentum spectrum before these cuts, bin by bin, yielding the particle-ID efficiencies versus momentum. To calculate the detection efficiency, the measured efficiency is folded in by randomly rejecting the corresponding fraction of Monte Carlo tracks in each momentum bin. The particle-ID ( $L_K > 0.1$ ,  $\mathcal{P}_K > 0.001$ ) efficiency for the kaons is derived in an analogous manner, except that the kaons are taken from  $D^*$  decays through the cascade process  $D^{*+} \rightarrow D^0\pi^+$ ,  $D^0 \rightarrow K^-\pi^+$ . A sample of 11000 such  $D^0 \rightarrow K^-\pi^+$  decays is obtained with an 8:1 signal-to-background ratio [14]. The particle-ID efficiency for protons is near 90% from 300 MeV/ $c^2$  to 1.1 GeV/ $c^2$  falling off to below 10% by 2.5 GeV/ $c^2$ . For kaons the particle-ID efficiency remains relatively flat at about 95%.

Using a Monte Carlo sample of  $\Lambda_c^+ \rightarrow p\phi$  decays, where the  $\Lambda_c^+$  fragmentation takes place according to the Lund JETSET Monte Carlo [15], the full detection efficiency is determined, with the particle-ID portion folded in as described above. For  $\Lambda_c^+ \rightarrow p\phi$ , the overall efficiency is  $0.178 \pm 0.004$  including the particle-ID efficiency which is  $0.425 \pm 0.011$ . For  $\Lambda_c^+ \rightarrow pK^-K^+$  and  $\Lambda_c^+ \rightarrow pK^-\pi^+$  the overall efficiencies are  $0.216 \pm 0.005$  and  $0.224 \pm 0.005$ , respectively.

Since for all the decay modes the requirement  $x_p > 0.5$  is applied, the relative branching ratio for each mode is found simply by dividing the corrected yields. Table I gives the details, listing only the statistical errors. The  $\phi \rightarrow K^-K^+$  branching ratio is explicitly included in the calculation of the  $\Lambda_c^+ \rightarrow p\phi$  branching ratio, and its uncertainty is included in the systematic errors.

The estimates for the main sources of systematic error include the  $\Lambda_c^+ \rightarrow p\phi$  and  $\Lambda_c^+ \rightarrow pK^-K^+$  signal shapes (7% and 11%, respectively) and background shapes (2% and 10%, respectively), particle-ID efficiency (6%), and the  $\Lambda_c^+ \rightarrow pK^-\pi^+$  fit (4%). In addition, for the  $\Lambda_c^+ \rightarrow p\phi$  mode, varying the  $\phi$  signal and sideband regions gives a 5% variation in the yield. Finally, there is a 1.8% contribution to the  $\Lambda_c^+ \rightarrow p\phi$  systematic error from the  $\phi \rightarrow K^-K^+$  branching ratio uncertainty. Thus we estimate 12% systematic error in  $B(p\phi)/B(pK\pi)$ , 17% in  $B(pKK)/B(pK\pi)$ , and 18% in  $B(p\phi)/B(pKK)$ . The final results appear in Table II, along with those from NA32 [9] and E687 [10] and theoretical predictions from Cheng and Tseng [5], Körner and Krämer [6], Żenczykowski [7], and Datta [8]. From Table I we also find  $B(\Lambda_c^+ \rightarrow pK^-K^+[\text{non-}\phi]) = 0.029 \pm 0.010 \pm 0.005$  for  $\Lambda_c^+ \rightarrow pK^-K^+$  decays not arising from  $\Lambda_c^+ \rightarrow p\phi$ .

In summary, we have observed the Cabibbo-suppressed decays  $\Lambda_c^+ \rightarrow p\phi$  and  $\Lambda_c^+ \rightarrow pK^-K^+$ . The results appear in Table II, which show that the phenomenological treatments of the  $\Lambda_c^+ \rightarrow p\phi$  decay rate agree within a factor of two or three with our result. Our measured branching ratio  $B(p\phi)/B(pKK)$  is consistent with the E687 upper limit, while our measurement of  $B(pKK)/B(pK\pi)$  differs from the E687 result by 1.7 sigma. Within the factorization approach using a  $1/N_c$  expansion, our result supports the validity of taking the large  $N_c$  limit in charm baryon decays.

## ACKNOWLEDGEMENTS

We gratefully acknowledge the effort of the CESR staff in providing us with excellent luminosity

and running conditions. This work was supported by the National Science Foundation, the U.S. Department of Energy, the Heisenberg Foundation, the Alexander von Humboldt Stiftung, the Natural Sciences and Engineering Research Council of Canada, and the A.P. Sloan Foundation.

---

- [1] Particle Data Group, L. Monatet *et al.*, Phys. Rev. D **50**, 1173 (1994).
- [2] Unless otherwise specified, reference to a state also implies refernce to the charge conjugate state.
- [3] A. J. Buras, J.-M. Gérard, and R. Rückl, Nuc. Phys. **B268**, 16 (1986).
- [4] M. Bauer, B. Stech, and M. Wirbel, Z. Phys. C **34**, 103 (1987).
- [5] H. Cheng and B. Tseng, Phys. Rev. D **46**, 1042 (1992).
- [6] J.G. Körner and M. Krämer, Z. Phys. C **55**, 659 (1992).
- [7] P. Żenczykowski, Phys. Rev. D **50**, 402 (1994).
- [8] A. Datta, UH-511-824-95, April 1995.
- [9] S. Barlag *et al.*, Z. Phys. C **48**, 29 (1990).
- [10] P.L. Frabetti *et al.*, Phys. Lett. B 314, 477-481 (1993).
- [11] Y. Kubota *et al.*, Nucl. Instrum. Methods **A320**, 66-113 (1992).
- [12] The Monte Carlo employs the CERN GEANT package:  
R. Brun *et al.*, GEANT 3.14, CERN DD/EE/84-1.
- [13] The quoted uncertainties in mass measurements refer to statistical error only.
- [14] The remaining background is removed by sideband subtraction.
- [15] T. Sjöstrand, Comp. Phys. Comm., **43** 367 (1987).



TABLE I. Calculation of the branching ratios for  $\Lambda_c^+ \rightarrow p\phi$  and  $\Lambda_c^+ \rightarrow pK^-K^+$  relative to  $\Lambda_c^+ \rightarrow pK^-\pi^+$  and  $\Lambda_c^+ \rightarrow pK^-K^+$ . The errors are statistical only.

Decay Mode:	$\Lambda_c^+ \rightarrow p\phi$	$\Lambda_c^+ \rightarrow pK^-K^+$	$\Lambda_c^+ \rightarrow pK^-\pi^+$
Raw Yield	$54 \pm 13$	$214 \pm 50$	$5683 \pm 138$
Efficiency	$0.178 \pm 0.004$	$0.216 \pm 0.005$	$0.224 \pm 0.005$
$B(\phi \rightarrow K^-K^+)$	$0.491 \pm 0.005$		
Corr. Yield	$618 \pm 138$	$991 \pm 233$	$25371 \pm 837$
$B/B(pK^-\pi^+)$	$0.024 \pm 0.006$	$0.039 \pm 0.009$	1
$B/B(pK^-K^+)$	$0.62 \pm 0.20$	1	

TABLE II. Final results on  $\Lambda_c^+ \rightarrow p\phi$  and  $\Lambda_c^+ \rightarrow pK^-K^+$ .

Ratio of interest:	$B(p\phi)/B(pK^-\pi^+)$	$B(p\phi)/B(pK^-K^+)$	$B(pK^-K^+)/B(pK^-\pi^+)$
This experiment	$0.024 \pm 0.006 \pm 0.003$	$0.62 \pm 0.20 \pm 0.12$	$0.039 \pm 0.009 \pm 0.007$
NA32	$0.04 \pm 0.03$		
E687		$< 0.58@90\%C.L.$	$0.096 \pm 0.029 \pm 0.010$
Cheng & Tseng	$0.045 \pm 0.011$		
Żenczykowski	0.023		
Datta	0.01		
Körner& Krämer	0.05		

3250795-005



FIG. 1. The decay  $\Lambda_c^+ \rightarrow p\phi$ .

FIG. 2. Invariant mass of inclusive  $pK^-K^+$  combinations passing all requirements. No  $\phi$  cut is applied. The region above 2.37  $\text{GeV}/c^2$ , where there is a large enhancement from  $\Lambda_c^+ \rightarrow pK^-\pi^+$  decays, is not included in the fit.




FIG. 3. Invariant mass of  $pK^-K^+$  combinations corresponding to  $K^-K^+$  mass in the  $\phi$  signal and sideband regions.




FIG. 4. Fit to  $K^-K^+$  mass from combinations belonging to the  $\Lambda_c^+$  signal and sideband regions. The region above 1.06 GeV/c<sup>2</sup> is not included in the fit because of  $K^{*0}$  feed-up when the  $\pi$  is misidentified as a  $K$ .




FIG. 5. Invariant mass of  $pK^-\pi^+$  combinations found in the same data sample. The  $\Lambda_c^+ \rightarrow pK^-\pi^+$  signal is used for normalization of the  $\Lambda_c^+ \rightarrow p\phi$  branching ratio.