

Centrality dependence of K^+ produced in Pb+Pb collisions at 158 GeV per nucleon

‡

Sonja Kabana for the NA52 collaboration:

G. Ambrosini¹, R. Arsenescu¹, C. Baglin³, H P Beck¹, K. Borer¹,
 A. Bussière³, K. Elsener², Ph. Gorodetzky⁵, J.P. Guillaud³, P. Hess¹,
 S. Kabana¹, R. Klingenberg¹, G Lehmann¹, T. Lindén⁴, K.D. Lohmann²,
 R. Mommsen¹, U. Moser¹, K. Pretzl¹, J. Schacher¹, F. Stoffel¹,
 R Spiwoaks¹, J. Tuominiemi⁴, M. Weber¹

¹ Laboratory for High Energy Physics, University of Bern, Sidlerstrasse 5,
 CH-3012 Bern, Switzerland,

² CERN, CH-1211 Geneva 23, Switzerland,

³ CNRS-IN2P3, LAPP Annecy, F-74941 Annecy-le-Vieux, France,

⁴ Dept. of Physics and Helsinki Institute of Physics, University of Helsinki, PO
 Box 9, FIN-00014 Helsinki, Finland,

⁵ PCC-College de France, 11 place Marcelin Berthelot, 75005 Paris, France

Abstract.

The NA52 collaboration searches for a discontinuous behaviour of charged kaons produced in Pb+Pb collisions at 158 A GeV as a function of the impact parameter, which could reveal a QGP phase transition. The K^+ yield is found to grow proportional to the number of participating ('wounded') nucleons N , above $N=100$. Previous NA52 data agree with the above finding and show a discontinuous behaviour in the kaon centrality dependence near $N=100$, marking the onset of strangeness enhancement -over e.g. p+A data at the same \sqrt{s} - in a chemically equilibrated phase.

1. Introduction

The quark-gluon plasma phase transition predicted by QCD [1] may occur and manifest itself in ultrarelativistic heavy ion collisions through discontinuities in the temperature and energy density dependence of relevant observables. A major example of such a discontinuity is seen in the J/Ψ to Drell Yan ratio [2]. The NA52 collaboration searches for discontinuities in strangeness production measuring charged kaons as a function of the impact parameter. Results from the 1995 NA52 run are published in [3]. We report here on new preliminary results from the 1998 run of the NA52 experiment, on K^+ at rapidity 4.1 and transverse momentum near 0 produced in Pb+Pb collisions at 158 A GeV. In this run a new electromagnetic lead/quartz fiber calorimeter (QFC) with improved acceptance and resolution [4] was used.

‡ Work supported in part by the Schweizerischer Nationalfonds.




Figure 1. Preliminary NA52 data from Pb+Pb collisions at 158 A GeV. (a) Calorimetric energy distribution in arbitrary units. (b): K^+ yield in arb. units per N, as a function of N. For comparison the 1995 NA52 data [3] are also shown. The 1998 data are scaled by 2.4 with respect to the 1995 data.

2. Results and discussion

For the kaon measurement we modified the 1998 set up of NA52 [5] by placing the target 0.6 m upstream of the calorimeter. The results have been corrected for empty target contributions. The number of participant nucleons N has been estimated from the energy measured with the calorimeter (figure 1, (a)) in the way described in [3]. Particle identification is described in [3] and references there. The positive kaon yield divided by N is independent of N, for $N > 100$ (figure 1, (b)) in agreement with previous NA52 results [3]. Assuming that N is proportional to the volume of the particle source, figure 1, (b) shows that the kaon number density exhibit a discontinuity, saturating above $N=100$. This indicates a transition to a phase characterized by a high degree of chemical equilibrium and enhancement [3] of kaons from the point $N=100$ on, corresponding to energy density $\epsilon \sim 1.3 \text{ GeV}/\text{fm}^3$ [6], near ϵ_c [1, 7]. This change marks the onset of strangeness enhancement seen in kaons in an equilibrated phase, which may be suggestive for a QCD phase transition, depending on the simultaneous appearance of thresholds in other signatures at the relevant ϵ values and their theoretical understanding [7, 2, 6].

References

- [1] E. Laermann, Nucl. Phys. A610 (1996) 1c.
- [2] M.C. Abreu et al. (NA50 coll.), CERN-EP-2000-013.
- [3] G. Ambrosini et al. (NA52 Coll.), N. J. of Phys. (1999) 1, 22, G. Ambrosini et al. (NA52 Coll.), N. J. of Phys. (1999) 1, 23, S. Kabana et al. (NA52 Coll.), Nucl. Phys. A 661 (99) 370c.
- [4] M. Weber et al (NA52 Coll.), CALOR97, World Scientific (1998) 151.
- [5] M. Weber et al (NA52 Coll.), these proceedings and Un. of Bern preprint BUHE-00-04.
- [6] S. Kabana, hep-ph-0004138 and hep-ph/0010228.
- [7] H. Satz, hep-ph/0007069, P.B. Munzinger, J. Stachel, nucl-th/0007059.