

MEASUREMENT OF BOSON SELF COUPLINGS AT LEP AND SEARCH FOR ANOMALIES

Martin Weber

III. Physikalisches Institut, Physikzentrum, RWTH Aachen, D-52056 Aachen, Germany



With center of mass energies up to 207 GeV of LEP II, massive W and Z bosons can be produced in pairs and jointly with photons. This allows to study boson-boson couplings. Since the W and Z bosons are unstable and decay into fermions, two- and four-fermion final states, accompanied possibly by photons, play an important role for these measurements. The couplings of the W to other bosons have been measured to be $g_1^Z = 0.990_{-0.024}^{+0.023}$, $\kappa_\gamma = 0.896_{-0.056}^{+0.058}$, and $\lambda_\gamma = -0.023_{-0.023}^{+0.025}$. They are in agreement with the Standard Model expectation of $g_1^Z = 1$, $\kappa_\gamma = 1$, and $\lambda_\gamma = 0$. No sign for couplings of three neutral bosons, parametrized by $f_{4,5}^{\gamma,Z}$ and $h_{1,2,3,4}^{\gamma,Z}$, and for anomalous couplings of four gauge bosons, parametrized by a_0 , a_n and a_c has been found.

1 Couplings of the W to other bosons

The $SU(2)_L \times U(1)_Y$ symmetry of the Standard Model predicts the pair production of W bosons through abelian and non-abelian graphs. On the left side of Fig. 1, the three Standard Model feynman diagrams for W pair production are shown.

As has been measured by the LEP experiments, all three diagrams are needed to describe the data. This can be seen on the right of Fig. 1. Using only the single abelian graph (the neutrino exchange) or neglecting the non-abelian Z exchange graph, data and theory disagree. But still the contribution of the graphs could differ from the Standard Model prediction, and therefore a more sophisticated method is performed to analyze the non-abelian gauge sector.

To study possible other contributions, the Lagrangian for the VWW vertex ($V=Z,\gamma$) can be written in the most general Lorentz invariant form¹

$$\begin{aligned}
 i\mathcal{L}^{WWV}/g_{WWV} &= g_1^V \left(W_{\mu\nu}^\dagger W^\mu V^\nu - W_\mu^\dagger V_\nu W^{\mu\nu} \right) + \kappa_V W_\mu^\dagger W_\nu V^{\mu\nu} \\
 &+ \frac{\lambda_V}{m_W^2} W_{\mu\nu}^\dagger W_\rho^\nu V^{\rho\mu} + \mathcal{C} + \mathcal{P} + \mathcal{CP},
 \end{aligned}$$




Figure 1: Feynman graphs (left) and measured cross-section (right) for the pair production of W bosons.

where C, P and CP-violating terms are not shown and assumed to vanish in the following discussion. To further reduce the parameter set from six to three free couplings, firstly $U(1)_{\text{em}}$ gauge invariance is required, fixing the charge of the W boson to $q_W = \pm 1$, which is equivalent to $g_1^\gamma = 1$. Secondly, the requirement of $SU(2)_L \times U(1)_Y$ symmetry of the Lagrangian leads to the two constraints $\kappa_Z = g_1^Z - (\kappa_\gamma - 1) \tan^2 \theta_W$ and $\lambda_Z = \lambda_\gamma$. The three parameters left are g_1^Z , κ_γ and λ_γ . In the Standard Model, their values are predicted to be $g_1^Z = 1$, $\kappa_\gamma = 1$ and $\lambda_\gamma = 0$. Often one finds in the literature also the differences to the Standard Model expectations: $\Delta g_1^Z = g_1^Z - 1$ and $\Delta \kappa_\gamma = \kappa_\gamma - 1$.

The couplings are not only accessible in W pair production, but also in single W and single photon production, which also involve the γWW vertex, as can be seen from Fig. 2. The W pair production is most sensitive to the couplings g_1^Z and λ_γ , and its sensitivity to κ_γ is comparable to the single W production, which in turn is most sensitive to κ_γ . From all processes, the single photon production is least sensitive.

Deviations from the couplings as they are predicted by the Standard Model would lead to changes of the total cross section, of the production and decay angles and of the average polarization of the bosons.

In the W pair production process, all information about production and decay is contained in five variables: The production angle θ_{W^-} of the W^- , the polar and azimuthal angles θ_1, ϕ_1 and θ_4, ϕ_4 of the decay products in the rest frame of the decaying W^- and W^+ relative to the W flight direction. If a W decays into a charged lepton and a neutrino, θ and ϕ are taken from the charged lepton, and if a W decays into two quarks, the angles from either quark modulo π




Figure 2: Other processes that are used in the determination of the VWW couplings.




Figure 3: Distribution of the production angle of the W boson and of the decay angles of the lepton in the rest frame of the decaying W.

are taken to compensate the missing charge determination. The distributions of $\cos \theta_1$, ϕ_1 and $\cos \theta_{W^-}$ are shown in Fig. 3 as they have been measured by the L3 experiment and together with the expectations for $g_1^Z = 0, 2$. From the shape of these distributions and the total rate, constraints on the value of the couplings are derived.

The shape of the $\cos \theta_{W^-}$ distribution shows stronger distortions than the shape of the $\cos \theta_1$ and ϕ_1 distributions, if the couplings are changed. Therefore, a reliable calculation of these distributions is necessary. Until recently, the theory error (2%) on the shape of the $\cos \theta_{W^-}$ distribution was double the size of the statistical accuracy ($\sim .1\%$). By using the predictions from the newly developed Monte Carlo generators YFSWW3² and RacoonWW³, a theory error of .5% on λ_γ has been achieved⁴. The two generators take into account $\mathcal{O}(\alpha)$ -corrections, i.e. diagrams with internal and external photon lines, in the Leading Pole Approximation (LPA) and the Double Pole Approximation DPA, respectively. Some example diagrams of these corrections are shown in Fig. 4.

By using the predictions from YFSWW3, measurements of the couplings are performed by each experiment, and combined with a log-likelihood method⁵. The likelihood curves of the combined fit are shown in Fig. 5. The measurement of κ_γ agrees within two standard deviations with the Standard Model, and both λ_γ and g_1^Z agree within one standard deviation with the Standard Model. The fitted values with the errors corresponding to $\Delta L = 0.5$ are:

$$g_1^Z = 0.990^{+0.023}_{-0.024} \quad \kappa_\gamma = 0.896^{+0.058}_{-0.056} \quad \lambda_\gamma = -0.023^{+0.025}_{-0.023}$$

For this combination, both the L3 and OPAL experiments did not submit the $e^+e^- \rightarrow W^+W^- \rightarrow q_1\bar{q}_2q_3\bar{q}_4$ channel. By adding these channels, the statistical accuracy of the measurement will improve. As far as systematic uncertainties are concerned, the $\mathcal{O}(\alpha)$ corrections are the largest correlated ones (± 0.04 on κ_γ , ± 0.01 on λ_γ and $^{+0.01}_{-0.02}$ on g_1^Z). For the result shown above, they have been set to the full difference between the Monte Carlo prediction with and




Figure 4: Some example diagrams for $\mathcal{O}(\alpha)$ corrections.




Figure 5: Result of the triple gauge coupling fit.

without $\mathcal{O}(\alpha)$ corrections. More refined numbers will be used in the future. Also, updates on the fits of higher dimensionality relating two or three couplings are planned.

2 Couplings of three neutral bosons

Couplings of three neutral bosons do not exist in the Standard Model. By imposing only Lorentz and $U(1)_{\text{em}}$ invariance, and for final states with equal bosons Bose symmetry, one ends up with possible anomalous vertices shown in Fig. 6. The corresponding Lagrangians^{1,6} describing these anomalous vertices are

$$\begin{aligned} \mathcal{L}_{NP}^{VZZZ} &= \frac{e}{m_Z^2} \left[- f_4^V (\partial_\mu V^{\mu\beta}) Z_\alpha (\partial^\alpha Z_\beta) + f_5^V (\partial^\sigma V_{\sigma\mu}) \tilde{Z}^{\mu\beta} Z_\beta \right] \\ \mathcal{L}_{NP}^{VZ\gamma} &= \frac{e}{m_Z^2} \left[- h_1^V (\partial^\sigma V_{\sigma\mu}) Z_\beta F^{\mu\beta} - h_3^V (\partial_\sigma V^{\sigma\rho}) Z^\alpha \tilde{F}_{\rho\alpha} \right. \\ &\quad \left. - \frac{h_2^V}{m_Z^2} [\partial_\alpha \partial_\beta (\square + m_V^2) V_\mu] Z^\alpha F^{\mu\beta} + \frac{h_4^V}{2m_Z^2} [(\square + m_V^2) \partial^\sigma V^{\rho\alpha}] Z_\sigma \tilde{F}_{\rho\alpha} \right], \end{aligned}$$

with $\tilde{V}_{\mu\nu} = 1/2 \epsilon_{\mu\nu\rho\sigma} V^{\rho\sigma}$ and $V = Z, \gamma$. The Lagrangians are of higher order than those for the gauge couplings of the W boson, so that one would expect either to detect deviations more easily with the W boson couplings or the scale of New Physics (which is artificially set to m_Z in the above formulae) to be close. The couplings f_4^V , h_1^V and h_2^V are CP violating, whereas the couplings f_5^V , h_3^V and h_4^V conserve CP. One interesting option for the future, which has not been followed yet, is to relate the couplings through $SU(2)_L \times U(1)_Y$ symmetry⁷. This relates the couplings from the $Z\gamma$ and from the ZZ final state in the following way: $f_5^V = h_3^V \tan \theta_W$ and $f_4^V = h_1^V \tan \theta_W$.

The measurement of the f couplings proceeds by selecting events from all visible ZZ final states and then reweighting the distributions for different values of the anomalous couplings $f_{4,5}^{Z,\gamma}$.




Figure 6: Couplings of three neutral bosons: Anomalous vertices.




Figure 7: ZZ production angle measured by DELPHI compared to the Standard Model prediction and $f_5^Z = \pm 1.5$.

In the presence of anomalous couplings, the total cross-section, the production angle of the Z boson and the average polarization of the Z bosons would change. In Fig. 7 the distribution of the Z boson production angle $\cos \theta_Z$ as predicted by the Standard Model and for $f_5^Z = \pm 1.5$ is compared to the data, as they have been measured by the DELPHI experiment.

Since in all LEP data no evidence for the presence of anomalous f couplings has been found, limits at the 95% confidence level are set. These limits are derived either one-dimensional by fixing all other couplings to zero, or two-dimensional by fitting couplings with the same CP behavior at the same time. The one-dimensional limits are⁸:

$$-0.17 < f_4^\gamma < 0.19 \quad -0.30 < f_4^Z < 0.28 \quad -0.34 < f_5^\gamma < 0.38 \quad -0.36 < f_5^Z < 0.38$$

For the h -couplings, events of the reactions $e^+e^- \rightarrow Z/\gamma^* \rightarrow q\bar{q}\gamma$ and $e^+e^- \rightarrow Z/\gamma^* \rightarrow \nu\bar{\nu}\gamma$ are selected. The photon energy E_γ , the angle $\cos \alpha_{\gamma\text{-jet}}$ between the photon and the nearest jet, and the photon production angle $\cos \theta_\gamma$ are sensitive to the anomalous couplings. In Fig. 8, distributions of these variables from the OPAL experiment are shown, for the Standard Model prediction and for $h_3^\gamma = \pm 0.5$. No evidence for anomalous h couplings has been found, and one- and two-dimensional limits are derived. The one-dimensional limits are⁸:

$$\begin{aligned} -0.056 < h_1^\gamma < 0.055 & \quad -0.045 < h_2^\gamma < 0.025 & \quad -0.130 < h_1^Z < 0.130 & \quad -0.078 < h_2^Z < 0.071 \\ -0.049 < h_3^\gamma < 0.008 & \quad -0.002 < h_4^\gamma < 0.034 & \quad -0.200 < h_3^Z < 0.070 & \quad -0.050 < h_4^Z < 0.120 \end{aligned}$$




Figure 8: Distributions of E_γ , $\cos \alpha_{\gamma\text{-jet}}$ and $\cos \theta_\gamma$ for the $q\bar{q}\gamma$ final state. Predictions from the Standard Model and for $h_3^\gamma = \pm 0.5$ are compared to the data.




Figure 9: Anomalous contributions to quartic gauge couplings.

3 Quartic boson self couplings

Starting from $U(1)_{\text{em}}$ gauge invariance and requiring a custodial $SU(2)_c$ symmetry, genuine quartic couplings (i.e. quartic couplings that are not introduced to counteract the trilinear gauge couplings to achieve $SU(2)_L \times U(1)_Y$ symmetry) arise through the Lagrangians⁹

$$\begin{aligned} \mathcal{L}_0 &= -\frac{e^2}{16} \frac{a_0^{W,Z}}{\Lambda^2} F^{\mu\nu} F_{\mu\nu} \vec{W}^\alpha \vec{W}_\alpha & WW\gamma\gamma, ZZ\gamma\gamma \\ \mathcal{L}_c &= -\frac{e^2}{16} \frac{a_c^{W,Z}}{\Lambda^2} F^{\mu\alpha} F_{\mu\beta} \vec{W}^\beta \vec{W}_\alpha & WW\gamma\gamma, ZZ\gamma\gamma \\ \mathcal{L}_n &= -\frac{e^2}{16} \frac{a_n}{\Lambda^2} \vec{W}_{\mu\alpha} \cdot (\vec{W}_\nu \times \vec{W}^\alpha) F^{\mu\nu} & WWZ\gamma \end{aligned}$$

The couplings a_0 and a_c conserve CP, the coupling a_n violates CP. Figure 9 shows the relationship between the vertices and the anomalous couplings. In principle, the couplings of the W could be different from the couplings of the Z, but they are assumed to coincide here.

These couplings are accessible either through boson fusion (e.g. in the reaction $e^+e^- \rightarrow \nu\bar{\nu}\gamma\gamma$ or through the production of three gauge bosons in the processes $e^+e^- \rightarrow W^+W^-\gamma$ or $e^+e^- \rightarrow ZZ\gamma$). Although boson fusion becomes important for Linear Collider energies, at LEP the three boson final states dominate the measurement.

In both cases, the presence of the anomalous couplings leads to different kinematics. The energy of the second highest energetic photon, which is expected to be larger for the quartic couplings, is sensitive to the presence of anomalous quartic couplings. No evidence for such couplings is found in the data, and the LEP combined limits set at 95% CL are⁸:

$$-0.02 < a_0/\Lambda^2 \times \text{GeV}^2 < 0.02 \quad -0.03 < a_c/\Lambda^2 \times \text{GeV}^2 < 0.05 \quad -0.17 < a_n/\Lambda^2 \times \text{GeV}^2 < 0.15$$

Acknowledgments

I would like to thank my colleagues from L3 and the convenors of the LEP four-fermion and gauge-coupling working groups for their support and many fruitful discussions.

References

1. K. Hagiwara, R. D. Peccei, D. Zeppenfeld and K. Hikasa, *Probing the weak boson sector in $e^+e^- \rightarrow W^+W^-$* , *Nucl. Phys. B* **282**, 253 (1987)
2. S. Jadach *et al.*, *Comp. Phys. Comm.* **140**, 432 (2001)
3. A. Denner *et al.*, *Nucl. Phys. B* **560**, 33 (1999)
4. R. Brunelière *et al.*, hep-ph/0201304, (2002)
5. J. Alcaraz, L3 internal note 2718, (2001)
6. G. J. Gounaris *et al.*, *Phys. Rev. D* **61**, 073013 (2000)
7. J. Alcaraz, *Phys. Rev. D* **65**, 075020 (2002)
8. The LEP Collaborations ALEPH, DELPHI, L3, OPAL and the LEP GC Working Group, *LEPEWWG/TGC/2001-03*, (2001)
9. G. Bélanger and F. Boudjema, *Phys. Lett. B* **288**, 210 (1992)