

# WEAK COUPLING PHASES

EFI 02-96

hep-ph/0207197

Presented at CERN CKM Workshop

13–16 February 2002

*J. L. Rosner*

*Enrico Fermi Institute and Department of Physics*

*University of Chicago, Chicago, IL, USA*

## Abstract

Recent results obtained from  $B$  decays on the phases of weak couplings described by the Cabibbo-Kobayashi-Maskawa (CKM) matrix are discussed, with particular emphasis on  $\alpha$  and  $\gamma = \pi - \beta - \alpha$ .

## 1 INTRODUCTION

The phases of Cabibbo-Kobayashi-Maskawa (CKM) matrix elements describing charge-changing weak couplings of quarks are fundamental quantities. They are sometimes described in terms of angles  $\alpha = \phi_2$ ,  $\beta = \phi_1$ , and  $\gamma = \phi_3$  in the unitarity triangle. Now that BaBar and Belle are converging on a value of  $\sin(2\beta)$ , attention has turned to ways of learning  $\alpha$  and  $\gamma = \pi - \beta - \alpha$ . This summary describes some recent work on the subject.

In Sec. 2 we discuss  $B^0 \rightarrow \pi^+\pi^-$  in the light of recent measurements at BaBar [1] and Belle [2] of time-dependent asymmetries. This work was performed in part in collaboration with M. Gronau [3, 4, 5] and in part with Z. Luo [6]. We then mention how to learn  $\gamma$  from various  $B \rightarrow K\pi$  decays (Sec. 3, collaboration with M. Gronau [3] and M. Neubert [7, 8]),  $2\beta + \gamma$  from  $B \rightarrow D^{(*)}\pi$  (Sec. 4, collaboration with D. Suprun and C.-W. Chiang [9]), and  $\alpha$  and  $\gamma$  from tree-penguin interference in  $B \rightarrow PP$ ,  $PV$  decays, where  $P$  is a light pseudoscalar and  $V$  a light vector meson (Sec. 5, collaboration with C.-W. Chiang [10]). Sec. 6 is a short guide to other recent work, while we summarize in Sec. 7.

## 2 $\alpha$ FROM $B^0 \rightarrow \pi^+\pi^-$

We regard  $\alpha, \gamma$  as uncertain to about  $\pi/4$ :  $126^\circ \geq \alpha \geq 83^\circ$ ,  $32^\circ \leq \gamma \leq 75^\circ$  [3], in accord with  $122^\circ \geq \alpha \geq 75^\circ$ ,  $37^\circ \leq \gamma \leq 80^\circ$  [11]. If  $B^0 \rightarrow \pi^+\pi^-$  were dominated by the “tree” amplitude  $T$  with phase  $\gamma = \text{Arg}(V_{ub}^*V_{ud})$ , the parameter  $\lambda_{\pi\pi} \equiv e^{-2i\beta}A(\overline{B}^0 \rightarrow \pi^+\pi^-)/A(B^0 \rightarrow \pi^+\pi^-)$  would be just  $e^{2i\alpha}$  and the indirect CP-violating asymmetry  $S_{\pi\pi} = 2\text{Im}\lambda_{\pi\pi}/(1 + |\lambda_{\pi\pi}|^2)$  would be  $\sin 2\alpha$ . Here

$$\frac{d\Gamma}{dt} \left\{ \begin{array}{l} B^0|_{t=0} \rightarrow f \\ \overline{B}^0|_{t=0} \rightarrow f \end{array} \right\} \propto e^{-\Gamma t} [1 \mp S_{\pi\pi} \sin \Delta m t \pm C_{\pi\pi} \cos \Delta m t] \quad , \quad (1)$$

$C_{\pi\pi} = (1 - |\lambda_{\pi\pi}|^2)/(1 + |\lambda_{\pi\pi}|^2)$ , and  $\Delta\Gamma \simeq \Delta m/200$  has been neglected. In the presence of non-zero  $\Delta\Gamma$  one can also measure  $A_{\pi\pi} = 2\text{Re}\lambda_{\pi\pi}/(1 + |\lambda_{\pi\pi}|^2)$ . Since  $|S_{\pi\pi}|^2 + |C_{\pi\pi}|^2 + |A_{\pi\pi}|^2 = 1$  one has  $|S_{\pi\pi}|^2 + |C_{\pi\pi}|^2 \leq 1$ . However, one also has a penguin amplitude  $P$  involving a  $\bar{b} \rightarrow \bar{d}$  loop transition involving contributions  $\sim V_{ud}^*V_{ub}$ ,  $V_{cd}^*V_{cb}$ , and  $V_{td}^*V_{tb} = -V_{ud}^*V_{ub} - V_{cd}^*V_{cb}$ . The decay amplitudes are then

$$A(B^0 \rightarrow \pi^+\pi^-) = -(|T|e^{i\delta_T}e^{i\gamma} + |P|e^{i\delta_P}), \quad A(\overline{B}^0 \rightarrow \pi^+\pi^-) = -(|T|e^{i\delta_T}e^{-i\gamma} + |P|e^{i\delta_P}), \quad (2)$$

where the strong phase difference  $\delta \equiv \delta_P - \delta_T$ . It will be convenient to define  $R_{\pi\pi} \equiv \overline{\mathcal{B}}(B^0 \rightarrow \pi^+\pi^-)/\overline{\mathcal{B}}(B^0 \rightarrow \pi^+\pi^-)_{\text{tree}}$ , where  $\overline{\mathcal{B}}$  refers to a branching ratio averaged over  $B^0$  and  $\overline{B}^0$ . One may use  $S_{\pi\pi}$  and  $C_{\pi\pi}$  to learn  $\alpha, \delta$ , resolving a discrete ambiguity with the help of  $R_{\pi\pi}$  [4]. Alternatively, one may directly use  $S_{\pi\pi}$ ,  $C_{\pi\pi}$ , and  $R_{\pi\pi}$  to learn  $\alpha, \delta$ , and  $|P/T|$  [5, 13].

Explicit expressions for  $R_{\pi\pi}$ ,  $S_{\pi\pi}$  and  $C_{\pi\pi}$  may be found in [4, 5]. In [4] we estimated  $|P/T| = 0.276 \pm 0.064$  (see also [12]), obtaining  $|P|$  from  $B^+ \rightarrow K^0\pi^+$  via (broken) flavor SU(3) and  $|T|$  from  $B \rightarrow \pi\ell\nu$ . Plotting  $C_{\pi\pi}$  against  $S_{\pi\pi}$  for various values of  $\alpha$  in the likely range, one obtains curves parametrized by  $\delta$  which establish a one-to-one correspondence between a pair  $(S_{\pi\pi}, C_{\pi\pi})$  and a pair  $(\alpha, \delta)$  as long as  $|\delta| \leq 90^\circ$ . However, if  $|\delta|$  is allowed to exceed about  $90^\circ$  these curves can intersect with one another, giving rise to a discrete ambiguity corresponding to as much as  $30^\circ$  uncertainty in  $\alpha$  when  $C_{\pi\pi} = 0$ . In this case, when  $\delta = 0$  or  $\pi$ , one has  $|\lambda_{\pi\pi}| = 1$  and  $S_{\pi\pi} = \sin 2(\alpha + \Delta\alpha)$ , where  $\tan(\Delta\alpha) = \pm(|P/T| \sin \gamma)/(1 \pm (|P/T| \cos \gamma))$  is typically  $\pm 15^\circ$ . One can resolve the ambiguity either by comparing the predicted  $R_{\pi\pi}$  with experiment (see [4] for details), or by comparing the allowed  $(\rho, \eta)$  region with that determined by other observables [11]. An example is shown in [3].

Once errors on  $R_{\pi\pi}$  are reduced to  $\pm 0.1$  (they are now about three times as large [4]), a distinction between  $\delta = 0$  and  $\delta = \pi$  will be possible when  $S_{\pi\pi} \simeq 0$ , as appears to be the case for BaBar [1]. For the Belle data [2], which suggest  $S_{\pi\pi} < 0$ , the distinction becomes easier; it becomes harder for  $S_{\pi\pi} > 0$ . With  $100 \text{ fb}^{-1}$  at each of BaBar and Belle, it will be possible to reduce  $\Delta|T|^2/|T|^2$  from its present error of 44% and  $\overline{\mathcal{B}}(B^0 \rightarrow \pi^+\pi^-)$  from its present error of 21% each to about 10% [6], which will go a long way toward this goal. In an analysis independent of  $|P/T|$  performed since the workshop, the somewhat discrepant BaBar and Belle values of  $S_{\pi\pi}$  and  $C_{\pi\pi}$ , when averaged, favor  $\alpha$  between about  $90^\circ$  and  $120^\circ$  (see Fig. 1 of [5]).

### 3 $\gamma$ from $B \rightarrow K\pi$

#### 3.1 $\gamma$ from $B^0 \rightarrow K^+\pi^-$ and $B^+ \rightarrow K^0\pi^+$

We mention some results of [3] on information provided by  $B^0 \rightarrow K^+\pi^-$  decays, which involve both a penguin  $P'$  and a tree  $T'$  amplitude. One can use the flavor-averaged branching ratio  $\overline{\mathcal{B}}$  and the CP asymmetry in these decays, together with  $P'$  information from the  $B^+ \rightarrow K^0\pi^+$  decay rate (assuming it is equal to the charge-conjugate rate, which must be checked) and  $T'$  information from  $B \rightarrow \pi\ell\nu$  and flavor SU(3), to obtain constraints on  $\gamma$ . One considers the ratio  $R \equiv [\overline{\mathcal{B}}(B^0 \rightarrow K^+\pi^-)/\overline{\mathcal{B}}(B^+ \rightarrow K^0\pi^+)][\tau_+/\tau_0]$ , where the  $B^+/B^0$  lifetime ratio  $\tau_+/\tau_0$  is about 1.07. Once the error on this quantity is reduced to  $\pm 0.05$  from its value of  $\pm 0.14$  as of February 2002, which should be possible with  $200 \text{ fb}^{-1}$  at each of BaBar and Belle, one should begin to see useful constraints arising from the value of  $R$ , especially if errors on the ratio  $r \equiv |T'/P'|$  can be reduced with the help of better information on  $|T'|$ .

#### 3.2 $\gamma$ from $B^+ \rightarrow K^+\pi^0$ and $B^+ \rightarrow K^0\pi^+$

One can use the ratio  $R_c \equiv 2\overline{\mathcal{B}}(BB^+ \rightarrow K^+\pi^0)/\overline{\mathcal{B}}(B^+ \rightarrow K^0\pi^+)$  to determine  $\gamma$  [3, 7, 8]. Given the values as of February 2002,  $R_c = 1.25 \pm 0.22$ ,  $A_c \equiv [\mathcal{B}(B^- \rightarrow K^-\pi^0) - \mathcal{B}(B^+ \rightarrow K^+\pi^0)]/\overline{\mathcal{B}}(B^+ \rightarrow K^0\pi^+) = -0.13 \pm 0.17$ , and  $r_c \equiv |T' + C'|/|p'| = 0.230 \pm 0.035$  (here  $C'$  is a color-suppressed amplitude, while  $p'$  is a penguin amplitude including an electroweak contribution), and an estimate [7, 8] of the electroweak penguin contribution, one finds  $\gamma \leq 90^\circ$  or  $\gamma \geq 140^\circ$  at the  $1\sigma$  level, updating an earlier bound [3]  $\gamma \geq 50^\circ$ . A useful determination would involve  $\Delta R_c = \pm 0.1$ , achievable with  $150 \text{ fb}^{-1}$  each at BaBar and Belle.

#### 4 $2\beta + \gamma$ FROM $B \rightarrow D^{(*)}\pi$

The “right-sign” (RS) decay  $B^0 \rightarrow D^{(*)-}\pi^+$ , governed by the CKM factor  $V_{cb}^*V_{ud}$ , and the “wrong-sign” (WS) decay  $\bar{B}^0 \rightarrow D^{(*)-}\pi^+$ , governed by  $V_{cd}^*V_{ub}$ , can interfere through  $B^0$ - $\bar{B}^0$  mixing, leading to information on the weak phase  $2\beta + \gamma$ . One must separate out the dependence on a strong phase  $\delta$  between the RS and WS amplitudes, measuring time-dependent observables

$$A_{\pm}(t) = (1 + R^2) \pm (1 - R^2) \cos \Delta mt, \quad B_{\pm}(t) = -2R \sin(2\beta + \gamma \pm \delta) \sin \Delta mt, \quad (3)$$

where  $R \equiv |\text{WS/RS}| = r|V_{cd}^*V_{ub}/V_{cb}^*V_{ud}| \simeq 0.02r$ , with  $r$  a parameter of order 1 which needs to be known better. In Ref. [9] we use the fact that  $R$  can be measured in the decay  $B^+ \rightarrow D^{*+}\pi^0$  to conclude that with 250 million  $B\bar{B}$  pairs one can obtain an error of less than  $\pm 0.05$  on  $\sin(2\beta + \gamma)$ , which is expected to be greater than about 0.89 in the standard model. Thus, such a measurement is not likely to constrain CKM parameters, but has potential for an interesting non-standard outcome.

#### 5 $\alpha$ and $\gamma$ FROM $B \rightarrow PP, PV$

Some other processes which have a near-term potential for providing information on tree-penguin interference (and hence on  $\alpha$  and  $\gamma$ ) are the following [10]: (1) the CP asymmetries in  $B^+ \rightarrow \pi^+\eta$  and  $\pi^+\eta'$ ; (2) rates in  $B^+ \rightarrow \eta'K^+$  and  $B^0 \rightarrow \eta'K^0$ ; (3) rates in  $B^+ \rightarrow \eta K^{*+}$  and  $B^0 \rightarrow \eta K^{*0}$ ; and (4) rates in  $B^+ \rightarrow \omega K^+$  and  $B^0 \rightarrow \omega K^0$ . Other interesting branching ratios include those for  $B^0 \rightarrow \pi^- K^{*+}$ ,  $B^0 \rightarrow K^+\rho^-$ ,  $B^+ \rightarrow \pi^+\rho^0$ ,  $B^+ \rightarrow \pi^+\omega$ , and  $B^{(+,0)} \rightarrow \eta' K^{*(+,0)}$ , with a story for each [10]. In order to see tree-penguin interference at the predicted level one needs to measure branching ratios at the level of  $\Delta\bar{B} = (1 - 2) \times 10^{-6}$ .

## 6 OTHER WORK

For other recent suggestions on measuring  $\alpha$  and  $\gamma$ , see the review of [14] and the contributions of [15] on the isospin triangle in  $B \rightarrow \pi\pi$  ( $\alpha$ ), [16, 17] on  $B^+ \rightarrow DK^+$  ( $\gamma$ ), [18] on  $B^0 \rightarrow DK_S$  ( $2\beta + \gamma$ ), [19] on  $B^0 \rightarrow K\pi$  ( $\gamma$ ), [20] on  $B^0 \rightarrow \pi^+\pi^-$  and  $B_s \rightarrow K^+K_-$  ( $\gamma$ ), and [21] on  $B^0 \rightarrow K^+\pi^-$  and  $B_s \rightarrow K^-\pi^+$  ( $\gamma$ ). These contain references to earlier work.

## 7 SUMMARY

CKM phases will be learned in many ways. While  $\beta$  is well-known now and will be better-known soon, present errors on  $\alpha$  and  $\gamma$  are about  $45^\circ$ . To reduce them to  $10^\circ$  or less, several methods will help. (1) Time-dependent asymmetries in  $B^0 \rightarrow \pi^+\pi^-$  already contain useful information. The next step will come when both BaBar and Belle accumulate samples of at least  $100 \text{ fb}^{-1}$ . (2) In  $B^0 \rightarrow \pi^+\pi^-$  an ambiguity between a strong phase  $\delta$  near zero and one near  $\pi$  (if the direct asymmetry parameter  $C_{\pi\pi}$  is small) can be resolved experimentally, for example by better measurement of the  $B^0 \rightarrow \pi^+\pi^-$  branching ratio and the  $B \rightarrow \pi\ell\nu$  spectrum. (3) Several  $B \rightarrow K\pi$  modes, when compared, can constrain  $\gamma$  through penguin-tree interference. This has been recognized, for example, in [11]. (4) The rates in several  $B \rightarrow PP, PV$  modes are sensitive to tree-penguin interference. One needs to measure branching ratios with errors less than  $2 \times 10^{-6}$  to see such effects reliably.

## ACKNOWLEDGEMENTS

I would like to thank C.-W. Chiang, M. Gronau, Z. Luo, M. Neubert, and D. Suprun for enjoyable collaborations, the workshop organizers for the opportunity to participate, and the Fermilab

Theory Group for hospitality. This work was supported in part by the United States Department of Energy under Grant No. DE FG02 90ER40560.

## References

- [1] BaBar Collaboration, B. Aubert et al., SLAC preprint SLAC-PUB-9229, hep-ex/0205082, presented at 37th Rencontres de Moriond on Electroweak Interactions and Unified Theories, Les Arcs, France, 9–16 Mar 2002.
- [2] Belle Collaboration, K. Abe et al., Phys. Rev. Lett. **89** (2002) 071801.
- [3] M. Gronau and J. L. Rosner, Phys. Rev. D **65** (2002) 013004.
- [4] M. Gronau and J. L. Rosner, Phys. Rev. D **65** (2002) 093012.
- [5] M. Gronau and J. L. Rosner, Phys. Rev. D **66** (2002), to be published.
- [6] Z. Luo and J. L. Rosner, Phys. Rev. D **65** (2002) 054027.
- [7] M. Neubert and J. L. Rosner, Phys. Lett. B **441** (1998) 403–409.
- [8] M. Neubert and J. L. Rosner, Phys. Rev. Lett. **81** (1998) 5076–5079.
- [9] D. A. Suprun, C.-W. Chiang, and J. L. Rosner, Phys. Rev. D **65** (2002) 054025.
- [10] C.-W. Chiang and J. L. Rosner, Phys. Rev. D **65** (2002) 074035.
- [11] A. Höcker, H. Lacker, S. Laplace, and F. Le Diberder, in *Proceedings of the 9th International Symposium on Heavy Flavor Physics*, Pasadena, California, 10–13 Sep 2001, AIP Conf. Proc. **618** (2002) 27.
- [12] M. Beneke, G. Buchalla, M. Neubert, and C. Sachrajda, Nucl. Phys. **B606** (2001) 245–321.
- [13] J. Charles, Phys. Rev. D **59** (1999) 054007.
- [14] R. Fleischer, in *Proceedings of the 9th International Symposium on Heavy Flavor Physics*, Pasadena, California, 10–13 Sep 2001, AIP Conf. Proc. **618** (2002) 266.
- [15] M. Gronau, D. London, N. Sinha, and R. Sinha, Phys. Lett. B **514** (2001) 315–320.
- [16] M. Gronau, Phys. Rev. D **58** (1998) 037301.
- [17] D. Atwood, I. Dunietz, and A. Soni, Phys. Rev. D **63** (2001) 036005.
- [18] B. Kayser and D. London, Phys. Rev. D **63** (2000) 116013.
- [19] A. J. Buras and R. Fleischer, Eur. Phys. J. C **16** (2000) 97–104.
- [20] R. Fleischer, Phys. Lett. B **459** (1999) 306–320.
- [21] M. Gronau and J. L. Rosner, Phys. Lett. B **482** (2000) 71–76.