

Energy Dependence of the Cronin Effect from Nonlinear QCD Evolution

Javier L. Albacete,^{1,2} Néstor Armesto,² Alex Kovner,^{2,3} Carlos A. Salgado,² and Urs Achim Wiedemann²

¹*Departamento de Física, Universidad de Córdoba, 14071 Córdoba, Spain*

²*Theory Division, CERN, CH-1211 Geneva 23, Switzerland*

³*Department of Mathematics and Statistics, University of Plymouth, Drake Circus, Plymouth PL4 8AA, United Kingdom*

(Received 14 July 2003; published 24 February 2004)

The nonlinear evolution of dense partonic systems has been suggested as a novel physics mechanism relevant for the dynamics of p - A and A - A collisions at collider energies. Here we study to what extent the description of Cronin enhancement in the framework of this nonlinear evolution is consistent with the recent observation in $\sqrt{s} = 200$ GeV d -Au collisions at the Relativistic Heavy Ion Collider. We solve the Balitsky-Kovchegov evolution equation numerically for several initial conditions encoding Cronin enhancement. We find that the properly normalized nuclear gluon distribution is suppressed at all momenta relative to that of a single nucleon. For the resulting spectrum of produced gluons in p - A and A - A collisions, the nonlinear QCD evolution is unable to generate a Cronin-type enhancement, and it quickly erases any such enhancement which may be present at lower energies.

DOI: 10.1103/PhysRevLett.92.082001

PACS numbers: 12.38.Bx

The observation that the ratio of particle yields in p - A and A - A , scaled by the number of collisions, exceeds unity in an intermediate transverse momentum range of a few GeV is commonly referred to as Cronin effect. The effect was first seen at lower fixed target energies [1] and has recently been confirmed in $\sqrt{s} = 200$ GeV d -Au collisions at Relativistic Heavy Ion Collider (RHIC) [2]. The current interest focuses mainly on comparing this Cronin *enhancement* in d -Au to the relative *suppression* of produced hadrons in Au-Au collisions at the same center of mass energy and in the same transverse momentum range [3]. The opposite trend of the two effects and their centrality dependence suggests that d -Au data may serve as an efficient benchmark measurement to distinguish between the two different physical mechanisms suggested for the relative suppression of hadron spectra in Au-Au collisions: initial state parton saturation [4] and final state jet quenching [5].

In particular, it has been suggested that, due to quantum evolution, saturation effects can account for the suppression of the high- p_T hadronic spectra in Au-Au collisions at RHIC above the saturation scale [4]. On the other hand, it is known that saturation models based on multiple scattering (the so-called Glauber-Mueller [6] or McLerran-Venugopalan [7] models) exhibit Cronin enhancement in p - A [8–10] and A - A [9,11]. In these models, quantum evolution is not implemented, and the saturation of low p_T gluons is the result of a redistribution of gluons in transverse phase space [12,13] which does not change the total number of gluons, thus resulting in a compensating enhancement at momenta just above the saturation momentum Q_s . These models are now being used to understand bulk properties of ultrarelativistic heavy ion collisions such as the multiplicity, rapidity distribution, and centrality dependence of particle production [14]. What is not fully understood is (i) whether such Cronin enhancement encoded in the initial condition of a nuclear

wave function persists in the nonlinear perturbative QCD evolution to higher energy and (ii) whether such Cronin enhancement can be generated by the nonlinear evolution itself. This Letter goes beyond earlier discussions [9–11] by providing the first complete (numerical) answer to these questions. We do not address other approaches to Cronin enhancement [15].

We start from the Balitsky-Kovchegov (BK) evolution equation [16,17], which describes the evolution of the forward scattering amplitude $N(\mathbf{r}, y)$ of a QCD dipole of transverse size $|\mathbf{r}|$ with rapidity Y and $y = (\alpha_s N_c / \pi) Y$,

$$\begin{aligned} \frac{dN(|\mathbf{r}|, y)}{dy} &= \frac{1}{2\pi} \int d^2z \frac{\mathbf{r}^2}{(\mathbf{r}-\mathbf{z})^2 \mathbf{z}^2} \\ &\times [N(|\mathbf{r}-\mathbf{z}|) + N(|\mathbf{z}|) - N(|\mathbf{r}|) \\ &\quad - N(|\mathbf{r}-\mathbf{z}|)N(|\mathbf{z}|)]. \end{aligned} \quad (1)$$

The unintegrated gluon distribution is related to the inclusive gluon distribution $\phi(k) \propto \{d[xG(x, k^2)]\}/(d^2k d^2b)$ and is given in terms of the dipole amplitude

$$\phi(k) = \int \frac{d^2r}{2\pi r^2} \exp\{i\mathbf{r} \cdot \mathbf{k}\} N(r). \quad (2)$$

In the following, we also use the modified gluon distribution

$$h(k) = k^2 \nabla_{\vec{k}}^2 \phi(k). \quad (3)$$

The two definitions coincide for the leading order perturbative distribution $\phi(k) \propto 1/k^2$, but are different in general, and especially at low momenta.

Using the second order Runge-Kutta algorithm [18], we solve the BK Eq. (1) numerically with 8000 equally spaced intervals in $\ln k$ space between -15 and 35 and a step $\Delta y = 0.0025$. The accuracy of this algorithm is better than 2% in the entire range of k discussed below.

We evolve two initial conditions given by the McLerran-Venugopalan [7] (MV) and Golec-Biernat-Wüsthoff [19] (GBW) model, respectively:

$$N_{\text{MV}}^{Q_s} = 1 - \exp\left[-\frac{Q_s^2 r^2}{4} \ln\left(\frac{1}{r^2 \Lambda_{\text{QCD}}^2} + e\right)\right], \quad (4)$$

$$N_{\text{GBW}}^{Q_s} = 1 - \exp\left[-\frac{Q_s^2 r^2}{4}\right], \quad (5)$$

where $\Lambda_{\text{QCD}} = 0.2 \text{ GeV}$. For momenta $k \geq O(1 \text{ GeV})$, the sensitivity on the infrared cutoff e is negligible. The amplitudes N_{MV} and N_{GBW} are similar for momenta of order Q_s , but differ strongly in their high k behavior; $\phi_{\text{GBW}}(k)$ decays exponentially while ϕ_{MV} has a power-like tail $1/k^2$.

Figure 1 shows the evolution of $h(k, y)$ and $\phi(k, y)$ for different initial conditions. The solutions for $h(k, y)$ quickly approach a universal solitonlike shape and do not change further except uniformly moving in k on the logarithmic plot. The position of the maximum is the evolved value of the saturation momentum $Q_s(y)$. The solutions for different initial conditions and different rapidities scale as a function of the scaling variable $\rho = k/Q_s(y)$. The shape of the initial condition affects only the value of the saturation momentum $Q_s(y)$, but not the shape of the




FIG. 1. Solutions of the BK equation. Upper left: $h(k)$ evolved (left to right) from $y = 0$ to 5 and 10 for different initial conditions: GBW with $Q_s^2 = 0.36 \text{ GeV}^2$ (solid lines), MV with $Q_s^2 = 4 \text{ GeV}^2$ (dashed lines), and MV with $Q_s^2 = 100 \text{ GeV}^2$ (dotted lines). Upper right: The same as upper left for $\phi(k)$. Lower left: The scaled function $h(\rho)$ versus $\rho = k/Q_s$ for $y = 4, 6, 8,$ and 10 , and the same initial conditions and conventions (lines cannot be distinguished). Lower right: Ratio of $h(y, \rho)/h(y, \rho = 1)$ over $h(y = 10, \rho)/h(y = 10, \rho = 1)$ for $y = 4$ (solid line), 6 (dashed line), 8 (dotted line), and 10 (dash-dotted line), and initial condition MV with $Q_s^2 = 4 \text{ GeV}^2$.

evolved function $h(\rho, y)$. The y dependence of $h(\rho, y)$ is very weak: The function evolves fast towards a scaling form $h(\rho)$. As the rapidity changes between $y = 4$ and $y = 10$, the ratio $h(\rho, y_1)/h(\rho, y_2)$ varies by at most 40% over 3 orders of magnitude of the scaling variable ρ . Similar behavior was found for ϕ (results not shown). This is consistent with previous numerical works [20,21].

To get a quantitative idea of the behavior of the scaling solution, we fitted the numerical solution of $\phi(\rho)$ to two analytical expressions: $s_1(\rho) = a\rho^{2(1-\lambda)}$ and $s_2(\rho) = a \ln(b\rho)\rho^{2(1-\lambda)}$ for $\rho > 5$. The functional form s_1 with $\lambda = 0.37$ and $\ln Q_s \propto y$ describes the scaling behavior of solutions of the linear Balitsky-Fadin-Kuraev-Lipatov (BFKL) equation [22]. It was argued in Ref. [23] that s_2 with the same value of λ and $\ln Q_s \propto \{[2\chi(\lambda)]/(1-\lambda)\}y - \{3/[2(1-\lambda)]\} \ln y$ accounts for the effects of nonlinearities in (1). We find that s_1 does not give an acceptable fit to $\phi(\rho)$ in any extended range of ρ . For values of ρ between 1 and 10^3 , the value of λ varies between 0.39 and 0.46. This is in contrast to the BFKL equation, where we find numerically that s_1 with $\lambda = 0.37$ does indeed approximate the solution over several orders of magnitude with very good accuracy (results not shown). On the other hand, for $5 < \rho < 1000$, s_2 gives a good fit with $\lambda = 0.32$. If, following [23], we fix $\lambda = 0.37$, the fit is still good.

To study the effect of the evolution on the Cronin enhancement, we consider two initial conditions, $N_{\text{MV}}^q(r)$ and $N_{\text{GBW}}^q(r)$ with $q^2 = 0.1 \text{ GeV}^2$ and $Q^2 = 2 \text{ GeV}^2$. Since $q \sim \Lambda_{\text{QCD}}$ and Q is of the order of the estimated saturation momentum for a gold nucleus [14], this choice mimics the gluon distributions of a proton and of a nucleus, respectively. At large transverse momenta, the ratio of the corresponding Fourier transforms is given by the ratio of the saturation momenta,

$$\frac{h^Q(k, y = 0)}{h^q(k, y = 0)} = \frac{Q^2}{q^2} = A^{1/3}. \quad (6)$$

This relation also holds for ϕ . As discussed in [9,10], these initial conditions exhibit Cronin enhancement, namely, $[h^Q(k, y = 0)]/[A^{1/3}h^q(k, y = 0)] > 1$ for $k \sim Q$. We solve the BK equation with these two initial conditions and construct the ratio $R(k, y) = h^Q(k, y)/A^{1/3}h^q(k, y)$ and the corresponding ratio for ϕ (see Fig. 2). The initial Cronin enhancement at rapidity $y = 0$ is seen to be wiped out very quickly by the evolution. Within less than half a unit of rapidity y , the ratios show uniform suppression for all values of transverse momentum. The observed behavior persists if different amounts of Cronin enhancement are included in the initial condition.

As seen in the lower panel of Fig. 2, the Cronin enhancement also disappears rapidly with rapidity when gluon distributions are evolved according to the linear BFKL equation. Qualitative differences between the BFKL and BK dynamics are only visible at momenta $k < Q_s$, where saturation effects are important. For larger




FIG. 2. Ratio of distributions ϕ and h in nucleus and proton, normalized to 1 at $k \rightarrow \infty$. Upper plots: BK evolution, with MV as initial condition with $Q_s^2 = 0.1 \text{ GeV}^2$ for p and 2 GeV^2 for A . Lines from top to bottom correspond to $y = 0, 0.05, 0.1, 0.2, 0.4, 0.6, 1, 1.4,$ and 2 . Lower plots: BFKL evolution, with MV as initial condition with $Q_s^2 = 4 \text{ GeV}^2$ for p and 100 GeV^2 for A . Lines from top to bottom correspond to $y = 0, 1,$ and 4 .

momenta k , the ratios are very similar for linear and nonlinear QCD evolution. We, thus, conclude that the wiping out of the initial enhancement is primarily driven by the linear BFKL dynamics which is contained in the BK equation as well.

For the evolved gluon distributions determined above, we have calculated the yield of produced gluons in p - A and A - A collisions at central rapidity according to the factorized expressions [24]

$$\frac{dN_{pA}}{dy d^2 p d^2 b} \propto \frac{1}{p^2} \int d^2 k h^q(y, k) h^Q(y, \mathbf{p} - \mathbf{k}), \quad (7)$$

$$\frac{dN_{AA}}{dy d^2 p d^2 b} \propto \frac{A^{2/3}}{p^2} \int d^2 k h^Q(y, k) h^Q(y, \mathbf{p} - \mathbf{k}). \quad (8)$$

From these spectra we compute the p - and y -dependent ratios

$$R_{pA} = \frac{\frac{dN_{pA}}{dy d^2 p d^2 b}}{A^{1/3} \frac{dN_{pp}}{dy d^2 p d^2 b}}, \quad R_{AA} = \frac{\frac{dN_{AA}}{dy d^2 p d^2 b}}{A^{4/3} \frac{dN_{pp}}{dy d^2 p d^2 b}}.$$

As seen in Fig. 3, the nonlinear BK evolution quickly wipes out any initial Cronin enhancement not only on the level of single parton distribution functions but also on the level of particle spectra. We performed several checks to establish that this behavior is generic. First, we checked the disappearance of Cronin enhancement by evolving different initial conditions corresponding to different initial amounts of enhancement. Second, we note that, in some calculations of gluon production in




FIG. 3. Ratios R_{pA} and R_{AA} of gluon yields in p - A (upper plot) and A - A (lower plot) for BK evolution, with MV as initial condition with $Q_s^2 = 0.1 \text{ GeV}^2$ for p and 2 GeV^2 for A . Lines from top to bottom correspond to $y = 0, 0.05, 0.1, 0.2, 0.4, 0.6, 1, 1.4,$ and 2 .

p - A , the gluon distribution ϕ [25] rather than h [24,26,27] enters the right-hand side of (7). Results using ϕ were found to be close to those shown in Fig. 3. Third, different calculations of (7) employ h defined in terms of the scattering amplitude either of the fundamental [24] dipole used here, or of the adjoint [10,26,27] one. By expressing the adjoint dipole amplitude in terms of the fundamental one, $N_{\text{adj}} = 2N - N^2$, we checked that our results remain qualitatively the same for the adjoint case. To summarize, the expressions (7) and (8) are subject to uncertainties as discussed in more detail in [9]. However, our conclusion about the disappearance of Cronin enhancement during QCD evolution is likely to persist in more refined ways of calculating particle spectra, since it is rooted directly in the rapidity dependence of gluon distributions [9].

We now comment on a recent formal argument which — in contrast to our numerical findings — suggests enhancement survives the nonlinear evolution. It is based on the observation that at very short distances $r \rightarrow 0$, $N(r)$ is not affected by evolution. Thus, the integral of ϕ over the transverse momentum is expected to be rapidity independent,

$$\int d^2 k \phi(k) = \frac{1}{r^2} N(r)|_{r=0}. \quad (9)$$

One thus obtains the sum rule

$$\int d^2 k \phi_A(k, y) = A^{1/3} \int d^2 k \phi_p(k, y), \quad (10)$$

valid for any rapidity, since it is satisfied by the initial condition ϕ_{MV} . Since the nonlinear evolution leads to the

depletion of the gluon distribution $\phi_A(k)$ relative to $A^{1/3}\phi_p(k)$ at low momenta, it must follow that in some range of momenta this effect is compensated by enhancement of ϕ_A . However, this argument breaks down since the quantity defined in (9) is infinite. As such, (10) relates only the (UV) divergent parts of the integrals, and carries no information about possible behavior at finite momentum. To be more specific, we use the scaling property $\phi(k, y) = \phi[k/Q_s(y)]$ of the solution of the BK equation established above. It is known that the ratio of the saturation momenta for any two solutions is preserved by the BK evolution [20–22]. For our two solutions representing a nucleus and a nucleon, this implies $[Q_s^A(y)]/[Q_s^p(y)] = A^{1/6}$. We now rewrite the sum rule (10) by regulating the divergent integrals with a large but finite UV cutoff $aQ_s^A(y)$,

$$\begin{aligned} \int_0^{a^2(Q_s^A)^2} d^2k \phi_A(k, y) &= (Q_s^A)^2 \int_0^{a^2} d^2\rho \phi(\rho) \\ &= A^{1/3}(Q_s^p)^2 \int_0^{a^2} d^2\rho \phi(\rho) \\ &= A^{1/3} \int_0^{a^2(Q_s^p)^2} d^2k \phi_p(k, y). \end{aligned} \quad (11)$$

The formal limit $a \rightarrow \infty$ recovers Eq. (10). However, since $Q_s^A \gg Q_s^p$, the regularized sum rule (11) is satisfied even if the nuclear distribution is suppressed relative to that of a single nucleon uniformly at all momenta. Thus, the sum rule (10) carries no information about either presence or absence of Cronin enhancement.

In summary, we have found that the nonlinear QCD evolution to high energy is very efficient in erasing any Cronin-type enhancement which may be present in the initial conditions. For “realistic” initial conditions, this disappearance occurs within half a unit of rapidity. We note that in our units the evolution from 130 to 200 GeV corresponds to $\delta y \simeq 0.1$ for $\alpha_s = 0.2$, and thus is not sufficient to completely eliminate an initial enhancement at central rapidity. For forward rapidity, δy is greater. The evolution to the LHC energy corresponds to $\delta y \sim 1$. Thus, the BK evolution suggests the reduction of the Cronin effect in d -Au for forward rapidities at RHIC and predicts its disappearance for p -A collisions at LHC.

We thank the CERN TH Division (J. L. A. and A. K.) and the INT Seattle for its hospitality, and the DOE for partial support (N. A., A. K., C. A. S., and U. A. W.). J. L. A. is supported by MECO of Spain, Grant No. AP2001-3333. Useful discussions with R. Baier, E. Iancu, D. Kharzeev, Y. Kovchegov, P. Jacobs, and L. McLerran are gratefully acknowledged. We also thank M. A. Braun for discussions and a numerical cross-check.

Note added.—After appearance of this work, Refs. [9,10] were revised. Reference [10] no longer invokes the sum rule argument criticized above, and both now agree with our main conclusion.

- [1] J. W. Cronin *et al.*, Phys. Rev. D **11**, 3105 (1975).
- [2] PHENIX Collaboration, S.S. Adler *et al.*, Phys. Rev. Lett. **91**, 072303 (2003); STAR Collaboration, J. Adams *et al.*, *ibid.* **91**, 072304 (2003); PHOBOS Collaboration, B. B. Back *et al.*, *ibid.* **91**, 072302 (2003); BRAHMS Collaboration, I. Arsene *et al.*, *ibid.* **91**, 072305 (2003).
- [3] PHENIX Collaboration, S.S. Adler *et al.*, Phys. Rev. Lett. **91**, 072301 (2003); STAR Collaboration, J. Adams *et al.*, *ibid.* **91**, 172302 (2003); PHOBOS Collaboration, B. B. Back *et al.*, Phys. Lett. B **578**, 297 (2004).
- [4] D. Kharzeev, E. Levin, and L. McLerran, Phys. Lett. B **561**, 93 (2003).
- [5] X. N. Wang, Phys. Lett. B **579**, 299 (2004).
- [6] A. H. Mueller, Nucl. Phys. **B335**, 115 (1990).
- [7] L. D. McLerran and R. Venugopalan, Phys. Rev. D **49**, 2233 (1994); **49**, 3352 (1994).
- [8] F. Gelis and J. Jalilian-Marian, Phys. Rev. D **67**, 074019 (2003).
- [9] R. Baier, A. Kovner, and U. A. Wiedemann, Phys. Rev. D **68**, 054009 (2003).
- [10] D. Kharzeev, Y.V. Kovchegov, and K. Tuchin, Phys. Rev. D **68**, 094013 (2003).
- [11] J. Jalilian-Marian, Y. Nara, and R. Venugopalan, Phys. Lett. B **577**, 54 (2003).
- [12] Y.V. Kovchegov, Phys. Rev. D **54**, 5463 (1996).
- [13] J. Jalilian-Marian, A. Kovner, L. D. McLerran, and H. Weigert, Phys. Rev. D **55**, 5414 (1997).
- [14] D. Kharzeev and M. Nardi, Phys. Lett. B **507**, 121 (2001); D. Kharzeev and E. Levin, Phys. Lett. B **523**, 79 (2001); D. Kharzeev, E. Levin, and M. Nardi, hep-ph/0111315.
- [15] E. Wang and X. N. Wang, Phys. Rev. C **64**, 034901 (2001); K. J. Eskola and H. Honkanen, Nucl. Phys. **A713**, 167 (2003); B. Z. Kopeliovich, J. Nemchik, A. Schafer, and A.V. Tarasov, Phys. Rev. Lett. **88**, 232303 (2002); A. Accardi, hep-ph/0212148; I. Vitev, Phys. Lett. B **562**, 36 (2003).
- [16] I. Balitsky, Nucl. Phys. **B463**, 99 (1996).
- [17] Y.V. Kovchegov, Phys. Rev. D **60**, 034008 (1999).
- [18] M. Braun, Eur. Phys. J. C **16**, 337 (2000).
- [19] K. Golec-Biernat and M. Wüsthoff, Phys. Rev. D **59**, 014017 (1999).
- [20] N. Armesto and M. A. Braun, Eur. Phys. J. C **20**, 517 (2001).
- [21] M. Lublinsky, Eur. Phys. J. C **21**, 513 (2001); K. Golec-Biernat, L. Motyka, and A. M. Stasto, Phys. Rev. D **65**, 074037 (2002).
- [22] E. Iancu, K. Itakura, and L. McLerran, Nucl. Phys. **A708**, 327 (2002).
- [23] A. H. Mueller and D. N. Triantafyllopoulos, Nucl. Phys. **B640**, 331 (2002).
- [24] M. A. Braun, Phys. Lett. B **483**, 105 (2000).
- [25] L. V. Gribov, E. M. Levin, and M. G. Ryskin, Phys. Rep. **100**, 1 (1983).
- [26] Y.V. Kovchegov and A. H. Mueller, Nucl. Phys. **B529**, 451 (1998).
- [27] Y.V. Kovchegov and K. Tuchin, Phys. Rev. D **65**, 074026 (2002).