

MEASUREMENTS OF σ_{tot} , $d\sigma_{\text{el}}/dt$, AND EVENT DISTRIBUTIONS
IN $\bar{p}p$ AND pp COLLISIONS AT $\sqrt{s} = 31, 53, \text{ AND } 63 \text{ GEV}$

CERN - Napoli - Pisa - Stony Brook Collaboration

M. Ambrosio, G. Anzivino, G. Barbarino, G. Carboni, V. Cavasinni,
T. Del Prete, P.D. Grannis, D. Lloyd Owen, M. Morganti,
G. Paternoster, S. Patricelli, F. Schiavo, and M. Valdata-Nappi

(Presented by G. Carboni)

Experiment R210 has been designed with the primary aim of measuring precisely $\sigma_{\text{tot}}(\bar{p}p)$ and $\sigma_{\text{tot}}(pp)$ over the ISR energy range. The experiment is also equipped with small-angle detectors which allow us to measure $d\sigma_{\text{el}}/dt$, and with a system of drift chambers and scintillation hodoscopes to measure charged multiplicities and angular distributions of emitted secondaries. Data have been collected at $\sqrt{s} = 31, 53, \text{ and } 63 \text{ GeV}$. Results at 53 GeV have already been published -- results at 31 and 63 GeV must still be considered preliminary.

The total cross-section is obtained by measuring simultaneously the total interaction rate and the ISR luminosity: $\sigma_{\text{tot}} = R_{\text{tot}}/L$. Excellent machine performance and accurate calibrations of the beam displacement scale allowed us to attain a better than 1% accuracy in pp runs. Our data reproduce well the old ISR results, except at $\sqrt{s} = 63 \text{ GeV}$, where, however, the agreement is good if we restrict the comparison to total-rate results only. We plan to collect more data at this energy in order to clarify this point. The difference $\Delta\sigma_{\text{tot}} = \sigma_{\text{tot}}(\bar{p}p) - \sigma_{\text{tot}}(pp)$ is positive over the range measured, showing conclusively that $\sigma_{\text{tot}}(\bar{p}p)$ increases in the ISR energy range. As expected from Regge phenomenology, $\Delta\sigma_{\text{tot}}$ behaves as s^{-2} . This result disfavors exotic possibilities, such as odderons, which would have a different s dependence. Both pp and $\bar{p}p$ data, moreover, favour a $\ln^2 s$ behaviour for σ_{tot} , and the extrapolation of this behaviour to the Collider agrees well with the result of UA4.

The elastic cross-section has only been analyzed at 53 GeV, and all the elastic cross-section parameters are the same for $\bar{p}p$ and pp , and consistent with geometrical scaling. Extrapolation of the elastic rate to measure the total cross-section via the optical theorem gives good agreement with the total-rate method.

As far as particle distributions are concerned, we focus our attention more on the comparison of $\bar{p}p$ and pp than on absolute numbers,

since most instrumental effects disappear in the comparison. In single-particle pseudorapidity distributions, a small excess (5%) is observed in the central region for $\bar{p}p$. Moreover, the average charged multiplicity $\langle n_{ch} \rangle$ is 2% higher for $\bar{p}p$ than for pp . Both $\bar{p}p$ and pp distributions satisfy KNO scaling fairly well.

A more interesting quantity is the difference of $\bar{p}p$ and pp topological cross-sections $\Delta\sigma_n$. The mean charged multiplicity of this distribution is 30-40% higher than $\langle n_{ch} \rangle$ for the individual reactions, and this effect occurs at all energies. The normalized form $\langle n \rangle \Delta\sigma_n / \Delta\sigma_{tot}$ is not fitted by the KNO function, but the distribution is similar to that obtained for annihilation at lower energy and for e^+e^- reactions.

A further difference in $\bar{p}p$ is the presence of an excess in the two-particle correlation function around 90° . The excess (roughly the same at the three energies measured) has a very short pseudorapidity range ($\Delta\eta = \pm 0.3$) compared to the classical short-range correlation ($\Delta\eta = \pm 1$). This effect depends on the multiplicity of the event, being present only for those multiplicities corresponding to the largest values of $\Delta\sigma_n$, suggesting that it is related to the "annihilation" mechanism, which still seems to be important at these energies.



MEASUREMENT OF σ_{TOT} , $\frac{d\sigma}{dt}_{EL}$ AND
EVENT DISTRIBUTIONS IN $p-p$ AND $\bar{p}-p$
COLLISIONS AT $\sqrt{s} = 31, 53, 63$ GeV

H. AMBROSIO, G. ANZIVINO, G. BARBARINO,
G. CARBONI, V. CAVASINNI, T. DEL PRETE,
P. GRANNIS, D. LLOYD OWEN, M. MORGANTI,
G. PATERNOSTER, S. PATRIZELLI, F. SCHIAVO,
M. VALDATA-NAPPI

ISR EXP'T R210 CERN-NAPOLI-PISA-STONY BROOK

PUBLISHED:

P.L. 108 B (1982) 145 } σ_{tot} , $\sqrt{s} = 53$ GeV
P.L. 113 B (1982) 87 }
(P.L. 113 B (1982) 347 } $\alpha \alpha / \alpha p$ ELASTIC)
P.L. 115 B (1982) 495 } ELASTIC, $\sqrt{s} = 53$ GeV



LUMINOSITY MONITORS:

- H5_L • H5_R
- H34_L • H34_R
- H34_L • TB_R

LEFT

RIGHT

$$T = HIT_L * HIT_R$$

$$HIT = (TB_A \cdot TB_B + H5_A \cdot H5_B + H4 \cdot H3 + H2 \cdot H1 + C0 \cdot C1)$$

TOTAL CROSS SECTION

$$\sigma_{TOT} = R_{TOT} / \mathcal{L}$$

- we measure ~ 95% of all collisions. extrapolate elastic and diffractive-like inelastic to correct for the loss

TRIGGER: (LEFT) * (RIGHT)

$$LEFT = H_1 \cdot H_2 + H_3 \cdot H_4 + H_5 + TB + E_2$$

- MEASURE: - TOF's between all left-right hodoscope pairs
- (θ, ϕ) for all charged particles
 - live time (Rate = N / τ_{LIVE})

- \mathcal{L}
- Choose a monitor, M, (hodoscope pair), and measure σ_M in Van der Meer's method

$$\sigma_M = \frac{1}{K} \int R_M(\delta) d\delta$$

and $\mathcal{L} = \frac{R_M}{\sigma_M}$

ISR disp. scale acc. $\pm 0.2\%$
(K. Potter)

-3-

In PP runs large imbalance in beam currents (~10 A p - 2-4 mA \bar{p})

large background from beam-gas & beam-pipe



Background subtraction performed by variation of luminosity (vertical steering):

$$\sigma_{tot} = \frac{\Delta R_{tot}}{\Delta L}$$


Beams at full overlap = background + signal
Beams separated = background

The luminosity monitor has to be free as possible from beam-gas background



-4-

VAN DER MEER CALIBRATION



EXTRAPOLATION TO R_{TOT}

- 1) ELASTIC: $\Delta\sigma_{EL} = \int_{\Delta\Omega} \frac{d\sigma}{dE} \Big|_{EL} \cdot \Delta t$
- EXPLOIT OPTICAL THEOREM TO FIX $\frac{d\sigma}{dE} \Big|_{E=0}$
 - $\Delta\sigma_{EL} \propto P_{ISR}^2$
 - RATHER INSENSITIVE TO THE VALUE OF ELASTIC SLOPE

- 2) INELASTIC: (SINGLE DIFFRACTIVE)
- FOR EVENTS WITH A PARTICLE ON ONE SIDE $\eta > 2^\circ$
- PLOT MAXIMUM t^* ($= P_{ISR}^2 \eta^2$) OF OPPOSITE SIDE.
- INTEGRATE OVER THE UNCOVERED $\Delta\Omega$

$\Delta\Omega = 10^{-4} \text{ sr}$
 $\theta_{MIN} = 0.2^\circ$

RUN CONDITIONS
IN IZ


DATE	I_p (A)	I_F (mA)	\mathcal{E} (cm ² sr)	$\int \mathcal{E} dt$ (cm ²)	\sqrt{S} (GeV)
10/81	10	3.0	4×10^{26}	4×10^{32}	53
4/82	12	3.4	10^{27}	4×10^{32}	63
5/82	11	4.2	10^{27}	7×10^{32}	31
6/82	12	4.0	$6 \cdot 10^{26}$	4×10^{32}	53

P-P runs taken immediately before
P-P runs.


Note: $1 \text{ mA} = 2 \times 10^{10} \text{ P}$

SUMMARY OF G_{TOT} RESULTS

G_{obs} (mb)	ΔG_{LL} (mb)	ΔG_{inec} (mb)	G_{TOT} (mb)	ΔG_{TOT} (mb)
39.3 ± 0.1	0.509 ± 0.03	0.231 ± 0.02	40.04 ± 0.20	2.3 ± 0.3
41.5 ± 0.3	0.588 ± 0.03	0.247 ± 0.02	42.33 ± 0.3	
40.75 ± 0.20	2.0 ± 0.1	0.51 ± 0.03	43.26 ± 0.2	1.44 ± 0.44
42.13 ± 0.4	2.05 ± 0.1	0.52 ± 0.03	44.7 ± 0.4	
41.37 ± 0.11	2.82 ± 0.14	0.50 ± 0.03	44.68 ± 0.22	0.57 ± 0.3
41.85 ± 0.30	2.87 ± 0.15	0.535 ± 0.03	45.25 ± 0.3	



FIT : $\sigma_{tot}(pp) = \sigma_0 + \gamma \cdot \ln^2(\sqrt{s})$
 $\sigma_{tot}(\bar{p}p) = \sigma_0 + \gamma \cdot \ln^2(\sqrt{s}) + \beta \cdot s^{-\alpha}$
 $1-\alpha = 0.58$





ELASTIC SCATTERING $|t| < 0.05 \text{ GeV}^2$

- 1) TRIGGER $\equiv T_{BL} * T_{BR}$
- 2) REJECT EVENTS WITH PARTICLES AT LARGE ANGLES
- 3) FIND COLLINEAR HITS

Bkgnd (NON-COLLINEAR EVENTS) $< 6\%_{TOT}$

MULTIPLE HITS OCCURRED IN 31% OF CASES. THE EVENT IS KEPT IF THE MOST COLLINEAR PAIR SATISFY 3).



$\sqrt{s} = 53 \text{ GeV}$	P-P	$\bar{P}-P$
$b \text{ (GeV}^{-2}\text{)}$	$13.09 \pm 0.37 \pm 0.21$	$13.92 \pm 0.37 \pm 0.22$
$\sigma_{EL} \text{ (}\mu\text{b)}$	$7.79 \pm 0.13 \pm 0.11$	$7.89 \pm 0.17 \pm 0.11$
$\sigma_{TOT}^{O.T.} \text{ (}\mu\text{b)}$	$43.34 \pm 0.29 \pm 0.13$	$44.86 \pm 0.44 \pm 0.13$
$\sigma_{EL} / \sigma_{TOT}$	$0.180 \pm 0.003 \pm 0.003$	$0.176 \pm 0.004 \pm 0.003$
$b / \sigma_{TOT}^2 \text{ (GeV}^{-2}\mu\text{b}^{-1}\text{)}$	$0.302 \pm 0.009 \pm 0.005$	$0.310 \pm 0.009 \pm 0.005$
$b' \text{ (GeV}^{-2}\text{)}$ ($0.09 < t < 1.0 \text{ GeV}^2$)	$10.34 \pm 0.19 \pm 0.06$	$10.68 \pm 0.20 \pm 0.06$

* COMBINED WITH TOTAL RATE METHOD

PSEUDORAPIDITY AND MULTIPLICITY DISTRIBUTIONS

SELECTION :

TRIGGER = $(T_8 + H_5 + H_{34})_L * (T_8 + H_5 + H_{34})_R$ } forward cone trigger
 > 90% of inelastic events

TWO SETS OF DATA :

- a) COUNTERS $|\eta| \leq 5.$
- b) CHAMBERS $|\eta| \leq 2.$

RAW DATA

NO CORRECTIONS FOR :

- SECONDARY INTERACTIONS
- δ -RAYS, CONVERSIONS
- BINNING
- k_0, Λ

FOCUS

DIFFERENCES in $\bar{p}p$ and pp in :

$$P_1(\eta) = \frac{1}{\sigma_I} \frac{d\sigma}{d\eta}$$

$$P_m(\eta) = \frac{1}{\sigma_m} \frac{d\sigma_m}{d\eta}$$

$$R_2(\eta_1, \eta_2) = \frac{P_2(\eta_1, \eta_2)}{P_1(\eta_1) \cdot P_1(\eta_2)} - 1.$$

σ_m








Fig. 3



MULTIPLICITY DISTRIBUTIONS $\sqrt{s} = 31$ GeV




$\bar{p}p - pp$

R210






● ($\bar{p}p - p\bar{p}$) THIS EXP. $\sqrt{S} = 53 \text{ GeV}$




MULTIPLICITY DISTRIBUTIONS

$\sqrt{S} = 31 \text{ GeV}$

○ pp
● $\bar{p}p$







TWO-BODY CORRELATIONS $\sqrt{s} = 31 \text{ GeV}$

COUNTERS
ALL MULTIPLICITIES


$$R_2(\eta_1, \eta_2) = \frac{\rho_2(\eta_1, \eta_2) - 1}{\rho_1(\eta_1)\rho_1(\eta_2)}$$

- PP
- $\bar{P}P$




INCLUSIVE TWO-BODY CORRELATIONS $\sqrt{s} = 31 \text{ GeV}$
(CHAMBERS) R210

$$|\eta| < 2.$$




$$R_2(\eta_1, \eta_2) = \frac{f_2(\eta_1, \eta_2)}{f_1(\eta_1) \cdot f_1(\eta_2)} - 1.$$

TWO-BODY CORRELATIONS $\sqrt{s} = 31$ GeV (INCLUSIVE)
(CHAMBERS)



TWO-BODY CORRELATIONS $\sqrt{s} = 31$ GeV (SEMI-INCLUSIVE)
(CHAMBERS) $|\eta| < 2$



CONCLUSIONS

A) Total cross-section

- 1) σ_{TOT} rising as $\ln^2 s$
- 2) $\Delta\sigma$ positive, but $\rightarrow 0$

$$\Delta\sigma \propto s^{\alpha-1} \quad \text{with } \alpha = 0.44.$$

B) Elastic cross-section

- 1) b rising with s ($\ln^2 s$)
- 2) geometrical scaling
 $\frac{b}{\sigma_{TOT}}, \frac{\sigma_{el}}{\sigma_{TOT}}$ indep. from s

C) Inelastic collisions

$$1) \frac{P_1(\bar{p}p)}{P_1(pp)} \begin{cases} \approx 1.05 & |\eta| \leq 2.5 \\ \leq 1 & |\eta| \geq 3 \end{cases}$$

$$2) R_2(\bar{p}p) > R_2(pp) \rightarrow \text{very short range correlations (0.3)}$$

KNO scaling violated KNO scaling OK

$$\downarrow \qquad \qquad \downarrow \qquad \downarrow \qquad \rightarrow \text{central region } (|\eta| < 1.5)$$

$$3) \langle n_{\Delta\sigma_n} \rangle > \langle n_{pp} \rangle, \langle n_{\bar{p}p} \rangle$$

↳ hint for the existence of "annihilation process" (similar to other very inelastic processes?)