

The Deuteron Spin-dependent Structure Function g_1^d and its First Moment

The COMPASS Collaboration

Abstract

We present a measurement of the deuteron spin-dependent structure function g_1^d based on the data collected by the COMPASS experiment at CERN during the years 2002–2004. The data provide an accurate evaluation for Γ_1^d , the first moment of $g_1^d(x)$, and for the matrix element of the singlet axial current, a_0 . The results of QCD fits in the next to leading order (NLO) on all g_1 deep inelastic scattering data are also presented. They provide two solutions with the gluon spin distribution function ΔG positive or negative, which describe the data equally well. In both cases, at $Q^2 = 3 (\text{GeV}/c)^2$ the first moment of ΔG is found to be of the order of $0.2 - 0.3$ in absolute value.

Keywords: Deep inelastic scattering; Spin; Structure function; QCD analysis; A1; g_1

(To be Submitted to Physics Letters B)

The COMPASS Collaboration

V.Yu. Alexakhin⁸⁾, Yu. Alexandrov¹⁸⁾, G.D. Alexeev⁸⁾, M. Alexeev²⁹⁾, A. Amoroso²⁹⁾, B. Badelek³⁰⁾, F. Balestra²⁹⁾, J. Ball²⁵⁾, J. Barth⁴⁾, G. Baum¹⁾, M. Becker²⁰⁾, Y. Bedfer²⁵⁾, C. Bernet²⁵⁾, R. Bertini²⁹⁾, M. Bettinelli¹⁹⁾, R. Birsa²⁸⁾, J. Bisplinghoff³⁾, P. Bordalo^{15,a)}, F. Bradamante²⁸⁾, A. Bressan²⁸⁾, G. Brona³⁰⁾, E. Burtin²⁵⁾, M.P. Bussa²⁹⁾, V.N. Bytchkov⁸⁾, A. Chapiro²⁷⁾, A. Cicuttin²⁷⁾, M. Colantoni^{29,b)}, A.A. Colavita²⁷⁾, S. Costa^{29,c)}, M.L. Crespo²⁷⁾, N. d'Hose²⁵⁾, S. Dalla Torre²⁸⁾, S. Das⁷⁾, S.S. Dasgupta⁶⁾, R. De Masi²⁰⁾, N. Dedek¹⁹⁾, D. Demchenko¹⁶⁾, O.Yu. Denisov^{29,d)}, L. Dhara⁷⁾, V. Diaz^{28,27)}, A.M. Dinkelbach²⁰⁾, S.V. Donskov²⁴⁾, V.A. Dorofeev²⁴⁾, N. Doshita^{2,21)}, V. Duic²⁸⁾, W. Dünnweber¹⁹⁾, A. Efremov⁸⁾, P.D. Eversheim³⁾, W. Eyrich⁹⁾, M. Faessler¹⁹⁾, P. Fauland¹⁾, A. Ferrero²⁹⁾, L. Ferrero²⁹⁾, M. Finger²²⁾, M. Finger jr.⁸⁾, H. Fischer¹⁰⁾, J. Franz¹⁰⁾, J.M. Friedrich²⁰⁾, V. Frolov^{29,d)}, R. Garfagnini²⁹⁾, F. Gautheron¹⁾, O.P. Gavrichtchouk⁸⁾, S. Gerassimov^{18,20)}, R. Geyer¹⁹⁾, M. Giorgi²⁸⁾, B. Gobbo²⁸⁾, S. Goertz^{2,4)}, A.M. Gorin²⁴⁾, O.A. Grajek³⁰⁾, A. Grasso²⁹⁾, B. Grube²⁰⁾, A. Guskov⁸⁾, F. Haas²⁰⁾, J. Hannappel^{4,16)}, D. von Harrach¹⁶⁾, T. Hasegawa¹⁷⁾, S. Hedicke¹⁰⁾, F.H. Heinsius¹⁰⁾, R. Hermann¹⁶⁾, C. Heß²⁾, F. Hinterberger³⁾, M. von Hodenberg¹⁰⁾, N. Horikawa^{21,e)}, S. Horikawa²¹⁾, I. Horn³⁾, C. Ilgner^{11,19)}, A.I. Ioukaev⁸⁾, I. Ivanchin⁸⁾, O. Ivanov⁸⁾, T. Iwata^{21,f)}, R. Jahn³⁾, A. Janata⁸⁾, R. Joosten³⁾, N.I. Jouravlev⁸⁾, E. Kabuß¹⁶⁾, D. Kang¹⁰⁾, B. Ketzer²⁰⁾, G.V. Khaustov²⁴⁾, Yu.A. Khokhlov²⁴⁾, Yu. Kisselev^{1,2)}, F. Klein⁴⁾, K. Klimaszewski³⁰⁾, S. Koblitz¹⁶⁾, J.H. Koivuniemi^{2,13)}, V.N. Kolosov²⁴⁾, E.V. Komissarov⁸⁾, K. Kondo^{2,21)}, K. Königsmann¹⁰⁾, I. Konorov^{18,20)}, V.F. Konstantinov²⁴⁾, A.S. Korentchenko⁸⁾, A. Korzenev^{16,d)}, A.M. Kotzinian^{8,29)}, N.A. Koutchinski⁸⁾, O. Kouznetsov⁸⁾, K. Kowalik³⁰⁾, D. Kramer¹⁴⁾, N.P. Kravchuk⁸⁾, G.V. Krivokhizhin⁸⁾, Z.V. Kroumchtein⁸⁾, J. Kubart¹⁴⁾, R. Kuhn²⁰⁾, V. Kukhtin⁸⁾, F. Kunne²⁵⁾, K. Kurek³⁰⁾, M.E. Ladygin²⁴⁾, M. Lamanna^{11,28)}, J.M. Le Goff²⁵⁾, M. Leberig^{11,16)}, A.A. Lednev²⁴⁾, A. Lehmann⁹⁾, J. Lichtenstadt²⁶⁾, T. Liska²³⁾, I. Ludwig¹⁰⁾, A. Maggiora²⁹⁾, M. Maggiora²⁹⁾, A. Magnon²⁵⁾, G.K. Mallot¹¹⁾, C. Marchand²⁵⁾, J. Marroncle²⁵⁾, A. Martin²⁸⁾, J. Marzec³¹⁾, L. Masek¹⁴⁾, F. Massmann³⁾, T. Matsuda¹⁷⁾, D. Matthiä¹⁰⁾, A.N. Maximov⁸⁾, W. Meyer²⁾, A. Mielech^{28,30)}, Yu.V. Mikhailov²⁴⁾, M.A. Moinester²⁶⁾, T. Nagel²⁰⁾, O. Nähle³⁾, J. Nassalski³⁰⁾, S. Neliba²³⁾, D.P. Neyret²⁵⁾, V.I. Nikolaenko²⁴⁾, K. Nikolaev⁸⁾, A.A. Nozdrin⁸⁾, V.F. Obraztsov²⁴⁾, A.G. Olshevsky⁸⁾, M. Ostrick^{4,16)}, A. Padee³¹⁾, P. Pagano²⁸⁾, S. Panebianco²⁵⁾, D. Panzieri^{29,b)}, S. Paul²⁰⁾, D.V. Peshekhonov⁸⁾, V.D. Peshekhonov⁸⁾, G. Piragino²⁹⁾, S. Platchkov^{11,25)}, J. Pochodzalla¹⁶⁾, J. Polak¹⁴⁾, V.A. Polyakov²⁴⁾, G. Pontecorvo⁸⁾, A.A. Popov⁸⁾, J. Pretz⁴⁾, S. Procureur²⁵⁾, C. Quintans¹⁵⁾, S. Ramos^{15,a)}, G. Reicherz²⁾, E. Rondio³⁰⁾, A.M. Rozhdestvensky⁸⁾, D. Ryabchikov²⁴⁾, V.D. Samoylenko²⁴⁾, A. Sandacz³⁰⁾, H. Santos¹⁵⁾, M.G. Sapozhnikov⁸⁾, I.A. Savin⁸⁾, P. Schiavon²⁸⁾, C. Schill¹⁰⁾, L. Schmitt²⁰⁾, W. Schroeder⁹⁾, D. Seeharsch²⁰⁾, M. Seimetz²⁵⁾, D. Setter¹⁰⁾, O.Yu. Shevchenko⁸⁾, H.-W. Siebert^{12,4)}, L. Silva¹⁵⁾, L. Sinha⁷⁾, A.N. Sissakian⁸⁾, M. Slunecka⁸⁾, G.I. Smirnov⁸⁾, F. Sozzi²⁸⁾, A. Srnka⁵⁾, F. Stinzing⁹⁾, M. Stolarski³⁰⁾, V.P. Sugonyaev²⁴⁾, M. Sulc¹⁴⁾, R. Sulej³¹⁾, V.V. Tchalishvili⁸⁾, S. Tessaro²⁸⁾, F. Tessarotto²⁸⁾, A. Teufel⁹⁾, L.G. Tkatchev⁸⁾, S. Trippel¹⁰⁾, G. Venugopal³⁾, M. Virius²³⁾, N.V. Vlassov⁸⁾, R. Webb⁹⁾, E. Weise^{3,10)}, Q. Weitzel²⁰⁾, R. Windmolders⁴⁾, W. Wislicki³⁰⁾, K. Zarembo³¹⁾, M. Zavertyaev¹⁸⁾, E. Zemlyanichkina⁸⁾, J. Zhao^{16,25)}, R. Ziegler³⁾, and A. Zvyagin¹⁹⁾

¹⁾ Universität Bielefeld, Fakultät für Physik, 33501 Bielefeld, Germany^{g)}

²⁾ Universität Bochum, Institut für Experimentalphysik, 44780 Bochum, Germany^{g)}

³⁾ Universität Bonn, Helmholtz-Institut für Strahlen- und Kernphysik, 53115 Bonn, Germany^{g)}

⁴⁾ Universität Bonn, Physikalisches Institut, 53115 Bonn, Germany^{g)}

-
- 5) Institute of Scientific Instruments, AS CR, 61264 Brno, Czech Republic^{h)}
 - 6) Burdwan University, Burdwan 713104, India^{j)}
 - 7) Matrivani Institute of Experimental Research & Education, Calcutta-700 030, India^{k)}
 - 8) Joint Institute for Nuclear Research, 141980 Dubna, Moscow region, Russia
 - 9) Universität Erlangen–Nürnberg, Physikalisches Institut, 91054 Erlangen, Germany^{g)}
 - 10) Universität Freiburg, Physikalisches Institut, 79104 Freiburg, Germany^{g)}
 - 11) CERN, 1211 Geneva 23, Switzerland
 - 12) Universität Heidelberg, Physikalisches Institut, 69120 Heidelberg, Germany^{g)}
 - 13) Helsinki University of Technology, Low Temperature Laboratory, 02015 HUT, Finland and University of Helsinki, Helsinki Institute of Physics, 00014 Helsinki, Finland
 - 14) Technical University in Liberec, 46117 Liberec, Czech Republic^{h)}
 - 15) LIP, 1000-149 Lisbon, Portugalⁱ⁾
 - 16) Universität Mainz, Institut für Kernphysik, 55099 Mainz, Germany^{g)}
 - 17) University of Miyazaki, Miyazaki 889-2192, Japan^{l)}
 - 18) Lebedev Physical Institute, 119991 Moscow, Russia
 - 19) Ludwig-Maximilians-Universität München, Department für Physik, 80799 Munich, Germany^{g)}
 - 20) Technische Universität München, Physik Department, 85748 Garching, Germany^{g)}
 - 21) Nagoya University, 464 Nagoya, Japan^{l)}
 - 22) Charles University, Faculty of Mathematics and Physics, 18000 Prague, Czech Republic^{h)}
 - 23) Czech Technical University in Prague, 16636 Prague, Czech Republic^{h)}
 - 24) State Research Center of the Russian Federation, Institute for High Energy Physics, 142281 Protvino, Russia
 - 25) CEA DAPNIA/SPhN Saclay, 91191 Gif-sur-Yvette, France
 - 26) Tel Aviv University, School of Physics and Astronomy, 69978 Tel Aviv, Israel^{m)}
 - 27) INFN Trieste and ICTP–INFN MLab Laboratory, 34014 Trieste, Italy
 - 28) INFN Trieste and University of Trieste, Department of Physics, 34127 Trieste, Italy
 - 29) INFN Turin and University of Turin, Physics Department, 10125 Turin, Italy
 - 30) Sołtan Institute for Nuclear Studies and Warsaw University, 00-681 Warsaw, Polandⁿ⁾
 - 31) Warsaw University of Technology, Institute of Radioelectronics, 00-665 Warsaw, Poland^{o)}
- a) Also at IST, Universidade Técnica de Lisboa, Lisbon, Portugal
 - b) Also at University of East Piedmont, 15100 Alessandria, Italy
 - c) deceased
 - d) On leave of absence from JINR Dubna
 - e) Also at Chubu University, Kasugai, Aichi, 487-8501 Japan
 - f) Also at Yamagata University, Yamagata, 992-8510 Japan
 - g) Supported by the German Bundesministerium für Bildung und Forschung
 - h) Supported by Czech Republic MEYS grants ME492 and LA242
 - i) Supported by the Portuguese FCT - Fundação para a Ciência e Tecnologia grants POCTI/FNU/49501/2002 and POCTI/FNU/50192/2003
 - j) Supported by DST-FIST II grants, Govt. of India
 - k) Supported by the Shailabala Biswas Education Trust
 - l) Supported by the Ministry of Education, Culture, Sports, Science and Technology, Japan; Daikou Foundation and Yamada Foundation
 - m) Supported by the Israel Science Foundation, founded by the Israel Academy of Sciences and Humanities
 - n) Supported by KBN grant nr 621/E-78/SPUB-M/CERN/P-03/DZ 298 2000 and nr 621/E-78/SPB/CERN/P-03/DWM 576/2003-2006
 - o) Supported by KBN grant nr 134/E-365/SPUB-M/CERN/P-03/DZ299/2000

The spin structure function g_1^d of the deuteron has been measured for the first time almost 15 years ago by the SMC experiment at CERN [1]. Since then, high accuracy measurements of g_1^d in the deep inelastic scattering (DIS) region have been performed at SLAC [2,3] and DESY [4]. Due to the relatively low incident energy, the DIS events collected in those experiments cover only a limited range of x for $Q^2 > 1 (\text{GeV}/c)^2$: $x > 0.015$ and $x > 0.03$, respectively. Further measurements covering the low x region were also performed at CERN (see [5] and references therein). Besides its general interest for the understanding of the spin structure of the nucleon, g_1^d is specially important because its first moment is directly related to the matrix element of the singlet axial vector current a_0 . A precise measurement of g_1^d can thus provide an evaluation of the fraction of nucleon spin carried by quarks, on the condition that the covered range extends far enough to low x to provide a reliable value of the first moment.

Here we present new results from the COMPASS experiment at CERN on the deuteron spin asymmetry A_1^d and the spin-dependent structure function g_1^d covering the range $1 (\text{GeV}/c)^2 < Q^2 < 100 (\text{GeV}/c)^2$ in the photon virtuality and $0.004 < x < 0.7$ in the Bjorken scaling variable. The data sample used in the present analysis was collected during the years 2002–2004 and corresponds to an integrated luminosity of about 2 fb^{-1} . Partial results based on the data collected during the first two years of the data taking have been published in Ref. [6]. At the time, the values of g_1^d were not precise enough, in particular at large x , to allow a meaningful evaluation of the first moment, Γ_1^d . The results presented here are based on a 2.5 times larger statistics and supersede those of Ref. [6]. We refer the reader to this reference for the description of the 160 GeV muon beam, the ^6LiD polarised target and the COMPASS spectrometer which remained basically unchanged in 2004.

The COMPASS data acquisition system is triggered by coincidence signals in hodoscopes, defining the direction of the scattered muon behind the spectrometer magnets, and by signals in the hadron calorimeters [7]. Triggers due to halo muons are eliminated by veto counters installed upstream of the target. Inclusive triggers, based on muon detection only, cover the full range of x and are dominant in the medium (x, Q^2) region. Semi-inclusive triggers, based on the muon energy loss and the presence of a hadron signal in the calorimeters, contribute mainly at low x and low Q^2 . Purely calorimetric triggers, based on the energy deposit in the hadron calorimeter without any condition on the scattered muon, account for most events at large Q^2 . The relative contributions of these three trigger types are shown in Fig. 1 as a function of x . The minimum hadron energy deposit required for the purely calorimetric trigger has been reduced to 10 GeV for the events collected in 2004. As a consequence, the contribution of this trigger now reaches 40% at large x , compared to 20% in 2002–2003 (Ref. [6]).

All events used in the present analysis require the presence of reconstructed beam muon and scattered muon trajectories defining an interaction point, which is located inside one of the target cells. The momentum of the incoming muon, measured in the beam spectrometer, is centered around 160 GeV/ c with an RMS of 8 GeV/ c for the Gaussian core. In the present analysis its value is required to be between 140 and 180 GeV/ c . In addition the extrapolated beam muon trajectory is required to cross entirely both target cells in order to equalize the fluxes seen by each of them. The scattered muon is identified by signals collected behind the hadron absorbers and (except for the purely calorimetric trigger) its trajectory must be consistent with the hodoscope signals defining the event trigger. For hadronic triggers, a second outgoing reconstructed track is required at the interaction point. The DIS events used in the present analysis are selected by cuts on the four-momentum transfer squared ($Q^2 > 1 (\text{GeV}/c)^2$) and the fractional energy of the virtual photon ($0.1 < y < 0.9$). The resulting sample consists of 89×10^6 events, out of which about 10% were obtained in 2002, 30% in 2003 and 60% in 2004. In order to extend the coverage of the low x region, we also analyse events in the interval $0.003 < x < 0.004$

selected in the same way but with a Q^2 cut lowered to $0.7 \text{ (GeV}/c)^2$. These events are included in the figures but not used in QCD calculations or moment estimation, in view of their low Q^2 .

During data taking the two target cells are polarised in opposite directions, so that the deuteron spins are parallel ($\uparrow\uparrow$) or antiparallel ($\uparrow\downarrow$) to the spins of the incoming muons. The spins are inverted every 8 hours by a rotation of the target magnetic field. The average beam and target polarisations are about -0.80 (-0.76 in 2002 and 2003) and ± 0.50 , respectively.

The cross-section asymmetry $A^d = (\sigma^{\uparrow\downarrow} - \sigma^{\uparrow\uparrow})/(\sigma^{\uparrow\downarrow} + \sigma^{\uparrow\uparrow})$, for antiparallel ($\uparrow\downarrow$) and parallel ($\uparrow\uparrow$) spins of the incoming muon and the target deuteron can be obtained from the numbers of events N_i collected from each cell before and after reversal of the target spins:

$$N_i = a_i \phi_i n_i \bar{\sigma} (1 + P_B P_T f A^d), \quad i = 1, 2, 3, 4, \quad (1)$$

where a_i is the acceptance, ϕ_i the incoming flux, n_i the number of target nucleons, $\bar{\sigma}$ the spin-averaged cross-section, P_B and P_T the beam and target polarisations and f the target dilution factor. The latter includes a corrective factor $\rho = \sigma_d^{1\gamma}/\sigma_d^{tot}$ [8] accounting for radiative events on the unpolarised deuteron and a correction for the relative polarisation of deuterons bound in ${}^6\text{Li}$ compared to free deuterons. Fluxes and acceptances cancel out in the asymmetry calculation on the condition that the ratio of the acceptances of the two cells is the same before and after spin reversal [9].

The longitudinal virtual-photon deuteron asymmetry, A_1^d , is defined via the asymmetry of absorption cross-sections of transversely polarised photons as

$$A_1^d = (\sigma_0^T - \sigma_2^T)/(2\sigma^T), \quad (2)$$

where σ_J^T is the γ^* -deuteron absorption cross-section for a total spin projection J and σ^T is the total transverse photoabsorption cross-section. The relation between A_1^d and the experimentally measured A^d is

$$A^d = D(A_1^d + \eta A_2^d), \quad (3)$$

where D and η depend on kinematics. The transverse asymmetry A_2^d has been measured at SLAC and found to be small [10]. In view of this, in our analysis, Eq. (3) has been reduced to $A_1^d \simeq A^d/D$. The virtual-photon depolarisation factor D depends on the ratio of longitudinal and transverse photoabsorption cross sections $R = \sigma^L/\sigma^T$. In the present analysis an updated parametrisation of R taking into account all existing measurements is used [11]. The tensor-polarised structure function of the deuteron has been measured by HERMES [12] and its effect on the measurement of the longitudinal spin structure was found to be negligible, which justifies the use of Eqs (1-3) in the present analysis.

In order to minimize the statistical error of the asymmetry, the kinematic factors f , D and the beam polarisation P_B are calculated event-by-event and used to weight events. A parametrisation of P_B as a function of the beam momentum is used, while for P_T an average value is used for the data sample taken between two consecutive target spin reversals. The obtained asymmetry is corrected for spin-dependent radiative effects according to Ref. [13]. The asymmetry is evaluated separately for inclusive and for hadronic events because the dilution factors and the radiative corrections to the asymmetry are different. This is because the correction due to radiative elastic and quasi-elastic scattering events only affects the inclusive sample.

It has been checked that the use of hadronic triggers does not bias the inclusive asymmetries. The most critical case is for the calorimetric trigger events at large x , where high-energy hadron production is limited by kinematics. This effect has been studied by Monte Carlo, using the program POLDIS [14]. DIS events were generated within the acceptance of the calorimetric

trigger and their asymmetry calculated analytically at the leading order. A selection based on the hadron requirements corresponding to the trigger was applied and the asymmetries for the selected sample compared to the original ones. The differences were found to be smaller than 0.001 in all intervals of x (Fig. 2) and thus negligible, so that inclusive and hadronic asymmetries can be safely combined for further analysis (see also the SMC analysis [5]).

The final values of $A_1^d(x, Q^2)$, obtained as weighted averages of the asymmetries in the inclusive and hadronic data sets, are listed in Table 1 with the corresponding average values of x and Q^2 . They are also shown as a function of x in Fig. 3 in comparison with previous results from experiments at CERN [5], DESY [4] and SLAC [2, 3]. The values of A_1^d confirm, with increased statistical precision, the observation made in Ref. [6] that the asymmetry is consistent with zero for $x < 0.03$. Values of A_1^d originating from experiments at different energies tend to coincide due to the very small Q^2 dependence of A_1^d at fixed x .

The systematic error of A_1^d contains multiplicative factors resulting from uncertainties on P_B and P_T , on the dilution factor f and on the ratio $R = \sigma^L/\sigma^T$ used to calculate the depolarisation factor D . When combined in quadrature, these errors result in a global scale uncertainty of 10% (Table 2). The other important contribution to the systematic error is due to false asymmetries which could be generated by instabilities in some components of the spectrometer. In order to minimize their effect, the values of A_1^d in each interval of x have been calculated for 184 subsamples, each of them covering a short period of running time and, therefore, ensuring similar detector operating conditions. An upper limit of the effect of detector instabilities has been evaluated by a statistical approach. The dispersion of the values of A_1^d around their mean agrees with the statistical error. There is thus no evidence for any broadening due to time dependent effects. Allowing the dispersion of A_1^d to vary within its two standard deviations we obtain an upper limit for the systematic error of A_1^d in terms of its statistical precision: $\sigma_{syst} < 0.4\sigma_{stat}$. This estimation accounts for the time variation effects of spectrometer components.

Several other searches for false asymmetries were performed. Data from the two target cells were combined in different ways in order to eliminate the physical asymmetry. Data obtained with different settings of the microwave frequencies, used to polarise the target by dynamic nuclear polarisation, were compared. No evidence was found for any significant apparatus induced asymmetry.

The longitudinal spin structure function is obtained as

$$g_1^d = \frac{F_2^d}{2x(1+R)} A_1^d, \quad (4)$$

where F_2^d is the spin-independent deuteron structure function. The values of g_1^d listed in the last column of Table 1 have been calculated with the F_2^d parametrisation of Ref. [5], which covers the range of our data, and the new parametrisation of R already used in the depolarisation factor. The systematic errors on g_1^d are obtained in the same way as for A_1^d , with an additional contribution from the uncertainty on F_2^d . The values of $x g_1^d(x)$ for the COMPASS data and, for comparison, the SMC results [5] moved to the Q^2 of the corresponding COMPASS point are shown in Fig. 4. The two curves on the figure represent the results of two QCD fits at NLO, described below, at the measured Q^2 of each data point.

The evaluation of the first moment $\Gamma_1^d(Q^2) = \int_0^1 g_1^d(x, Q^2) dx$ requires the evolution of all g_1 measurements to a common Q_0^2 . This is done by using a fitted parametrisation $g_1^{fit}(x, Q^2)$, so that

$$g_1(x, Q_0^2) = g_1(x, Q^2) + \left[g_1^{fit}(x, Q_0^2) - g_1^{fit}(x, Q^2) \right]. \quad (5)$$

We have used several fits of g_1 from the Durham data base [15]: Blümlein-Böttcher [16], GRSV

[17] and LSS05 [18], and we have chosen $Q_0^2 = 3 (\text{GeV}/c)^2$ as reference Q^2 because it is close to the average Q^2 of the COMPASS DIS data. The three parametrisations are quite similar in the range of the COMPASS data and have been averaged. The resulting values of $g_1^N = (g_1^p + g_1^n)/2$ are shown as open squares in Fig. 5. For clarity we now use g_1^N instead of g_1^d because the correction for the D-wave state of the deuteron has been applied:

$$g_1^N(x, Q^2) = g_1^d(x, Q^2)/(1 - 1.5\omega_D) \quad (6)$$

with $\omega_D = 0.05 \pm 0.01$ [19]. It can also be seen in Fig. 5 that the curve representing the average of the three fits does not reproduce the trend of our data for $x < 0.02$ and therefore cannot be used to estimate the unmeasured part of g_1^N at low x .

In view of this, we have performed a new NLO QCD fit of all g_1 data at $Q^2 > 1 (\text{GeV}/c)^2$ from proton, deuteron and ^3He targets, including the COMPASS data. The deuteron data are from Refs [2–5], the proton data from Refs [2, 4, 5, 20, 21] and the ^3He data from Refs [22–25].

In order to optimise the use of the COMPASS data in this fit, all x bins of Table 1, except the last one, have been subdivided into three Q^2 intervals (Fig. 6). The number of COMPASS data points used in the fit is thus 43, out of a total of 230.

The fit is performed in the $\overline{\text{MS}}$ renormalisation and factorisation scheme and requires parametrisations of the quark singlet spin distribution $\Delta\Sigma(x)$, non-singlet distributions $\Delta q_3(x)$, $\Delta q_8(x)$ and the gluon spin distribution $\Delta G(x)$. These distributions are given as an input at a reference $Q^2 (=Q_0^2)$ which is set to $3 (\text{GeV}/c)^2$ and evolved according to the DGLAP equations. The resulting values of $g_1(x, Q^2)$ are calculated for the (x_i, Q_i^2) of each data point and compared to the experimental values.

The input parametrisations are written as

$$\Delta F_k = \eta_k \frac{x^{\alpha_k} (1-x)^{\beta_k} (1+\gamma_k x)}{\int_0^1 x^{\alpha_k} (1-x)^{\beta_k} (1+\gamma_k x) dx}, \quad (7)$$

where ΔF_k represents each of the polarised parton distribution functions (PDF) $\Delta\Sigma$, Δq_3 , Δq_8 and ΔG , and η_k is the integral of ΔF_k . The moments, η_k , of the non-singlet distributions Δq_3 and Δq_8 are fixed by the baryon decay constants (F+D) and (3F–D) respectively [26], assuming $\text{SU}(3)_f$ flavour symmetry. The linear term γx is used only for the singlet distribution, in which case the exponent β_G is fixed because it is poorly constrained by the data. This leaves 10 parameters in the input distributions. In addition, the normalisation of E155 proton data is allowed to vary within the limits quoted by the authors of Ref. [21].

The optimal values of the parameters are obtained by minimizing the sum

$$\chi^2 = \sum_{i=1}^{N=230} \frac{[g_1^{fit}(x_i, Q_i^2) - g_1^{exp}(x_i, Q_i^2)]^2}{[\sigma(x_i, Q_i^2)]^2}. \quad (8)$$

Here the errors σ are the statistical ones for all data sets, except for the proton data of E155 where the uncorrelated part of the systematic error on each point is added in quadrature to the statistical one. In order to keep the parameters in their physical range, the polarised strange sea distribution $\Delta s(x) + \Delta \bar{s}(x) = (1/3)(\Delta\Sigma(x) - \Delta q_8(x))$ is calculated at every step and required to satisfy the positivity condition $|\Delta s(x)| \leq s(x)$ at all Q^2 values. A similar condition is imposed on the gluon spin distribution $\Delta G(x)$. The unpolarised distributions $s(x)$ and $G(x)$ used in this test are taken from the MRST parametrisation [27]. This procedure leads to asymmetric errors on the parameters when the fitted value is close to the allowed limit.

The fits have been performed with two different programs: the first one uses the DGLAP evolution equations for the spin structure functions [28], the other one, referred to in [29], uses the evolution of moments. Both programs give consistent values of the fitted PDFs and similar χ^2 -probabilities. Each program yields two solutions, one solution with $\Delta G > 0$, the other one with $\Delta G < 0$ (Fig. 5). The fitted distributions of $g_1^N(x)$ differ at low x but are both compatible with the data. The two additional data points at $x < 0.004$ and $Q^2 > 0.7$ (GeV/c)², not used in the fit, have too large statistical errors to provide a discrimination between the two solutions. The values of the parameters obtained in the fits with positive and negative ΔG are listed in Table 3 with their statistical errors and will be discussed below.

The integral of g_1^N in the measured region is obtained from the experimental values evolved to a fixed Q^2 and averaged over the two fits. Taking into account the contributions from the fits in the unmeasured regions at low and high x we obtain (Table 4):

$$\Gamma_1^N(Q^2=3 \text{ (GeV/c)}^2) = 0.050 \pm 0.003(\text{stat.}) \pm 0.002(\text{evol.}) \pm 0.005(\text{syst.}). \quad (9)$$

The second error accounts for the difference in Q^2 evolution between the two fits. The systematic error is the dominant one and mainly corresponds to the 10% scale uncertainty resulting from the errors on the beam and target polarisations and on the dilution factor.

For comparison, the SMC result was [5]

$$\Gamma_{1,SMC}^N(Q^2=10 \text{ (GeV/c)}^2) = 0.021 \pm 0.007(\text{stat.}) \pm 0.014(\text{evol.}) \pm 0.003(\text{syst.}). \quad (10)$$

while our result evolved to $Q^2 = 10$ (GeV/c)² is $0.051 \pm 0.003(\text{stat.}) \pm 0.002(\text{evol.}) \pm 0.005(\text{syst.})$. The difference between these two results reflects the fact that the COMPASS data do not support the fast decrease of $g_1^d(x, Q_0^2 = 3 \text{ (GeV/c)}^2)$ at low x which was assumed in the SMC analysis, and thus force the fit to be different. In the COMPASS analysis, the part of Γ_1^N obtained from the measured region represents 98% of the total value. This correction of only 2% has to be compared to a correction of about 50% with respect to the measured value in case of the SMC analysis [5].

Γ_1^N is of special interest because it gives access to the matrix element of the singlet axial current a_0 which, except for a possible gluon contribution, measures the quark spin contribution to the nucleon spin. At NLO, the relation between Γ_1^N and a_0 reduces to

$$\Gamma_1^N(Q^2) = \frac{1}{9} \left(1 - \frac{\alpha_s(Q^2)}{\pi} + \mathcal{O}(\alpha_s^2) \right) \left(a_0(Q^2) + \frac{1}{4} a_8 \right). \quad (11)$$

From the COMPASS result on Γ_1^N (Eq. (9)) and taking the value of a_8 from hyperon β decay, assuming $SU(3)_f$ flavour symmetry ($a_8 = 0.585 \pm 0.025$ [26]), one obtains with the value of α_s evolved from the PDG value $\alpha_s(M_z^2) = 0.1187 \pm 0.005$:

$$a_0(Q^2=3 \text{ (GeV/c)}^2) = 0.35 \pm 0.03(\text{stat.}) \pm 0.05(\text{syst.}). \quad (12)$$

The quoted systematic error accounts for the error from the evolution and for the experimental systematic error, combined in quadrature.

The relation between Γ_1^N and a_0 can also be rewritten in order to extract the value of the matrix element a_0 in the limit $Q^2 \rightarrow \infty$. Here we will follow a notation of Ref. [30] introducing a ‘‘hat’’ for the coefficient C_1^S and a_0 at this limit:

$$\Gamma_1^N(Q^2) = \frac{1}{9} \hat{C}_1^S(Q^2) \hat{a}_0 + \frac{1}{36} C_1^{NS}(Q^2) a_8.$$

The coefficients \hat{C}_1^S and C_1^{NS} have been calculated in perturbative QCD up to the third order in $\alpha_s(Q^2)$ [30]:

$$\begin{aligned}\hat{C}_1^S(Q^2) &= 1 - 0.33333\left(\frac{\alpha_s}{\pi}\right) - 0.54959\left(\frac{\alpha_s}{\pi}\right)^2 - 4.44725\left(\frac{\alpha_s}{\pi}\right)^3 \\ C_1^{NS}(Q^2) &= 1 - \left(\frac{\alpha_s}{\pi}\right) - 3.5833\left(\frac{\alpha_s}{\pi}\right)^2 - 20.2153\left(\frac{\alpha_s}{\pi}\right)^3.\end{aligned}$$

With α_s evolved at the same order, one obtains

$$\hat{a}_0 = 0.33 \pm 0.03(stat.) \pm 0.05(syst.). \quad (13)$$

Combining this value with a_8 , the first moment of the strange quark spin distribution in the limit $Q^2 \rightarrow \infty$ is found to be

$$(\Delta s + \Delta \bar{s})_{Q^2 \rightarrow \infty} = \frac{1}{3}(\hat{a}_0 - a_8) = -0.08 \pm 0.01(stat.) \pm 0.02(syst.). \quad (14)$$

Previous fits of g_1 , not including the COMPASS data, found a positive $\Delta G(x)$ and a fitted function $g_1^d(x)$ becoming negative for $x \lesssim 0.025$ at $Q^2 = 3 \text{ (GeV}/c)^2$, as shown by the dotted line in Fig. 5. The new COMPASS data do not reveal any evidence for a decrease of the structure function at limit $x \rightarrow 0$. For our fit the data are still compatible with a positive ΔG , as shown by the full line in Fig. 5. However in this case an unexpected dip appears at $x \simeq 0.25$. The origin of this dip is related to the shape of the fitted $\Delta G(x)$, shown in Fig. 7 (left). Indeed, the gluon spin distribution must be close to zero at low x , to avoid pushing g_1^d down to negative values, and is also strongly limited at higher x by the positivity constraint $|\Delta G(x)| < G(x)$. The whole distribution is thus squeezed in a narrow interval around the maximum at $x \simeq \alpha_G/(\alpha_G + \beta_G) \simeq 0.25$.

In contrast, the fit with negative ΔG reproduces very well the COMPASS low x data with a much smoother distribution of $\Delta G(x)$ (dashed line on Fig. 5) and without approaching the positivity limit (Fig. 7, right). The $(1 + \gamma x)$ factor in the singlet quark distribution is not used in this case because it does not improve the confidence level of the fit.

Comparing the fitted parameters for ΔG positive and negative (Table 3), we observe that the parameters of the non-singlet distributions $\Delta q_3(x)$ and $\Delta q_8(x)$ are practically identical. The value of η_Σ is slightly larger in the fit with $\Delta G < 0$, as could be expected since in this case $\Delta \Sigma(x)$ remains positive over the full range of x :

$$\eta_\Sigma(Q^2=3 \text{ (GeV}/c)^2) = 0.28 \pm 0.01(stat.) (\Delta G > 0), \quad (15)$$

$$\eta_\Sigma(Q^2=3 \text{ (GeV}/c)^2) = 0.32 \pm 0.01(stat.) (\Delta G < 0). \quad (16)$$

We remind that in \overline{MS} scheme η_Σ is identical to the matrix element a_0 .

The singlet moment derived from the fits to all g_1 data is thus:

$$\eta_\Sigma(Q^2=3 \text{ (GeV}/c)^2) = 0.30 \pm 0.01(stat.) \pm 0.02(evol.). \quad (17)$$

Here we have taken the difference between the fits as an estimate of the systematic error and do not further investigate other contributions related to the choice of the QCD scale or the PDF parametrisations. The singlet moment obtained with COMPASS data alone (Eq. (12)) is slightly above this value and its statistical error is larger by a factor of 3. As stated before, the main uncertainty on the COMPASS result is due to the 10% normalisation uncertainty from the

beam and target polarisations and from the dilution factor. The fact that the COMPASS data are on average slightly above the world average can already be detected by a comparison of the measured g_1^d values to the curves fitted to the world data (Fig. 5). Hence a_0 derived from the COMPASS value of Γ_1^N is found to be slightly larger than η_Σ .

The polarised strange quark distributions, obtained from the difference between $\Delta\Sigma(x)$ and $\Delta q_8(x)$ are shown in Fig. 8. They are negative and concentrated in the highest x region, compatible with the constraint $|\Delta s(x)| < s(x)$. This condition is indeed essential in the determination of the Δq_8 parameters which otherwise would be poorly constrained.

Although the gluon distributions strongly differ in the two fits, the fitted values of their first moments are both small and about equal in absolute value $|\eta_G| \approx 0.2 - 0.3$. In Fig. 9 the existing direct measurements of $\Delta G/G$ [31–33] are shown with the distributions of $\Delta G(x)/G(x)$ derived from our fits with $G(x)$ taken from Ref. [27]. The HERMES value is positive and 2σ away from zero. The measured SMC point is too unprecise to discriminate between positive or negative ΔG . The published COMPASS point, which has been obtained from a partial data sample corresponding to about 40% of the present statistics, is almost on the $\Delta G > 0$ curve but is only 1.3σ away from the $\Delta G < 0$ one, so that no preference for any of the curves can be given so far. It should also be noted that the measured values of $\Delta G/G$ have all been obtained in leading order QCD analyses.

In summary, we have measured the deuteron spin asymmetry A_1^d and its longitudinal spin-dependent structure function g_1^d with improved precision at $Q^2 > 1$ (GeV/c)² over the range $0.004 < x < 0.70$. The g_1^d values are consistent with zero for $x < 0.03$. The measured values have been evolved to a common Q^2 by a new fit of the world g_1 data, and the first moment Γ_1^N has been evaluated at $Q^2 = 3$ (GeV/c)² with a statistical error smaller than 0.003. From Γ_1^N we have derived the matrix element of the singlet axial current \hat{a}_0 in the limit $Q^2 \rightarrow \infty$. With COMPASS data alone, at the order α_s^3 , it has been found that $\hat{a}_0 = 0.33 \pm 0.03(stat.) \pm 0.05(syst.)$ and the first moment of the strange quark distribution $(\Delta s + \Delta \bar{s})_{Q^2 \rightarrow \infty} = -0.08 \pm 0.01(stat.) \pm 0.02(syst.)$. We also observe that the fit of world g_1 data at NLO yields two solutions with either $\Delta G(x) > 0$ or $\Delta G(x) < 0$, which equally well describe the present data. In both cases, the first moment of $\Delta G(x)$ is of the order of $0.2 - 0.3$ in absolute value at $Q^2 = 3$ (GeV/c)² but the shapes of the distributions are very different.

Acknowledgements

We gratefully acknowledge the support of the CERN management and staff and the skill and effort of the technicians of our collaborating institutes. Special thanks are due to V. Anosov and V. Pesaro for their technical support during the installation and the running of this experiment. This work was made possible by the financial support of our funding agencies.

References

- [1] SMC Collaboration, B. Adeva *et al.*, Phys. Lett. B **302** (1993) 533.
- [2] E143 Collaboration, K. Abe *et al.*, Phys. Rev. D **58** (1998) 112003.
- [3] E155 Collaboration, P. L. Anthony *et al.*, Phys. Lett. B **463** (1999) 339.
- [4] HERMES Collaboration, A. Airapetian *et al.*, Phys. Rev. D **71** (2005) 012003.
- [5] SMC Collaboration, B. Adeva *et al.*, Phys. Rev. D **58** (1998) 112001.
- [6] COMPASS Collaboration, E. S. Ageev *et al.*, Phys. Lett. B **612** (2005) 154.
- [7] C. Bernet *et al.*, Nucl. Instrum. Methods A **550** (2005) 217.
- [8] A. A. Akhundov *et al.*, Fortsch. Phys. **44** (1996) 373.
- [9] SMC Collaboration, D. Adams *et al.*, Phys. Rev. D **56** (1997) 5330.
- [10] E155 Collaboration, P. L. Anthony *et al.*, Phys. Lett. B **553** (2003) 18.
- [11] E143 Collaboration, K. Abe *et al.*, Phys. Lett. B **452** (1999) 194.
- [12] HERMES Collaboration, A. Airapetian *et al.*, Phys. Rev. Lett. **95** (2005) 242001.
- [13] I. V. Akushevich and N. M. Shumeiko, J. Phys. G **20** (1994) 513.
- [14] A. Bravar, K. Kurek and R. Windmolders, Comput. Phys. Commun. **105** (1997) 42.
- [15] The Durham HEP Databases, <http://durpdg.dur.ac.uk/HEPDATA/pdf.html>
- [16] J. Blümlein and H. Böttcher, Nucl. Phys. B **636** (2002) 225.
- [17] M. Glück, E. Reya, M. Stratmann and W. Vogelsang, Phys. Rev. D **63** (2001) 094005.
- [18] E. Leader, A. V. Sidorov and D. B. Stamenov, Phys. Rev. D **73** (2006) 034023.
- [19] R. Machleidt *et al.*, Phys. Rep. **149** (1987) 1.
- [20] EMC Collaboration, J. Ashman *et al.*, Nucl. Phys. B **328** (1989) 1.
- [21] E155 Collaboration, P. L. Anthony *et al.*, Phys. Lett. B **493** (2000) 19.
- [22] E142 Collaboration, P. L. Anthony *et al.*, Phys. Rev. D **54** (1996) 6620.
- [23] E154 Collaboration, K. Abe *et al.*, Phys. Rev. Lett. **79** (1997) 26.
- [24] JLAB/Hall A Collaboration, X. Zheng *et al.*, Phys. Rev. Lett. **92** (2004) 012004.
- [25] HERMES Collaboration, K. Ackerstaff *et al.*, Phys. Lett. B **404** (1997) 383.
- [26] Y. Goto *et al.*, Phys. Rev. D **62** (2000) 034017.
- [27] A. D. Martin *et al.*, Eur. Phys. J. C4 (1998) 463.
- [28] SMC Collaboration, B. Adeva *et al.*, Phys. Rev. D **58** (1998) 112002.
- [29] A. N. Sissakian, O. Yu. Shevchenko and O. N. Ivanov, Phys. Rev. D **70** (2004) 074032.
- [30] S. A. Larin *et al.*, Phys. Lett. B **404** (1997) 153.
- [31] SMC Collaboration, B. Adeva *et al.*, Phys. Rev. D **70** (2004) 012002.
- [32] HERMES Collaboration, A. Airapetian *et al.*, Phys. Rev. Lett. **84** (2000) 2584.
- [33] COMPASS Collaboration, E. S. Ageev *et al.*, Phys. Lett. B **633** (2006) 25.
- [34] F. James, MINUIT, CERN Program Library Long Writeup D506.

x range	$\langle x \rangle$	$\langle Q^2 \rangle$ [(GeV/c) ²]	A_1^d	g_1^d
0.0030–0.0035	0.0033	0.78	$0.003 \pm 0.009 \pm 0.004$	$0.090 \pm 0.240 \pm 0.107$
0.0035–0.0040	0.0038	0.83	$-0.004 \pm 0.007 \pm 0.003$	$-0.097 \pm 0.183 \pm 0.082$
0.004–0.005	0.0046	1.10	$0.004 \pm 0.009 \pm 0.004$	$0.082 \pm 0.210 \pm 0.089$
0.005–0.006	0.0055	1.22	$0.003 \pm 0.007 \pm 0.003$	$0.062 \pm 0.146 \pm 0.062$
0.006–0.008	0.0070	1.39	$-0.002 \pm 0.005 \pm 0.002$	$-0.034 \pm 0.086 \pm 0.036$
0.008–0.010	0.0090	1.61	$-0.010 \pm 0.006 \pm 0.003$	$-0.139 \pm 0.078 \pm 0.035$
0.010–0.020	0.0141	2.15	$0.002 \pm 0.004 \pm 0.002$	$0.017 \pm 0.033 \pm 0.014$
0.020–0.030	0.0244	3.18	$0.003 \pm 0.006 \pm 0.003$	$0.017 \pm 0.035 \pm 0.015$
0.030–0.040	0.0346	4.26	$0.009 \pm 0.008 \pm 0.004$	$0.041 \pm 0.035 \pm 0.016$
0.040–0.060	0.0487	5.80	$0.017 \pm 0.008 \pm 0.004$	$0.054 \pm 0.026 \pm 0.012$
0.060–0.100	0.0765	8.53	$0.058 \pm 0.009 \pm 0.007$	$0.121 \pm 0.019 \pm 0.014$
0.100–0.150	0.121	12.6	$0.095 \pm 0.013 \pm 0.011$	$0.123 \pm 0.017 \pm 0.014$
0.150–0.200	0.171	17.2	$0.123 \pm 0.020 \pm 0.014$	$0.103 \pm 0.016 \pm 0.012$
0.200–0.250	0.222	21.8	$0.183 \pm 0.028 \pm 0.021$	$0.106 \pm 0.016 \pm 0.012$
0.250–0.350	0.290	28.3	$0.216 \pm 0.030 \pm 0.024$	$0.077 \pm 0.011 \pm 0.009$
0.350–0.500	0.405	39.7	$0.343 \pm 0.049 \pm 0.038$	$0.055 \pm 0.008 \pm 0.006$
0.500–0.700	0.566	55.3	$0.626 \pm 0.112 \pm 0.075$	$0.027 \pm 0.005 \pm 0.003$

Table 1: Values of A_1^d and g_1^d with their statistical and systematical errors as a function of x with the corresponding average values of x and Q^2 . The minimum Q^2 cut is 1 (GeV/c)² except for the first two points where it is lowered to 0.7 (GeV/c)². These two data points are shown on the figures as complementary information but were not used in the fits.

Multiplicative variables error, ΔA_1^{mult}	Beam polarization	dP_B/P_B	5%
	Target polarization	dP_T/P_T	5%
	Depolarization factor	$dD(R)/D(R)$	2 – 3 %
	Dilution factor	df/f	6 %
	Total		$\Delta A_1^{mult} \simeq 0.1A_1$
Additive variables error, ΔA_1^{add}	Transverse asymmetry	$\eta/\rho \cdot \Delta A_2$	$10^{-4} - 5 \cdot 10^{-3}$
	Radiative corrections	ΔA_1^{RC}	$10^{-4} - 10^{-3}$
	False asymmetry	A_{false}	$< 0.4 \cdot \Delta A_1^{stat}$

Table 2: Decomposition of the systematic error of A_1 into multiplicative and additive variables contributions.

$\Delta G > 0$			$\Delta G < 0$		
	Prog. Ref. [28]	Prog. Ref. [29]		Prog. Ref. [28]	Prog. Ref. [29]
η_Σ	0.276 ± 0.013	0.288 ± 0.011	η_Σ	0.321 ± 0.009	$0.329^{+0.009}_{-0.008}$
α_Σ	$-0.285^{+0.073}_{-0.085}$	$-0.187^{+0.072}_{-0.065}$	α_Σ	$1.39^{+0.15}_{-0.14}$	1.40 ± 0.12
β_Σ	$3.61^{+0.26}_{-0.24}$	$3.81^{+0.25}_{-0.18}$	β_Σ	$4.09^{+0.29}_{-0.27}$	$4.10^{+0.24}_{-0.23}$
γ_Σ	$-16.6^{+1.6}_{-1.8}$	$-15.8^{+1.4}_{-1.0}$	γ_Σ	-	-
η_G	$0.263^{+0.038}_{-0.062}$	$0.194^{+0.012}_{-0.097}$	η_G	$-0.31^{+0.10}_{-0.14}$	$-0.181^{+0.042}_{-0.031}$
α_G	$6.15^{+0.58}_{-0.76}$	$9.9^{+1.0}_{-0.74}$	α_G	$0.39^{+0.64}_{-0.48}$	0.39 ± 0.17
β_G	20 (fixed)	30 (fixed)	β_G	$13.8^{+7.8}_{-5.3}$	$16.1^{+1.3}_{-4.0}$
α_3	$-0.221^{+0.028}_{-0.027}$	$-0.217^{+0.027}_{-0.027}$	α_3	-0.212 ± 0.027	$-0.208^{+0.027}_{-0.026}$
β_3	$2.43^{+0.11}_{-0.10}$	$2.40^{+0.11}_{-0.10}$	β_3	$2.44^{+0.11}_{-0.10}$	2.40 ± 0.10
α_8	$0.36^{+0.19}_{-0.44}$	$0.43^{+0.11}_{-0.41}$	α_8	0.42 ± 0.16	$0.347^{+0.071}_{-0.095}$
β_8	$3.37^{+0.63}_{-1.07}$	$3.51^{+0.42}_{-0.99}$	β_8	$3.53^{+0.56}_{-0.53}$	$3.31^{+0.30}_{-0.34}$
χ^2/ndf	233/219	234/219	χ^2/ndf	247/219	248/219

Table 3: Values of the parameters obtained from the QCD analysis at $Q^2 = 3 \text{ (GeV}/c)^2$ in fits with $\Delta G > 0$ and $\Delta G < 0$ with the two programs. The quoted errors correspond to one σ and have been obtained from the MINOS analysis [34]. The strongly asymmetric errors obtained for some parameters are due to the positivity constraints applied in the fits.

Range in x	COMPASS data evolved to $Q^2 = 3 \text{ (GeV}/c)^2$ using			
	fits of		COMPASS fits (prog. [28])	
	BB [16]	LSS [18]	$\Delta G > 0$	$\Delta G < 0$
[0.004, 0.7]	0.0455	0.0469	0.0479	0.0508
[0.7, 1]	0.0014	0.0008	0.0011	0.0010
[0, 0.004]	-0.0040	-0.0029	-0.0009	0.0004
[0, 1]	0.0430	0.0448	0.0481	0.0522

Table 4: Contributions to $\Gamma_1^N(Q^2 = 3 \text{ (GeV}/c)^2)$ from different kinematic regions. The values in the first line are the COMPASS results evolved according to different fits and integrated over the measured x range. The second and third lines show the corresponding high and low x extrapolations.




Figure 1: Fraction of inclusive, semi-inclusive, and calorimetric triggers as a function of x . Events are counted with the weight they carry in the asymmetry calculation.




Figure 2: Difference between asymmetries for inclusive and hadronic Monte-Carlo events in the kinematic range covered by the purely calorimetric trigger.




Figure 3: The asymmetry $A_1^d(x)$ as measured in COMPASS and previous results from SMC [5], HERMES [4], SLAC E143 [2] and E155 [3] at $Q^2 > 1$ (GeV/c) 2 . The SLAC values of g_1/F_1 have been converted to A_1 and the E155 data corresponding to the same x have been averaged over Q^2 . Only statistical errors are shown with the data points. The shaded areas show the size of the COMPASS systematic errors.




Figure 4: Values of $xg_1^d(x)$. The COMPASS points are given at the $\langle Q^2 \rangle$ where they were measured. The SMC points have been moved to the Q^2 of the corresponding COMPASS points. Only statistical errors are shown with the data points. The shaded band at the bottom shows the COMPASS systematic error. The curves show the results of QCD fits with $\Delta G > 0$ and $\Delta G < 0$.




Figure 5: The COMPASS values of g_1^N evolved to $Q^2 = 3 (\text{GeV}/c)^2$. The open triangles at low x correspond to $Q^2 > 0.7 (\text{GeV}/c)^2$, the other symbols to $Q^2 > 1 (\text{GeV}/c)^2$. Results of QCD fits are shown by curves. In addition to our fits ($\Delta G > 0$ and $\Delta G < 0$) the curve obtained with three published polarised PDF parameterizations (Blümlein and Böttcher, GRSV and LSS05) [15] is shown. These parameterizations lead almost to the same values of $g_1^N(x, Q^2 = 3 (\text{GeV}/c)^2)$ and have been averaged. For clarity the data points evolved with different fits are shifted in x with respect to each other. Only statistical errors are shown.




Figure 6: Values of A_1^d as a function of Q^2 in intervals of x . The solid lines show the results of fits to a constant.




Figure 7: Gluon distribution $x\Delta G(x)$ corresponding to the fits with $\Delta G > 0$ (left) and $\Delta G < 0$ (right) obtained with the program of Ref. [28]. The dashed, solid and dotted lines correspond to $Q^2 = 1.5, 3$ and 10 (GeV/c)^2 , respectively. The unpolarised distributions $\pm x G(x)$ which were used in the fit as constrains for the polarised ones are shown for $Q^2 = 1.5$ and 3 (GeV/c)^2 .




Figure 8: Strange quark distribution $x\Delta s(x)$ corresponding to the fits with $\Delta G > 0$ (left) and $\Delta G < 0$ (right) obtained with the program of Ref. [28]. The dashed, solid and dotted lines correspond to $Q^2 = 1.5, 3$ and 10 (GeV/c)^2 , respectively. The unpolarised distributions $\pm x s(x)$ are shown for $Q^2 = 1.5$ and 3 (GeV/c)^2 .




Figure 9: Distribution of the gluon polarisation $\Delta G(x)/G(x)$ at $Q^2 = 3 \text{ (GeV/c)}^2$ for the fits with $\Delta G > 0$ and $\Delta G < 0$ obtained with the program of Ref. [28]. The error bands correspond to the statistical error on $\Delta G(x)$ at a given x . The unpolarised gluon distribution is taken from the MRST parametrisation [27]. The three data points show the measured values from SMC [31], HERMES [32] and COMPASS [33]. Two error bars are associated to each data point, one corresponding to the statistical precision and the other one to the statistical and systematic errors added in quadrature. The horizontal bar on each point shows the x -range of measurement.