An Intro to Game Theory

15-451 Avrim Blum 12/02/03

Plan for Today

- 2-Player Zero-Sum Games (matrix games)
  - Minimax optimal strategies

test material

not test material

- General-Sum Games (bimatrix games)
  - notion of Nash Equilibrium
- Proof of existence of Nash Equilibria
  - using Brouwer's fixed-point theorem
- do FCEs at end...

2-Player Zero-Sum games

- Two players R and C. Zero-sum means that what's good for one is bad for the other.
- Game defined by matrix with a row for each of R's options and a column for each of C's options. Matrix tells who wins how much.
- E.g., matching pennies / penalty shot / hide-a-coin:

	Left	Right
Left	(-1,1)	(1,-1)
Right	(1,-1)	(-1,1)

shooter loses and goalie wins

shooter wins and goalie loses

An algorithmic example

Sorting three items (A,B,C):

- Compare two of them. Then compare 3rd to larger of 1st two. If we're lucky it's larger, else need one more comparison.

	largest is: C	B	A ← adversary
(A,B) first	2	3	3
(A,C) first	3	2	3
(B,C) first	3	3	2

(payoff to adversary)

Minimax-optimal strategies

- Minimax optimal strategy is a (randomized) strategy that has the best worst-case expected gain. [maximizes the minimum]
- I.e., it's the thing to play if your opponent knows you well.
- Same as our notion of a randomized strategy with a good worst-case bound.

Minimax-optimal strategies

Sorting three items (A,B,C): Compare two of them. Then compare 3rd to larger of 1st two. Minimax optimal cost is 2+(2/3).

	largest is: C	B	A ← adversary
(A,B) first	2	3	3
(A,C) first	3	2	3
(B,C) first	3	3	2

(payoff to adversary)

Minimax-optimal strategies

- E.g., matching pennies / penalty shot / hide-a-coin

	Left	Right
Left	(-1,1)	(1,-1)
Right	(1,-1)	(-1,1)

Minimax optimal for both players is 50/50. Gives expected gain of 0. Any other is worse.

Minimax-optimal strategies

- E.g., penalty shot with goalie who's weaker on the left.

	Left	Right
Left	(0,0)	(1,-1)
Right	(1,-1)	(-1,1)

Minimax optimal for both players is (2/3,1/3). Gives expected gain 1/3. Any other is worse.

Minimax Theorem (von Neumann 1928)

- Every 2-player zero sum game has a unique value V.
- Minimax optimal strategy for R guarantees R's expected gain at least V.
- Minimax optimal strategy for C guarantees R's expected gain at most V.

Counterintuitive: against an optimal opponent, it doesn't hurt to reveal your randomized strategy. (Borel had proved for symmetric 5x5 but thought was false for larger games)

Matrix games and Algorithms

- Gives a useful way of thinking about guarantees on algorithms.
- Think of rows as different algorithms, columns as different possible inputs.
- M(i,j) = cost of algorithm i on input j.

Of course matrix is HUGE. But helpful conceptually.

Matrix games and Algs

- What is a deterministic alg with a good worst-case guarantee?
  - A row that does well against all columns.
- What is a lower bound for deterministic algorithms?
  - Showing that for each row i there exists a column j such that M(i,j) is bad.
- How to give lower bound for randomized algs?
  - Give randomized strategy for adversary that is bad for all i.

E.g., hashing



- Rows are different hash functions.
- Cols are different sets of items to hash.
- M(i,j) = #collisions incurred by alg i on set j.
  [alg is trying to minimize]
- For any row, can reverse-engineer a bad column.
- Universal hashing is a randomized strategy for row player.

One more example

1-card poker in a 3-card deck {J,Q,K}:

	[FP,FP,CB]	[FP,CP,CB]	[FB,FP,CB]	[FB,CP,CB]
[PF,PF,PC]				
[PF,PF,B]				
[PF,PC,PC]				
[PF,PC,B]				
[B,PF,PC]				
[B,PF,B]				
[B,PC,PC]				
[B,PC,B]				

Minimax-optimal strategy

- Minimax optimal for 1st player is:
  - If hold J, then 5/6 PassFold and 1/6 Bet.
  - If hold Q, then ½ PassFold and ½ PassCall.
  - If hold K, then ½ PassCall and ½ Bet.
- Note the bluffing and underbidding… (Minimax for 2nd player has this too)
- Minimax value of game is –1/18 for 1st player and 1/18 for 2nd.
- See Chvatal, *Linear Programming*. Chap 15. (Remember can solve for minimax with LP)

General-sum games

- In general-sum (bimatrix) games, can have win-win and lose-lose situations.
- E.g., "what side of road to drive on?":

	Left	Right
Left	(1,1)	(-1,-1)
Right	(-1,-1)	(1,1)

General-sum games

- In general-sum (bimatrix) games, can have win-win and lose-lose situations.
- E.g., "which movie should we go to?":

	MatRev	loveactually
MatRev	(8,2)	(0,0)
loveactually	(0,0)	(2,8)

No longer a unique "value" to the game.

General-sum games

- Economists use as models of interaction.
- E.g., pollution / prisoner's dilemma:
  - (imagine pollution controls cost $4 and improve everyone's environment by $3)

	pollute	don't pollute
pollute	(-1,-1)	(2,-2)
don't pollute	(-2,2)	(1,1)

Need to add extra incentives to get desired behavior.

Nash Equilibrium

- A Nash Equilibrium is a stable pair of strategies (could be randomized).
- Stable means that neither player has incentive to deviate on their own.
- E.g., "what side of road to drive on":

	Left	Right
Left	(1,1)	(-1,-1)
Right	(-1,-1)	(1,1)

NE are: both left, both right, or both 50/50.

Nash Equilibrium

- A Nash Equilibrium is a stable pair of strategies (could be randomized).
- Stable means that neither player has incentive to deviate.
- E.g., "which movie to go to":

	MatRev	loveactually
MatRev	(8,2)	(0,0)
loveactually	(0,0)	(2,8)

NE are: both MR, both la, or (80/20,20/80)

Existence of NE

- Nash (1950) proved: any general-sum game must have at least one such equilibrium.
  - Might require randomized strategies (called "mixed strategies")
- This also yields minimax thm as a corollary.
  - Pick some NE and let V = value to row player in that equilibrium.
  - Since it's a NE, neither player can do better even knowing the (randomized) strategy their opponent is playing.
  - So, they're each playing minimax optimal.

Existence of NE

- Proof will be non-constructive.
- Unlike case of zero-sum games, we know of no polynomial-time algorithm for finding Nash Equilibria in general-sum games.
- Notation:
  - Assume an nxn matrix.
  - Use $(p_1,\ldots,p_n)$ to denote mixed strategy for row player, and $(q_1,\ldots,q_n)$ to denote mixed strategy for column player.

Proof

- We'll start with Brouwer's fixed point theorem.
  - Let S be a compact convex region in R^n and let $f:S \rightarrow S$ be a continuous function.
  - Then there must exist $x \in S$ such that $f(x)=x$.
  - x is called a "fixed point" of f.
- Simple case: S is the interval [0,1].
- We will care about:
  - $S = \{(p,q)$: p,q are legal probability distributions on $1,\ldots,n\}$. I.e., $S = \text{simplex}_n \times \text{simplex}_n$

Proof (cont)

- S = {(p,q): p,q are mixed strategies}.
- Want to define f(p,q) = (p',q') such that:
  - f is continuous. This means that changing p or q a little bit shouldn't cause p' or q' to change a lot.
  - Any fixed point of f is a Nash Equilibrium.

Try #1

- What about f(p,q) = (p',q') where p' is best response to q, and q' is best response to p?
- Problem: not continuous:
  - E.g., matching pennies. If p = (0.51, 0.49) then q' = (1,0). If p = (0.49,0.51) then q' = (0,1).

	Left	Right
Left	(-1,1)	(1,-1)
Right	(1,-1)	(-1,1)

Try #1

- What about f(p,q) = (p′,q′) where p′ is best response to q, and q′ is best response to p?
- Problem: also not necessarily well-defined:
  - E.g., if p = (0.5,0.5) then q′ could be anything.

	Left	Right
Left	(-1,1)	(1,-1)
Right	(1,-1)	(-1,1)

Instead we will use…

- f(p,q) = (p′,q′) such that:
  - q′ maximizes [(expected gain wrt p) - $||q-q'||^2$]
  - p′ maximizes [(expected gain wrt q) - $||p-p'||^2$]



Note: quadratic + linear = quadratic.

Instead we will use…

- f(p,q) = (p′,q′) such that:
  - q′ maximizes [(expected gain wrt p) - $||q-q'||^2$]
  - p′ maximizes [(expected gain wrt q) - $||p-p'||^2$]



Note: quadratic + linear = quadratic.

Instead we will use…

- f(p,q) = (p′,q′) such that:
  - q′ maximizes [(expected gain wrt p) - $||q-q'||^2$]
  - p′ maximizes [(expected gain wrt q) - $||p-p'||^2$]



Note: quadratic + linear = quadratic.

Instead we will use…

- f(p,q) = (p′,q′) such that:
  - q′ maximizes [(expected gain wrt p) - $||q-q'||^2$]
  - p′ maximizes [(expected gain wrt q) - $||p-p'||^2$]

- f is well-defined and continuous since quadratic has unique maximum and small change to p,q only moves this a little.
- Also fixed point = NE. (even if tiny incentive to move, will move little bit).
- So, that's it!