

Particle acceleration at relativistic shock waves ... and gamma-ray bursts

Martin Lemoine

Institut d'Astrophysique de Paris
CNRS, Université Pierre & Marie Curie



Outline:

1. Particle acceleration and relativistic collisionless shocks
2. Microphysics of gamma-ray burst afterglows

Relativistic Fermi acceleration - small scale turbulence



Test particle picture:


→ particles gain energy by bouncing across the shock front,

exploiting the convective electric fields : $\delta E = -\frac{v}{c} \times \delta B$

→ if $\gamma_{sh} \gg 1$, advection beats acceleration unless particles

scatter in small-scale turbulence $\lambda_B < r_g$, $\delta B > B$ and $r_g < \lambda_B \delta B/B$

(r_g gyroradius of accelerated particles, λ_B length scale of δB)
 (ML et al. 06, Niemiec et al. 06, Pelletier et al. 09)




$$\Leftarrow \sigma < \epsilon_B^2 (\lambda \omega_p / c)^2$$


weak magnetization!

PIC simulations:

(e.g. Spitkovsky 08, Nishikawa et al. 09, Martins et al. 09, Sironi & Spitkovsky 09, 11, 13, Haugbolle 11)




Phase diagram for relativistic shock acceleration



Caveats and open questions



Most PIC simulations have not converged to a stationary state! (Keshet et al. 09)



$$\varepsilon_B = \delta B^2 / (16\pi \gamma_{sh}^2 n_u m c^2)$$

→ Keshet et al. 09: **time = $10^4 \omega_p^{-1} \Leftrightarrow \sim 0.1\%$ of a dynamical timescale for a GRB!**

→ theoretical extrapolation is needed!

Main open questions:

→ phase space still largely unexplored... **mildly relativistic shocks = terra incognita**

→ high energy particles stream further away and modify the precursor: how?

→ other instabilities on larger (MHD?) scales?


→ acceleration at magnetized shocks, e.g. PWNe up to $\gamma_e \sim 10^9$?

Relativistic Fermi acceleration - unmagnetized limit



PIC simulations:

(e.g. Spitkovsky 08, Nishikawa et al. 09, Martins et al. 09, Sironi & Spitkovsky 09, 11, 13, Haugbolle 11)



→ supra-thermal particles stream ahead of the shock and excite plasma instabilities (Weibel/filamentation, two-stream, current-driven etc.), which build δB ...

→ δB builds a magnetic barrier ($\sim 10\%$ of equipartition) which mediates the shock transition...

→ δB on c/ω_p scales provides the scattering required for acceleration...

→ δB provides the turbulence in which particles radiate (?) (Medvedev & Loeb 99)

Maximum energy



Maximum energy:


- scattering in small scale turbulence $\lambda_B < r_g$ is not as efficient as Bohm...
- max energy for electrons by comparing $t_{\text{acc}} \sim t_{\text{scatt}}$ to synchrotron loss, with $t_{\text{scatt}} \sim r_g^2 / (\lambda_B c)$ and $\lambda_B \sim 10 c / \omega_p$, implies a maximum synchrotron photon energy:
(e.g. Kirk & Reville 10, Plotnikov et al. 13, Wang et al. 13, Sironi et al. 13):

$$\epsilon_{\gamma, \text{max}} \simeq 2 \text{ GeV } E_{54}^{1/4} \epsilon_{B,-2}^{1/2} \lambda_1^{2/3} n_0^{-1/12} t_{\text{obs},2}^{-3/4}$$

- **long-lived GeV emission on 1000sec can result from synchrotron afterglow**
(Kumar & Barniol-Duran 09, 10, Ghisellini et al. 10)
- ... photons above 10GeV result from IC interactions...** (Wang et al. 13)

in GRB130427A:

two spectral components with $\epsilon_{\gamma, \text{max}} \sim \text{GeV}$ at 100-1000 sec for the synchrotron afterglow...



Evolution of turbulence in GRB blast waves



Theory vs observations/phenomenology:

→ comparison between theory, PIC sims. and GRB phenomenology overall satisfactory:

electrons are heated to $\gamma_{\min} = \gamma_{\text{sh}} m_p/m_e \sim 10^5$,

to near equipartition $\epsilon_e \sim 0.1-0.5$... with a power-law tail of index $s \sim -2.2$


magnetized turbulence is excited up to $\epsilon_B \sim 0.01$ (canonical value !?)

→ actually, a long-standing notorious problem for ϵ_B : turbulence lies on plasma scales c/ω_{pi} , and should decay on 100's of c/ω_{pi} , whereas observations probe the width of the blast, many orders of magnitude beyond...

⇒ origin of the magnetisation of GRB blast waves? (e.g. Gruzinov 99, Gruzinov & Waxman 99)

$$\frac{l_{\text{blast}}}{c/\omega_{pi}} \approx 10^7 t_{\text{obs},2}^{5/8}$$

$$\frac{l_{\text{cool}}}{c/\omega_{pi}} \approx 10^7 t_2^{9/8} E_{53}^{-3/8} n_{-3}^{-1/8} \epsilon_{B,-2}^{-1} \epsilon_{e,-0.3}^{-1} \frac{\gamma_{\min}}{\gamma_e}$$



Spitkovsky 08


Evolution of turbulence in GRB blast waves



(ref. frame:
shocked plasma)

how does the turbulence evolve
with distance to shock?

damping or additional source of turbulence?
e.g. Gruzinov & Waxman 99, Medvedev & Loeb 99,
Chang et al. 08, Keshet et al. 09, ML 13



micro-instabilities associated
with the shock structure:
typically on plasma scales c/ω_{pi}


Microphysics of turbulence evolution:

→ through Landau damping, δB is expected to decay as power-law in proper time t
(Chang et al. 08, ML 14):

(t : comoving time since injection through the shock $\sim 3x$ distance to the shock)

$$\frac{d \langle \delta B^2 \rangle}{dt} = -2 \int \frac{d^3 k}{(2\pi)^3} \gamma_k |\delta B_k|^2 \quad \text{with (linear) damping rate } \gamma_k \simeq \frac{4}{\pi} \frac{k^3 c^3}{\omega_p^2}$$


$$\Rightarrow \epsilon_B(t) \propto k_{\max}^{n_B+3}(t) \propto t^{-(n_B+3)/3}$$



General picture



Blast wave geometry:



GRB orders of magnitude (comoving frame):

radius for afterglow: $R \sim 10^{17}$ cm

Lorentz factor: $\gamma_b \sim 100$


B field: $B_{\text{ISM}} \sim 1 \mu\text{G}$ ($\Leftrightarrow \epsilon_{B,\text{ISM}} \sim 10^{-9}$)

blast width: $R / (\gamma_b c) \sim 10^7 \omega_{\text{pi}}^{-1}$

gyration: $t_L \sim \epsilon_{B,-2}^{-1/2} (\gamma_e / \gamma_{\text{min}}) \omega_{\text{pi}}^{-1}$

cooling: $t_{\text{synch}} \sim 10^7 \dots (\gamma_e / \gamma_{\text{min}})^{-1} \omega_{\text{pi}}^{-1}$

PIC simulations: $\sim 10\,000 \omega_{\text{pi}}^{-1}$



acceleration zone $\sim 100 \omega_{\text{pi}}^{-1}$

\Rightarrow particles get "instantaneously" accelerated to a power-law then cool in microturbulence...

Synchrotron spectra in decaying microturbulence




Example:

slowly decaying turbulence, $\alpha_t = -0.8$,
 $t_{\text{obs}} = 100 \text{ sec}$, $n = 10^{-3} \text{ cm}^{-3}$,
 $E = 10^{53} \text{ ergs}$, with inverse Compton
 losses, $Y=3$

$$\delta B^2(t) \sim \delta B^2(t=0) [t/(100c/\omega_{\text{pi}})]^{\alpha_t}$$

vs

homogeneous turbulence, $\varepsilon_B = 10^{-2}$



→ decaying turbulence leaves a strong signature in the spectral flux $F_\nu(t_{\text{obs}})$:
modifies slopes and characteristic frequencies...

General trend: (for $-1 < \alpha_t < 0$)

→ flux F_ν at ν comes from electrons with γ_e : $v_p(\gamma_e) = \nu \dots$


→ $v_p \propto \gamma_e^2$ and $t_{\text{synch}} \propto \gamma_e^{-1}$ imply that **low frequencies are produced in regions of low magnetic field, high frequencies are produced in regions of strong magnetic field...**

Confrontation to observations



... *synchrotron emission of shock accelerated electrons in decaying micro-turbulence nicely reproduces the afterglows and >100MeV extended emissions of GRBs...* (ML et al. 13)


GRB 090902B



$E_{ej} \sim 1.6 \cdot 10^{54} \text{ erg}$
 $n \sim 0.012 \text{ cm}^{-3}$
 $\epsilon_e \sim 0.50, p \sim 2.3$
 $\alpha_t \sim -0.44 \pm 0.10$

$$\epsilon_B = 0.01 \left[t / \left(100 \omega_{pi}^{-1} \right) \right]^{\alpha_t}$$

GRB 090323




$E_{ej} \sim 5.4 \cdot 10^{54} \text{ erg}$
 $n \sim 8.4 \times 10^{35} \text{ r}^{-2} \text{ cm}^{-3}$
 $\epsilon_e \sim 0.29, p \sim 2.5$
 $\alpha_t \sim -0.54 \pm 0.09$

Confrontation to observations



... synchrotron emission of shock accelerated electrons in decaying micro-turbulence nicely reproduces the afterglows and >100MeV extended emissions of GRBs... (ML et al. 13)


GRB 090328



$E_{ej} \sim 0.73 \times 10^{54}$ erg
 $n \sim 1.5 \times 10^{35} \text{ r}^{-2} \text{ cm}^{-3}$
 $\epsilon_e \sim 0.18, p \sim 2.5$
 $\alpha_t \sim -0.48 \pm 0.11$

$$\epsilon_B = 0.01 \left[t / \left(100 \omega_{pi}^{-1} \right) \right]^{\alpha_t}$$

GRB 110731A



$E_{ej} > \sim 6 \times 10^{54}$ erg
 $n < \sim 0.1 \times 10^{35} \text{ r}^{-2} \text{ cm}^{-3}$
 $\epsilon_e < \sim 0.04, p \sim 2.1$
 $\alpha_t < \sim -0.35 \pm 0.20$



→ a simple solution, which reconciles data and theory, for the problem of the origin of magnetization in GRB blast waves:

→ **synchrotron radiation takes place in the partially decayed Weibel turbulence, which is self-generated at the (ultra-relativistic, unmagnetized) collisionless shock**

→ **4 GRBs seen in radio, optical, X-ray through >100MeV point to a consistent net decay power law of the magnetic field downstream of the shock:**

$$\epsilon_B \propto t^{\alpha_t} \quad \text{with: } -0.5 < \alpha_t < -0.4 \quad (t \text{ indicates } 3x \text{ distance}/c \text{ to shock})$$

→ values for ϵ_B **do not agree** with other estimates by Cenko et al. for 090902B, 090323, 090328, or with Ackermann et al. (Fermi Coll.) for 110731A:

difference: these works do not account for >100MeV emission...

... so 3 constraints for 4 parameters...

degeneracy implies that $\epsilon_B \sim 0.01$ in these works is a choice rather than a result!

→ is this even more general? What about earlier determinations of ϵ_B ?

Does the canonical value $\epsilon_B \sim 0.01$ hold at all?



Particle acceleration at relativistic shock waves is intimately connected to the self-generation of turbulence...

→ shock physics in mildly relativistic regime, high or low magnetization, less ideal conditions remain to be worked out...

→ a clearer view in the past decade thanks to PIC simulations (+theory!), especially at low magnetization

A microphysical solution for the origin of magnetization in GRB blast waves:

→ **synchrotron radiation takes place in the partially decayed Weibel turbulence, which is self-generated at the (ultra-relativistic, unmagnetized) collisionless shock**

→ a broad turbulence power spectrum at the shock leads to a power-law decay:

$$\delta B^2(t) \simeq \delta B^2(t=0) \left[t / \left(100 \omega_{\text{pi}}^{-1} \right) \right]^{\alpha_t}$$

→ **4 GRBs seen in radio, optical, X-ray through >100MeV point to a net decay power law of the magnetic field downstream of the shock: $-0.5 < \alpha_t < -0.4$**