OBTAINING EXACT INTERPOLATION MULTIVARIATE POLYNOMIAL BY APPROXIMATION*

Yong FENG · Jingzhong ZHANG · Xiaolin QIN · Xun YUAN

Received: 3 September 2008 ©2006 Springer Science + Business Media, Inc.

Abstract In some fields such as Mathematics Mechanization, automated reasoning and Trustworthy Computing etc., exact results are needed. Symbolic computations are used to obtain the exact results. Symbolic computations are of high complexity. In order to improve the situation, exactly interpolating methods are often proposed for the exact results and approximate interpolating methods for the approximate ones. In this paper, we study how to obtain exact interpolation polynomial with rational coefficients by approximate interpolating methods.

Key words Numerical approximate computation, symbolic-numerical computation, continued fraction, multivariate interpolation, Vandermonde determinant.

1 Introduction

Some fields such as automated reasoning and trustworthy computing etc., need exact results, and symbolic computations are used to obtain the exact results. Symbolic computations are principally exact and stable. However, they have the disadvantage of intermediate expression swell. Numerical computations have not the problem, however only give approximate results. In recent two decades, numerical methods are applied in the field of symbolic computations. In 1985, Kaltofen presented an algorithm for performing the absolute irreducible factorization, and suggested to perform his algorithm by floating-point numbers, then the factor obtained is an approximate one. After then, numerical methods have been studied to get approximate factors of a polynomial^[1-6]. In the meantime, numerical methods are applied to get approximate greatest

Yong FENG

School of Computer Science and Engineering, University of Electronic Science and Technology of China, Chengdu 610054, China; Laboratory for Automated Reasoning and Programming, Chengdu Institute of Computer Applications, Chinese Academy of Sciences, Chengdu 610041, China.Email:yongfeng@casit.ac.cn. Jingzhong ZHANG

Laboratory for Automated Reasoning and Programming, Chengdu Institute of Computer Applications, Chinese Academy of Sciences, Chengdu 610041, China.

Xiaolin QIN

Laboratory for Automated Reasoning and Programming, Chengdu Institute of Computer Applications, Chinese Academy of Sciences, Chengdu 610041, China.Email: qinxl811028@163.com.

Xun YUAN

Laboratory for Automated Reasoning and Programming, Chengdu Institute of Computer Applications, Chinese Academy of Sciences, Chengdu 610041, China.

^{*}This research is supported by China 973 Project NKBRPC-2004CB318003, the Knowledge Innovation Program of the Chinese Academy of Sciences KJCX2-YW-S02, and the National Natural Science Foundation of China(Grant NO. 10771205).

common divisors of approximate polynomials^[7-10], to compute functional decompositions^[11], to test primality^[12] and to find zeroes of a polynomial^[13]. In 2000, Corless et al. applied numerical methods in implicitization of parametric curves, surfaces and hypersurfaces^[14], and the resulting implicit equation is still an approximate one. In [15], Chèze et al. discussed how to obtain an exact absolute polynomial factorization from its approximate one, which only involves recovering an integral coefficient from its approximation.

Interpolation methods as an efficient numerical method have been proverbially used to compute resultants and determinants, etc. And approximate interpolation methods are still used to get the approximate ones^[16–18]. In order to obtain exact results, people usually use exact interpolation methods to meliorate intermediate expression swell problem arising from symbolic computations ^[19–23]. In fact, these are not approximate numerical computations but big number computations, which are also exact computations and only improve intermediate expression swell problem. Recently, Zhang et al. proposed an algorithm to recover the exact rational number from its approximate computations. In this paper, we discuss how the errors of support points affect that of coefficients of the interpolation multivariate polynomial. The algorithm can be carried out in parallel computers. Compared with exact interpolation method, the efficiency of the our algorithm is higher when the scale of problem is larger. This paper provide a way to obtain exact results by numerical computations.

The remainder of the paper is organized as follows. Section 2 gives a review of modified continued fraction method, by which an exact rational number can be obtained from its approximation, and gives a review of Kronecker product of two matrix. Section 3 first proposes estimation of the error to ensure the exact interpolation polynomial to be obtained, and then presents an algorithm to obtain an exact polynomial from its approximation for univariate and multivariate polynomial over rational number field, respectively. Section 4 gives some experimental results. The final section makes conclusions.

2 Preliminaries

In general, obtaining exact polynomial by approximate computations consists of two steps. First compute approximate polynomial within an error control by approximate numerical methods, and then recover the exact coefficients of the approximate polynomial by continued fraction method. In this section, we review the main results to be used in this paper.

A continued fraction representation of a real number x is one of the forms:

$$a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \cdots}}},$$
(1)

where a_0 is an integer and a_1, a_2, a_3, \cdots are positive integers. One can abbreviate the above continued fraction as $[a_0; a_1, a_2, \cdots]$. Truncating the above continued fraction representation of a number x early yields a rational number which is in a certain sense the best possible rational approximation. We call $[a_0; a_1, \cdots, a_n]$ the n-th convergent of $[a_0; a_1, a_2, \cdots]$. By continued fraction representation, the relation between a rational number and its approximation was disclosed as follows^[24].

Theorem 1 Let n_0/n_1 be a reduced rational number and r its approximation. Assume that n_0, n_1 are positive integers and $N \ge \max\{n_1, 2\}$. We have the continued fraction representation

 $r = [b_0, b_1, \dots, b_M]$ and $|r - n_0/n_1| < 1/(2N^2)$. Then, $n_0/n_1 = [b_0, b_1, \dots, b_L]$, where either M = L or M > L. In the case of M > L, for any positive integer $L < t \le M$, the denominator of rational number $g = [b_0, b_1, \dots, b_t]$ is greater than N.

Based on the theorem above, an algorithm for obtaining the exact number was designed as follows^[24]:

Algorithm 1 Input: a nonnegative floating-point number r and a positive number N; Output: a rational number b.

Step 1: Set i = 0, $h_{-1} = 1$, $h_{-2} = 0$, $k_{-1} = 0$, $k_{-2} = 1$, and convert r to a rational number and assign it to tem;

Step 2: Get integral part of tem and assign it to a, assign its remains to b;

Step 3: Compute $h_i = a * h_{i-1} + h_{i-2}$ and $k_i = a * k_{i-1} + k_{i-2}$. If $k_i > N$, then goto Step 6;

Step 4: Set i := i + 1;

Step 5: If $b \neq 0$, set $tem = \frac{1}{b}$ and go to Step 2;

Step 6: Compute h_{i-1}/k_{i-1} and assign it to b;

Step 7: return b.

It follows from theorem 1 that once an bound N on the denominator of a positive rational number is estimated, we can obtain the exact rational number as follows. Compute its approximation with the error less than $1/(2N^2)$, and then recover the rational number from its approximation by algorithm 1.

Now, let us give a brief review on the Kronecker product of two matrix. Denoted by $\mathbf{M}_{m,n}(\mathbb{F})$ the set of all m by n matrices over field \mathbb{F} , and abbreviate $\mathbf{M}_{n,n}(\mathbb{F})$ to $\mathbf{M}_n(\mathbb{F})$. The Kronecker product of $\mathbf{A} = [a_{i,j}] \in \mathbf{M}_{m,n}(\mathbb{F})$ and $\mathbf{B} \in \mathbf{M}_{p,q}(\mathbb{F})$ is denoted by $\mathbf{A} \otimes \mathbf{B}$ and is defined to be the block matrix

$$\mathbf{A} \otimes \mathbf{B} = \begin{pmatrix} a_{11}\mathbf{B} & \cdots & a_{1n}\mathbf{B} \\ \vdots & \ddots & \vdots \\ a_{m1}\mathbf{B} & \cdots & a_{mn}\mathbf{B} \end{pmatrix} \in \mathbf{M}_{mp,nq}(\mathbb{F}).$$
(2)

The Kronecker products has many properties^[25]. Here we mention two properties, one of which is as follows:

Theorem 2 (mixed-product) Let $\mathbf{A} \in \mathbf{M}_{m,n}(\mathbb{F})$, $\mathbf{B} \in \mathbf{M}_{p,q}(\mathbb{F})$, $\mathbf{C} \in \mathbf{M}_{n,k}(\mathbb{F})$, and $\mathbf{D} \in \mathbf{M}_{q,r}(\mathbb{F})$. Then $(\mathbf{A} \otimes \mathbf{B})(\mathbf{C} \otimes \mathbf{D}) = \mathbf{A}\mathbf{C} \otimes \mathbf{B}\mathbf{D}$.

Another property is concerned with the eigenvalues of the Kronecker product of two square complex matrices.

Theorem 3 Let $\mathbf{A} \in \mathbf{M}_n$, and $\mathbf{B} \in \mathbf{M}_m$. Denote by $\sigma(\mathbf{A})$ the all eigenvalues of matrix \mathbf{A} . If $\lambda \in \sigma(\mathbf{A})$ and $\mathbf{x} \in \mathbb{C}^n$ is a corresponding eigenvector of \mathbf{A} , and if $\mu \in \sigma(\mathbf{B})$ and $\mathbf{y} \in \mathbb{C}^m$ is a corresponding eigenvector of \mathbf{A} , and $\mathbf{x} \otimes \mathbf{y} \in \mathbb{C}^{nm}$ is a corresponding eigenvector of $\mathbf{A} \otimes \mathbf{B}$ and $\mathbf{x} \otimes \mathbf{y} \in \mathbb{C}^{nm}$ is a corresponding eigenvector of $\mathbf{A} \otimes \mathbf{B}$. Every eigenvalues of $\mathbf{A} \otimes \mathbf{B}$ arise as such a product of eigenvalues of \mathbf{A} and \mathbf{B} .

From theorem 3 and the fact that the determinant of any matrix is equal to the product of its eigenvalues, it follows that $[^{26}]$.

Theorem 4 Let $\mathbf{A} \in \mathbf{M}_n$ and $\mathbf{B} \in \mathbf{M}_m$ be given. Then $\det(\mathbf{A} \otimes \mathbf{B}) = (\det \mathbf{A})^m (\det \mathbf{B})^n$. Thus, $\mathbf{A} \otimes \mathbf{B}$ is nonsingular if and only if both \mathbf{A} and \mathbf{B} are nonsingular.

In this paper, we need to solve an equation whose coefficient matrix is the Kronecker product. The following theorem shows us how to solve the problem^[25].

Theorem 5 Let \mathbb{F} denote a field. Matrices $\mathbf{A} \in \mathbf{M}_{m,n}(\mathbb{F})$, $\mathbf{B} \in \mathbf{M}_{q,p}(\mathbb{F})$, and $\mathbf{C} \in \mathbf{M}_{m,q}(\mathbb{F})$ are given and assume $\mathbf{X} \in \mathbf{M}_{n,p}(\mathbb{F})$ to be unknown. With matrix \mathbf{X} , associate the vector $\operatorname{vec}(\mathbf{X}) \in \mathbb{F}$ defined by

$$\operatorname{vec}(\mathbf{X}) = [a_{11}, \cdots a_{n1}, a_{12}, \cdots, a_{n2}, \cdots, a_{1p}, \cdots, a_{np}]^T.$$

Then, the following equation:

$$(\mathbf{B} \otimes \mathbf{A}) \operatorname{vec}(\mathbf{X}) = \operatorname{vec}(\mathbf{C}) \tag{3}$$

is equivalent to matrix equation:

$$\mathbf{A}\mathbf{X}\mathbf{B}^T = \mathbf{C}.\tag{4}$$

Obviously, equation (4) is equivalent to the system of equations

$$\begin{cases} \mathbf{A}\mathbf{Y} = \mathbf{C} \\ \mathbf{B}\mathbf{X}^T = \mathbf{Y}^T. \end{cases}$$
(5)

3 Algorithms

3.1 Univariate Interpolation Polynomial

Let $f(x) = \sum_{i=0}^{n} a_i x^i$ be an univariate rational polynomial with degree *n*. Its approximate support points are $\{(x_i, f_i)\}$ for $i = 0, 1, \dots, n$, which means that

$$f(x_i) = f_i$$
, for $i = 0, 1, \cdots, n$

where x_i are called interpolation nodes and values f_i interpolation datum. We can construct polynomial f(x) from its support points by interpolation method. Polynomial interpolation is a classical numerical method. It is studied very well for univariate polynomials. In general, interpolation problem is essentially to solve the following equation:

$$\begin{pmatrix} 1 & x_0 & x_0^2 & \cdots & x_0^n \\ 1 & x_1 & x_1^2 & \cdots & x_1^n \\ \cdots & \cdots & \cdots & \cdots \\ 1 & x_n & x_n^2 & \cdots & x_n^n \end{pmatrix} \begin{pmatrix} a_0 \\ a_1 \\ a_2 \\ \vdots \\ a_n \end{pmatrix} = \mathbf{U}_{n+1}(x_0, \cdots, x_n) \mathbf{a} = \begin{pmatrix} f_0 \\ f_1 \\ f_2 \\ \vdots \\ f_n \end{pmatrix},$$
(6)

where

$$\mathbf{U}_{n+1}(x_0,\cdots,x_n) = \begin{pmatrix} 1 & x_0 & x_0^2 & \cdots & x_0^n \\ 1 & x_1 & x_1^2 & \cdots & x_1^n \\ 1 & x_2 & x_2^2 & \cdots & x_2^n \\ 1 & x_3 & x_3^2 & \cdots & x_n^n \\ \cdots & \cdots & \cdots & \cdots \\ 1 & x_n & x_n^2 & \cdots & x_n^n \end{pmatrix}, \quad \mathbf{a} = \begin{pmatrix} a_0 \\ a_1 \\ a_2 \\ a_3 \\ \cdots \\ a_n \end{pmatrix},$$
(7)

Matrix $\mathbf{U}_{n+1}(x_0, \dots, x_n)$ is called Vandermonde matrix and its determinant $\mathbf{V}_{n+1}(x_0, \dots, x_n) = \det \mathbf{U}_{n+1}(x_0, \dots, x_n) = \prod_{0 \le i < j \le n} (x_j - x_i)$ is called Vandermonde determinant. Let $\mathbf{D}_n(i, j)$

denote the determinant of the submatrix of matrix $\mathbf{U}_{n+1}(x_0, \cdots, x_n)$ resulting from deletion of row i + 1 and column j + 1. $\mathbf{D}_n(i, j)$ is often called generalized Vandermonde determinant. How do we compute it? The follow theorem shows us do it^[27].

Theorem 6 Let $\delta_k(x_1, x_2, \dots, x_n) = \sum_{1 \le i_1 < i_2 < \dots < i_k \le n} x_{i_1} x_{i_2} \cdots x_{i_k}$ be elementary symmetric polynomial with degree k. Then

 $\mathbf{D}_{n}(i,j) = \delta_{n-j}(x_{0}, \cdots, x_{i-1}, x_{i+1}, \cdots, x_{n}) \mathbf{V}_{n}(x_{0}, \cdots, x_{i-1}, x_{i+1}, \cdots, x_{n}).$

From theorem 6, it follows that the solution of equation (6)

$$a_j = \frac{\sum_{i=0}^n (-1)^{i+j} \mathbf{D}_n(i,j) f_i}{\mathbf{V}_{n+1}(x_0, x_1, \cdots, x_n)}.$$
(8)

Theoretically, we can compute the interpolation polynomial after choosing n + 1 distinct points x_0, x_1, \dots, x_n and then obtaining their corresponding exact interpolation values f_0, f_1, \dots, f_n . However, in practice, we often get the approximate function values of f(x), denoted by $\tilde{f}_0, \tilde{f}_1, \dots, \tilde{f}_n$. So an approximate interpolation polynomial $\tilde{f}(x) = \sum_{j=0}^n \tilde{a}_j x^j$ is only produced. The following theorem gives an error estimation of coefficients of the approximate interpolation polynomial.

Theorem 7 Let $\varepsilon = \max\{|f_j - \tilde{f}_j|, j = 0, \dots, n\}, \lambda_x = \min\{|x_j - x_i|, i \neq j\}$ and $M = \max\{1, \max\{|x_j|, j = 0, \dots, n\}\}.$

$$\max\{|a_j - \tilde{a}_j|, j = 0, \cdots, n\} < \frac{\varepsilon}{\lambda_x^n} (n+1) \binom{n}{\lfloor n/2 \rfloor} M^n, \tag{9}$$

where $\lfloor x \rfloor$ stands for the greatest integer which is less than or equal to x, and $\binom{n}{\lfloor n/2 \rfloor}$ the number of $\lfloor n/2 \rfloor$ -combinations of an n-element set.

Before giving the proof of theorem 7, we introduce two lemmas:

Lemma 1 Let $\delta_k(x_1, x_2, \dots, x_n) = \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq n} x_{i_1} x_{i_2} \cdots x_{i_k}$ be elementary symmetric polynomial with degree k. Then

$$\sum_{j=0}^{n} \delta_k(\overbrace{x_0, x_1, \cdots, x_{j-1}, x_{j+1}, \cdots, x_n}^{n \text{ points}}) = (n+1-k)\delta_k(\overbrace{x_0, x_1, \cdots, x_j, \cdots, x_n}^{n+1 \text{ points}}).$$
(10)

Proof: Every term $x_{i_0}x_{i_1}\cdots x_{i_{k-1}}$ of $\delta_k(x_0, x_1, \cdots, x_j, \cdots, x_n)$ does not appear in $\delta_k(x_0, x_1, \cdots, x_{i-1}, x_{i+1}, \cdots, x_n)$ with $i = i_l$ for $l = 0, 1, \cdots, k-1$, and appears in the other cases. Therefore, term $x_{i_0}x_{i_1}\cdots x_{i_{k-1}}$ appears (n+1-k) times on the left hand side of equation (10). So, the lemma is finished.

Another lemma is concerned with binomial coefficients as follows^[28]:

Lemma 2 For n a positive integer, the largest of the binomial coefficients

$$\binom{n}{0}, \binom{n}{1}, \binom{n}{2}, \cdots, \binom{n}{n}$$

is $\binom{n}{\lfloor n/2 \rfloor}$, where $\lfloor x \rfloor$ stands for the greatest integer which is less than or equal to x.

Now we turn to give the proof of theorem 7. **Proof:** From equation (8), we have

$$\begin{aligned} |a_{j} - \tilde{a}_{j}| &= \left| \frac{\sum_{i=0}^{n} (-1)^{i+j} \mathbf{D}_{n}(i,j)(f_{i} - \tilde{f}_{i})}{\mathbf{V}_{n+1}(x_{0}, x_{1}, \cdots, x_{n})} \right| \\ &\leq \sum_{i=0}^{n} \frac{|\mathbf{D}_{n}(i,j)||(f_{i} - \tilde{f}_{i})|}{|\mathbf{V}_{n+1}(x_{0}, x_{1}, \cdots, x_{n})|} &\leq \varepsilon \sum_{i=0}^{n} \frac{|\mathbf{D}_{n}(i,j)|}{|\mathbf{V}_{n+1}(x_{0}, x_{1}, \cdots, x_{n})|} \\ &\leq \varepsilon \sum_{i=0}^{n} \frac{|\delta_{n-j}(x_{0}, \cdots, x_{i-1}, x_{i+1}, \cdots, x_{n})||\mathbf{V}_{n}(x_{0}, \cdots, x_{i-1}, x_{i+1}, \cdots, x_{n})|}{|\mathbf{V}_{n+1}(x_{0}, x_{1}, \cdots, x_{n})|} \\ &\leq \frac{\varepsilon}{\lambda_{x}^{n}} \sum_{i=0}^{n} |\delta_{n-j}(\widetilde{x_{0}, \cdots, x_{i-1}, x_{i+1}, \cdots, x_{n})|, \end{aligned}$$

noticing equation (10) yields

$$\begin{aligned} |a_j - \tilde{a}_j| &\leq \frac{\varepsilon}{\lambda_x^n} (n+1-n+j) \delta_{n-j} (\overbrace{|x_0|, \cdots, |x_{i-1}|, |x_i|, |x_{i+1}|, \cdots, |x_n|}^{n+1}) \\ &\leq \frac{\varepsilon}{\lambda_x^n} (j+1) \binom{n+1}{n-j} M^{n-j} \\ &= \frac{\varepsilon}{\lambda_x^n} (j+1) \frac{(n+1)!}{(n-j)!(j+1)!} M^{n-j} \\ &= \frac{\varepsilon}{\lambda_x^n} (n+1) \frac{n!}{(n-j)!j!} M^{n-j}. \end{aligned}$$

From lemma 2, it follows that

$$\begin{aligned} |a_j - \tilde{a}_j| &\leq \frac{\varepsilon}{\lambda_x^n} (n+1) \frac{n!}{(n - \lfloor n/2 \rfloor)! \lfloor n/2 \rfloor!} M^{n-j} \\ &\leq \frac{\varepsilon}{\lambda_x^n} (n+1) \frac{n!}{(n - \lfloor n/2 \rfloor)! \lfloor n/2 \rfloor!} M^n \\ &= \frac{\varepsilon}{\lambda_x^n} (n+1) \binom{n}{\lfloor n/2 \rfloor} M^n. \end{aligned}$$

The proof is completed.

And now, let us discuss how to recover the exact polynomial from its approximate interpolation polynomial.

Let n denote the degree of f(x) and N an upper bound of absolute values of denominators of its coefficients. Note that f(x) is unknown, so we should estimate an upper bound N and its degree n in advance. We may calculate them by the form of the given expression of polynomial f(x), and should obtain as less bound as possible. The less bound we obtain, the less the amount of computation is for obtaining approximate interpolation polynomial. Once an upper bound N and n are gotten, we choose n + 1 interpolate nodes x_0, x_1, \dots, x_n and calculate

$$\varepsilon = \frac{\lambda_x^n}{2(n+1)\binom{n}{\lfloor n/2 \rfloor} M^n N^2}.$$
(11)

Then, compute the values $\tilde{f}_i \approx f(x_i)$ for $i = 0, 1, \dots, n$ with an error less than ε . By interpolation method, we compute the approximate interpolation polynomial $\tilde{f}(x)$ with coefficient

error less than $1/(2N^2)$. Finally, use algorithm 1 to obtain the exact polynomial f(x) from its approximate polynomial $\tilde{f}(x)$. In summary, the algorithm is as follows:

Algorithm 2 Input: an expression f(x) which result is a polynomial; Output: a polynomial g(x) such that g(x) = f(x).

Step 1: By the structure of the expression, estimate an upper bound on the degree of f(x) and an upper bound on absolute values of denominators of its coefficients, denoted by n and N respectively;

Step 2: Choose distinct points x_0, x_1, \dots, x_n ; Compute $\lambda_x = \min\{|x_j - x_i|, i \neq j\}$ and $M = \max\{1, \max\{|x_i|, i = 0, \dots, n\}\}.$

Step 3: Compute ε in formula (11);

Step 4: By numerical method, approximately compute the values of f(x) at the points x_0, x_1, \dots, x_n with an error less than ε and denote the corresponding values by $\tilde{f}_i \approx f(x_i)$, for $i = 0, 1, \dots, n$;

Step 5: By interpolate method, obtain approximate interpolation polynomial $\tilde{f}(x)$;

Step 6: Call algorithm 1 to recover the exact coefficients from the coefficients of $\tilde{f}(x)$ one by one. Denote the exact polynomial by g(x);

Step 7: return g(x).

The correctness of the algorithm above is shown as follows: From step 1 to step 4, we obtain the interpolate datum \tilde{f}_i with an error less than ε . By theorem 7, The errors of the coefficients of the approximate polynomial are all less than $1/(2N^2)$, which ensure the exact coefficients of the polynomial to obtain from its approximation by algorithm 1.

3.2 Multivariate Interpolation Polynomial

For simplicity, we first consider the bivariate interpolate problem, and then generalize the results to the case of multivariate interpolation. Let $f(x, y) = \sum_{i,j} a_{ij} x^i y^j$ be a polynomial with rational coefficients, and n, m be the bounds on the degree of f(x, y) in x, y respectively. We choose the interpolation nodes (x_i, y_j) $(i = 0, \dots, n; j = 0, \dots, m)$, and obtain the values of f(x, y), denoted by $f_{ij} \in \mathbb{R}$ $(i = 0, \dots, n; j = 0, \dots, m)$. The set of monomials is ordered as follows:

$$\{x_i y_j | i = 0, \cdots, n; j = 0, \cdots, m\}$$

= $\{1, y, \cdots, y^m, x, xy, \cdots, xy^m, \cdots, x^n, x^n y, \cdots, x^n y^m\},$

and the interpolation nodes in the corresponding order is as follows:

$$\{(x_i, y_j)|i = 0, \cdots, n; j = 0, \cdots, m\} = \{(x_0, y_0), (x_0, y_1), \cdots, (x_0, y_m), (x_1, y_0), (x_1, y_1), \cdots, (x_1, y_m), \cdots, (x_n, y_0), \cdots, (x_n, y_m)\}.$$

Note that the above ordering is reverse to conventional lexicographic order. Of course it is a lexicographic order under assumption of ordering $0 > 1 > 2 > \cdots$ and y > x. They do not

coincide with our convention, so the above ordering is called reverse lexicographic order in this paper. Let

$$\mathbf{A} = \begin{pmatrix} a_{00} & \cdots & a_{0m} \\ \cdots & \cdots & \cdots \\ a_{n0} & \cdots & a_{nm} \end{pmatrix}, \quad \mathbf{F} = \begin{pmatrix} f_{00} & \cdots & f_{0m} \\ \cdots & \cdots & \cdots \\ f_{n0} & \cdots & f_{nm} \end{pmatrix},$$

the bivariate interpolate problem can be expressed as to solve the following equation:

$$\mathbf{M}\mathrm{vec}(\mathbf{A}^T) = \mathrm{vec}(\mathbf{F}^T),\tag{12}$$

where the coefficients matrix $\mathbf{M} = \mathbf{U}_x \otimes \mathbf{U}_y$. \mathbf{U}_x and \mathbf{U}_y are Vandermonde matrix, i.e.,

$$\mathbf{U}_{x} = \begin{pmatrix} 1 & x_{0} & x_{0}^{2} & \cdots & x_{0}^{n} \\ 1 & x_{1} & x_{1}^{2} & \cdots & x_{1}^{n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_{n} & x_{n}^{2} & \cdots & x_{n}^{n} \end{pmatrix}, \quad \mathbf{U}_{y} = \begin{pmatrix} 1 & y_{0} & y_{0}^{2} & \cdots & y_{0}^{m} \\ 1 & y_{1} & y_{1}^{2} & \cdots & y_{1}^{m} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & y_{m} & y_{m}^{2} & \cdots & y_{m}^{m} \end{pmatrix}.$$

As said in section 2, in practice we often only get the approximate interpolation values $\tilde{f}_{ij} \approx f(x_i, y_j)$. Of course, the interpolate polynomial obtained is an approximation. The following theorem gives an estimation error:

Theorem 8 Set $f_{ij} = f(x_i, y_j)$ and denote by \tilde{f}_{ij} the approximation of f_{ij} . $\tilde{f}(x, y) = \sum_{ij} \tilde{a}_{ij} x^i y^j$ is the interpolate polynomial from interpolate datum \tilde{f}_{ij} . Let $\varepsilon = \max\{|f_{ij} - \tilde{f}_{ij}| : i = 0, \dots, n; j = 0, \dots, m\}$, $\lambda_x = \min\{|x_j - x_i| : i \neq j\}$, $\lambda_y = \min\{|y_j - y_i| : i \neq j\}$ and $M_x = \max\{1, \max\{|x_j| : j = 0, \dots, n\}\}$, $M_y = \max\{1, \max\{|y_j| : j = 0, \dots, m\}\}$. Then

$$\max_{i,j} |a_{ij} - \tilde{a}_{ij}| \le \frac{(m+1)(n+1)M_x^n M_y^m}{\lambda_x^n \lambda_y^m} \binom{n}{\lfloor n/2 \rfloor} \binom{m}{\lfloor m/2 \rfloor} \varepsilon.$$

Proof: Let $\tilde{\mathbf{A}} = (\tilde{a}_{ij})$ and $\tilde{\mathbf{F}} = (\tilde{f}_{ij})$. From equation (12), it holds that

$$\mathbf{M}\mathrm{vec}((\tilde{\mathbf{A}} - \mathbf{A})^T) = \mathrm{vec}((\tilde{\mathbf{F}} - \mathbf{F})^T),$$

by theorem 5, the above equation is equivalent to the following equation:

$$\mathbf{U}_{y}(\mathbf{\tilde{A}} - \mathbf{A})^{\mathbf{T}} \mathbf{U}_{\mathbf{x}}^{T} = (\mathbf{F} - \mathbf{\tilde{F}})^{T}.$$

Thus, it is equivalent to

$$\mathbf{U}_{\boldsymbol{y}}\mathbf{Z} = (\mathbf{F} - \tilde{\mathbf{F}})^T \tag{13a}$$

$$\mathbf{U}_{\mathbf{x}}(\tilde{\mathbf{A}} - \mathbf{A}) = \mathbf{Z}^T \tag{13b}$$

Where $\mathbf{Z} = (z_{ij})$. Matrix equation (13a) is equivalent to

$$\mathbf{U}_{y}\mathbf{Z}_{.i} = (\mathbf{F}_{i.} - \tilde{\mathbf{F}}_{i.})^{T}, \quad i = 1, \cdots m + 1$$
(14)

where \mathbf{Z}_{i} stands for the *i*-th column of \mathbf{Z} and \mathbf{F}_{i} the *i*-th row of matrix \mathbf{F} . By theorem 7, for every *i*, it holds that

$$\max_{j=0}^{m} |z_{ji}| < \frac{\varepsilon}{\lambda_y^m} (m+1) \binom{m}{\lfloor m/2 \rfloor} M_y^m.$$

Hence, we conclude that

$$\max_{i,j} |z_{ji}| < \frac{\varepsilon}{\lambda_y^m} (m+1) \binom{m}{\lfloor m/2 \rfloor} M_y^m.$$

Let

$$\delta = \frac{\varepsilon}{\lambda_y^m} (m+1) \binom{m}{\lfloor m/2 \rfloor} M_y^m,$$

argue equation (13b) in the same way as do above, we deduce that

$$\max_{i,j} |a_{ij} - \tilde{a}_{ij}| \le \frac{(m+1)(n+1)M_x^n M_y^m}{\lambda_x^n \lambda_y^m} \binom{n}{\lfloor n/2 \rfloor} \binom{m}{\lfloor m/2 \rfloor} \varepsilon_{ij}$$

The proof is finished.

Now, After having a relation between error of approximate coefficients and that of interpolate datum, we establish an algorithm to obtain an exact polynomial from its approximation as follows:

Algorithm 3 Input: an expression f(x, y) which result is a polynomial; Output: a polynomial g(x, y) such that g(x, y) = f(x, y).

Step 1: By the structure of the expression, estimate an upper bounds on the degrees of f(x, y) in x and y, denoted by n and m respectively. Estimate an upper bound on absolute values of denominators of its coefficients, denoted by N;

Step 2: Choose interpolate nodes $(x_i, y_j), (i = 0, 1, \dots, n; j = 0, 1, \dots, m)$. Compute $\lambda_x = \min_{j \neq i} \{ |x_j - x_i| \}, \ \lambda_y = \min_{j \neq i} \{ |y_j - y_i| \}$ and $M_x = \max\{1, \max_{i=0}^n |x_i| \}, M_y = \max\{1, \max_{i=0}^m |y_i| \}$.

Step 3: Compute ε :

$$\varepsilon = \frac{\lambda_x^n \lambda_y^m}{2(n+1)(m+1)\binom{n}{\lfloor n/2 \rfloor}\binom{m}{\lfloor m/2 \rfloor}M_x^n M_y^m N^2};$$

Step 4: By numerical method, compute the approximate values of f(x, y) at the points $(x_i, y_j)(i = 0, 1, \dots, n; j = 0, 1, \dots, m)$ with an error less than ε , and denote the corresponding values by $\tilde{f}_{ij} \approx f(x_i, y_j)$, $(i = 0, 1, \dots, n; j = 0, 1, \dots, m)$;

Step 5: By interpolate method, obtain approximate interpolation polynomial $\tilde{f}(x,y)$;

Step 6: Call algorithm 1 to recover the exact coefficients from the coefficients of $\tilde{f}(x,y)$ one by one. Denote the exact polynomial by g(x,y);

Step 7: return g(x, y).

As for generalization of the above results to the case of multivariate polynomials r > 2, we assert that the situation is completely analogous to the bivariate case.

Let $f(X_1, \dots, X_r) = \sum a_{i_1 \dots i_r} X_1^{i_1} \dots X_r^{i_r}$ be a polynomial in X_1, \dots, X_r , and let n_i be a bound on the degree of $f(X_1, \dots, X_r)$ in X_i $(i = 1, \dots, r)$. Denotes by N an upper bound on absolute values of denominators of coefficients of polynomial $f(X_1, \dots, X_r)$. Choose interpolation nodes $(x_{1i_1}, x_{2i_2} \dots, x_{ri_r})$ $(i_k = 0, 1, \dots, n_k)$, where $k = 1, \dots, r$. Set $f_{i_1 i_2 \dots i_r} =$ $f(x_{1i_1}, x_{2i_2}, \dots, x_{ri_r})$ and denote by $\tilde{f}_{i_1 i_2 \dots i_r}$ the approximation of $f_{i_1 i_2 \dots i_r}$, by which we compute an approximate interpolation polynomial $\tilde{f}(X_1, X_2, \dots, X_r) = \sum \tilde{a}_{i_1 i_2 \dots i_r} X_1^{i_1} X_2^{i_2} \dots X_r^{i_r}$. Thus we have the following theorem: **Theorem 9** Let $\varepsilon = \max_{i_1, i_2, \dots, i_r} |f_{i_1 i_2 \dots i_r} - \tilde{f}_{i_1 i_2 \dots i_r}|, \ \lambda_k = \min_{i \neq j} |x_{kj} - x_{ki}|, \ and \ M_k = \max\{1, \max_{j=0}^{n_k} |x_{kj}|\}.$ Then

$$\max_{i_1, i_2, \cdots, i_r} |a_{1i_1, 2i_2, \cdots, ri_r} - \tilde{a}_{1i_1, 2i_2, \cdots, ri_r}| \le \varepsilon \prod_{k=1}^r \frac{(n_k + 1)M_k^{n_k} \binom{n_k}{\lfloor n_k/2 \rfloor}}{(\lambda_k^{n_k})}.$$
(15)

Proof: By the Reverse Lexicographic order, the interpolation problem comes down to solving the following equation:

$$(\mathbf{U}_{X_1}\otimes\mathbf{U}_{X_2}\cdots\otimes\mathbf{U}_{X_r})\mathbf{a}=\mathbf{F},$$

where **a** is a column vector consisting of elements $a_{i_1i_2\cdots i_r}$ in the reverse lexicographic order and **F** is a column vector consisting of elements $f_{i_1i_2\cdots i_r}$ in the reverse lexicographic order. Thus we have the following equation:

$$(\mathbf{U}_{X_1} \otimes \mathbf{U}_{X_2} \cdots \otimes \mathbf{U}_{X_r})(\mathbf{a} - \tilde{\mathbf{a}}) = (\mathbf{F} - \tilde{\mathbf{F}}),$$
(16)

where $\tilde{\mathbf{a}}$ and $\tilde{\mathbf{F}}$ are column vectors respectively consisting of elements $\tilde{a}_{i_1i_2\cdots i_r}$ and $f_{i_1i_2\cdots i_r}$ in the reverse lexicographic order. Equation (16) is equivalent to the following equation:

$$((\mathbf{U}_{X_1}\otimes\mathbf{U}_{X_2}\cdots\otimes\mathbf{U}_{X_{r-1}})\otimes\mathbf{U}_{X_r})(\mathbf{a}-\mathbf{\tilde{a}})=(\mathbf{F}-\mathbf{\tilde{F}}).$$

Hence, by recursion and as did in the proof of theorem 8, we can deduce that inequality (15) holds. The proof is finished.

Based on the above theorem, algorithm 3 is generalized to the case of multivariate interpolation as follows:

Algorithm 4 Input: an expression $f(X_1, \dots, X_r)$ which result is a polynomial; Output: a polynomial $g(X_1, \dots, X_r)$ such that $g(X_1, \dots, X_r) = f(X_1, \dots, X_r)$.

Step 1: By the structure of the expression, estimate bounds on the degrees of $f(X_1, \dots, X_r)$ in X_i , denoted by $n_i (i = 1, \dots, r)$. Estimate an upper bound on absolute values of denominators of its coefficients, denoted by N;

Step 2: Choose interpolation nodes $(x_{1i_1}, \dots, x_{ri_r}), (i_k = 0, 1, \dots, n_k; k = 0, 1, \dots, r).$ Compute $\lambda_k = \min_{j \neq i} \{ |x_{kj} - x_{ki}| \}$ and $M_k = \max\{1, \max_{i=0}^n |x_{ki}| \}$ for $k = 1, \dots, r$.

Step 3: Compute ε :

$$\varepsilon = \frac{\prod_{k=1}^{r} (\lambda_k^{n_k})}{2N^2 \prod_{k=1}^{r} (n_k + 1) M_k^{n_k} \binom{n_k}{\lfloor n_k/2 \rfloor}};$$
(17)

Step 4: By numerical method, compute the approximate values of $f(X_1, \dots, X_r)$ at the points $(x_{1i_1}, \dots, x_{ri_r})(i_k = 0, 1, \dots, n_k; k = 0, 1, \dots, r)$ with an error less than ε and denote the corresponding values by $\tilde{f}_{i_1\dots i_r} \approx f(x_{1i_1}, \dots, x_{ri_r}), (i_k = 0, 1, \dots, n_k; k = 0, 1, \dots, r)$;

Step 5: By interpolate method, obtain approximate interpolation polynomial $f(X_1, \dots, X_r)$;

Step 6: Call algorithm 1 to recover the exact coefficients from the coefficients of $\tilde{f}(X_1, \dots, X_r)$ one by one. Denote the exact polynomial by $g(X_1, \dots, X_r)$;

Step 7: return $g(X_1, \cdots, X_r)$.

The correctness of the above algorithm is ensured by theorem 8 and theorem 1. The proof can be given similarly to that in algorithm 3.

4 Experimental Results

Our algorithms are implemented in *Maple*. The following examples run in the platform of Maple11 and PIV3.0G,512M RAW. The first three examples illuminate how to obtain exact interpolation polynomials. Example 4 tests our algorithm for more variables and higher degree interpolation multivariate polynomials.

Example 1 Let f(x) be an unknown univariate rational polynomial. Assume that we know in advance the degree of f(x): n = 8 and an upper bound of absolute values of denominators of its coefficients N = 181. According to theorem7, computing exact interpolation polynomial f(x)as follows: Choose floating-point interpolation nodes [0.5001, 1.5003, 3.1201, 2.314, 4.02, 5.23, 6, 6.8,7.2001; Calculate $\lambda_x = 0.4001$, M=7.2001, Compute $\varepsilon = 0.2202 \times 10^{-17}$. Compute the approximate interpolate datum $\tilde{f}_i \approx f(x_i)$ such that $|f(x_i) - \tilde{f}_i| < \varepsilon$, for $i = 0, \dots, 8$, and use interpolation method to obtain approximate interpolation polynomial $\tilde{f}(x)$:

 $\begin{array}{l} .333333333333333333327598x^8 - .0277777777777775996307x^7 + .6666666666666666664376920x^6 \\ - .07222222222064098893x^5 + 1.958333333326967612x^4 + 1.750000000015123646x^3 \\ - .241666666666870050587x^2 + .17500000000137107868x + 1.49999999999996690437. \end{array}$

Then we recover the exact coefficients of interpolation polynomial by algorithm 1:

$$f(x) = \frac{1}{3}x^8 - \frac{1}{36}x^7 + \frac{2}{3}x^6 - \frac{13}{180}x^5 + \frac{47}{24}x^4 + \frac{7}{4}x^3 - \frac{29}{120}x^2 + \frac{7}{40}x + \frac{3}{2}.$$

Example 2 Let f(x, y) be an unknown bivariate rational polynomial. Assume that we know in advance the degrees of f(x, y) in x and y: n = 3, m = 3, and an upper bound of absolute values of denominators of its coefficients N = 13. According to theorem 8, computing exact interpolation polynomial f(x, y) as follows: Choose floating-point interpolation nodes X = [0.1, 0.5, 1.2, 1.3], Y = [0.2, 0.8, 2.1, 2.6], and calculate $\lambda_x = 0.1, \lambda_y = 0.5, M_x = 1.3, M_y = 2.6$; Compute $\varepsilon = 0.6659 \times 10^{-10}$. Compute the approximate interpolate datum \tilde{f}_{ij} such that $|f(x_i, y_j) - \tilde{f}_{ij}| < \varepsilon$, for $i = 0, \dots, 3, j = 0, \dots, 3$, and use interpolation method to obtain approximate interpolation polynomial $\tilde{f}(x, y)$ as follows:

 $-.2500000301x^3y^2 - .2500000111x^2y^3 + .08333326696x^2y - .08333336318xy^2 - .1666666674y^3 - .2499999755x^2 + .2500000031y^2 - .5000000114x - .500000029y.$

Then we recover the exact coefficients of interpolation polynomial by algorithm 1:

$$f(x,y) = -\frac{1}{4}x^3y^2 - \frac{1}{4}x^2y^3 + \frac{1}{12}x^2y - \frac{1}{12}xy^2 - \frac{1}{6}y^3 - \frac{1}{4}x^2 + \frac{1}{4}y^2 - \frac{1}{2}x - \frac{1}{2}y$$

Example 3 Let f(x, y, z) be an unknown tri-variate rational polynomial. Assume that we know in advance the degrees of f(x, y, z) in x, y, z: n = 3, m = 2, l = 3, and an upper bound of absolute values of denominators of its coefficients N = 231. According to theorem 9, computing exact interpolation polynomial f(x, y, z) as follows: Choose floating-point interpolation nodes X = [0.5, 1.2, 1.8, 2.7], Y = [0.3, 1.5, 2.2], Z = [0.6, 1.8, 2, 2.8], and calculate $\lambda_x = 0.6, \lambda_y = 0.7, \lambda_z = 0.4, M_x = 2.7, M_y = 2.2, M_z = 2.8$; Compute $\varepsilon = 0.735 \times 10^{-10}$. Compute the approximate interpolate datum \tilde{f}_{ijk} such that $|f(x_i, y_j, z_k) - \tilde{f}_{ijk}| < \varepsilon$, for $i = 0, \dots, 3, j = 0, \dots, 2, k = 0, \dots, 3$, and use interpolation method to obtain approximate interpolation polynomial $\tilde{f}(x, y, z)$ as follows:

$$\begin{array}{l} -.08333331999x^3y^2 + .1666668454x^2y^2z - .004629485562xy^2z^2 - .02777770742xyz^3 \\ -.05555579922xy^2z - .3333337487xyz^2 + .08333353340x^2y - .1666659870xyz \\ +.01388866836yz + .08333327541z^2 + .5000001136x + .1666667565y. \end{array}$$

Then we recover the exact coefficients of interpolation polynomial by algorithm 1:

$$f(x, y, z) = -\frac{1}{12}x^3y^2 + \frac{1}{6}x^2y^2z - \frac{1}{216}xy^2z^2 - \frac{1}{36}xyz^3 - \frac{1}{18}xy^2z - \frac{1}{3}xyz^3 - \frac{1}{18}xy^2z - \frac{1}{3}xyz^2 + \frac{1}{12}x^2y - \frac{1}{6}xyz + \frac{1}{72}yz + \frac{1}{12}z^2 + \frac{1}{2}x + \frac{1}{6}y.$$

Example 4 This example tests algorithm 4 by computing determinants, and the results are shown in table 1, where '?' represents that the computation is not finished. Table 1 shows that our algorithm is more efficient than exact interpolation algorithm when scale of problem is larger.

matrix sizes	variates	terms	error control	time(sec)	exact in- terp(sec)
3×3	3	210	0.1679e - 26	1.766	1.531
3×3	4	343	0.1829e - 30	5.578	3.837
4×4	4	256	0.5441e - 19	2.078	1.734
5×5	3	816	0.4471e - 37	5.837	4.241
6×6	3	2774	0.3781e - 45	147.452	67.731
7×7	4	5657	0.4731e - 50	258.129	347.372
8×8	4	8847	0.4251e - 55	657.221	899.372
9×9	3	15367	0.5882e - 63	2074.771	2744.305
10×10	4	23371	0.7321e - 61	5721.491	8737.707
11×11	4	40721	0.6781e - 73	9287.527	?

Table 1: Comparison between our algorithm and exact interpolation

5 Conclusions

The paper presents a new method for obtaining exact interpolation multivariate polynomial with rational coefficients from its approximate interpolation. The exact results can be obtained by our algorithm as long as we estimate the upper bounds on the degrees in every variable and upper bound N on absolute values of denominators of its coefficients. As we know, many complicate polynomials computation can be completed by exact interpolation method, so they can also be completed by our algorithm, i.e., by approximately numerical computation.

The paper proposes a way to get an exact polynomial by approximate numerical computation. Like exact interpolation methods, this algorithm is a parallel method. The accuracy control ε in formula (17) is an exponential function in degrees of polynomial, and is a polynomial function in the upper bound 1/N. Experimental results show that our method is more efficient than the exact interpolation methods when scale of a problem gets larger.

References

- Robert M. Corless, Mark W. Giesbrecht, et al, Towards factoring bivariate approximate polynomials, In Proc. ISSAC 2001, ACM press, pp.85-92.
- Huang Y., Wu W., Stetter H., and Zhi L. Pseudofactors of multivariate polynomials. In Proc. ISSAC00(2000), ACM Press, pp.161-168.
- [3] Mou-Yan, Z., and Unbehausen, R. Approximate factorization of multivariable polynomials. Signal Proces. 14(1988), pp.141-152.
- [4] Sasaki, T., Suzuki, M., et al., Approximate factorization of multivariate polynomials and absolute irreducibility testing. Japan J. Indust. Appl. Math. 8 (1991), pp.357-375.
- [5] Tateaki Sasaki, Approximate multivariate polynomial factorization based on zero-sum relations. In Proc. ISSAC2001, ACM Press, pp.284-291.
- [6] Sasaki, T., Saito T., and Hilano, T., Analysis of approximate factorization algorithm. Japan J. Indust. Appl. Math.9 (1992), pp.351-368.
- [7] Corless R. M., Gianni P.M., Trager B.M. and Watt, S.M. The singular value decomposition for polynomial systems. In International Symposium on Symbolic and Algebraic Computation (Montreal, Canada, 1995), A Levelt, Ed., ACM pp.195-207.
- [8] Beckermann, B., and Labahn, G., When are two polynomials relatively prime? Journal of Symbolic Computation 26 (1998), pp.677-689.
- [9] Karmarka N., and Lakshman Y. N., Approximate polynomial greatest common divisors and nearest singular polynomials. In Proceeding of ISSAC 1996, ACM, pp.35-42.
- [10] Robert M. Corless, Stephen M. Watt, and Lihong Zhi, QR Factoring to Compute the GCD of Univariate Approximate Polynomials IEEE Transactions on Signal Processing, 52(12),(2004), pp.3394-3402.
- [11] Robert M. Corless, MarkW. Giesbrecht, et al, Approximate polynomial decomposition. In proceeding of ISSAC 1999, S.S. Dooley, Ed., ACM pp.213-220.
- [12] Galligo A., and Watt S. M. A numerical absolute primality test for bivariate polynomials In proceeding of ISSAC 1997. W.Küchlin, Ed. ACM, pp.217-224.
- [13] Greg Reid , and Lihong Zhi, Solving Nonlinear Polynomial System via Symbolic-Numeric Elimination Method, In Proceedings of international conference on polynomial system solving,(2004), pp.50-53.
- [14] Corless, R.M., Giesbrecht, M.W., et al, Numerical implicitization of parametric hypersurfaces with linear algebra. In proceeding of AISC2000, LNAI 1930, pp.174-183.
- [15] G.Chèze, A. Galligo. From an approximate to an exact absolute polynomial factorization. J. Symbolic Comput., 41(6),(2006),pp.682-696.
- [16] Erich Kaltofen, Lakshman Yagati, Improved Sparse Mutivariate Polynomial Interpolation Algorithms. In Symbolic Algebraic Comput. Internat. Symp. ISSAC '88 Proc. [-17], pages 467-474, 1988.
- [17] Manocha D, Efficient Algorithms for multipolynomial Resultant. The Computer Journal, 1993, 36(5) 485-496.
- [18] M.Giesbrecht, G. Labahn and W-s Lee, Symbolic-numeric Sparse Interpolation of Multivariate Polynomials, Proceedings of ISSAC'06, Genoa, Italy, ACM Press, (2006) 116-123.
- [19] A Marco, JJ Martinez, Parallel computation of determinants of matrices with polynomial entries. J. Symbolic Comput. 37(2004), No.6,749-760.
- [20] Tushar Saxena, Efficient Variable Elimination using Resultants, Doctoral Thesis, Computer Science Division, State University of New York at Albany. 1997.
- [21] Yong Feng, Yaohui Li, Checking RSC Criteria for Extended Dixon Resultant by Interpolation Method, Proceedings of 7-th International Symposium on Symbolic and Numeric Algorithms for Scientific Computing, Timisoara, Romania, September 25-29, 2005, IEEE Computer Society press pages 48-51.
- [22] Yong Feng, Yaohui Li, Computing extended Dixon resultant by interpolation method. Proceedings of international workshop on Symbolic-Numeric Computation. Xi'an China, July 19-21, 2005, pp. 177-187.

- [23] Carl de Boor, Polynomial Interpolation in Several Variables, in Studies in Computer Science (in Honor of Samuel D. Conte), R. DeMillo and J. R. Rice (eds.), P87C119, Plenum Press New York. 1994.
- [24] Jingzhong Zhang, YongFeng, Obtaining Exact Value by Approximate Computations. Science in China Series A: Mathematics Vol. 50, No.9, 1361-1368, Sep. 2007.
- [25] R.A., Johnson, C. R., Topics in Matrix Analysis, Cambridge University Press, Cambridge, 1991.
- [26] John W. BREWER, Kronecker products and matrix calculus in system theory, IEEE Transactions on Circuits and Systems, 1978.
- [27] Xianke Zhang, Puhua Xu, Higher Algebra(second edition) (in Chinese), Beijing: Tsinghua University Press, 2004.
- [28] Richard A. Brualdi, Introductory Combinatorics. North-Holland Publishing Company, NewYork Oxford Amsterdam, 1977.