Stellar Archaeology in the Galactic halo with the Ultra-Faint Dwarfs: VI. Ursa Major II¹

M. Dall'Ora², Karen Kinemuchi³, Vincenzo Ripepi², Christopher T. Rodgers⁴, Gisella Clementini⁵, Luca Di Fabrizio⁶, Horace A.Smith⁷, Marcella Marconi², Ilaria Musella², Claudia Greco⁸, Charles A.Kuehn⁷, Márcio Catelan⁹, Barton J.Pritzl¹⁰, Timothy C. Beers¹¹

ABSTRACT

We present a B,V color-magnitude diagram (CMD) of the Milky Way dwarf satellite Ursa Major II (UMa II), spanning the magnitude range from $V \sim 15$ to $V \sim 23.5$ mag and extending over a 18×18 arcmin² area centered on the galaxy. Our photometry goes down to about 2 magnitudes below the galaxy's

¹Based on data collected at the 1.52 m telescope of the INAF-Osservatorio Astronomico di Bologna, Loiano, Italy, at the 2.3 m telescope of the Wyoming Infrared Observatory (WIRO) at Mt. Jelm, Wyoming, USA, and at the 1.8 m Perkins telescope of the Lowell Observatory, at Anderson Mesa, Flagstaff, Arizona, USA.

²INAF, Osservatorio Astronomico di Capodimonte, Napoli, Italy, dallora@na.astro.it, ripepi@na.astro.it, marcella@na.astro.it, ilaria@na.astro.it

³NASA-Ames Research Center/Bay Area Environmental Research Institute Mail Stop 244-30, P.O. Box 1, Moffett Field, CA 94035-0001; karen.kinemuchi@nasa.gov

 $^{^4 \}rm University$ of Wyoming, Department of Physics & Astronomy, Laramie, WY 82071, USA; crodgers@uwyo.edu

⁵INAF, Osservatorio Astronomico di Bologna, Bologna, Italy; gisella.clementini@oabo.inaf.it

⁶INAF, Centro Galileo Galilei & Telescopio Nazionale Galileo, S. Cruz de La Palma, Spain; difabrizio@tng.iac.es

⁷Department of Physics and Astronomy, Michigan State University, East Lansing, MI 48824, USA; smith@pa.msu.edu, kuehncha@msu.edu, beers@pa.msu.edu

 $^{^8{\}rm Observatoire}$ de Geneve, 51, ch. Des Maillettes, CH-1290 Sauverny, Switzerland; claudia.greco@obs.unige.ch

⁹Pontificia Universidad Católica de Chile, Departamento de Astronomía y Astrofísica, Santiago, Chile; mcatelan@astro.puc.cl

 $^{^{10}\}mbox{Department}$ of Physics and Astronomy, University of Wisconsin Oshkosh, Oshkosh, WI 54901, USA; pritzlb@uwosh.edu

¹¹Department of Physics & Astronomy and JINA: Joint Institute for Nuclear Astrophysics, Michigan State University, East Lansing, MI 48824, USA

main sequence turn-off, that we detected at $V \sim 21.5$ mag. We have discovered a bona-fide RR Lyrae variable star in UMa II, which we use to estimate a conservative dereddened distance modulus for the galaxy of $(m-M)_0=17.70\pm0.04\pm0.12$ mag, where the first error accounts for the uncertainties of the calibrated photometry, and the second reflects our lack of information on the metallicity of the star. The corresponding distance to UMa II is $34.7^{+0.6}_{-0.7}(^{+2.0}_{-1.9})$ kpc. Our photometry shows evidence of a spread in the galaxy subgiant branch, compatible with a spread in metal abundance in the range between Z=0.0001 and Z=0.001. Based on our estimate of the distance, a comparison of the fiducial lines of the Galactic globular clusters (GCs) M68 and M5 ([Fe/H]= -2.27 ± 0.04 dex and -1.33 ± 0.02 dex, respectively), with the position on the CMD of spectroscopically confirmed galaxy members, may suggest the existence of stellar populations of different metal abundance/age in the central region of UMa II.

Subject headings: galaxies: dwarf—galaxies: individual (Ursa Major II)—stars: distances—stars: variables: other—techniques: photometric

1. Introduction

In the Λ -Cold Dark Matter (Λ -CDM) paradigm, larger galaxies are hierarchically assembled by merging of small cold dark matter fragments (Diemand et al. 2007; Springel 2005), as also recently confirmed by the high-resolution simulations "Aquarius" (Lunnan et al. 2011; Springel et al. 2008). The idea is attractive when applied to our Galaxy, since it echoes the pioneering scenario suggested by Searle & Zinn (1978) (SZ), in which the Milky Way (MW) was successively built up from small substructures. Indeed, first attempts to link the SZ scenario with the cosmological simulations foresaw the assembling of the Galactic halo starting from a number of satellites, and producing a number of tidal streams that should be (and are in fact) observed (Bullock et al. 2001). The survivors of such a process should have the observational characteristics of the present day dwarf spheroidal galaxies (dSphs), which are old, metal-poor and gas deficient systems.

However, according to the Λ -CDM theory, the expected number of surviving fragments of the accretion process is one or two orders of magnitude larger than the observed number of MW "bright" dSph companions. It was also pointed out that the present-day "bright" dwarfs may be the *wrong* remnants, as they could be instead tidal dwarf galaxies, with a non-primordial origin (e.g. Metz et al. 2008). In any case this so-called "missing satellites problem" (Moore et al. 1999; Klypin et al. 1999) has represented, so far, a major issue of the

comparison between theoretical expectations and observational evidence ¹. The discrepancy was partially alleviated in the past few years by the discovery of several "ultra-faint" dwarf galaxies (UFDs) surrounding the MW, on the basis of the analysis of the Sloan Digital Sky Survey (SDSS) data (e.g., Koposov et al. 2009, and references therein). The UFDs bring the number of known MW dSph satellites to ≈ 27 (e.g. Mateo 1998; Belokurov et al. 2010 and references therein). Since the sky coverage of the SDSS is only $\sim 1/4$ of the celestial sphere, basic statistical arguments suggest that tens of these faint MW satellites are still undiscovered, thus further narrowing the gap between theoretical expectations and observational evidence unless, as suggested by Metz et al. (2009), the distribution of these faint systems is not uniform and follows instead a disc around the MW. Whatever the case, in order to be "proper" remnants of the primordial Galactic halo contributors the observed satellites should host old stellar populations with properties compatible with those of the MW halo. In particular, they should contain stars as metal-poor as [Fe/H] < -3.0 or -4.0 dex ("extremely metal-poor stars", Beers & Christlieb 2005) and RR Lyrae stars with pulsation properties conforming to the Oosterhoff dichotomy (Oosterhoff 1939) observed for field and cluster MW RR Lyrae variables (Catelan 2009, and references therein). Regarding both aspects the "bright" MW dSph satellites do not seem to be the possible primordial contributors to the Galactic halo, as spectroscopic studies (e.g. Helmi et al. 2006) show that there are very few extremely metal-poor stars in most of the "bright" MW dSph satellites (Tolstoy et al. 2009, and reference therein) and, on the other hand, these galaxies are generally classified as "Oosterhoff-intermediate" (e.g. Carina: Dall'Ora et al. 2003; Fornax: Bersier & Wood 2002) because their fundamental-mode RR Lyrae stars have average periods $(\langle P_{ab} \rangle)$ intermediate between the Oosterhoff I (OoI; $\langle P_{ab} \rangle = 0.55$ days) and II (OoII; $\langle P_{ab} \rangle$ = 0.65 days) types observed for the MW field and cluster variables. The Ursa Minor dSph $(\langle P_{ab}\rangle = 0.638 \text{ days})$ and the Sagittarius $(\langle P_{ab}\rangle = 0.574 \text{ days})$ are the only exceptions to this generally accepted "rule" among the "bright" dwarfs (Smith, Catelan, & Clementini 2009. and references therein).

The UFDs, instead, seem to possess both proper metal abundances, as shown by a number of spectroscopic studies (e.g., Simon & Geha 2007; Kirby et al. 2008; Frebel et al. 2010), as well as compatible properties of stellar populations and variable stars, as shown by our long-term project to monitor the UFD variable stars (see Moretti et al. 2009, for a summary). In brief, the five galaxies we have studied so far, namely, Bootes I (Dall'Ora et al. 2006), Canes Venatici I (CVn I; Kuehn et al. 2008), Canes Venatici II (CVn II; Greco et al. 2008), Coma Berenices (Coma; Musella et al. 2009), and Leo IV (Moretti et al. 2009), contain RR

¹However, it is important to stress that cosmological simulations refer to cold dark matter minihaloes, which may or may not host a stellar population (Bullock et al. 2001)

Lyrae stars with pulsation periods suggesting an Oo II classification, with the exception being CVn I, which appears to be Oosterhoff-intermediate. In this respect, we remark that both total luminosity and global metallicity of CVn I make this object more likely a "classical" dSph than a UFD (see e.g. Fig. 5 of Kirby et al. 2008). Continuing our study of variable stars in the Galactic UFDs in this paper we present results for Ursa Major II (UMa II).

UMa II (R.A.= $08^h 51^m 30^s$, decl.= $+63^\circ 07' 48''$, J2000, Zucker et al. 2006) was initially discovered as a candidate star cluster in the Galactic halo by Grillmair (2006), and only afterwards recognized as a dwarf galaxy by Zucker et al. (2006). According to Simon & Geha (2007) and Muñoz et al. (2010) UMa II appears to be a very elongated and extended object, likely undergoing tidal disruption. This suggestion is supported by the existence of a velocity gradient along the major axis of 8.4 ± 1.4 km s^{-1} between eastern and western sides of the galaxy (Simon & Geha 2007). This gradient shows the same direction of elongation as discussed in Muñoz et al. (2010). The galaxy was proposed to be the progenitor of the so-called Orphan Stream (Fellhauer et al. 2007). Zucker et al. (2006) derived a half light radius of $r_h \sim 13'$ for the galaxy, and highlighted that the central part of UMa II breaks up into three distinct clumps. Subsequently, Muñoz et al. (2010) on the basis of deep, widefield Canada-France-Hawaii Telescope photometry, showed that this apparent clumping is due to the poor statistics of the bright stars, and disappears when deeper data are used. Moreover, they showed that UMa II is more extended than previously thought. The same conclusion was reached by Newberg et al. (2010), from the analysis of the Sloan Extension for Galactic Understanding and Exploration (SEGUE, Yanny et al. 2009) spectroscopic and SDSS/SEGUE photometric data. These findings seem to rule out the link between UMaII and the Orphan Stream. Spectroscopic studies disclosed the presence of very metal-poor stars in the galaxy, with $[Fe/H] \sim -3.0$ dex, and detailed abundance patterns consistent with those observed in the Galactic halo (Kirby et al. 2008; Frebel et al. 2010). Martin et al. (2008) derived new structural parameters for UMaII, adopting a slightly different position for the galaxy centre and a flatter shape with a slightly larger r_h , compared to Zucker et al. (2006). In the following we use the structural parameters derived by Zucker et al. (2006). However, our conclusions would not be affected whether we used Martin et al.'s parameters, instead.

2. Observations and Data Reduction

Time-series observations of UMa II were obtained between January and March 2007 with three different telescopes. At the Wyoming Infrared Observatory (WIRO) 2.3m telescope, using the WIRO Prime Focus Camera, we obtained 40 images in Johnson V, 12 images in Johnson B, and 22 images in Cousins I. At the 1.8m Perkins telescope of the Lowell

Observatory, we used the Perkins Re-Imaging SysteM (PRISM) instrument in photometry mode to obtain 12 V, 4 B and 4 I images of UMa II. Five nights at the Loiano Observatory 1.5m telescope provided 14 B and 15 V images using the Bologna Faint Object Spectrograph and Camera (BFOSC). The WIRO observations cover the largest area with a field of view (FOV) of 17.8' × 17.8' at the pixel scale of 0.55 arcsec/pix, pointed on the center coordinates of UMa II derived by Zucker et al. (2006). The FOVs at the Lowell and Loiano telescopes are slightly smaller, 13.65' × 13.65' (0.39 arcsec/pix) and 13' × 12.6' (0.58 arcsec/pix), respectively. The total area covered by our observations is 18.38' × 18.26'. This corresponds to the central clump of UMa II, while the other two clumps reported by Zucker et al. (2006) are both slightly outside the field covered by our observations. The log of observations is provided in Table 1.

Datasets from each observatory were pre-reduced using standard IRAF² techniques (bias subtraction and flat fielding). Since we lack a photometric calibration of the I-band data, observations in this band were not considered in the subsequent analysis and only the B, V data are presented in the paper. Point-spread function (PSF) photometry was performed with the DAOPHOT IV/ALLFRAME (Stetson 1987, 1994) package. After an accurate evaluation of the PSF of each individual frame, a reference image was built by averaging all the available frames and a source catalogue was extracted from the stacked image. The source list was then passed to ALLFRAME, which performed homogeneous PSF photometry simultaneously on all images, thus producing b and v instrumental magnitude catalogues for each telescope. Typical internal photometric errors were of the order of 0.01-0.02 mag at the horizontal branch (HB) level, on the averaged catalog, while the signal-to-noise ratio greatly changes between the individual exposures, due to the different telescope sizes and very different exposure times adopted at the various sites.

To calibrate our photometry, additional B,V observations of UMa II, along with standard fields centered on the open clusters NGC 188 and NGC 7790, selected from the publicly available standard stars archive maintained by P. B. Stetson 3 , were obtained on a photometric night in 2008 January, at the 3.5m Telescopio Nazionale Galileo in La Palma, Canary Islands. We used the standard La Palma extinction coefficients 4 to derive the following calibration equations:

²IRAF is distributed by the National Optical Astronomical Observatory, which is operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation

³http://www4.cadc-ccda.hia-iha.nrc-cnrc.gc.ca/community/STETSON/standards/

⁴http://www.ast.cam.ac.uk/~dwe/SRF/camc_extinction.html

$$B = b + (0.058 \pm 0.007)(B - V) + (26.400 \pm 0.020)$$
$$V = v + (0.082 \pm 0.014)(B - V) + (26.141 \pm 0.010)$$

where B,V and b,v are the standard and the instrumental magnitudes, respectively ⁵. The r.m.s. of these equations is of ~ 0.02 mag in both bands. These calibrations were used to define secondary standards in the field of UMa II. In particular, we selected 25 isolated local standards with photometric errors less than 0.05 mag which were visually inspected to remove any non-stellar objects. The remaining stars cover a range in color of 0.1 < B-V < 1.7 mag. Since the three telescopes define slightly different photometric systems, the individual datasets were calibrated separately, and for each filter a final master catalogue was obtained by averaging the measurements of the individual telescopes. Typical uncertainties in calibrating our dataset on the TNG local standards were in the range 0.01-0.03 mag. Figure 1 shows the photometric internal error as a function of the calibrated magnitude, both in the B and V bands.

3. Identification of Variable Stars and distance to the galaxy

For each star calibrated time-series photometry was obtained firstly by shifting the instrumental magnitudes to the WIRO instrumental photometric system, and then calibrating the WIRO-aligned data to the Johnson photometric system. We omitted the relative color term dependence since the relative color terms were small enough in the observed color range of the RR Lyrae stars $(0.2 < (B - V)_0 < 0.5 \text{ mag})$ to be considered negligible for our purposes. Search and identification of candidate variable stars were performed with two different methods: 1) the Stetson (1993) variability index, which compares the spread of individual measurements with the intrinsic photometric error; 2) an ad hoc procedure in which we first computed the Fourier transforms (in the Schwarzenberg-Czerny 1996 formulation) of the stars having at least 12 measurements in each photometric band, and then we averaged these transforms to estimate the noise and calculated the signal-to-noise ratios. Results from B and V photometries were cross-correlated, and all stars with S/N> 4 in both photometric bands were visually inspected, for a total of ~ 200 candidates. In particular, we checked all the stars around the galaxy's horizontal branch and some of the SX Phoenicis type. The light

⁵The instrumental magnitude for a generic band X is defined as $m_X = -2.5 \log(F_X) - K_X X$, where F_X is the flux normalized to 1sec, K_X is the extinction coefficient, and X is the airmass.

curves of the candidate variables were inspected by eye. We confirmed only one candidate variable, whose multiband light curves were analyzed with the Graphical Analyzer of Time Series (GRaTiS, Clementini et al. 2000) package, obtaining a classification as RR Lyrae with pulsation period of 0.6593 days. We note that the time window of our data would imply an accuracy on the second decimal place of the derived period, however, we have used a period value to the forth decimal place to fold the light curves of the RR Lyrae star, since this allowed us to significantly reduce the rms scatter of the truncated Fourier series best fitting the data. The star's time series data are provided in Table 2 and the pulsation characteristics (period, epoch of maximum light, amplitudes of the V, B light variations, and intensity-averaged mean magnitudes) are summarized in Table 3. The internal accuracy of the average magnitudes of the RR Lyrae star is of \pm 0.02 mag, as estimates from the rms scatter of the models best fitting the light curves. For the amplitudes we list both the values of the difference between minimum and maximum light data points, and the values of the truncated Fourier series best fit models of the light curves (in brackets). Corresponding uncertainties in the amplitudes are of the order of 0.1 mag. Light curves are shown in Figure 2. Figure 3 shows the position of the only RR Lyrae in UMa II on the period-amplitude diagram defined by the RR Lyrae stars of the Bootes I, CVn II, Coma, and Leo IV UFDs. The pulsation period and the position on the period-amplitude diagram both suggest the similarity of UMa II to Oosterhoff type II systems. This classification is unchanged if we round the period to two decimal places. Figure 4 shows the CMD of UMa II with the RR Lyrae star marked by a cyan filled circle. The B-V color of the RR Lyrae star $(\langle B \rangle - \langle V \rangle$ = 0.22 mag) appears to be slightly too blue for the star pulsation period. We will discuss this further at the end of the section.

We have estimated the distance to UMa II from its RR Lyrae, by adopting the calibration of the absolute V magnitude of the RR Lyrae stars as a function of the metallicity provided by Clementini et al. (2003) and Gratton et al. (2004), who derived $\Delta M_V(RR)/\Delta[Fe/H]=0.214\pm0.047$ mag/dex from a sample of 100 RR Lyrae stars in the Large Magellanic Cloud (LMC) spanning the metallicity range from [Fe/H]=-2.12 to -0.27 dex, and the RR Lyrae zero-point $M_V=0.59$ mag at [Fe/H]=-1.5 dex, proposed by Cacciari & Clementini (2003). We explicitly note that, selecting a brighter zero-point of $M_V=0.54$ mag at [Fe/H]=-1.5 dex, which is in agreement with the LMC distance modulus of $(m-M)_0=18.52$ mag (Clementini et al. 2003), the kernel of our analysis is not changed. Adopting a metallicity of $[Fe/H]=-2.44\pm0.06$ dex for UMa II (Kirby et al. 2008), an average apparent magnitude of the UMa II HB of $\langle V_{\rm HB} \rangle=18.39$ mag, as derived from the galaxy RR Lyrae star, and a reddening of E(B-V)=0.096 mag from the Schlegel et al. (1998) maps, we obtain a true distance modulus for the galaxy of $(m-M)_0=17.70\pm0.04$ mag, corresponding to $34.7^{+0.6}_{-0.7}$ kpc. The quoted uncertainty takes into account only the uncertainties of the

intrinsic photometry, of the photometric calibration, and of the slope and zeropoint of the RR Lyrae M_V - [Fe/H] relationship. However, as pointed out by our refereee, the UMa II red giants exhibit a considerable spread in metal abundance, $(\sigma_{\text{Fe/H}}=0.57 \text{ dex}, \text{ according})$ to Kirby et al. 2008), that we should take into account since we lack a direct estimate of metallicity for the RR Lyrae star. This translates into an additional uncertainty of ± 0.12 mag on the distance modulus, which thus becomes $(m - M)_0 = 17.70 \pm 0.04 \pm 0.12$ mag, corresponding to $34.7^{+0.6}_{-0.7}(^{+2.0}_{-1.9})$ kpc. We also note that, since the $M_V - [Fe/H]$ relationship is extrapolated for metallicities below [Fe/H]=- 2.1 dex and there are claims that it may not be unique and/or linear (e.g. Caputo et al. 2000), this error might still be an underestimate. Our distance agrees within the uncertainties with the value obtained by Zucker et al. (2006) $((m-M)_0 = 17.5 \pm 0.3 \text{ mag})$ based on isochrone-fitting of the main sequence and turn-off regions. Our distance of 34.7 kpc also agrees with the predictions of the heliocentric distance to UMa II by Fellhauer et al. (2007). These authors simulated the orbit of UMa II obtaining for the galaxy a distance of 34 kpc, which taken at the face value would support their suggestion that UMa II is the progenitor of the Orphan Stream. However, the N-body simulations by Sales et al. (2008) show that there is no obvious link between the Orphan Stream and the orbit of UMa II.

As mentioned previously the RR Lyrae star shows a rather blue B-V color. We do not have a clear explanation for this occurence, as the image of the star does not seem to be contaminated by any companion, and the good agreement with the HB ridgeline of M68 as well as with the theoretical models by Pietrinferni et al. (2006) (see below) both in the V, B-V and B, B-V planes, seem to support the notion that the star is not affected by blending with a hot companion or by deceptive photometric errors. On the other hand, our B light curve is based only on six different epochs, as the 12 exposures obtained at the WIRO telescope span a very short time interval and contribute in fact only 2 distinct epochs, thus the derived $\langle B \rangle$ magnitude may be somewhat uncertain. However, this uncertainty does not affect significantly our distance estimate. Our time-series photometry was referred to the WIRO data, with photometric transformations to the Johnson standard system given by

$$B = b + 0.115(b - v) + 3.477$$
$$V = v - 0.098(b - v) + 3.223$$

which translate into the V-(B-V) transformation

$$V = v - 0.080(B - V) + 3.342$$

Therefore, a shift as large as 0.4 mag of the star B-V color (thus pushing the variable on the "red" side of the Instability Strip), would affect the derived $\langle V \rangle$ luminosity by ~ 0.03 mag at most.

4. The Stellar Population(s) of the UMa II UFD

Figure 4 shows our CMD of UMa II, where to minimize the contamination by non-point sources we only plot objects with DAOPHOT parameters $\chi \leq 1.4$ and $-0.3 \leq sharpness \leq$ +0.3 and also considered only stars within the galaxy's half light radius $(r_h \sim 13 \text{ arcmin})$ as worked out from the preliminary values of distance, $d \sim 30$ kpc, and linear half light radius $r_h \sim 120 \,\mathrm{pc}$, published by Zucker et al. 2006). UMa II appears to be quite elongated with an ellipticity of ~ 0.5 , therefore we adopted the azimuthally averaged value of the half light radius (Zucker et al. 2006). We explicitly note that UMa II shows a rapidly varying ellipticity, with values ranging from $\epsilon = 0.40$ in the center to $\epsilon = 0.73$ near its half light radius (Muñoz et al. 2010), but this does not affect our conclusions about the stellar populations present in the galaxy. The RR Lyrae star we have discovered in the galaxy is marked by a cyan filled circle, while the cyan filled triangle shows the position of star SDSSJ084947.6+630830, which was suspected to be an RR Lyrae variable by Simon & Geha (2007) on the basis of the observed radial velocity and spectral type variations. Unfortunately, this star falls outside our field of view. Nevertheless, we show its position on our CMD, after transforming the SDSS g, r magnitudes to the Johnson standard system with the relations by Fukugita et al. (1996), as it appears to be consistent with the RR Lyrae instability strip, thus supporting the Simon & Geha (2007) hypothesis. Blue, red, magenta and cyan open circles mark spectroscopically confirmed members of UMa II according to Martin et al. (2007), Simon & Geha (2007), Kirby et al. (2008), and Frebel et al. (2010), respectively. No correction for contamination by Galactic field stars was made, as our attempt to compare the PPM-Extended (PPMX) proper motions catalog (Roeser et al. 2010) of the spectroscopically confirmed UMaII members with all the other stars present in the field did not show a clear separation.

UMa II is characterized by a very low average metal abundance \langle [Fe/H] \rangle =-2.44 dex, an internal metallicity spread as large as \sim 0.6 dex (Kirby et al. 2008), and α element abundances in agreement with those observed in the Galactic halo ([α/Fe] = 0.4 dex (Frebel et al. 2010, and references therein). In Figure 4 we have overplotted to the UMa II CMD the ridgelines of the Galactic GCs M68 (blue dashed line, from Walker 1994) and M5 (blue solid line, from Sandquist et al. 1996), because these two GCs well encompass the metallicity range found in the galaxy. Indeed, M68 is a metal-poor cluster, with $[Fe/H] = -2.27 \pm 0.04$

dex (Carretta et al. 2009), and more recent determinations claim an even lower metallicity of $[Fe/H] = -2.44 \pm 0.1$ dex (Simmons et al. 2011), hence very similar to the average metallicity of UMaII; M5 is a metal-intermediate cluster, with $[Fe/H] = -1.33 \pm 0.02$ dex (Carretta et al. 2009). The ridgelines were properly shifted to account for differences in distance and reddening. For M68 we used the RR Lyrae mean apparent V magnitude $\langle V(RR) \rangle = 15.60$ mag by Walker (1994) and a reddening of E(B-V) = 0.05 mag, as available in the compilation by Harris (1996). With these values we get an apparent distance modulus of 15.11 mag, by adopting the RR Lyrae magnitude-metallicity calibration as above. For M5, we have adopted $(m-M)_0 = 14.44$ mag from Coppola et al. (2011) and E(B-V) = 0.03 mag (Harris 1996).

Our RR Lyrae-based distance to UMa II is supported by the good match of the M68 ridgelines and the UMaII subgiant branch (SGB), and by the very satisfactory alignment, along both the M68 and M5 HBs, of three bona-fide UMaII stars, one of them being a spectroscopically confirmed member. The good match between few stars located at V in the range from 21.0 to 21.5 mag and B-V color between 0.4-0.5 and 0.7-0.8 mag, two of which are spectroscopically confirmed members of UMaII with the M5 ridgeline seems to suggest the presence of a second SGB. This is not surprising, since the quoted spectroscopic works clearly demonstrate the occurrence of stellar populations with different metallicities in this UFD. Moreover, the galaxy bona-fide main sequence appears to have an observed width of ≈ 0.2 mag in the range of luminosities between $V \approx 22$ and $V \approx 23$ mag, while the width expected in that range due to the photometric errors alone would be of the order of $\sigma_{B-V} \approx 0.1 \text{ mag}$ (see Fig. 1). In the following, we will refer to the stars close to the M68 and M5 ridgelines as "blue" and "red" populations, respectively. To check if the blue and red population are also compatible with a spread in the age, we have compared in Figure 5 the CMD of UMa II with the BaSTI (Bag of Stellar Tracks and Isochrones) α -enhanced isochrones by Pietrinferni et al. (2006). Following the recent α -enhanced abundances measured by Frebel et al. (2010), we have adopted for the blue population the α -enhanced models with Z = 0.0001 and $[\alpha/Fe] = 0.4$ dex, corresponding to an iron-over-hydrogen ratio of [Fe/H] = -2.62 dex and a global metallicity of [M/H] = -2.27 dex, and ages of 12, 13, and 14 Gyr. For the red population we have used the α -enhanced models with Z=0.001and $\left[\alpha/Fe\right] = 0.4$ dex, corresponding to an iron-over-hydrogen ratio of $\left[Fe/H\right] = -1.62$ dex and a global metallicity of [M/H] = -1.27 dex, with ages of 11, 12, 13 and 14 Gyr, since M5 is thought to be ~ 1 Gyr younger than M68 (see Marín-Franch et al. 2009). We stress that our selection of the isochrones is entirely based on the measured metallicities. It is also important to note that UMaII hosts a very metal-poor stellar component with |Fe/H| = -3 dex, (Frebel et al. 2010). Unfortunately, the BaSTI database does not contain such low metallicities. We also explicitly note that the spectroscopically confirmed member of UMa II at $V \approx 19.8$ mag and $(B-V) \approx 1$ mag has a measured metallicity of [Fe/H] = -1.8 dex (SDSSJ085117.07+630347.3, Martin et al. 2007) and appears to be definitely too red for its metallicity, being redder than spectroscopically confirmed members close to the M5 ridgeline and the Z=0.001 isochrones. The star could be contaminated by a red companion, even if the values of its DAOPHOT structural parameters χ and sharpness do not support this explanation. Strangely, this star is not present in the 2MASS catalog, even if stars of similar luminosity are reported.

The isochrone fitting appears to be quite satisfactory (see Fig. 5) but, as commented by the referee, the small statistics does not allow us to prove the existence of two distinct stellar populations, with different age and/or metallicity in UMa II. It is worth to mention that Zucker et al. (2006) suggested the presence of an age/metallicity spread in UMa II that could be accounted for by an an intermediate-metallicity stellar population with age of at least 10 Gyr. We do not recover such a spread, but our data marginally suggest a gap between the metal-poor and the metal-intermediate SGBs. If this gap is not due to the poor statistics, UMaII may have experienced distinct bursts of star formation during its star formation history, similar to those found in the Carina dwarf galaxy (Monelli et al. 2003), for example. However, an alternative, intriguing explanation has been proposed by Frebel & Bromm (2010), who suggested that in the poor baryonic matter content of the UFDs, pristine (i.e. Population III) core collapse (and maybe pair-instability) supernova events may have suppressed further star formation and inhibited a homogeneous mixing of the interstellar medium. According to this model, stars born close to the core collapse SNe progenitors would be chemically enriched from the supernova yields, while more distant stars would not. Therefore, the stellar populations in UMa II could have the same age but different metallicity as arising from differently enriched material. Alternatively, several Population III SNes could have enriched the environment but, given the shallow potential well of the UFDs, part of the enriched gas was lost (Frebel 2011). In either the case, this kind of galaxies should be regarded as "one-shot events". Our photometric data do not allow us to distinguish between the two possible scenarios since the red population is reasonably well described by isochrones in the range between 11 and 14 Gyrs.

In order to verify whether the stellar populations show a different spatial distribution in the field of view covered by our observations, we have used the ridgelines of M68 and M5 to select stars possibly compatible with a blue and a red population, and plotted them on the galaxy map with different colors. This is shown in Figures 6 and 7. In particular, Figure 6 illustrates our selection of the two components by displaying as blue and red filled circles the sources, within the UMaII half light radius, which lie within ± 0.05 mag in B-V for $V \leq 21.5$ mag and ± 0.10 mag in B-V for V > 21.5 mag from the ridgelines of M68 and M5, respectively. While the selection is rather straightforward for $V \leq 21.5$ mag, at fainter

magnitudes the two components cannot be totally separated. We have marked in green the sources with V>21.5 mag which could belong to either population. Figure 7 shows the spatial location of these blue-, red-, and green-coded populations, in the FOV covered by our observations, for sources with $V\leq 21.5$ mag (upper panel) and for all sources (lower panel). The upper panel of Figure 7 shows that sources brighter than V=21.5 mag lying closer to the M68 ridgeline (blue filled circles) are larger in number and seem to be more concentrated towards the center of UMa II than sources closer to the M5 ridgeline (red filled circles), but this could be an effect due to the poor statistics. On the other hand, the lower panel of Figure 7 shows that sources fainter than V=21.5 mag, which could belong either to the M68-like or to the M5-like population (green filled circles), do not show any obvious pattern.

5. Conclusions

We have presented the first Johnson V, B - V CMD of the UMa II UFD galaxy and reported the detection of a bona-fide RR Lyrae star in the galaxy, from which we have derived a true distance modulus of $(m-M)_0 = 17.70 \pm 0.04(\pm 0.12)$ mag $(d=34.7^{+0.6}_{-0.7}(^{+2.0}_{-1.9})$ kpc). The error quoted in the brackets takes into account our lack of knowledge of the metallicity of the RR Lyrae star. The pulsation period of this variable (P=0.6593 days) and its position in the period-amplitude (Bailey) diagram suggests the similarity of UMaII with Oosterhoff type II systems. A comparison of the observed CMD with the location of UMa II spectroscopically confirmed member stars (taken from the catalogs of Martin et al. 2007; Simon & Geha 2007; Kirby et al. 2008; Frebel et al. 2010) and with the fiducial lines of the Galactic GCs M68 and M5 is compatible with a spread in age/metallicity of the stellar content of the galaxy. The inner brightness profile of UMaII, as suggested by Muñoz et al. (2010) on the basis of deep CFHT photometry, could be due to the presence of multiple stellar generations, as also postulated in McConnachie et al. (2007). However, the comparison with the isochrones gives only a coarse estimate of the age of the stellar populations, and no definite conclusion can be reached about a possible spread in age. On the other hand, when only the comparison with the M68 and M5 ridgelines is considered, the hypothesis of an age difference of ~ 1 Gyr is favored. Since the field covered by our photometry is smaller than the fields observed by the Martin et al. (2007), Simon & Geha (2007), Kirby et al. (2008), and Frebel et al. (2010) studies, our conclusions apply mainly to the central part of UMaII monitored by our data. A deeper, more accurate and more spatially extended photometry is therefore highly desided to unveil all the secrets of the UMa II UFD.

We thank an anonymous referee, for helpful comments, which greatly improved both

the scientific accuracy and the readability of this paper. We warmly thank Evan Kirby and Joshua Simon for providing us identification and individual metallicities for member stars of the UMaII UFD. PRISM was developed with support from NSF (AST-0079541, K.A. Janes, P.I.) with additional support from Boston University and Lowell Observatory. We also thank Benjamin Kelly and Laura Portscheller for having joined us in the first phase of this project. We acknowledge financial contribution from the Italian PRIN MUR 2007 "Multiple stellar populations in globular clusters: census, characterization and origin" P.I.: G. Piotto, from COFIS ASI-INAF I/016/07/0, and from the agreement ASI-INAF I/009/10/0. HAS thanks the NSF for support under grants AST-0607249 and AST-0707756. MC acknowledges support by the Chilean Ministry for the Economy, Development, and Tourisms Programa Iniciativa Científica Milenio through grant P07-021-F, awarded to The Milky Way Millennium Nucleus; by the BASAL Center for Astrophysics and Associated Technologies (PFB-06); by the FONDAP Center for Astrophysics (15010003); by Proyecto Fondecyt Regular #1110326; and by Proyecto Anillo ACT-86.T.C.B. acknowledges partial funding of this work from grants PHY 02-16783 and PHY 08-22648: Physics Frontier Center/Joint Institute for Nuclear Astrophysics (JINA), awarded by the U.S. National Science Foundation.

REFERENCES

Beers, T. C., & Christlieb, N. 2005, ARA&A, 43, 531

Belokurov, V., et al. 2010, ApJ, 712, L103

Bersier, D, & Wood, P. R. 2002, AJ, 123, 840

Bullock, J. S., Kravtsov, A. V., & Weinberg, D. H. 2001, ApJ, 548, 33

Cacciari, C., & Clementini, G. 2003, in Stellar Candles for the Extragalactic Distance Scale, ed. D. Alloin & W.Gieren (Berlin: Springer), 105

Caputo, F., Castellani, V., Marconi, M., & Ripepi, V. 2000, MNRAS, 316, 819

Carretta, E., Bragaglia, A., Gratton, R., D'Orazi, V., & Lucatello, S. 2009, A&A, 505, 139

Catelan, M. 2009, Ap&SS, 320, 261.

Clement, C. M., & Rowe, J. 2000, AJ, 120, 2579

Clementini, G., et al. 2000, AJ, 120, 2054

Clementini, G., Gratton, R.G., Bragaglia, A., Carretta, E., Di Fabrizio, L., & Maio, M. 2003, AJ, 125, 1309

Coppola, G., Dall'Ora, M., Ripepi, V., et al. 2011, MNRAS, 416, 1056

Dall'Ora, M., et al. 2003, AJ, 126, 197

Dall'Ora, M., et al. 2006, ApJ, 653, L109

Diemand, J., Kuhlen, M., & Madau, P. 2007, ApJ, 667, 859

Fellhauer, M., et al. 2007, MNRAS, 375, 1171

Frebel, A. 2011, arXiv:1108.4692

Frebel, A., & Bromm, V. 2010, arXiv:1010.1261

Frebel, A., Simon, J. D., Geha, M., & Willman, B. 2010, ApJ, 708, 560

Fukugita, M., Ichikawa, T., Gunn, J. E., Doi, M., Shimasaku, K., & Schneider, D. P. 1996, AJ, 111, 1748

Gratton, R. G., Bragaglia, A., Clementini, G., Carretta, E., Di Fabrizio, L., Maio, M., & Taribello, E. 2004, A&A, 421, 937

Greco, C., Dall'Ora, M., Clementini, G., et al. 2008, ApJ, 675, L73

Grillmair, C. J. 2006, ApJ, 645, L37

Harris, W. E. 1996, AJ, 112, 1487

Helmi, A., et al. 2006, ApJ, 651, L121

Kirby, E. N., Simon, J. D., Geha, M., Guhathakurta, P., & Frebel, A. 2008, ApJ, 685, L43

Klypin, A., Kravtsov, A. V., Valenzuela, O., & Prada, F. 1999, ApJ, 522, 82

Koposov, S. E., et al. 2009, ApJ, 696, 2179

Kuehn, C., et al. 2008, ApJ, 674, L81

Lunnan, R., Vogelsberger, M., Frebel, A., Hernquist, L., Lidz, A., & Boylan-Kolchin, M. 2011, arXiv:1105.2293

Marín-Franch, A., et al. 2009, ApJ, 694, 1498

Martin, N. F., de Jong, J. T. A., & Rix, H.-W. 2008, ApJ, 684, 1075

Martin, N. F., Ibata, R. A., Chapman, S. C., Irwin, M., & Lewis, G. F. 2007, MNRAS, 380, 281

Mateo, M. L. 1998, ARA&A, 36, 435

McConnachie, A. W., Peñarrubia, J., & Navarro, J. F. 2007, MNRAS, 380, L75

Metz, M., Kroupa, P., & Jerjen, H. 2009, MNRAS, 394, 2223

Metz, M., Kroupa, P., & Libeskind, N. I. 2008, ApJ, 680, 287

Monelli, M., et al. 2003, AJ, 126, 218

Moore, B., Ghigna, S., Governato, F., Lake, G., Quinn, T., Stadel, J., & Tozzi, P. 1999, ApJ, 524, L19

Moretti, M. I., et al. 2009, ApJ, 699, L125

Muñoz, R. R., Geha, M., & Willman, B. 2010, AJ, 140, 138

Musella, I., et al. 2009, ApJ, 695, L83

Newberg, H. J., Willett, B. A., Yanny, B., & Xu, Y. 2010, ApJ, 711, 32

Oosterhoff, P. T. 1939, The Observatory, 62, 104

Pietrinferni, A., Cassisi, S., Salaris, M., & Castelli, F. 2006, ApJ, 642, 797

Roeser, S., Demleitner, M., & Schilbach, E. 2010, AJ, 139, 2440

Sales, L. V., et al. 2008, MNRAS, 389, 1391

Sandquist, E. L., Bolte, M., Stetson, P. B., & Hesser, J. E. 1996, ApJ, 470, 910

Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, ApJ, 500, 525

Schwarzenberg-Czerny, A. 1996, ApJ, 460, L107

Searle, L., & Zinn, R. 1978, ApJ, 225, 357

Simmons, S. K., Sneden, C., Preston, G., Thompson, I., Sobeck, J., & Shectman, S. 2011, American Astronomical Society Meeting Abstracts, 217, #153.13

Simon, J. D., & Geha, M. 2007, ApJ, 670, 313

Smith, H.A., Catelan, M., & Clementini, G. 2009, in Stellar Pulsation: Challenges for Theory and Observation, eds. J. A. Guzik, P. A. Bradley, AIPC, 1170, 179 press

Springel, V. 2005, MNRAS, 364, 1105

Springel, V., Wang, J., Vogelsberger, M., et al. 2008, MNRAS, 391, 1685

Stetson, P. B. 1987, PASP, 99, 191

Stetson, P. B. 1993, IAU Colloq. 136: Stellar Photometry - Current Techniques and Future Developments, 291

Stetson, P. B. 1994, PASP, 106, 250

Tolstoy, E., Hill, V., & Tosi, M. 2009, ARA&A, 47, 371

Walker, A.R. 1994, AJ, 108, 555

Yanny, B., et al. 2009, AJ, 137, 4377

Zucker, D. B., et al. 2006, ApJ, 650, L41

This preprint was prepared with the AAS $\mbox{\sc IAT}_{\mbox{\sc EX}}\mbox{\sc X}$ macros v5.2.

Table 1: Observing Log. See text for details on the individual instruments.

Night	Instrument	Filter	Exposure	N. of images	Median seeing	
			(sec)		(arcsec)	
2007-01-16	WIRO	V	10	4	3.4	
2007-01-17	WIRO	V	20	30	2.6	
2007-01-17	WIRO	V	60	6	2.6	
2007-01-17	WIRO	В	30	12	2.2	
2007-02-22	PRISM	V	300	12	2.0	
2007-02-22	PRISM	В	600	4	2.1	
2007-03-12	BFOSC	V	1200	3	1.6	
2007-03-12	BFOSC	В	1200	3	1.6	
2007-03-13	BFOSC	V	1200	3	1.6	
2007-03-13	BFOSC	В	1200	3	1.4	
2007-03-14	BFOSC	V	1200	2	1.6	
2007-03-14	BFOSC	В	1200	2	1.3	
2007-03-15	BFOSC	V	1200	2	1.5	
2007-03-15	BFOSC	В	1200	3	1.5	
2007-03-16	BFOSC	V	1200	3	1.6	
2007-03-16	BFOSC	V	1200	2	1.4	

Table 2: B, V Photometry of the RR Lyrae star detected in UMa II

HJD	В	HJD	V	
(-2454117)	(mag)	(-2454116)	(mag)	
0.756437	18.27	0.900689	18.92	
0.757167	18.27	0.902199	18.81	
0.758047	18.28	0.902819	18.87	
0.758747	18.27	0.903249	18.91	
0.759427	18.29	0.903679	18.75	
0.761087	18.29	1.750797	17.97	
0.832037	18.69	1.752317	17.98	
0.832677	18.69	1.752877	17.98	
0.833317	18.70	1.753467	18.00	
0.834037	18.68	1.753997	18.00	
0.834687	18.69	1.754927	18.00	
0.835567	18.71	1.781567	18.15	
36.764746	19.19	1.782127	18.18	
36.814344	19.23	1.782657	18.13	
36.841012	18.49	1.783177	18.17	
36.887420	17.97	1.783707	18.17	

Table 2 is published in its entirety in the electronic edition of the Journal. A portion is shown here for guidance regarding its form and content.

Table 3: Identification and properties of the RR Lyrae star detected in UMaII.

Name	α (2000)	δ (2000)	Туре	P (days)	Epoch (-2450000)	⟨V⟩ (mag)	N_{V}	⟨B⟩ (mag)	N_B	A _V (mag)	A _B (mag)
V1	08:50:37.43	+63:10:10.0	RRab	0.6593	4153.8779	18.39	53	18.61	16	$\geq 1.09(1.22)$	$\geq 1.27(1.38)$

Mean values are intensity-averaged mean magnitudes.




Fig. 1.— Photometric internal errors as a function of the calibrated B (left) and V (right) magnitudes.




Fig. 2.— B,V light curves of the lonely bona-fide RR Lyrae variable, star V1, detected in the UMa II dSph. See text for details.




Fig. 3.— Period-amplitude diagram in the V band. The solid lines show the positions of the Oo I and Oo II Galactic GCs, according to Clement & Rowe (2000). Different symbols correspond to RR Lyrae stars identified in five of the ultra-faint SDSS dwarfs studied so far for variability (namely Bootes I, CVn II, Coma, Leo IV, and UMa II, from the present study.) For UMa II we have plotted the amplitude of the truncated Fourier series best fitting the V data (A_V =1.22 mag). Results are unchanged if we use instead the amplitude corresponding to the difference of the observed minimum and maximum light data points (A_V =1.09 mag).




Fig. 4.— V, B-V Color-Magnitude Diagram of UMa II with superimposed the ridgelines of the Galactic globular clusters M68 (blue dashed line) and M5 (red dashed line). Large blue, red, magenta and cyan open symbols identify spectroscopically confirmed members of UMa II according to Martin et al. (2007), Simon & Geha (2007), Kirby et al. (2008), and Frebel et al. (2010), respectively. The cyan filled circle marks the galaxy's bona-fide RR Lyrae star found in our study, while the cyan filled triangle shows a suspected RR Lyrae in Simon & Geha (2007) study. This star falls outside the FOV of our observations, its magnitudes are taken from Simon & Geha (2007) and correspond to random phase values.




Fig. 5.— Isochrones from Pietrinferni et al. (2006) overplotted to the galaxy CMD. We adopted Z=0.0001 and $[\alpha/Fe]=0.4$ and ages of 12, 13, and 14 Gyrs (blue solid lines), and Z=0.001 and $[\alpha/Fe]=0.4$ and ages of 11, 12, 13, and 14 Gyrs (red solid lines), respectively.




Fig. 6.— V, B-V Color-Magnitude Diagram of UMa II with the ridgelines of the Galactic GCs M68 (blue solid line) and M5 (red solid line) superimposed, which we have used to select stars belonging to the two different populations that we suggest to exist in UMa II. Blue and red dots mark the stars within ± 0.05 mag in B-V for $V \leq 21.5$ mag, and ± 0.10 mag in B-V for V > 21.5 mag from the ridgelines of M68 and M5, respectively. Green dots show the sources with V > 21.5 mag which could belong to either populations. Large open circles identify spectroscopically confirmed members of UMa II according to Martin et al. (2007), Simon & Geha (2007), and Kirby et al. (2008). The filled triangle marks the galaxy's bona-fide RR Lyrae star we detected in our study.




Fig. 7.— Upper panel: Map of sources with $V \leq 21.5$ mag, within the UMa II half light radius, which we consider to belong to the galaxy according to the comparison with the ridgelines of M68 (blue filled circles) and M5 (red filled circles), or with spectroscopically confirmed membership (black open circles). The symbol sizes are proportional to the star's brightness. Lower panel: Map of all sources within the UMa II half light radius, which we consider to belong to the galaxy either according to the comparison with the ridgelines of M68 (blue filled circles) and M5 (red filled circles), or with spectroscopically confirmed membership (black open circles). Marked in green are sources with V > 21.5 mag which could belong to either populations. In both panels the RR Lyrae star is marked by a cyan filled triangle.