

SHARP HEAT KERNEL ESTIMATES IN THE FOURIER-BESSEL SETTING FOR A CONTINUOUS RANGE OF THE TYPE PARAMETER

ADAM NOWAK AND LUZ RONCAL

ABSTRACT. The heat kernel in the setting of classical Fourier-Bessel expansions is defined by an oscillatory series which cannot be computed explicitly. We prove qualitatively sharp estimates of this kernel. Our method relies on establishing a connection with a situation of expansions based on Jacobi polynomials and then transferring known sharp bounds for the related Jacobi heat kernel.

1. INTRODUCTION

Let J_ν denote the Bessel function of the first kind and order $\nu > -1$, and let $\{\lambda_{n,\nu} : n \geq 1\}$ be the sequence of successive positive zeros of J_ν . The Fourier-Bessel heat kernel is given by the oscillating sum

$$(1) \quad G_t^\nu(x, y) = 2(xy)^{-\nu} \sum_{n=1}^{\infty} \exp(-t\lambda_{n,\nu}^2) \frac{J_\nu(\lambda_{n,\nu}x)J_\nu(\lambda_{n,\nu}y)}{|J_{\nu+1}(\lambda_{n,\nu})|^2}.$$

The numbers $\lambda_{n,\nu}^2$ and the functions $u \mapsto \sqrt{2}u^{-\nu}J_\nu(\lambda_{n,\nu}u)/|J_{\nu+1}(\lambda_{n,\nu})|$ occurring here are the eigenvalues and the eigenfunctions, respectively, of the Bessel differential operator $-\frac{d^2}{du^2} - \frac{2\nu+1}{u} \frac{d}{du}$ considered on the interval $(0, 1)$. It is well known that this kernel gives the solution of the initial-value problem for the Bessel heat equation on $(0, 1)$ with Dirichlet boundary condition at the right endpoint.

Our main result is the following qualitatively sharp description of $G_t^\nu(x, y)$.

Theorem A. *Assume that $\nu > -1$. Given any $T > 0$, there exist positive constants C, c_1 and c_2 , depending only on ν and T , such that*

$$\begin{aligned} & \frac{1}{C} [t + xy]^{-\nu-1/2} \frac{(1-x)(1-y)}{t + (1-x)(1-y)} \frac{1}{\sqrt{t}} \exp\left(-c_1 \frac{(x-y)^2}{t}\right) \\ & \leq G_t^\nu(x, y) \\ & \leq C [t + xy]^{-\nu-1/2} \frac{(1-x)(1-y)}{t + (1-x)(1-y)} \frac{1}{\sqrt{t}} \exp\left(-c_2 \frac{(x-y)^2}{t}\right), \end{aligned}$$

uniformly in $x, y \in [0, 1]$ and $0 < t \leq T$. Moreover,

$$C^{-1} \exp(-t\lambda_{1,\nu}^2) (1-x)(1-y) \leq G_t^\nu(x, y) \leq C \exp(-t\lambda_{1,\nu}^2) (1-x)(1-y),$$

uniformly in $x, y \in [0, 1]$ and $t \geq T$.

2010 Mathematics Subject Classification: primary 42C05; secondary 35K08, 60J35.

Key words and phrases: Fourier-Bessel expansions, heat kernel, Bessel process, transition density.

The first-named author was supported in part by MNiSW Grant N N201 417839. Research of the second-named author supported by the grant MTM2012-36732-C03-02 from Spanish Government.

The main contents of Theorem A are of course the short time bounds. Proving them by a direct analysis of the heavily oscillating series in (1) is practically impossible. Note that the Bessel function J_ν is transcendental in general, and can be expressed by means of elementary functions only if the index ν is half-integer, i.e. $\nu = k/2$ for some integer k . Furthermore, the zeros of J_ν are known explicitly only when $\nu = \pm 1/2$. Note also that the order of magnitude of the numbers $\lambda_{n,\nu}^2$ appearing in the exponential factor is n^2 as $n \rightarrow \infty$; in particular, the asymptotic distribution of $\lambda_{n,\nu}^2$ is not linear.

The behavior of the Fourier-Bessel heat kernel does not seem to have been studied before, except for our previous paper [4]. More precisely, in [4, Theorem 3.3] we derived qualitatively sharp estimates for $G_t^\nu(x, y)$ under the assumption that ν is a half-integer not less than $-1/2$ (actually, this restriction is essential only for the short time bounds, see [4, Theorem 3.7]). We also conjectured analogous estimates for general $\nu > -1$, see [4, Conjecture 4.3], which are now confirmed by Theorem A. However, the methods we use in the present paper are much different from those applied in [4]. Here the main argument is based on a relation we establish between the Fourier-Bessel setting and the framework related to Jacobi ‘functions’. This makes it possible to transfer qualitatively sharp estimates of the Jacobi heat kernel obtained recently by Coulhon, Kerkycharian and Petrushev [2] and independently by Nowak and Sjögren [5]. We remark that actually in [2] heat kernel estimates were established in a very general setting but, nevertheless, they do not apply directly to the Fourier-Bessel case.

Our principal motivation for investigating $G_t^\nu(x, y)$ comes from an interest in harmonic analysis related to Fourier-Bessel expansions, see the references in [4, Section 1]. Additional motivation emerges from the probabilistic interpretation of the Fourier-Bessel heat kernel, see [3, Section I]. Namely, it is a transition probability density for the time scaled Bessel process $X_{2t}^{\nu+1/2}$ on $(0, \infty)$ (with reflecting barrier at $x = 0$ when $-1 < \nu < 0$, see [1, Appendix I, Section 21]) killed upon leaving the interval $(0, 1)$. We also note that for $\nu = d/2 - 1$, $d \geq 1$, the kernel $G_t^\nu(x, y)$ provides a fundamental solution to the standard heat equation in the d -dimensional Euclidean unit ball with Dirichlet boundary condition and a radial initial condition.

There are two simple cases occurring already in Fourier’s works, at least implicitly. These are $G_t^{-1/2}(x, y)$ and $G_t^{1/2}(x, y)$, and the two kernels can be written by means of non-oscillating series, see [4, Section 4]. The argument is based on the periodized Gauss-Weierstrass kernel and simple boundary-value problems for the classical heat equation in an interval. No more elementary representations in these cases seem to be possible, and this suggests that a general closed explicit formula for $G_t^\nu(x, y)$ does not exist. The estimates in Theorem A are therefore a natural and desirable substitute for an exact expression.

2. FOURIER-BESSEL AND JACOBI SETTINGS

Given $\nu > -1$, consider the Bessel differential operator

$$L^\nu = -\frac{d^2}{dx^2} - \frac{2\nu + 1}{x} \frac{d}{dx}$$

on the interval $(0, 1)$. It is well known that L^ν is symmetric and nonnegative on $C_c^2(0, 1) \subset L^2(d\mu_\nu)$, where μ_ν is a power measure in the interval $[0, 1]$ given by

$$d\mu_\nu(x) = x^{2\nu+1} dx.$$

A classical orthonormal basis in $L^2(d\mu_\nu)$ consisting of eigenfunctions of L^ν is the Fourier-Bessel system $\{\phi_n^\nu : n \geq 1\}$ (cf. [8, Chapter XVIII]) defined by

$$\phi_n^\nu(x) = \frac{\sqrt{2}}{|J_{\nu+1}(\lambda_{n,\nu})|} x^{-\nu} J_\nu(\lambda_{n,\nu}x), \quad n \geq 1, \quad x \in [0, 1];$$

here the value $\phi_n^\nu(0)$ is understood in the limiting sense. We have

$$L^\nu \phi_n^\nu = \lambda_{n,\nu}^2 \phi_n^\nu, \quad n \geq 1.$$

The self-adjoint extension of L^ν associated with the Fourier-Bessel system is defined naturally in $L^2(d\mu_\nu)$ via its spectral decomposition as

$$\tilde{L}^\nu f = \sum_{n=1}^{\infty} \lambda_{n,\nu}^2 \langle f, \phi_n^\nu \rangle_{d\mu_\nu} \phi_n^\nu,$$

on the domain

$$\text{Dom } \tilde{L}^\nu = \left\{ f \in L^2(d\mu_\nu) : \sum_{n=1}^{\infty} |\lambda_{n,\nu}^2 \langle f, \phi_n^\nu \rangle_{d\mu_\nu}|^2 < \infty \right\}.$$

The semigroup generated by $-\tilde{L}^\nu$ has the integral representation

$$\exp(-t\tilde{L}^\nu) f(x) = \int_0^1 G_t^\nu(x, y) f(y) d\mu_\nu(y),$$

where the kernel is given by (1).

For technical reasons, we also need to consider a closely related setting emerging from incorporating the measure μ_ν into the system $\{\phi_n^\nu\}$. In this way we derive the Fourier-Bessel system $\{\psi_n^\nu : n \geq 1\}$,

$$\psi_n^\nu(x) = x^{\nu+1/2} \phi_n^\nu(x), \quad n \geq 1, \quad x \in [0, 1];$$

the value $\psi_n^\nu(0)$ is understood in the limiting sense, in particular $\psi_n^\nu(0) = \infty$ for $\nu < -1/2$. For each $\nu > -1$ this system is an orthonormal basis in $L^2(dx)$ (here and elsewhere dx stands for Lebesgue measure in the interval $[0, 1]$) consisting of eigenfunctions of the differential operator

$$(2) \quad \mathbb{L}^\nu = -\frac{d^2}{dx^2} - \frac{1/4 - \nu^2}{x^2};$$

more precisely, we have

$$\mathbb{L}^\nu \psi_n^\nu = \lambda_{n,\nu}^2 \psi_n^\nu, \quad n \geq 1.$$

The operator \mathbb{L}^ν is symmetric and nonnegative on $C_c^2(0, 1) \subset L^2(dx)$. Similarly as in the previous setting, there is a natural self-adjoint extension of \mathbb{L}^ν in $L^2(dx)$, denoted by $\tilde{\mathbb{L}}^\nu$, whose spectral decomposition is given by the ψ_n^ν . The associated heat semigroup $\exp(-t\tilde{\mathbb{L}}^\nu)$ possesses an integral representation (with integration against dx), and the relevant kernel is

$$(3) \quad \mathbb{K}_t^\nu(x, y) = (xy)^{\nu+1/2} G_t^\nu(x, y);$$

this identity is understood in the limiting sense whenever $x = 0$ or $y = 0$.

Finally, we introduce the Jacobi setting based on Jacobi trigonometric ‘functions’, see [5, Section 2], scaled to the interval $[0, 1]$. Let $P_k^{\alpha,\beta}$, $k = 0, 1, 2, \dots$, be the classical Jacobi polynomials with type parameters $\alpha, \beta > -1$, as defined in Szegő’s monograph [7]. Define

$$\Phi_k^{\alpha,\beta}(x) = c_k^{\alpha,\beta} \left(\sin \frac{\pi x}{2} \right)^{\alpha+1/2} \left(\cos \frac{\pi x}{2} \right)^{\beta+1/2} P_k^{\alpha,\beta}(\cos \pi x), \quad k \geq 0, \quad x \in [0, 1],$$

with

$$c_k^{\alpha,\beta} = \left(\frac{\pi(2k + \alpha + \beta + 1)\Gamma(k + \alpha + \beta + 1)\Gamma(k + 1)}{\Gamma(k + \alpha + 1)\Gamma(k + \beta + 1)} \right)^{1/2},$$

where for $k = 0$ and $\alpha + \beta = -1$ the product $(2k + \alpha + \beta + 1)\Gamma(k + \alpha + \beta + 1)$ must be replaced by $\Gamma(\alpha + \beta + 2)$. The values $\Phi_k^{\alpha,\beta}(0)$ and $\Phi_k^{\alpha,\beta}(1)$ are understood in the limiting sense and they may be infinite for some α and β . Then the system $\{\Phi_k^{\alpha,\beta} : k \geq 0\}$ is an orthonormal basis in $L^2(dx)$. Moreover, each $\Phi_k^{\alpha,\beta}$ is an eigenfunction of the differential operator

$$\mathbb{J}^{\alpha,\beta} = -\frac{d^2}{dx^2} - \frac{\pi^2(1/4 - \alpha^2)}{4\sin^2(\pi x/2)} - \frac{\pi^2(1/4 - \beta^2)}{4\cos^2(\pi x/2)},$$

and we have

$$\mathbb{J}^{\alpha,\beta}\Phi_k^{\alpha,\beta} = \Lambda_k^{\alpha,\beta}\Phi_k^{\alpha,\beta}, \quad k \geq 0, \quad \text{where} \quad \Lambda_k^{\alpha,\beta} = \pi^2 \left(k + \frac{\alpha + \beta + 1}{2} \right)^2.$$

Thus $\mathbb{J}^{\alpha,\beta}$ has a natural self-adjoint extension in $L^2(dx)$ given by

$$\tilde{\mathbb{J}}^{\alpha,\beta} f = \sum_{k=0}^{\infty} \Lambda_k^{\alpha,\beta} \langle f, \Phi_k^{\alpha,\beta} \rangle \Phi_k^{\alpha,\beta}$$

on the domain

$$\text{Dom } \tilde{\mathbb{J}}^{\alpha,\beta} = \left\{ f \in L^2(dx) : \sum_{k=0}^{\infty} |\Lambda_k^{\alpha,\beta} \langle f, \Phi_k^{\alpha,\beta} \rangle|^2 < \infty \right\}.$$

The semigroup generated by $-\tilde{\mathbb{J}}^{\alpha,\beta}$ in $L^2(dx)$ has the integral representation

$$\exp(-t\tilde{\mathbb{J}}^{\alpha,\beta})f(x) = \int_0^1 \mathbb{G}_t^{\alpha,\beta}(x, y) f(y) dy,$$

where the Jacobi heat kernel is defined by the oscillating series

$$\mathbb{G}_t^{\alpha,\beta}(x, y) = \sum_{k=0}^{\infty} \exp(-t\Lambda_k^{\alpha,\beta}) \Phi_k^{\alpha,\beta}(x) \Phi_k^{\alpha,\beta}(y).$$

Recently, qualitatively sharp estimates of $\mathbb{G}_t^{\alpha,\beta}(x, y)$ were obtained in [2] for $\alpha, \beta > -1$ and independently in [5], with completely different methods and under the restriction $\alpha, \beta \geq -1/2$. The result below is a consequence of [2, Theorem 7.2] (see also [5, Theorem A] and the beginning of [5, Section 4]), the relation between various Jacobi heat kernels [5, (3)] and a simple scaling argument.

Theorem 2.1 ([2, 5]). *Assume that $\alpha, \beta > -1$. Given any $T > 0$, there exist positive constants C, c_1 and c_2 , depending only on α, β and T , such that*

$$\begin{aligned} & \frac{1}{C} \left[\frac{xy}{t+xy} \right]^{\alpha+1/2} \left[\frac{(1-x)(1-y)}{t+(1-x)(1-y)} \right]^{\beta+1/2} \frac{1}{\sqrt{t}} \exp\left(-c_1 \frac{(x-y)^2}{t}\right) \\ & \leq \mathbb{G}_t^{\alpha,\beta}(x, y) \\ & \leq C \left[\frac{xy}{t+xy} \right]^{\alpha+1/2} \left[\frac{(1-x)(1-y)}{t+(1-x)(1-y)} \right]^{\beta+1/2} \frac{1}{\sqrt{t}} \exp\left(-c_2 \frac{(x-y)^2}{t}\right), \end{aligned}$$

uniformly in $x, y \in [0, 1]$ and $0 < t \leq T$, where the expressions in the lower and upper estimates are understood in the limiting sense whenever x or y is an endpoint of $[0, 1]$.

These estimates are crucial for our reasoning proving Theorem A.

3. PROOF OF THEOREM A

We first focus on showing the short time bounds, which constitute the main part of Theorem A. Our argument relies on relating the Fourier-Bessel heat kernel $\mathbb{K}_t^\nu(x, y)$ with the Jacobi heat kernel $\mathbb{G}_t^{\alpha, \beta}(x, y)$ and then transferring the estimates of Theorem 2.1. This will be achieved by proving that the generator of the Fourier-Bessel semigroup $\{\exp(-t\tilde{\mathbb{L}}^\nu)\}$ is a ‘slight’ perturbation of the Jacobi generator with the parameters of type $\alpha = \nu$ and $\beta = 1/2$. Then the desired relation will follow by the so-called Trotter product formula.

A close connection between the generators is suggested by a relation at the level of differential operators. Observe that \mathbb{L}^ν and $\mathbb{J}^{\nu, 1/2}$ differ only by a zero order term,

$$\mathbb{L}^\nu - \mathbb{J}^{\nu, 1/2} = \left(\frac{1}{4} - \nu^2\right) \left[\frac{\pi^2}{4 \sin^2 \frac{\pi x}{2}} - \frac{1}{x^2} \right] =: H^\nu(x).$$

Moreover, the difference function $H^\nu(x)$ is continuous and bounded on $[0, 1]$, with the value at $x = 0$ understood in the limiting sense. In the sequel we will use repeatedly, sometimes implicitly, the bound

$$|H^\nu(x)| \leq |H^\nu(1)| = \left| \frac{1}{4} - \nu^2 \right| \left(\frac{\pi^2}{4} - 1 \right), \quad x \in [0, 1].$$

This follows by analyzing the function in square brackets above. Indeed, its limit value at $x = 0$ is positive, and its derivative is positive in $(0, 1)$, as easily verified by means of the inequality

$$\sin^3 u > u^3 \cos u, \quad 0 < u < \frac{\pi}{2}.$$

The latter can readily be checked by using the estimates

$$\sin u > u - \frac{u^3}{6}, \quad \cos u < 1 - \frac{u^2}{2} + \frac{u^4}{24}, \quad u > 0,$$

coming from Taylor expansions of the two trigonometric functions.

An essential point of the whole method is the coincidence of domains, which is stated in the following.

Theorem 3.1. *For each $\nu > -1$,*

$$\text{Dom } \tilde{\mathbb{L}}^\nu = \text{Dom } \tilde{\mathbb{J}}^{\nu, 1/2}.$$

To prove this we will need a preparatory result.

Lemma 3.2. *Let $\nu > -1$. Then*

$$\langle \tilde{\mathbb{L}}^\nu \psi_n^\nu, \Phi_k^{\nu, 1/2} \rangle = \langle \psi_n^\nu, \mathbb{L}^\nu \Phi_k^{\nu, 1/2} \rangle, \quad n \geq 1, \quad k \geq 0,$$

where \mathbb{L}^ν is the differential operator given by (2).

Proof. We shall use the divergence form of \mathbb{L}^ν ,

$$\mathbb{L}^\nu f(x) = -x^{-\nu-1/2} D \left(x^{2\nu+1} D \left(x^{-\nu-1/2} f(x) \right) \right),$$

and integrate by parts. Denote

$$\mathcal{I} = \langle \tilde{\mathbb{L}}^\nu \psi_n^\nu, \Phi_k^{\nu, 1/2} \rangle = \int_0^1 \mathbb{L}^\nu \psi_n^\nu(x) \Phi_k^{\nu, 1/2}(x) dx.$$

Clearly, the integral here converges since $\mathbb{L}^\nu \psi_n^\nu$ and $\Phi_k^{\nu,1/2}$ are in $L^2(dx)$. We have

$$\begin{aligned}
\mathcal{I} &= - \int_0^1 x^{-\nu-1/2} D \left(x^{2\nu+1} D(x^{-\nu-1/2} \psi_n^\nu(x)) \right) \Phi_k^{\nu,1/2}(x) dx \\
&= -x^{\nu+1/2} D(x^{-\nu-1/2} \psi_n^\nu(x)) \Phi_k^{\nu,1/2}(x) \Big|_0^1 + \int_0^1 D(x^{-\nu-1/2} \psi_n^\nu(x)) D(x^{-\nu-1/2} \Phi_k^{\nu,1/2}(x)) x^{2\nu+1} dx \\
&= -x^{\nu+1/2} D(x^{-\nu-1/2} \psi_n^\nu(x)) \Phi_k^{\nu,1/2}(x) \Big|_0^1 + x^{\nu+1/2} \psi_n^\nu(x) D(x^{-\nu-1/2} \Phi_k^{\nu,1/2}(x)) \Big|_0^1 \\
&\quad - \int_0^1 \psi_n^\nu(x) x^{-\nu-1/2} D \left(x^{2\nu+1} D(x^{-\nu-1/2} \Phi_k^{\nu,1/2}(x)) \right) dx \\
&\equiv \mathcal{I}_1 + \mathcal{I}_2 + \mathcal{I}_3.
\end{aligned}$$

Since $\mathcal{I}_3 = \langle \psi_n^\nu, \mathbb{L}^\nu \Phi_k^{\nu,1/2} \rangle$, it remains to ensure that $\mathcal{I}_1 = \mathcal{I}_2 = 0$.

From the definition of $\Phi_k^{\alpha,\beta}$ it is clear that

$$\Phi_k^{\nu,1/2}(x) = \mathcal{O}(x^{\nu+1/2}), \quad x \rightarrow 0^+, \quad \text{and} \quad \Phi_k^{\nu,1/2}(x) = \mathcal{O}(1-x), \quad x \rightarrow 1^-.$$

The estimate $|\phi_n^\nu(x)| \leq C(1-x)$, see the proof of [4, Theorem 3.7], holds uniformly in $x \in (0, 1)$ and implies

$$\psi_n^\nu(x) = \mathcal{O}(x^{\nu+1/2}), \quad x \rightarrow 0^+, \quad \text{and} \quad \psi_n^\nu(x) = \mathcal{O}(1-x), \quad x \rightarrow 1^-.$$

Further, using $D(z^{-\nu} J_\nu(z)) = -z^{-\nu} J_{\nu+1}(z)$ (cf. [8, Chapter III, Section 3.2]), we see that

$$D(x^{-\nu-1/2} \psi_n^\nu(x)) = c x^{-\nu} J_{\nu+1}(\lambda_{n,\nu} x) = \begin{cases} \mathcal{O}(x), & x \rightarrow 0^+ \\ \mathcal{O}(1), & x \rightarrow 1^- \end{cases},$$

where c does not depend on x . Moreover, by the differentiation rule for Jacobi polynomials (cf. [7, (4.21.7)]),

$$\frac{d}{du} P_k^{\alpha,\beta}(u) = \frac{1}{2}(k + \alpha + \beta + 1) P_{k-1}^{\alpha+1,\beta+1}(u), \quad k \geq 0,$$

(here we use the convention that $P_{-1}^{\alpha,\beta} = 0$) it follows that

$$\begin{aligned}
D(x^{-\nu-1/2} \Phi_k^{\nu,1/2}(x)) &= c_1 \sin(\pi x) P_{k-1}^{\nu+1,3/2}(\cos(\pi x)) \left(\frac{\sin \frac{\pi x}{2}}{x} \right)^{\nu+1/2} \cos \frac{\pi x}{2} \\
&\quad + c_2 P_k^{\nu,1/2}(\cos(\pi x)) \left(\frac{\sin \frac{\pi x}{2}}{x} \right)^{\nu+1/2} \sin \frac{\pi x}{2} \\
&\quad + c_3 P_k^{\nu,1/2}(\cos(\pi x)) \frac{\frac{\pi x}{2} \cos \frac{\pi x}{2} - \sin \frac{\pi x}{2}}{x^2} \left(\frac{\sin \frac{\pi x}{2}}{x} \right)^{\nu-1/2} \cos \frac{\pi x}{2},
\end{aligned}$$

with c_1, c_2, c_3 independent of x . Therefore

$$D(x^{-\nu-1/2} \Phi_k^{\nu,1/2}(x)) = \begin{cases} \mathcal{O}(x), & x \rightarrow 0^+ \\ \mathcal{O}(1), & x \rightarrow 1^- \end{cases}.$$

Altogether, these facts imply $\mathcal{I}_1 = \mathcal{I}_2 = 0$. The lemma follows. \square

Proof of Theorem 3.1. We demonstrate that $\text{Dom } \tilde{\mathbb{J}}^{\nu,1/2} \subset \text{Dom } \tilde{\mathbb{L}}^\nu$. Then the other inclusion follows by self-adjointness of the two operators.

Assume that $f \in \text{Dom } \tilde{\mathbb{J}}^{\nu,1/2}$ and let

$$\mathcal{S} := \sum_{n=1}^{\infty} |\lambda_{n,\nu}^2 \langle f, \psi_n^\nu \rangle|^2.$$

We must show that \mathcal{S} is finite. We have

$$\mathcal{S} = \sum_{n=1}^{\infty} |\langle f, \tilde{\mathbb{L}}^\nu \psi_n^\nu \rangle|^2 = \sum_{n=1}^{\infty} \left| \sum_{k=0}^{\infty} \langle f, \Phi_k^{\nu,1/2} \rangle \langle \Phi_k^{\nu,1/2}, \tilde{\mathbb{L}}^\nu \psi_n^\nu \rangle \right|^2.$$

Applying now Lemma 3.2 we can write

$$\begin{aligned} \mathcal{S} &= \sum_{n=1}^{\infty} \left| \sum_{k=0}^{\infty} \langle f, \Phi_k^{\nu,1/2} \rangle \langle (\mathbb{J}^{\nu,1/2} + H^\nu) \Phi_k^{\nu,1/2}, \psi_n^\nu \rangle \right|^2 \\ &= \sum_{n=1}^{\infty} \left| \left\langle \sum_{k=0}^{\infty} \Lambda_k^{\nu,1/2} \langle f, \Phi_k^{\nu,1/2} \rangle \Phi_k^{\nu,1/2}, \psi_n^\nu \right\rangle + \left\langle \sum_{k=0}^{\infty} \langle f, \Phi_k^{\nu,1/2} \rangle \Phi_k^{\nu,1/2}, \psi_n^\nu H^\nu \right\rangle \right|^2 \\ &= \sum_{n=1}^{\infty} \left| \langle \tilde{\mathbb{J}}^{\nu,1/2} f, \psi_n^\nu \rangle + \langle f H^\nu, \psi_n^\nu \rangle \right|^2. \end{aligned}$$

This together with Parseval's identity implies

$$\mathcal{S} \leq 2 \|\tilde{\mathbb{J}}^{\nu,1/2} f\|_{L^2(dx)}^2 + 2 \|f H^\nu\|_{L^2(dx)}^2 < \infty$$

and consequently $f \in \text{Dom } \tilde{\mathbb{L}}^\nu$. \square

We now recall the Trotter product formula, see [6, Chapter VIII, Section 8]. Let A and B be (possibly unbounded) self-adjoint operators on a Hilbert space \mathcal{H} . If $A + B$ is essentially self-adjoint on $\text{Dom } A \cap \text{Dom } B$, and A and B are bounded from below, then

$$\exp(-t(A+B))h = \lim_{m \rightarrow \infty} \left(\exp(-tA/m) \exp(-tB/m) \right)^m h, \quad h \in \mathcal{H}, \quad t \geq 0.$$

Specifying this general situation to our context we take $\mathcal{H} = L^2(dx)$, $A = \tilde{\mathbb{J}}^{\nu,1/2}$ and choose B to be the multiplication operator by H^ν . Then, in view of Theorem 3.1, $A + B = \tilde{\mathbb{L}}^\nu$. Moreover, the Trotter product formula applies and since $|H^\nu(x)| \leq |H^\nu(1)|$, $x \in [0, 1]$, it implies

$$e^{-t|H^\nu(1)|} \exp(-t\tilde{\mathbb{J}}^{\nu,1/2})f \leq \exp(-t\tilde{\mathbb{L}}^\nu)f \leq e^{t|H^\nu(1)|} \exp(-t\tilde{\mathbb{J}}^{\nu,1/2})f, \quad 0 \leq f \in L^2(dx),$$

for each $t \geq 0$. Taking now into account that the Jacobi and the Fourier-Bessel heat kernels are continuous functions of their arguments (see [5, Section 2], [4, Section 2]), we infer that

$$e^{-t|H^\nu(1)|} \mathbb{G}_t^{\nu,1/2}(x, y) \leq \mathbb{K}_t^\nu(x, y) \leq e^{t|H^\nu(1)|} \mathbb{G}_t^{\nu,1/2}(x, y), \quad x, y \in [0, 1], \quad t > 0.$$

This combined with Theorem 2.1 and (3) justifies the short time bounds stated in Theorem A.

The long time estimates were proved in [4, Theorem 3.7], under the assumption that the threshold T is sufficiently large. However, given any $0 < T_0 < T$, the already proved short time bounds imply existence of a constant $C > 0$ such that

$$C^{-1}(1-x)(1-y) \leq G_t^\nu(x, y) \leq C(1-x)(1-y), \quad x, y \in [0, 1], \quad T_0 \leq t \leq T,$$

and the desired conclusion follows.

The proof of Theorem A is complete.

Remark 3.3. *A slightly more detailed analysis reveals that for $\nu \in [-1/2, 1/2]$*

$$e^{-tH^\nu(1)}\mathbb{G}_t^{\nu,1/2}(x,y) \leq \mathbb{K}_t^\nu(x,y) \leq e^{-tH^\nu(0)}\mathbb{G}_t^{\nu,1/2}(x,y) \leq \mathbb{G}_t^{\nu,1/2}(x,y), \quad x, y \in [0,1], \quad t > 0,$$

and for $\nu \notin [-1/2, 1/2]$

$$\mathbb{G}_t^{\nu,1/2}(x,y) \leq e^{-tH^\nu(0)}\mathbb{G}_t^{\nu,1/2}(x,y) \leq \mathbb{K}_t^\nu(x,y) \leq e^{-tH^\nu(1)}\mathbb{G}_t^{\nu,1/2}(x,y), \quad x, y \in [0,1], \quad t > 0.$$

Notice that $\mathbb{K}_t^{\pm 1/2}(x,y) = \mathbb{G}_t^{\pm 1/2,1/2}(x,y)$.

REFERENCES

- [1] A. N. Borodin, P. Salminen, *Handbook of Brownian motion: facts and formulae*, 2nd ed., Birkhäuser Verlag, Basel-Boston-Berlin, 2002.
- [2] T. Coulhon, G. Kerkycharian, P. Petrushev, *Heat kernel generated frames in the setting of Dirichlet spaces*, J. Fourier Anal. Appl. 18 (2012), 995–1066.
- [3] S. Karlin, J. McGregor, *Classical diffusion processes and total positivity*, J. Math. Anal. Appl. 1 (1960), 163–183.
- [4] A. Nowak, L. Roncal, *On sharp heat and subordinated kernel estimates in the Fourier-Bessel setting*, Rocky Mountain J. Math. (to appear). [arXiv:1111.5700](https://arxiv.org/abs/1111.5700)
- [5] A. Nowak, P. Sjögren, *Sharp estimates of the Jacobi heat kernel*, preprint 2011. [arXiv:1111.3145](https://arxiv.org/abs/1111.3145)
- [6] M. Reed, B. Simon, *Methods of modern mathematical physics. I. Functional analysis*, Second edition. Academic Press, Inc., New York, 1980.
- [7] G. Szegő, *Orthogonal polynomials*, Fourth Edition, Amer. Math. Soc. Colloq. Publ. 23, Amer. Math. Soc., Providence, R. I., 1975.
- [8] G. N. Watson, *A treatise on the theory of Bessel functions*, Cambridge University Press, Cambridge, 1966.

INSTYTUT MATEMATYCZNY, POLSKA AKADEMIA NAUK, ŚNIADECKICH 8, 00-956 WARSZAWA, POLAND
E-mail address: adam.nowak@impan.pl

DEPARTAMENTO DE MATEMÁTICAS Y COMPUTACIÓN, UNIVERSIDAD DE LA RIOJA, 26004 LOGROÑO, SPAIN
E-mail address: luz.roncal@unirioja.es