The warm, the excited, and the molecular gas: GRB 121024A shining through its star-forming galaxy. \star

M. Friis¹, A. De Cia², T. Krühler^{3,4}, J. P. U. Fynbo⁴, C. Ledoux³, P. M. Vreeswijk², D. J. Watson⁴, D. Malesani⁴, J. Gorosabel^{5,6,7}†, R. L. C. Starling⁸, P. Jakobsson¹, K. Varela⁹, K. Wiersema⁸, A. P. Drachmann⁴, A. Trotter^{10,11}, C. C. Thöne⁵, A. de Ugarte Postigo^{5,4}, V. D'Elia^{12,13}, J. Elliott⁹, M. Maturi¹⁴, P. Goldoni¹⁵, J. Greiner⁹, J. Haislip¹⁰, L. Kaper¹⁶, F. Knust⁹, A. LaCluyze¹⁰, B. Milvang-Jensen⁴, D. Reichart¹⁰, S. Schulze^{17,18}, V. Sudilovsky⁹, N. Tanvir⁸, S. D. Vergani¹⁹ †Deceased ¹ Centre for Astrophysics and Cosmology, Science Institute, University of Iceland, Dunhagi 5, 107 Reykjavík, Iceland

 2 Department of Particle Physics and Astrophysics, Faculty of Physics, Weizmann Institute of Science, 76100, Rehovot, Israel

 3 European Southern Observatory, Alonso de Córdova 3107, Casilla 19001, Santiago 19, Chile

- ⁴Dark Cosmology Centre, Niels Bohr Institute, University of Copenhagen, Juliane Maries Vej 30, 2100 Copenhagen, Denmark
- 5 Instituto de Astrofísica de Andalucía (IAA-CSIC), Glorieta de la Astronomía s/n, 18008 Granada, Spain
- ⁶Unidad Asociada Grupo Ciencia Planetarias UPV/EHU-IAA/CSIC, Departamento de F´ısica Aplicada I, E.T.S. Ingenieria,
- Universidad del Pais Vasco UPV/EHU, Alameda de Urquijo s/n, 48013 Bilbao, Spain
- 7 Ikerbasque, Basque Foundation for Science, Alameda de Urquijo 36-5, 48008 Bilbao, Spain
- ⁸Department of Physics and Astronomy, University of Leicester, University Road, Leicester LE1 7RH, UK
- 9 Max-Planck-Institut für extraterrestrische Physik, Giessenbachstraße 1, 85748 Garching, Germany
- 10 Department of Physics and Astronomy, University of North Carolina at Chapel Hill, Campus Box 3255, Chapel Hill, NC 27599, USA
- 11 Department of Physics, NC A&T State University, 1601 E. Market St, Greensboro, NC 27411, USA
- 12 INAF/Rome Astronomical Observatory, via Frascati 33, I-00040 Monteporzio Catone (Roma), Italy
- ¹³ASI-Science Data Center, Via del Politecnico snc, I-00133 Rome, Italy
- 14 Zentrum für Astronomie der Universität Heidelberg, Institut für Theoretische Astrophysik, Philosophenweg 12, 69120 Heidelberg, Germany

 15 APC, Astroparticule et Cosmologie, Universite Paris Diderot, CNRS/IN2P3, CEA/Irfu, Observatoire de Paris, Sorbonne Paris Cité,

10 Rue Alice Domon et Leonie Duquet, F-75205 Paris, Cedex 13, France

¹⁶ Anton Pannekoek Institute for Astronomy, University of Amsterdam, Science Park 904, 1098 XH, Amsterdam, The Netherlands ¹⁷Millennium Institute of Astrophysics, Casilla 306, Santiago 22, Chile

 18 Instituto de Astrofísica, Facultad de Física, Pontifícia Universidad Catolica de Chile, Casilla 306, Santiago 22, Chile

¹⁹Laboratoire GEPI, Observatoire de Paris, CNRS-UMR8111, Université Paris Diderot, 5 place Jules Janssen, F-92195 Meudon, France

ABSTRACT

We present the first reported case of the simultaneous metallicity determination of a gamma-ray burst (GRB) host galaxy, from both afterglow absorption lines as well as strong emission-line diagnostics. Using spectroscopic and imaging observations of the afterglow and host of the long-duration Swift GRB 121024A at $z = 2.30$, we give one of the most complete views of a GRB host/environment to date. We observe a strong damped $Ly\alpha$ absorber (DLA) with a hydrogen column density of $\log N(\text{H I}) = 21.88 \pm 0.10, \text{H}_2$ absorption in the Lyman-Werner bands (molecular fraction of $\log(f) \approx -1.4$; fourth solid detection of molecular hydrogen in a GRB-DLA), the nebular emission lines H α , H β , [O II], [O III] and [N II], as well as metal absorption lines. We find a GRB host galaxy that is highly star-forming $(SFR \sim 40 M_{\odot} \text{ yr}^{-1})$, with a dust-corrected metallicity along the line of sight of $\text{[Zn/H]}_{\text{corr}} = -0.6 \pm 0.2$ $([O/H] \sim -0.3$ from emission lines), and a depletion factor $[\text{Zn/Fe}] = 0.85 \pm 0.04$. The molecular gas is separated by 400 km s^{-1} (and 1-3 kpc) from the gas that is photoexcited by the GRB. This implies a fairly massive host, in agreement with the derived stellar mass of $log(M_*/M_{\odot}) = 9.9_{-0.3}^{+0.2}$. We dissect the host galaxy by characterising its molecular component, the excited gas, and the line-emitting star-forming regions. The extinction curve for the line of sight is found to be unusually flat $(R_V \sim 15)$. We discuss the possibility of an anomalous grain size distributions. We furthermore discuss the different metallicity determinations from both absorption and emission lines, which gives consistent results for the line of sight to GRB 121024A.

Key words: Galaxies: abundances – gamma-ray burst: individual: GRB 121024A

1 INTRODUCTION

The study of gamma-ray burst (GRB) afterglows has proven to be a powerful tool for detailed studies of the interstellar medium (ISM) of star-forming galaxies, out to high redshifts (e.g. [Vreeswijk et al. 2004](#page-17-0); [Prochaska et al. 2007;](#page-17-1) [Ledoux et al. 2009;](#page-16-0) [Sparre et al. 2014](#page-17-2)). With quickly fading emission spanning the entire electromagnetic spectrum, GRB afterglows offer a unique opportunity to probe the surrounding environment. The intrinsic spectrum of the afterglow is well fitted with simple power-law segments, so the imprints of the intergalactic medium (IGM) as well as the ISM surrounding the burst are relatively easy to distinguish from the afterglow in the observed spectrum. Moreover, with absorption and emission-line analysis it is possible to determineparameters such as H i column density, metallicity, dust depletion, star-formation rate (SFR) and kinematics of the GRB host galaxy.

Metallicity is a fundamental parameter for characterising a galaxy and it holds important information about its history. Metallicity might also play a crucial role in the GRB production mechanism. For GRB hosts, the metallicity is measured either from hydrogen and metal absorption lines,

or by using diagnostics based on the fluxes of strong nebular emission lines, calibrated in the local Universe. Different calibrations are in use leading to some discrepancy (e.g. [Kudritzki et al. 2012](#page-16-1)), and the different diagnostics have their strengths and weaknesses (e.g. less sensitive to reddening, multiple solutions, or more sensitive at high metallicities). The absorption lines probe the ISM along the line of sight, while the nebular line diagnostics determine the integrated metallicity of the H ii regions of the host. For GRB damped Ly α absorbers (GRB-DLAs, $N(H I) > 2 \times 10^{20}$ cm⁻² [Wolfe et al. 2005](#page-17-3)), a direct comparison of metallicity from the two methods is interesting because it can either provide a test of the strong-line methods or alternatively allow a measurement of a possible offset in abundances in H ii regions and in the ISM. So far, this comparison has only been carried out for a few galaxy counterparts of DLAs found in the line of sight of background QSOs (QSO-DLAs, e.g. [Bowen et al. 2005](#page-16-2); Péroux et al. 2012; [Noterdaeme et al.](#page-17-5) [2012](#page-17-5); [Fynbo et al. 2013](#page-16-3); [Jorgenson & Wolfe 2014](#page-16-4)). To our knowledge, a comparison for GRB-DLAs has not been reported before. For both emission and absorption measurements to be feasible with current instrumentation, the observed host needs to be highly star-forming, to have strong nebular lines, and at the same time be at a redshift high enough for the $Ly\alpha$ transition to be observed (at redshifts higher than $z \approx 1.5$ the Ly α absorption line is redshifted into the atmospheric transmission window). GRB 121024A is a $z = 2.30$ burst hosted by a highly star-forming galaxy. We measure abundances of the GRB host galaxy in absorption and compare them with the metallicity determined by strong-line diagnostics using observed nebular lines from [O II], [O III], [N II] and the Balmer emission lines.

Apart from the absorption features from metal lines, we also detect the Lyman-Werner bands of molecular hydrogen. Molecular hydrogen is hard to detect in absorption, because

[⋆] Based on observations carried out under prog. ID 090.A-0088(B) with the X-shooter spectrograph installed at the Cassegrain focus of the Very Large Telescope (VLT), Unit 2 – Kueyen, operated by the European Southern Observatory (ESO) on Cerro Paranal, Chile. Also used are observations made with the Nordic Optical Telescope, operated by the Nordic Optical Telescope Scientific Association, and the Gran Telescopio Canarias (program GTC67-13B), both at the Observatorio del Roque de los Muchachos, La Palma, Spain, of the Instituto de Astrofisica de Canarias. HAWK-I imaging used is part of the program 092.A-0076(B). This work made use of data supplied by the UK Swift Science Data Centre at the University of Leicester.

it requires high S/N and mid-high resolution. As long duration GRBs $(t_{obs} > 2 s)$ are thought to be associated with the death of massive stars (e.g. [Hjorth et al. 2003;](#page-16-5) [Stanek et al.](#page-17-6) [2003](#page-17-6); [Sparre et al. 2011](#page-17-7); [Cano 2013](#page-16-6); [Schulze et al. 2014](#page-17-8)), they are expected to be found near regions of active star formation, and hence molecular clouds. In spite of this, there are very few detections of molecular absorption towards GRBs (see e. g [Tumlinson et al. 2007](#page-17-9)). [Ledoux et al. \(2009](#page-16-0)) found that this is likely due to the low metallicities found in the systems observed with high resolution spectrographs $(R = \lambda/\Delta\lambda \gtrsim 40000)$. Typically, mid/high-resolution spectroscopy at a sufficient S/N is only possible for the brighter sources. As is the case for QSO-DLAs, lines of sight with H² detections will preferentially be metal-rich and dusty. The observed spectra are therefore UV-faint and difficult to observe (GRB 080607 is a striking exception, where observations were possible thanks to its extraordinarily intrinsic luminosity and rapid spectroscopy, see [Prochaska et al. 2009](#page-17-10)). Now with X-shooter [\(Vernet et al. 2011](#page-17-11)) on the Very Large Telescope (VLT) we are starting to secure spectra with sufficient resolution to detect H_2 for fainter systems resulting in additional detections (Krühler et al. 2013; [D'Elia et al.](#page-16-8) [2014](#page-16-8)).

Throughout this paper we adopt a flat ΛCDM cosmology with $H_0 = 71 \text{ km s}^{-1}$ and $\Omega_M = 0.27$, and report 1 σ errors (3 σ limits), unless otherwise indicated. Reference solar abundances are taken from [Asplund et al. \(2009\)](#page-16-9), where either photospheric or meteoritic values (or their average) are chosen according to the recommendations of [Lodders et al.](#page-17-12) [\(2009](#page-17-12)). Column densities are in cm[−]² . In Sect. [2](#page-2-0) we describe the data and data reduction used in this paper, in Sect. [3](#page-3-0) we present the data analysis and results, which are then discussed in Sect. [4.](#page-13-0)

2 OBSERVATIONS AND DATA REDUCTION

On 2012 October 24 at 02:56:12 UT the Burst Alert Telescope (BAT, [Barthelmy et al. 2005](#page-16-10)) onboard the Swift satellite [\(Gehrels et al. 2004](#page-16-11)) triggered on GRB 121024A. The X-Ray Telescope (XRT) started observing the field at 02:57:45 UT, 93 seconds after the BAT trigger. About one minute after the trigger, Skynet observed the field with the PROMPT telescopes located at CTIO in Chile and the 16" Dolomites Astronomical Observatory telescope (DAO) in Italy [\(Reichart et al. 2005\)](#page-17-13) in filters g', r', i', z' and BRi . Approximately 1.8 hours later, spectroscopic afterglow measurements in the wavelength range of 3000\AA to 25000\AA were acquired (at 04:45 UT), using the cross-dispersed, echelle spectrograph X-shooter [\(Vernet et al. 2011](#page-17-11)) mounted at ESO's VLT. Then at 05:53 UT, 3 hours after the burst, the Gamma-Ray burst Optical/NIR Detector (GROND, [Greiner et al. 2007,](#page-16-12) [2008](#page-16-13)) mounted on the 2.2 m MPG/ESO telescope at La Silla Observatory (Chile), performed followup optical/NIR photometry simultaneously in g', r', i', z' and JHK. About one year later (2013 November 07), VLT/HAWK-I imaging of the host was acquired in the J $(07:02:13 \text{ UT})$ and $K (06:06:47 \text{ UT})$ band. To supplement these, B, R and i band imaging was obtained at the Nordic Optical Telescope (NOT) at 2014 January 06 (i) and February 10 (R) and 19 (B) . Gran Telescopio Canarias (GTC) observations in the q and z band were optioned on 2014

Table 1. X-shooter observations

$t_{\rm obs}$ (UT) ^a	$(\min)^b$ $t_{\rm GRB}$	$t_{\rm exp.}$ (s)	Mean Airmass	Seeing
04:47:01	116	600	1.23	$0\frac{\prime}{6} - 0\frac{\prime}{7}$
04:58:35	127	600	1.19	$0\frac{7}{6} - 0\frac{7}{7}$
05:10:12	139	600	1.16	$0\frac{7}{6} - 0\frac{7}{7}$
05:21:46	151	600	1.13	$0\frac{7}{6} - 0\frac{7}{7}$

^a Start time of observation on October 24, 2012.

 b Mid-exposure time in seconds since GRB trigger.</sup>

Table 2. X-shooter resolution

Arm	Slit	$R = \lambda / \Delta \lambda$
NIR.	$0''\!\!.9$	6800
VIS	0''9	13000
UVB	$1\rlap{.}^{\prime\prime}$ (0	7100

February 28. For an overview see Tables [1,](#page-2-1) [2](#page-2-2) and [3.](#page-3-1) Linear and circular polarisation measurements for the optical afterglow of GRB 121024A have been reported in [Wiersema et al.](#page-17-14) [\(2014](#page-17-14)).

2.1 X-shooter NIR/Optical/UV Spectroscopy

The X-shooter observation consists of four nodded exposures with exposure times of 600 s each, taken simultaneously by the ultraviolet/blue (UVB), visible (VIS) and near-infrared (NIR) arms. The average airmass was 1.18 with a median seeing of ∼0". The spectroscopy was performed with slitwidths of 1".0, 0".9 and 0".9 in the UVB, VIS and NIR arms, respectively. The resolving power $R = \lambda/\Delta\lambda$ is determined from telluric lines to be $R = 13000$ for the VIS arm. This is better than the nominal value due to the very good seeing. Following [Fynbo et al. \(2011](#page-16-14)) we then infer $R = 7100$ and $R = 6800$ for the UVB and NIR arms, see Table [2](#page-2-2) for an overview.

X-shooter data were reduced with the ESO/X-shooter pipeline version 2.2.0 [\(Goldoni et al. 2006](#page-16-15)), rectifying the data on an output grid with a dispersion of 0.15 Å/pixel in the UVB, $0.13 \text{ Å}/\text{pixel}$ in the VIS and $0.5 \text{ Å}/\text{pixel}$ in the NIR arm. The wavelength solution was obtained against arc-lamp frames in each arm. Flux-calibration was performed against the spectrophotometric standard GD71 observed during the same night. We further correct the flux-calibrated spectra for slit-losses by integrating over filter curves from GROND photometry shifted to X-shooter observation times (assuming a slope of $\alpha = 0.8$). For the UVB arm, only the g' band photometry is available, which covers the DLA (see Sect. [3.1\)](#page-3-2), making this calibration less secure. Wavelengths are plotted in vacuum and corrected for heliocentric motion.

2.2 NOT, GTC and VLT/HAWK-I imaging

To derive physical parameters of the host of GRB 121024A via stellar population synthesis modelling, we obtained latetime photometry from VLT/HAWK-I, NOT and GTC. Exposure times and seeing can be found in Table [3.](#page-3-1)

J and K band images were observed with HAWK-I on

^a Time since the GRB trigger (observer's time frame).

For the afterglow measurements time is given in hours, while for the host galaxy, it is shown in days.

the Yepun (VLT-UT4) telescope at the ESO Paranal Observatory in Chile. HAWK-I is a near-infrared imager with a pixel scale of $0''$ 106/pix and a total field of view of $7'$: $5 \times 7'$: 5 . B, R and i images were obtained with the ALFOSC optical camera on the NOT. The photometric calibration was carried out by observing the standard star GD71 at a similar airmass to the GRB field. g' and z' -band host galaxy images were taken with the 10.4m GTC. The images were acquired with the OSIRIS instrument which provides an unvignetted field of view of $7/8 \times 7/8$ and a pixel scale of 0".25/pix [\(Cepa et al. 2000](#page-16-16)). Images were taken following a dithering pattern. The z'-band images were defringed by subtracting an interference pattern which was constructed based on the dithered individual frames. The photometric calibration was carried out by observing the standard star SA95-193 [\(Smith et al. 2002\)](#page-17-15). NOT and GTC are located at the observatory of Roque de los Muchachos, La Palma, Spain.

All images were dark-subtracted and flat-fielded using IRAF standard routines.

2.3 GROND and Skynet Photometry

GROND data was reduced using standard IRAF tasks [\(Tody 1997](#page-17-16); Krühler et al. 2008). The afterglow image was fitted using a general point spread function (PSF) model obtained from bright stars in the field. The optical images in g', r', i', z' were calibrated against standard stars in the SDSS catalogue, with an accuracy of ± 0.03 mag. The NIR magnitudes were calibrated using stars of the 2MASS catalogue, with an accuracy of ± 0.05 mag. Skynet obtained images of the field of GRB 121024A on 2012 October 24-25 with four $16''$ telescopes of the PROMPT array at CTIO, Chile, and the 16′′ DAO in Italy. Exposures ranging from 5 to 160 s were obtained in the $BVRI$ (PROMPT) and g', r', i' (DAO) bands, starting at 02:57:07 UT $(t = 55 s \text{ since the})$ GRB trigger) and continuing until $t = 7.3$ h on the first night, and continuing from $t = 20.7 - 25.5$ h on the second night. Bias subtraction and flat-fielding were performed via Skynet's automated pipeline. Post-processing occurred in Skynet's guided analysis pipeline, using both custom and IRAF-derived algorithms. Differential aperture photometry was performed on single and stacked images, with effective exposure times of 5s to 20 min on the first night, and up to ∼4 h on the second night. Photometry was calibrated to the catalogued B, V, g', r', i' magnitudes of five APASS DR7 stars in the field, with g', r', i' magnitudes transformed to RI using transformations obtained from prior observations of Landoldt stars (Henden, A. et al., in preparation). The Skynet magnitudes can be seen in Appendix [A.](#page-18-0)

3 ANALYSIS AND RESULTS

3.1 Absorption Lines

The most prominent absorption feature is the $Ly\alpha$ line. We plot the spectral region in Fig. [1.](#page-5-0) Over-plotted is a Voigt-profile fit to the strong $Ly\alpha$ absorption line yielding $\log N(H I) = 21.88 \pm 0.10$. The error takes into account the noise in the spectrum, the error on the continuum placement and background subtraction at the core of the saturated lines. Table [4](#page-4-0) shows the metal absorption lines identified in the spectrum. To determine the ionic column densities of the metals, we model the identified absorption lines with a number of Voigt-profile components, as follows. We use the Voigt-profile fitting software $VPFIT^1$ $VPFIT^1$ version 9.5 to model the absorption lines. We first normalise the spectrum around each line, fitting featureless regions with zero- or first-order polynomials. To remove the contribution of atmospheric absorption lines from our Voigt-profile fit, we compare the observed spectra to a synthetic telluric spectrum. This telluric spectrum was created following [Smette et al. \(2010](#page-17-17)) as described by [De Cia et al. \(2012](#page-16-18)) and assuming a precipitable

GRB 121024A 5

Table 4. Ionic column densities of the individual components of the line profile. The transitions used to derive column densities are reported in the second column. Transitions marked in bold are those unblended and unsaturated lines that we use to determine the lineprofile decomposition. The results for the ground state and excited levels are listed in the top and bottom part of the table, respectively. Velocities given are with respect to the [O III] λ 5007 line (z = 2.3015). (b) / (s) indicate that the line is blended / saturated. The error on the redshifts of each component is 0.0001

Component	Transition	\mathbf{a}	$\mathbf b$	\mathbf{C}	$\mathbf d$	\mathbf{e}
\boldsymbol{z}		2.2981	2.2989	2.3017	2.3023	2.3026
$b~(\rm km\,s^{-1})$		26	21	20	22	35
$v~(\rm km~s^{-1})$		-264	-191	64	118	145
log(N)						
MgI	λ 1827, λ 2026(b)	13.97 ± 0.05	13.57 ± 0.05	$<13.4\,$	13.57 ± 0.07	$<13.4\,$
Al III	λ 1854(s), λ 1862(s)					
Si _{II}	λ 1808(s)					
$\mathrm{S}\:\textsc{ii}$	λ 1253(s)					
Ca _{II}	λ 3934, λ 3969	13.25 ± 0.16^a	12.50 ± 0.16^b	12.20 ± 0.16	11.90 ± 0.16	11.20 ± 0.16
Cr II	λ 2056, λ 2062(b), λ 2066	13.47 ± 0.05	13.48 ± 0.05	13.39 ± 0.05	13.67 ± 0.05	13.34 ± 0.09
MnII	λ 2576, λ 2594, λ 2606	13.15 ± 0.05	13.07 ± 0.05	12.71 ± 0.05	13.21 ± 0.05	12.93 ± 0.05
Fe II	λ 1611, λ 2260, λ 2249	15.15 ± 0.05	15.09 ± 0.05	14.81 ± 0.05	15.27 ± 0.05	15.12 ± 0.05
Ni II	λ 1345, λ 1454, λ 1467.3, λ 1467.8, λ 1709	13.91 ± 0.10	13.88 ± 0.10	13.95 ± 0.09	14.17 ± 0.10	13.73 ± 0.29
Zn II	λ 2026(b), λ 2062(b)	13.14 ± 0.05	13.05 ± 0.05	12.50 ± 0.08	13.40 ± 0.05	12.19 ± 0.40
Component		α	β			
\boldsymbol{z}		2.2981	2.2989			
$b~(\rm km\,s^{-1})$		28	30			
$v~(\rm km~s^{-1})$		-264	-191			
log(N)						
$\rm Fe\,\textsc{ii}^*$	λ 2389, λ 2396(b)	13.25 ± 0.05	13.16 ± 0.05			
Fe II**	λ 2396(b), λ 2405(b), λ 2607	12.92 ± 0.05	12.80 ± 0.05			
Fe $II***$	λ 2405(b), λ 2407, λ 2411(b)	12.63 ± 0.05	12.58 ± 0.07			
Fe $II***$	λ 2411(b), λ 2414, λ 2622	12.53 ± 0.06	12.61 ± 0.05			
$Fe II*****$	λ 1559, λ 2360	13.95 ± 0.08	13.68 ± 0.13			
$NiII**$	λ 2166, λ 2217, λ 2223	13.43 ± 0.05	13.47 ± 0.05			
$Si II*$	λ 1309, λ 1533, λ 1816 ^e	14.98 ± 0.11^c	14.39 ± 0.05^d			

^a redshift: 2.2979, *b*-value: 30 km s^{-1} , see main text

^b redshift: 2.2989, b-value: 23 km s^{-1} , -

^c redshift: 2.2979, *b*-value: 20 km s^{-1} , –

d redshift: 2.2987, b-value: 30 km s^{-1} , -

^e The column density of the α component of Si_{II}* has been determined solely from the λ 1816 line.

Table 5. Total column densities (summed among individual velocity components and including excited levels) and abundances with respect to H and Fe.

Ion	$\log(N/\text{cm}^{-2})_{\text{tot}}$	$\log(N/\text{cm}^{-2})_{\text{a}+\text{b}}$	$\log(N/\text{cm}^{-2})_{c+d+e}$	$[X/H]_{\rm tot}$	[X/Fe]	$[X/Fe]_{a+b}$	$[X/Fe]_{c+d+e}$
H _I	21.88 ± 0.10						
MgI	< 14.31	14.11 ± 0.03	< 13.86				
Al III	>14.11						
Si II	>16.35			>-1.0	> 0.53		
$\mathrm{S}\,\mathrm{II}$	> 15.90			>-1.1	> 0.46		
Ca _{II}	13.37 ± 0.12	13.32 ± 0.13^a	12.40 ± 0.12	-2.9 ± 0.2	-1.29 ± 0.13	-0.97 ± 0.14^a	-2.02 ± 0.11
Cr _{II}	14.18 ± 0.03	13.78 ± 0.04	13.97 ± 0.03	-1.3 ± 0.1	$0.22 + 0.05$	0.18 ± 0.05	0.24 ± 0.04
MnII	13.74 ± 0.03	13.41 ± 0.04	13.47 ± 0.03	-1.6 ± 0.1	$-0.01 + 0.05$	0.03 ± 0.05	-0.04 ± 0.04
Fe II	15.82 ± 0.05	15.45 ± 0.05	15.58 ± 0.03	-1.6 ± 0.1			
Ni II	14.70 ± 0.06	14.33 ± 0.05	14.47 ± 0.06	-1.4 ± 0.1	0.17 ± 0.08	0.02 ± 0.08	0.16 ± 0.06
Zn II	13.74 ± 0.03	13.40 ± 0.03	13.47 ± 0.04	-0.7 ± 0.1	0.85 ± 0.06	0.88 ± 0.05	0.83 ± 0.05

 a Different a and b broadening parameter and redshift for CaII, see Sect. [3.1](#page-3-2)

Figure 1. The UVB spectrum centred on the damped $Ly\alpha$ absorption line at the GRB host galaxy redshift. For clarity purposes, the spectrum has been smoothed with a median filter with a sliding window width of 3 pixels. A neutral hydrogen column density fit $(\log N(H I) = 21.88 \pm 0.10)$ to the damped Ly α line is shown with a solid line (red), while the 1σ errors are shown with the shaded area (also red). In blue is shown the velocity range of the metal absorption lines. The dashed line shows the continuum placement, while the grey line near the bottom shows the error spectrum.

water-vapour column of 2.5 mm. We systematically reject from the fit the spectral regions affected by telluric features at a level of > 1 per cent^{[2](#page-5-1)}. None of the absorption lines that we include are severely affected by telluric lines. The resulting column densities are listed in Tables [4](#page-4-0) and [5](#page-4-1) for lines arising from ground-state and excited levels, respectively. We report formal $1-\sigma$ errors from the Voigt-profile fitting. We note that these do not include the uncertainty on the continuum normalisation, which can be dominant for weak lines (see e.g. [De Cia et al. 2012](#page-16-18)). We hence adopt a minimum error of 0.05 dex to account for this uncertainty. The error on the redshifts of each component is 0.0001. The Voigt-profile fits to the metal lines are shown in Figs. [2](#page-6-0) and [3.](#page-7-0)

The fit to the absorption lines from ground-state levels is composed of five components (a-e). We consider the redshift of the [O III] λ 5007 emission-line centroid $z = 2.3015$, as the reference zero-velocity. Components 'a' through 'e' are shifted -264 , -191 , 64, 118 and 145 km s⁻¹, respectively. Given the resolution of the instrument of 23 km s^{-1} (VIS arm), the individual components are blended, and therefore the profile decomposition is not unequivocal. However, regardless of the properties (and numbers) of the individual components, they are clearly divided into two well separated groups: $a+b$ and $c+d+e$. When forcing more components to the fit of each group, the resultant total column density are consistent with the previous estimate for each of the two groups. We stress that the resultant b-values are not physical, but likely a combination of smaller unresolved components. First we determine redshift z and broadening parameter b (purely turbulent broadening) of the individual components of the line profile, by considering only a mastersample of unblended and unsaturated lines (shown in bold

in Table [4\)](#page-4-0), with b and z tied among transitions of different ions. Values for z and b were then frozen for the rest of the absorption lines, and the column densities were fitted. We report $3-\sigma$ lower and upper limits for the saturated and undetected components, respectively. For the saturated lines Al III, Si II and S II we do not report column densities from the Voigt-profile fit, but instead from the measured equivalent widths (EWs), converted to column densities assuming a linear regime. For these, we only report the total column density for all the components together.

At the H_I column density that we observe, we expect most elements to be predominantly in their singly ionised state [\(Wolfe et al. 2005](#page-17-3)). We hence expect much of the Mg to be in Mg ii (for this reason we do not report the abun-dance of Mg_I in Table [5\)](#page-4-1). Ca_{II} seems to have a different velocity composition than the rest of the lines. One possibility is that Ca ii may extend to a slightly different gas phase, as its ionisation potential is the lowest among the observed lines (less than $1 \text{ Ryd} = 13.6 \text{ eV}$). Alternatively, since the Ca_{II} lines are located in the NIR arm, a small shift in the wavelength solution with respect to the VIS arm could cause the observed difference. However, a positive comparison between the observed and synthetic telluric lines rules out any shift in the wavelength calibration. We have allowed z and b to have different values for the two Ca ii lines. This resulted in a slightly different $a+b$ component, but the same $c+d+e$ component as for the rest of the sample.

The fine-structure lines show a different velocity profile composed only of two components, α and β , see Table [4.](#page-4-0) The redshift of α and β are the same as for component a and b found for the resonance lines (but different broadening parameters). Remarkably, no fine-structure lines are detected at the position of components $c+d+e$. The Si II^* lines are poorly fitted when tied together with the rest of the fine-structure lines, so we allow their z and b values to vary freely. These components are then referred to as γ and δ, which are quite similar to components α and β, respec-tively, see Fig. [2.](#page-6-0) The column density for component γ of the stronger $Si II^*$ line appears strongly saturated, so only the λ1816 line has been used to determine the column density in this component.

The total ionic column densities (summed over individual components and including excited levels when necessary) are given in Table [5.](#page-4-1) We also report the column densities of the groups of component a+b and c+d+e, which are well resolved from each other, unlike the individual components. Our first metallicity estimate is from Zn, as this element is usually not heavily depleted into dust (see e.g. [Pettini et al. 1994](#page-17-18)). We derive $[Zn/H]=-0.7\pm0.1$ (the other non-refractory elements Si and S are saturated, but the limits we find are consistent). This is in agreement with the value reported in [Cucchiara et al. \(2014](#page-16-19)).

We note that high ionisation lines from Si iv as well as C iv are detected, but are highly saturated, see Fig. [4.](#page-8-0)

3.2 Dust Depletion

Refractory elements, such as Fe, Ni, and Cr, can be heavily depleted into dust grains (e.g. [Savage & Sembach 1996](#page-17-19); [Ledoux et al. 2002b](#page-16-20), De Cia et al. in prep.), and thus can be missing from the gas-phase abundances. A first indicator of the level of depletion in the ISM is the relative abundance

² This procedure does not aim at reproducing the observed telluric spectrum, but simply reject suspect telluric lines from the Voigt-profile fit.

Figure 2. Velocity profiles of the metal resonance lines. Black lines show the normalised spectrum, with the associated error indicated by the dashed line at the bottom. The Voigt-profile fit to the lines is marked by the red line, while the single components of the fit are displayed in several colours (vertical dotted lines mark the centre of each component). The decomposition of the line-profile was derived by modelling only the underlined transitions. The oscillator strength 'f', is labelled in each panel. Saturated lines have not been fitted with a Voigt-profile, so for these we show only the spectrum. See online version for colours.

[Zn/Fe] (referred to as the depletion factor), because Zn is marginally if not at all depleted into dust grains, and its nucleosynthesis traces Fe. We measure $[Zn/Fe] = 0.85 \pm 0.06$. This value is among the highest for QSO-DLAs, but typical at the observed metallicity of $[Zn/H] = -0.7 \pm 0.1$ (e.g. [Noterdaeme et al. 2008](#page-17-20), De Cia et al. in prep.). Following [De Cia et al. \(2013](#page-16-21)) we calculate a column density of Fe in dust-phase of $log N(Fe)_{dust} = 16.74 \pm 0.17$ and a dust-corrected metallicity of $[Zn/H]_{\text{corr}} = -0.6 \pm 0.2$, indicating that even Zn is mildly depleted in this absorber, by ~ 0.1 dex. This is not surprising given the level of depletion, as also discussed by [Jenkins \(2009](#page-16-22)).

We also compare the observed abundances of a variety of metals (namely Zn, S, Si, Mn, Cr, Fe, and Ni) to the depletion patterns of a warm halo (H), warm disk+halo (DH), warm disk (WD) and cool disk (CD) types of environments, as defined in [Savage & Sembach \(1996\)](#page-17-19). These are fixed depletion patterns observed in the Galaxy and calculated assuming that Zn is not depleted into dust grains. We fit the observed abundances to the depletion patterns using the method described in [Savaglio \(2001\)](#page-17-21). We find that none of the environments are completely suitable to describe the observed abundances. The fits to cool- and warm-disk pat-terns are displayed in Fig. [5](#page-8-1) (χ^2_{ν} =1.18 and 1.58, respectively, with 4 degrees of freedom). For the cool disk the lower limit on the Si column density is not very well reproduced, while the fit for the warm disk overestimates the Mn abundance. The real scenario could be somewhere in between these two environments. Alternatively, the actual depletion pattern is different than what has been observed by [Savage & Sembach](#page-17-19) [\(1996](#page-17-19)), or there are some nucleosynthesis effects which we cannot constrain for our case.

Another quantity that is very useful to derive from the observed dust depletion is the dust-to-metals ratio $(DTM,$ normalised by the Galactic value). Constraining the DTM distribution on a variety of environments can indeed shed light on the origin of dust (e.g. [Mattsson et al. 2014](#page-17-22)). Based on the observed [Zn/Fe] and following [De Cia et al. \(2013](#page-16-21)),

Figure 3. The same as Fig. [2,](#page-6-0) but for fine-structure lines. Telluric features are highlighted in yellow.

we calculate $DTM = 1.01 \pm 0.03$, i.e. consistent with the Galaxy. From the depletion-pattern fit described above we derive similar, although somewhat smaller, $DTM = 0.84 \pm$ 0.02 (CD) and $DTM = 0.89 \pm 0.02$ (WD). These values are in line with the distribution of the DTM with metallicity and metal column densities reported by [De Cia et al. \(2013](#page-16-21)), and are also consistent with those of [Zafar & Watson \(2013](#page-17-23)). Following [Zafar & Watson \(2013](#page-17-23)), we calculate $D T M = 0.1$ now based on the dust extinction A_V that we model from the SED fit (Sect. [3.9\)](#page-11-0). Due to the small amount of reddening in the SED, this $DTM(A_V)$ value is a factor of 10 lower than expected at the metal column densities observed. This will be discussed further in Sect. [4.3.](#page-13-1)

At the metallicity of GRB 121024A (\sim 1/3 solar), it is not possible to draw further conclusions on the dust origin based on the DTM . Both models of pure stellar dust production and those including dust destruction and grain growth in the ISM converge to high (Galactic-like) DTM values at metallicities approaching solar [\(Mattsson et al. 2014](#page-17-22)).

3.3 Distance between GRB and Absorbing Gas

The most likely origin of the fine-structure transitions observed in the a+b $(\alpha+\beta)$ component, is photo-excitation by UV photons from the GRB afterglow itself (see e.g [Prochaska et al. 2006;](#page-17-24) [Vreeswijk et al. 2007](#page-17-25)). Assuming the afterglow to be the only source of excitation, we model the population of the different levels of Fe and Ni, closely following [Vreeswijk et al. \(2013\)](#page-17-26). Using an optical light curve to estimate the luminosity of the afterglow, we can then determine how far the excited gas must be located from the GRB site, for the afterglow to be able to excite these levels. We model the total column density from component a+b $(\alpha+\beta)$ of all observed levels (ground state and excited states) of Ni_{II} and Fe_{II}. We input the optical light curve from Skynet, see Fig. [6](#page-8-2) and Tables in Appendix [A,](#page-18-0) which is extrapolated to earlier times using the power-law decay observed. We use the broadeningparametersb derivedfrom the Voigt-profile fits, and the sameatomic parameters (see [Vreeswijk et al. 2013](#page-17-26)).

The best fit (see Fig. [7\)](#page-9-0) is obtained with a distance

Figure 4. High ionisation lines. These lines are highly saturated. See Fig. [2](#page-6-0) for details.

Figure 5. The dust-depletion pattern fit for a cold disk (red solid curve) and a warm disk (red dashed curve) to the observed abundances measured from absorption-line spectroscopy (diamonds and arrows, for the constrained and 3σ limits, respectively).

Figure 6. GRB-afterglow light curve from the Skynet instruments, used as input for the population modelling. The legend gives the instrument and observational band. The black arrow indicates the starting point of the X-shooter observations. Observations started 55 s after the GRB trigger. See online version for colours.

of 590 ± 100 pc between the cloud and burst, and a cloud size of $\langle 333 \text{ pc } (1\sigma)$. The resultant fit is rather poor $(\chi^2/\text{d.o.f} = 40.6/4)$. As can be seen in Fig. [2](#page-6-0) and [3,](#page-7-0) the column densities of the ground level of Ni ii (as probed by Ni II $\lambda\lambda\lambda$ 1709, 1454, 1467) and 5th excited level of Fe II (as probed by Fe ii 5s $\lambda\lambda$ 1559, 2360) are not very well constrained due to the observed spectrum having a low S/N ratio near those features. The formal errors from the Voigt profile fit are likely an underestimate of the true error for these column densities. This, in turn, results in the χ^2 of the excitation model fit being overestimated. Furthermore the lack of spectral time series means the resultant parameters are not well constrained. For the c+d+e component we are able to set a lower limit of 1.9 kpc on the distance to the burst using Fe II, and 3.5 kpc using Si II (3σ) . Since Si II is saturated, we use the EW to determine the column density, but that only gives the total value of all components together. Hence, for the c+d+e component we fitted using VPFIT and compared the total column density with what we get from the EWs. After establishing that both methods yield the same result, we feel confident in using the column density of $log N(Si II)_{c+d+e} > 15.99$ together with a detection limit $\log N(\text{Si II*})_{c+d+e}$ < 12.80 on the 1265 Å line, as this is the strongest of the $Si II^*$ lines. The lack of vibrationallyexcited H² in the spectra, see below, is in agreement with a distance $\gg 100$ pc, see [Draine \(2000](#page-16-23)).

3.4 Molecular Hydrogen

We detect Lyman- and Werner-band absorption lines of molecular hydrogen at redshift $z = 2.3021$ (corresponding to metal-line component $C + d + e$ " in rotational levels J=0, 1, 2 and 3, see Fig. [8.](#page-10-0) The fitting and analysis of the molecular hydrogen transition lines follow [Ledoux et al.](#page-16-20) $(2002b, 2003)$ $(2002b, 2003)$ $(2002b, 2003)$ $(2002b, 2003)$ and Krühler et al. (2013) . We performed a

Figure 7. Best-fit model for the excited-level populations of the a+b $(\alpha + \beta)$ column densities of Fe II (top panel) and Ni II (bottom). Black lines show the fit to the resonance level. For Fe II, from the lower levels and up, excited-level population are shown with red, green, purple, orange and blue. For NiII the red line shows the first excited level, while the blue line shows the second. Open circles show the actual values from Voigt-profile fits. See online version for colours.

Voigt-profile fit of lines mainly from the Lyman bands L0- 0 up to L3-0, as these are found in the less noisy part of the spectrum (a few $J = 2$ and 3 lines from the Lyman bands L4-0 and L5-0 were also fitted). $J = 0$ and 1 lines are strong and fairly well constrained by the presence of residual flux around them, hinting at damping wings in $L \geq 1$. Given the low spectral resolution of the data and the possibility of hidden saturation, we tested a range of Doppler parameters. The estimated H_2 column densities, log $N(\text{H}_2)$ are given in Table [6](#page-9-1) for Doppler parameter values of $b = 1$ and 10 km s^{-1} resulting in log $N(\text{H}_2) = 19.8 \text{--} 19.9$. Using the column density of neutral hydrogen for component $c+d+e'$ of $log N(H I) = 21.6$, calculated assuming the same Zn metallicity for the two main velocity components $('a+b')$ and $c+d+e'$, this results in a molecular fraction in the order of log $f \sim -1.4$, where $f \equiv 2N(\text{H}_2)/(N(\text{H I})+2N(\text{H}_2)).$ For the component 'a+b' at redshift $z \sim 2.2987$, we report $\log N(\text{H}_2) < 18.9$ as a conservative upper limit on detection. A more detailed analysis is not possible because of the high noise-level. The implications of this detection are discussed in Sect. [4.4.](#page-14-0)

We searched for vibrationally-excited H_2 by cross-

Table 6. Estimated column densities for H_2 for broadening parameter values $b = 10$ and $b = 1$ (km s⁻¹).

Rotational level	$\log(N({\rm H}_2)/{\rm cm}^{-2})$			
	$b = 10 \,\mathrm{km\,s^{-1}}$ $b = 1 \,\mathrm{km\,s^{-1}}$			
$J=0$	19.7	19.7		
$J=1$	19.2	19.3		
$J=2$	16.1	18.3		
$J=3$	16.0	18.2		
Total	19.8	19.9		

correlating the observed spectrum with a theoretical model from [Draine \(2000](#page-16-23)) and [Draine & Hao \(2002](#page-16-25)) similar to the procedure outlined in Krühler et al. (2013) . There is no evidence for H^{*}₂ in our data, neither through the crosscorrelation nor for individual strong transitions, and we set an upper limit of 0.07 times the optical depth of the input model. This approximately corresponds to $\log N(\text{H}_2^*)$ < 15.7. A column density of H_2^* as high as seen in e.g. GRBs 120815A or 080607 [\(Sheffer et al. 2009](#page-17-27)) would have been clearly detected in our data.

We furthermore note that CO is not detected. We set a conservative limit of $log N(CO) < 14.4$, derived by using four out of the six strongest CO AX bandheads with the lowest 6 rotational levels of CO. The wavelength range of the other two bandheads are strongly affected by metal lines, and thus do not provide constraining information.

3.5 Emission Lines

In the NIR spectrum, we detect H α , H β , the [O ii] $\lambda\lambda$ 3727, 3729 doublet, [N II] λ 6583 (highest-redshift [N II] detection published for a GRB host) and the two [O III] $\lambda\lambda$ 4959, 5007. Table [7](#page-11-1) shows the fluxes (extinction-corrected, see Sect. [3.8\)](#page-11-2). The reported fluxes are derived from Gaussian fits, with the background tied between the [O III] doublet and $H\beta$, and between $H\alpha$ and [N II], assuming a slope of the afterglow spectrum of 0.8. [O ii] is intrinsically a doublet, so we fit a double Gaussian with a fixed wavelength spacing based on the wavelength of the rest-frame lines. Using the GROND photometry, we estimate a slit-loss correction factor of 1.25 ± 0.10 . Fig. [9](#page-11-3) shows the emission-line profiles, the 2D as well as the extracted 1D spectrum. The figure shows a Gaussian fit to the lines, after subtracting the PSF for the continuum (done by fitting the spectral trace and PSF as a function of wavelength locally around each line, see [Møller 2000](#page-17-28), for details). For the weaker [N II], a formal χ^2 minimisation is done by varying the scale of a Gaussian with fixed position and width. The noise is estimated above and below the position of the trace (marked by a horizontal dotted line in Fig. [9\)](#page-11-3). We assign the zero-velocity reference at the redshift of the $[O III]$ λ 5007 line. For the weaker $[N II]$ line, we fix the Gaussian-profile fit to be centred at this zero-velocity.

3.6 Star-Formation Rate

The SFR can be derived from the emission line fluxes of $H\alpha$ and [O II]. Using conversion factors from [Kennicutt \(1998](#page-16-26)), but converted from a Salpeter initial mass function (IMF) to Chabrier [\(Treyer et al. 2007](#page-17-29)), we report extinction corrected

Figure 8. X-shooter spectrum showing Lyman- and Werner-band absorption. The shaded area shows the synthetic spectrum from a fit with Doppler parameter $b = 1 \text{ km s}^{-1}$, while the blue-dotted line shows the fit for $b = 10 \text{ km s}^{-1}$. J0 marks the transitions from the J = 0 rotational level, and likewise for higher J. See online version for colours.

(see Sect. [3.8\)](#page-11-2) values of $SFR_{H\alpha} = 42 \pm 11 \,\mathrm{M}_{\odot} \,\mathrm{yr}^{-1}$ from the H α flux and a SFR_[O II] = 53 ± 15 M $_{\odot}$ yr⁻¹ derived from [O ii]. For a comparison with results from the stellar population synthesis modelling see Sect. [3.10.](#page-12-0)

3.7 Metallicity from Emission Lines

We determine the gas-phase metallicity of the GRB host galaxy using the strong-line diagnostics R_{23} (using the ratio ([O II] $\lambda\lambda$ 3727 + [O III] $\lambda\lambda$ 4959, 5007)/H β), O3N2 (using

 $([O II]/H\beta)/([N II]/H\alpha)$ and N2 (using [N II]/H α ; for a discussion of the different diagnostics see e. g. [Kewley & Ellison](#page-16-27) [2008](#page-16-27)). Note that different metallicity calibrators give different values of metallicity. R_{23} appears to be consistently higher than $O3N2$ and N2. The R_{23} diagnostic has two branches of solutions, but the degeneracy can be broken using the ratios $[N II]/H\alpha$ or $[N II]/[O II]$. In our case $[N II]/H\alpha = 0.09 \pm 0.02$ and $[N II]/[O II] = 0.13 \pm 0.03$, which places the R_{23} solution on the upper branch (though not far from the separation). Because of the large difference in wavelength of the emission lines used for R_{23} , this method is sen-

Table 7. Measured emission-line fluxes

Transition	Wavelength ^a	$Flux^b$	Width ^c	Redshift
[O II]	3726.03, 3728.82	14.5 ± 1.2	\overline{d}	2.3015^e
$H\beta$	4861.33	7.4 ± 0.4	$218 + 12$	2.3012
[O III]	4958.92	9.0 ± 0.4	$194 + 28$	2.3017
[O III]	5006.84	27.2 ± 0.7	$192 + 7$	2.3010
$H\alpha$	6562.80	21.0 ± 1.5	$279 + 17$	2.3010
[NII]	6583.41	$1.9 + 0.7$	\sim 140	2.3015^{f}
	the contract of			

 a Wavelengths in air in units of \AA .

^b Extinction corrected flux in units of 10^{-17} erg s⁻¹ cm⁻².

^c FWHM of line (after removing instrumental broadening) in units of km s−¹ . Errors do not include uncertainty in continuum. d [O II] is intrinsically a doublet, which is not fully resolved here, so we do not give the width.

 e Calculated using a weighted wavelength average of 3727.7 Å. f The Gaussian fit shown of [N II] has a redshift frozen to that of the [O III] λ 5007 line.

sitive to the uncertainty on the reddening. Both O3N2 and N2 use lines that are close in wavelength, so for these we expect the reddening to have a negligible effect. Instead, they then both depend on the weaker [N II] line, which has not got as secure a detection. We derive $12 + \log(O/H) = 8.6 \pm 0.2$ for R₂₃ [\(McGaugh 1991](#page-17-30)), $12 + \log(O/H) = 8.2 \pm 0.2$ for O3N2 and $12 + \log(O/H) = 8.3 \pm 0.2$ for N2 (both from [Pettini & Pagel 2004](#page-17-31)). The errors include the scatter in the relations (these values are from [Kewley & Ellison 2008,](#page-16-27) and references therein), though the scatter in N2 is likely underestimated. See Sect. [4.1](#page-13-2) for a comparison with absorptionline metallicity.

3.8 Balmer Decrement

The ratio of the Balmer lines $H\alpha$ and $H\beta$ can be used to estimate the dust extinction. We use the intrinsic ratio found $I(H\alpha)/I(H\beta) = 2.86$ [\(Osterbrock 1989](#page-17-32)), for starforming regions (and case B recombination, meaning photons above 13.6 eV are not re-absorbed), where we expect GRBs to occur. The ratio we measure is 2.98 which, assuming the extinction law of Calzetti et al. $(2000)^3$ $(2000)^3$, results in $E(B - V) = 0.04 \pm 0.09$ mag. We note that adopting a different extinction law (from e.g. [Pei 1992](#page-17-33)) results in the same reddening correction within errors, because there is little difference within the wavelength range of the Balmer lines.

3.9 Broad-Band Spectral Energy Distribution

We fitted the broad-band afterglow data from XRT and GROND (without the g' -band, due to possible DLA contamination), where simultaneous data exist (11 ks after the trigger). The fit was perform within the ISIS software [\(Houck & Denicola 2000\)](#page-16-29) following the method of [Starling et al. \(2007](#page-17-34)). The XRT data were extracted using

Figure 9. Emission lines detected from the GRB 121024A host. Each panel shows the 2D spectrum after continuum PSF subtraction on top. The bottom part shows the extracted 1D spectrum. The blue line shows the Gaussian fit to the line profile. The abscissa shows the velocity dispersion with respect to the [O III] λ5007 reference frame. The [N ii] spectrum has been smoothed and binned differently than the other lines, and the fit has been performed with the Gaussian profile centre frozen at 0 km s−¹ with respect to the reference frame, as indicated with the dashed line in the figure. [O ii] has been fit as a doublet for the flux estimate.

Swift tools. We use single and broken power-law models. For the broken power-law, we tie thetwo spectral slopes to a fixed difference of 0.5. Such spectral feature is known as the cooling break of GRB afterglows (e.g. [Sari 1998](#page-17-35)), and is observed to be the best-fit model for most burst [Zafar et al.](#page-17-36) [\(2011](#page-17-36)), with the exception of GRB 080210 [\(Zafar et al. 2011](#page-17-36); [De Cia et al. 2011](#page-16-30)). We fit with two absorbers, one Galactic fixed at $N(\text{H})_{\text{X}}^{\text{Gal}} = 7.77 \times 10^{20} \text{ cm}^{-2}$ [\(Willingale et al.](#page-17-37) [2013](#page-17-37)), and one intrinsic to the host galaxy^{[4](#page-11-5)} An SMC dustextinction model (the average extinction curve observed in

⁴ We assume solar metallicity, not to provide a physical description of the absorbers, but purely to let $N(H)_X$ conform to the

³ We use the [Calzetti et al. \(2000\)](#page-16-28) law, which is an attenuation law for star burst galaxies, where the [Pei \(1992](#page-17-33)) laws are relevant for lines of sight towards point-sources inside galaxies where light is lost due to both absorption and scattering out of the line of sight.

Figure 10. NIR-to-X-ray spectral energy distribution and model for the afterglow at 11 ks after the trigger. The solid red line shows the model. g'-band magnitude is not included in the fit, due to possible contribution from the $Ly\alpha$ transition.

Table 8. Best fit parameters from the broad-band spectral energy distribution, for a single power-law SMC absorption model.

$N(H)_X$	$(1.2^{+0.8}_{-0.6}) \times 10^{22}$ cm ⁻²
$E(B-V)$	0.03 ± 0.02 mag
Power-law index β	0.90 ± 0.02

the Small Magellanic Cloud) was used for the host, while the reddening from the Milky Way was fixed to $E(B-V)$ = 0.123 [\(Schlegel et al. 1998\)](#page-17-38). A single power-law is preferred statistically $(\chi^2 / d.o.f = 1.07)$, see Fig. [10,](#page-12-1) but the two models give similar results.

The best fit parameters for the single power-law SMC absorption model are $N(H)_X = (1.2^{+0.8}_{-0.6}) \times 10^{22}$ cm⁻² and $E(B - V) = 0.03 \pm 0.02$ mag at a redshift of $z = 2.298$, and a power-law index of $\beta = 0.90 \pm 0.02$ (90 per cent confidence limits), see Table [8.](#page-12-2) LMC and MW (the average extinction curves observed in the Large Magellanic Cloud and the Milky Way) model fits result in the same values within errors. For a discussion on the extinction see Sect. [4.3.](#page-13-1)

3.10 Stellar Population Synthesis Modelling

Using our photometry of the host, see Table [3,](#page-3-1) we perform stellar population synthesis modelling of the host galaxy. We use a grid of stellar evolution models with different star formation timescales, age of stellar population and extinction, to compute theoretical magnitudes and compare them to the observed photometry. For the model input, we assume stellar models from [Bruzual & Charlot \(2003](#page-16-31)), based on an IMF from [Chabrier \(2003](#page-16-32)) and a Calzetti dust attenuation law [\(Calzetti et al. 2000\)](#page-16-28). Table [9](#page-12-3) lists the galaxy parameters resulting from the best-fit to the HAWK-I, NOT and GTC data. The best fit is obtained with a $\chi^2 = 8$ for the 7 data points used in the modelling. Most of the contribution to the χ^2 comes from the B-band observations. This

standard solar reference. The reference solar abundances used are from [Wilms et al. \(2000](#page-17-39)).

Table 9. Host galaxy parameters from stellar population synthesis modelling

Starburst age (Myr)	~ 250
Extinction (mag)	$0.15 + 0.15$
$M_{\rm B}$	-22.1 ± 0.2
$log(M_{*}/M_{\odot})$	$9.9^{+0.2}_{-0.3}$
$SFR (M_{\odot} yr^{-1})$	40^{+80}_{-25}

Table 10. Equivalent widths of intervening systems.

data point lies $\approx 3\sigma$ above the best-fit and the g-band measurement, which probes a very similar wavelength range. The reported value of the SFR takes into account the uncertainty in the dust attenuation, and thus has large error bars. We observe a significant Balmer break, which is well fit with star-burst ages between 50 and 500 Myr. The SFR of $\sim 40 \,\mathrm{M}_{\odot} \,\mathrm{yr}^{-1}$ is consistent with the results from Sect. [3.6.](#page-9-2)

3.11 Kinematics

The X-shooter spectrum contains information both on the kinematics of the absorbing gas along the line-of-sight to the location of the burst inside the host galaxy, as well as kinematics of the emitting gas in H ii regions probed by the emission lines. The emission lines have a full-width-at-halfmaximum (FWHM) of around 210 km s^{-1} from a Gaussian fit, see Table [7.](#page-11-1) We do not observe signs of rotation in the 2 dimensional spectrum. One possibility is that the galaxy could be dominated by velocity dispersion, as observed for galaxies of similar mass and properties, Förster et al. (2009). The velocity width that encloses 90% of the optical depth (as defined by [Ledoux et al. 2006\)](#page-16-34) is 460 km s⁻¹ based on the SiII λ 1808 line. This is consistent with the correlation between absorption line width and metallicity for GRB host galaxies of [Arabsalmani et al. \(2015\)](#page-16-35). The velocity for each absorption component, with respect to the emission lines, is given in Table [4.](#page-4-0) The characteristics of different gas components are discussed in Sect. [4.5.](#page-15-0)

3.12 Intervening Systems

We identify three intervening systems along the line of sight, at redshifts $z = 2.0798$, $z = 1.959$, and $z = 1.664$. Table [10](#page-12-4) lists the observed lines along with the measured EWs. Furthermore only at $z = 1.959$ do we observe the Ly α line, but blended with a Si iv line. The intervening systems will not be discussed further in this work.

4 DISCUSSION AND IMPLICATIONS

4.1 Abundance Measurements from Absorption and Emission Lines

The metallicity of GRB hosts is usually determined either directly through absorption line measurements, or via the strong-line diagnostics using nebular-line fluxes. The two methods probe different physical regions; the ISM of the host galaxy along the line of sight as opposed to the ionised star-forming H ii regions emission - weighted over the whole galaxy. Hence, the two methods are not necessarily expected to yield the same metallicity, see for instance [Jorgenson & Wolfe \(2014](#page-16-4)). The line of sight towards the GRB is expected to cross star-forming regions in the GRB host. Thus, the absorption and emission lines may probe similar regions. Local measurements from the solar neighbourhood show a concurrence of the two metallicities in the same region, see e. g. [Esteban et al. \(2004](#page-16-36)). Only a few cases where measurements were possible using both methods have been reported for QSO-DLAs (see e.g. [Bowen et al.](#page-16-2) [2005](#page-16-2); [Jorgenson & Wolfe 2014\)](#page-16-4) but never for GRB-DLAs. The challenge is that the redshift has to be high enough $(z \ge 1.5)$ to make Ly α observable from the ground, while at the same time the host has to be massively star-forming to produce sufficiently bright emission lines. Furthermore, the strong-line diagnostics are calibrated at low redshifts, with only few high redshift cases available (see for instance [Christensen et al. 2012](#page-16-37)).

The spectrum of GRB 121024A has an observable $Ly\alpha$ line as well as bright emission lines. We find that the three nebular line diagnostics R_{23} , O3N2 and N2 all find a similar oxygen abundance of $12 + \log(O/H) = 8.4 \pm 0.4$. Expressing this in solar units we get a metallicity of $\text{[O/H]} \sim -0.3$ (or slightly lower if we disregard the value found from the R²³ diagnostic, given that we cannot convincingly distinguish between the upper and lower branch). This is indeed consistent with the absorption line measurement from the low-depletion elements (dust-corrected value) $[Zn/H]_{\text{corr}} =$ -0.6 ± 0.2 , though the large uncertainty in the strong-line diagnostics hinders a more conclusive comparison.

[Krogager et al. \(2013](#page-16-38)) find a slightly lower metallicity from absorption lines in the spectrum of quasar Q2222- 0946, compared to the emission-line metallicity. However, this is easily explained by the very different regions probed by the nebular lines (6 kpc above the galactic plane for this quasar) and the line of sight, see also P \acute{e} roux et al. (2012). QSO lines of sight intersect foreground galaxies at high impact parameters, while the metallicity probed with GRB-DLAs are associated with the GRB host galaxy. Interestingly, [Noterdaeme et al. \(2012](#page-17-5)) find different values for the metallicities, even with a small impact parameter between QSO and absorber, for a QSO-DLAs. A comparison of the two metallicities is also possible for Lyman break galaxies (LBGs), see for instance [Pettini et al. \(2002](#page-17-40)) for a discussion on the metallicity of the galaxy MS 1512–cB58. They find that the two methods agree for a galaxy with an even larger velocity dispersion in the absorbed gas than observed here $($ ∼ 1000 km s⁻¹). The line of sight toward GRB 121024A crosses different clouds of gas in the host galaxy, as shown by the multiple and diverse components of the absorption-line profiles. The gas associated with component a+b is photoexcited, indicating that it is the closest to the GRB. Given

the proximity, the metallicity of this gas could be representative of the GRB birth site. Assuming the GRB exploded in an H ii region, the emission- and absorption-metallicities are expected to be similar, though if other H ii region are dominating the brightness, the GRB birth sight might contribute only weakly to the emission line-flux, see Sect [4.5.](#page-15-0) Building a sample of dual metallicity measurements will increase our understanding of the metallicity distribution and evolution in galaxies.

4.2 The Mass-Metallicity Relation at $z \sim 2$

Having determined stellar mass, metallicity and SFR of the GRB host, we can investigate whether the galaxy properties are consistent with the mass-metallicity relation at the observed redshift. Appropriate for a redshift of $z \approx 2$, we use equation 5 from [Mannucci et al. \(2010\)](#page-17-41):

$$
12 + \log(O/H) = 8.90 + 0.47 \times (\mu_{0.32} - 10)
$$

where $\mu_{0.32} = \log(M_*[M_{\odot}]) - 0.32 \times \log(SFR_{H\alpha}[M_{\odot}]yr^{-1}).$ Using the stellar mass from Sect. [3.10](#page-12-0) and the SFR from $H\alpha$ we find an equivalent metallicity of $12 + \log(O/H) = 8.6 \pm 0.2$ $([O/H] = -0.1 \pm 0.2)$. The error does not include a contribution from the scatter in the relation, and is hence likely underestimated. This value is consistent with the metallicity derived from the emission lines, but given the large uncertainty this is perhaps not that illustrative. Instead, we use the mass-metallicity relation determined in [Christensen et al. \(2014\)](#page-16-39) for QSO-DLAs for absorption-line metallicities remodelled to GRBs by [Arabsalmani et al.](#page-16-35) [\(2015](#page-16-35)). This results in a metallicity $[M/H] = -0.3 \pm 0.2$, not including scatter from the relation, and using the mean impact parameter of 2.3 kpc calculated in [Arabsalmani et al.](#page-16-35) [\(2015](#page-16-35)). This is consistent with the dust-corrected metallicity of $[Zn/H]_{\text{corr}} = -0.6 \pm 0.2$.

4.3 Grey Dust Extinction?

We determine the dust extinction/attenuation of the host galaxy of GRB 121024A both from the Balmer decrement (Sect. [3.8\)](#page-11-2) and a fit to the X-ray and optical spectral energy distribution (SED, see Sect. [3.9\)](#page-11-0), as well as from the stellar population synthesis modelling (Sect. [3.10\)](#page-12-0). The first method determines the attenuation of the host H ii regions (from the X-shooter spectrum alone), while the SED fitting probes the extinction along the line of sight (using XRT+GROND data). The stellar population synthesis modelling models the host attenuation as a whole (using host photometry). All methods determine the amount of extinction/attenuation by comparing different parts of the spectrum with known/inferred intrinsic ratios, and attribute the observed change in spectral form to dust absorption and scattering. We find values that agree on a colour index $E(B - V) \sim 0.04$ mag. This value is small, but falls within the range observed for GRB-DLA systems. However, low A_V 's are typically observed for the lowest metallicities. For our case we would expect a much higher amount of reddening at our determined H_I column density and metallicity. Using the metallicity of $[Zn/H]_{\text{corr}} = -0.6 \pm 0.2$, column density $log N(H I) = 21.88 \pm 0.10$, dust-to-metal ratio $DTM= 1.01 \pm 0.03$ (see Sect. 3.2), and a reference Galactic dust-to-metal ratio $A_{V,Gal}/N_{(H,Gal)} = 0.45 \times 10^{-21}$ mag cm² [\(Watson 2011](#page-17-42)), we expect an extinction of $A_V = 0.9 \pm$ 0.3 mag (De Cia et al. in prep. and [Savaglio, Fall & Fiore](#page-17-43) [2003](#page-17-43)). This is incompatible with the determined reddening, as it would require $R_V > 15$ (R_V for the Galaxy is broadly in the range 2–5). For the Balmer decrement and SED fitting we have examined different extinction curves (MW and LMC besides the SMC) and we have tried fitting the SED with a cooling break, neither option changing the extinction significantly. In an attempt to test how high a fitted reddening we can achieve, we tried fitting the SED with a lower Galactic $N(H I)$ and reddening. While keeping reasonable values (it is unphysical to expect no Galactic extinction at all), and fitting with the break, the resulting highest colour index is $E(B - V) \sim 0.06$ mag. This is still not compatible with the value derived from the metallicity, so this difference needs to be explained physically.

One possibility to consider is that the host could have a lower dust-to-metals ratio, and hence we overestimate the extinction we expect from the metallicity. However, we see no sign of this from the relative abundances, see Sect. [3.2.](#page-5-2) The metallicity is robustly determined from Voigt-profile fits and EW measurement of several lines from different elements (including a lower limit from the none Fe-peak element Si). The lines are clearly observed in the spectra, see Fig. [2,](#page-6-0) and the metallicity that we find is consistent with the mass-metallicityrelation.

To examine the extinction curve, we perform a fit to the XRT (energy range: 0.3–10 keV) X-ray data alone and extrapolate the resultant best-fit power-law to optical wavelengths. We try both a single power law, as well as a broken power law with a cooling break in the extrapolation. The latter is generally found to be the best model for GRB extinction in optical fit (e.g. [Zafar et al. 2011;](#page-17-36) [Greiner et al.](#page-16-40) [2011](#page-16-40); [Schady et al. 2012](#page-17-44)). We calculate the range of allowed A_{λ} by comparing the X-shooter spectrum to the extrapolation, within the 90 % confidence limit of the best-fit photon index, and a break in between the two data sets (X-ray and optical). The resulting extinction does not redden the afterglow strongly, so we cannot constrain the total extinction very well directly from the SED. However, the optical spectroscopy indicates a high metal column density. The strong depletion of metals from the gas phase supports the presence of dust at $A_V = 0.9 \pm 0.3$ mag (see Sect [3.2\)](#page-5-2). Fixing A_V at this level, allows us to produce a normalised extinction curve (Fig. [11\)](#page-14-1). This extinction curve is very flat, much flatter than any in the local group [\(Fitzpatrick & Massa 2005](#page-16-41)), with an $R_V > 9$.

The most likely physical scenario that can explain this shape of the curve is grey dust. If the dust extinction is 'grey', i.e., has a much weaker dependence on wavelength than in the local extinction laws, then a given visual extinction will be much less apparent in the SED ('flat' extinction curve) and thus underestimated in our analysis. Such grey extinction corresponds to larger R_V and is physically interpreted with large grain sizes. A weak wavelength dependance in the extinction for GRBs has been suggested before. [Savaglio & Fall \(2004](#page-17-45)) reported a MW-like depletion pattern, but a very low reddening in the SED. [Gao, Jiang & Li](#page-16-42) [\(2008](#page-16-42)) likewise claim a grey extinction law for GRB lines of sight determined by comparing observed spectra to intrin-

Figure 11. Extinction curves for the line of sight to GRB 121024A. We plot the extinction law assuming $A_V = 0.9 \pm$ 0.3 mag as expected from the measured metallicity, H_I column density and dust-to-metal ratio. The solid black curve shows the extinction curve for a broken power law with $A_V = 0.9$, while the grey-shaded area corresponds to the A_V error-space. Likewise, the extinction curve for a single power law is plotted with the dashed black curve, and the hatched area displays the errorspace. Over-plotted, in colours, are extinction curves from Pei (1992).

sic ones (by extrapolating from X-rays), arguing for grain growth through coagulation in the dense molecular clouds surrounding GRBs. The larger grains have an extinction that is less dependent on wavelength, because of the contribution of their physical cross-section to the opacity. Preferable destruction of the smaller grains by the GRB would be another possibility, but is unlikely in our case, because the absorbing gas is far from the GRB. We note that the other GRB-DLAs with molecular-hydrogen detection show the expected amount of reddening (using a standard extinction curve), though anomalies do exist in GRB observations. The most notable example to date is reported by [Perley et al.](#page-17-46) [\(2008](#page-17-46)) for GRB 061126. As for the GRB 121024A afterglow, they observe a very flat optical–to–X-ray spectral index, arguing for large quantities of grey dust, or a separate origin of the optical and X-ray afterglow. To fit the extinction curve for GRB 061126, an $R_V \sim 10$ is needed. We find an R_V even higher than this, making this case even more extreme than previously observed. We refer to future work on this problem (Friis et al., in prep), as a deeper analysis is beyond the scope of this paper.

4.4 Molecular Hydrogen in GRB-DLAs

The lack of detection of molecular hydrogen towards GRBs has puzzled astronomers (see e. g [Tumlinson et al. 2007](#page-17-9)), given that long GRBs are associated with active star formation, and hence are expected to show signatures of molecular clouds. Compared to QSO-DLA line of sights then, we would expect the presence of H_2 to be more common for GRB-DLAs, because the QSO-DLA line of sights have a higher probability to intersect the outskirts/halo of the intervening galaxy, where we would anticipate a low molecular con-

tent. Recently, a number of H_2 detections in GRB afterglows have been reported [\(Prochaska et al. 2009](#page-17-10); Krühler et al. [2013](#page-16-7); [D'Elia et al. 2014](#page-16-8)), making GRB121024A the fourth definite case. This detection supports the emerging picture that dust has played a major role in biasing past observations against molecular detection (e.g. [Ledoux et al. 2009](#page-16-0)). Molecules are thought to form on the surface of dust grains, and once formed, shielded from Lyman-Werner photons by the grains. Krühler et al. (2013) suggest that it is likely this connection that is responsible for the low number of H_2 detections towards GRB-DLAs. The high dust column density makes the GRB afterglow UV-faint, preventing highresolution and high S/N spectroscopy, which is needed to identify the presence of molecular gas. Thus, the lack of H2 detections in most GRB-DLAs can be explained with an observational bias. They illustrate this argument by investigating the metallicity, $N(H I)$ and dust depletion parameter space, showing that the GRB-DLAs with unsuccessful molecular searches fall outside the region where we would expect detections (with the only exception being GRB 050829A). This argument is supported by the observed $log N(H I)$, metallicity and depletion factor of GRB 121024A, which lies inside the parameter space where molecular detections are expect.

The high level of dust depletion observed in this GRB-DLA (see Sect. [3.2\)](#page-5-2), is consistent with molecular detec-tions in QSO-DLAs [\(Noterdaeme et al. 2008;](#page-17-20) Krühler et al. 2013), where there is a strong preference for H₂-bearing DLAs to have significant depletion factors. The dependence on the total neutral hydrogen column density is weak (although intrinsically-weak molecular lines are better constrained in strong DLAs), whereas the parameter that seems to determine whether H_2 is detected, is the column density of iron locked into dust. The $\log N(\text{Fe})_{\text{dust}}$ that we measure is 2 dex higher than the column density above which a significant presence of molecules has been observed in QSO-DLAs [\(Noterdaeme et al. 2008\)](#page-17-20). Indeed [De Cia et al. \(2013](#page-16-21)) studied log $N(\text{Fe})_{\text{dust}}$ and concluded that GRB hosts are promising sites for molecular detections.

[D'Elia et al. \(2014\)](#page-16-8) find a molecular fraction for GRB 120327A of $log(f)$ between -7 and -4 with a depletion factor of Zn/Fe = 0.56 \pm 0.14, while for the dustier line of sight towards GRB 120815A Krühler et al. (2013) reports a value of $\log(f) = -1.14 \pm 0.15$ ([Zn/Fe] = 1.01 \pm 0.10). For GRB 121024A we find intermediate values, although with the high noise-level the numbers are consistent with those reported for GRB 120815A. For GRB 080607 [Prochaska et al.](#page-17-10) [\(2009](#page-17-10)) only report limits on both the molecular fraction and the Zn+Fe column densities. Although the sample is too small to infer anything statistically, it appears that the H_2 detection criteria in GRB afterglows follow the trend observed for QSO-DLAs. For a fair estimate of the molecular fraction, the column densities of both H_2 and H_1 should be constrained for individual velocity components, while this is hardly the case for H i. Recent work (e.g. [Balashev et al.](#page-16-43) [2015](#page-16-43)) indicates that the molecular fraction in QSO-DLAs can possibly be much higherthan the line-of-sight average values usually measured.

4.5 Gas kinematics: dissecting the host components

One of the striking features of the metal absorption-line profiles observed towards GRB 121024A is that they consist of two widely separated groups of velocity components $(a+b \text{ and } c+d+e, \text{ see Sect. 3.1}).$ The separation is of about 340 km s⁻¹, which lies at the high end of the velocity distribution of [Møller et al. \(2013](#page-17-47)) and [Ledoux et al. \(2006](#page-16-34)). The latter is for QSO-DLAs, however [Arabsalmani et al. \(2015](#page-16-35)) showed that GRB-DLAs follow the velocity-metallicity distribution of QSO-DLAs. This suggests that either the two components belong to separate galaxies (see for instance [Savaglio et al. 2012,](#page-17-48) on the GRB 090323 systems), or that this galaxy is fairly massive compared to the average GRB host of $\sim 10^9$ M⊙ [\(Savaglio et al. 2009](#page-17-49); Castro Cerón et al. [2010](#page-16-44), but see also [Perley et al. 2013](#page-17-50) and [Hunt et al. 2014](#page-16-45)). The scenario with separate galaxies is disfavoured, because the two absorption components show very similar relative abundances (see Table [4\)](#page-4-0) and also because the emission lines are centred in between the two absorption components (unlike for GRBs 050820A and 060418; see [Chen 2012](#page-16-46)). Thus, a likely possibility is that the two absorption components are probing different regions within the host. This is in agree-ment with the mass found in Sect. [3.10,](#page-12-0) of almost $10^{10} M_{\odot}$.

Furthermore, the blue $(a+b)$ and the red $(c+d+e)$ absorption components are associated with gas at different physical conditions. On one hand, Fe ii, Ni ii and Si ii finestructure lines are detected only in the blue component. These lines are photo-excited by the GRB radiation at a distance of ∼600 pc. On the other hand, H² molecules are detected in the red component only, indicating a gas that is not disturbed by the GRB (at a distance of minimum 3.5 kpc). Through absorption-line spectroscopy at the X-shooter resolution, these two different gas components could be located inside the host (with respect to the GRB) and characterised.

The observed emission component (arbitrarily set at $v = 0$) traces the brightest star-forming regions. Since GRBs tend to reside around the brightest star-forming regions in their host [\(Fruchter et al. 2006](#page-16-47)), one might expect to observe absorption components at velocities close to that of the emission as the line of sight passes through this gas. However, the gas around the GRB can be photo-ionised out to hundreds of parsecs (e.g. [Ledoux et al. 2009](#page-16-0); [Vreeswijk et al.](#page-17-26) [2013](#page-17-26)); it is thus highly unlikely that the optical/UV absorption lines are probing the actual GRB environment. For GRB 121024A, this is further supported by the fact that the a,b component is located ∼600 pc away from the GRB. Given that giant molecular clouds have a maximum radius of \sim 200 pc [\(Murray 2011\)](#page-17-51), the a,b component is undoubtedly unrelated to the GRB surroundings.

Although GRBs are most often associated with the brightest star-forming regions, this is not always the case. GRB 980425 (e.g. Michałowski et al. 2009) is an example where the star-forming region in which the GRB occurred is quite faint compared to the larger and brighter star-forming regions in the host. A potentially similar scenario could hold for GRB 121024A as well, in which case the burst should not be identified with $v = 0$. In this situation, the possible interpretations of the kinematics would be different and lead to other geometric setups compared to those conceivable were the GRB localised close to $v = 0$. It should also be noted that

we have not discussed transverse motion which could complicate the interpretation even further. Finally, since the host is most likely an irregular galaxy, indicating a 3D perturbed environment without a rotating disk, we find it appropriate not to draw further conclusions.

While the available data do not allow us to discriminate between possible scenarios, this work demonstrates how powerful GRB afterglow observations can be to start dissecting individual building-block components of star-forming galaxies at $z \sim 2$ and above. This is especially true once we have gathered enough data to compile a statistical sample; see (e.g. [Fox et al. 2008](#page-16-48)) for previous work on VLT/UVES data and Fynbo et al. (in prep) for upcoming VLT/Xshooter results on a large afterglow sample.

ACKNOWLEDGMENTS

MF acknowledges support from the University of Iceland Research fund. ADC acknowledges support by the Weizmann Institute of Science Dean of Physics Fellowship and the Koshland Center for Basic Research. The Dark Cosmology Centre is funded by the DNRF. RLCS is supported by a Royal Society Dorothy Hodgkin Fellowship. JPUF acknowledge support from the ERC-StG grant EGGS-278202. CCT is supported by a Ramón y Cajál fellowship. The research activity of J. Gorosabel, CCT, and AdUP is supported by Spanish research project AYA2012-39362-C02-02. Ad.U.P. acknowledges support by the European Commission under the Marie Curie Career Integration Grant programme (FP7- PEOPLE-2012-CIG 322307). SS acknowledges support from CONICYT-Chile FONDECYT 3140534, Basal-CATA PFB-06/2007, and Project IC120009 "Millennium Institute of Astrophysics (MAS)" of Iniciativa Científica Milenio del Ministerio de Economía, Fomento y Turismo. Part of the funding for GROND (both hardware as well as personnel) was generously granted from the Leibniz-Prize to Prof. G. Hasinger (DFG grant HA 1850/28-1). We thank Alain Smette for providing the telluric spectrum, and the referee for very constructive feedback.

REFERENCES

- Arabsalmani M., Møller P., Fynbo J. P. U., Christensen L., Freudling W., Savaglio S., Zafar T., 2015, MNRAS, 446, 990
- Asplund M., Grevesse N., Sauval A. J., Scott P., 2009, ARA&A, 47, 481
- Barthelmy S.D. et al., 2005, SSRv, 120, 143
- Balashev S. A., Noterdaeme P., Klimenko V. V., Petitjean P., Srianand R., Ledoux C., Ivanchik A. V., Varshalovich D. A., 2015, A&A, 575, 8
- Bowen D. V., Jenkins E. B., Pettini M., Tripp T. M., 2005, ApJ, 635, 880
- Bruzual G., Charlot S., 2003, MNRAS, 344, 1000
- Calzetti D., Armus L., Bohlin R. C., Kinney A. L., Koorn-
- neef J., Storchi-Bergmann T., 2000, ApJ, 533, 682
- Cano Z., 2013, MNRAS, 434, 1098
- Castro Cerón J. M. et al., 2010, ApJ, 721, 1919
- Cepa J. et al., 2000, SPIE, 4008, 623
- Chabrier G., 2003, PASP, 115, 763
- Chen H.-W., 2012, MNRAS, 419, 3039
- Christensen L. et al., 2012, MNRAS, 427, 1973
- Christensen L., Møller P., Fynbo J. P. U., Zafar T., 2014, MNRAS, 445, 225
- Cucchiara A., Fumagalli M., Rafelski M., Kocevski D., Prochaska J. X., Cooke R. J., Becker G. D., 2014, [arXiv:1408.3578](http://arxiv.org/abs/1408.3578)
- De Cia A. et al., 2011, MNRAS, 412, 2229
- De Cia A. et al., 2012, A&A, 545, 64
- De Cia A., Ledoux C., Savaglio S., Schady P., Vreeswijk P. M., 2013, A&A, 560, 88
- D'Elia V. et al., 2014, A&A, 564, 38
- Elíasdóttir A. et al., 2009, ApJ, 697, 1725
- Draine B. T., 2000, ApJ, 532, 273
- Draine B. T., Hao L., 2002, ApJ, 569, 780
- Esteban C., Peimbert M., García-Rojas J., Ruiz M. T.,
- Peimbert A., Rodríguez M., 2004, MNRAS, 355, 229
- Fitzpatrick E. L., Massa D. 2005, AJ, 130, 1127
- Fox A. J., Ledoux C., Vreeswijk P. M., Smette A., Jaunsen A. O., 2008, A&A, 491, 189
- Förster S. et al., 2009, ApJ, 706, 1364
- Fruchter A. S. et al., 2006, Nature, 441, 463
- Fynbo J. P. U. et al., 2009, ApJS, 185, 526
- Fynbo J. P. U. et al., 2011, MNRAS, 413, 2481
- Fynbo J. P. U. et al., 2013, MNRAS, 436, 361
- Gao J., Jiang B. W., Li A., 2008, ApJ, 707, 89
- Gehrels N. et al., 2004, ApJ, 611, 1005
- Goldoni P., Royer F., François P., Horrobin M., Blanc G., Vernet J., Modigliani A., Larsen J., 2006, in SPIE conf. ser., 6269
- Greiner J. et al., 2007, The Messenger, 130, 12
- Greiner J. et al., 2008, PASP, 120, 405
- Greiner J. et al., 2011, A&A, 526, 30
- Haehnelt M. G., Steinmetz M., Rauch M., 1998, ApJ, 495, 647
- Hjorth J. et al., 2003, Nature, 423, 847
- Houck J. C., Denicola L.A., 2000, ASPC, 216, 519
- Hunt L. K. et al., 2014, A&A, 565, 112
- Jakobsson P. et al., 2006, A&A, 460, L13
- Jenkins E. B., 2009, ApJ, 700, 1299
- Jorgenson R., Wolfe A. M., 2014, ApJ, 785, 16
- Kennicutt Jr. R. C., 1998, ARA&A, 36, 189
- Kewley L. J., Ellison S. L., 2008, ApJ, 681, 1183
- Krogager. J.-K. et al., 2013, MNRAS, 433, 3091
- Krühler T. et al., 2008, ApJ, 685, 376
- Krühler T. et al., 2012, A&A, 546, 8
- Krühler T. et al., 2013, A&A, 557, 18
- Kudritzki et al., 2012, ApJ, 747, 15
- Levesque E. M., Berger E., Kewley L. J., Bagley M. M., 2010, AJ, 139, 694
- Ledoux C., Petitjean P., Bergeron J., Wampler E. J., Srianand R., 1998, A&A, 337, 51
- Ledoux C., Bergeron J., Petitjean P., 2002a, A&A, 385, 802
- Ledoux C., Srianand R., Petitjean P., 2002b, A&A, 392, 781
- Ledoux C., Petitjean P., Srianand R., 2003, MNRAS, 346, 209
- Ledoux C., Petitjean P., Fynbo J. P. U., Møller P., Srianand R., 2006, A&A, 457, 71
- Ledoux C., Vreeswijk P. M., Smette A., Fox A. J., Petitjean P., Ellison S. L., Fynbo J. P. U., Savaglio S., 2009, A&A, 506, 661
- Lodders K., Palme H., Gail H. P., 2009, Landolt Börnstein, 44
- Mannuci F., Cresci G., Maiolino R., Marconi A., Gnerucci A., 2010, MNRAS, 408, 2115
- Mattsson L., De Cia A., Andersen A.C., Zafar T., 2014, MNRAS, 440, 1562
- McGaugh S. S., 1991, ApJ, 380, 140
- Melioli C., Brighenti F., D'Ercole A., De Gouveia Dal Pino E. M., 2008, MNRAS, 388, 573
- Michałowski M. J. et al., 2009, ApJ, 693, 347
- Morton D. C., 2003, ApJs, 149, 205
- Møller P., 2000, The Messenger, 99, 31
- Møller P., Fynbo J. P. U., Ledoux C., Nilsson K. K., 2013, MNRAS, 430, 2680
- Murray N., 2011, ApJ, 729, 133
- Noterdaeme P., Ledoux C., Petitjean P., Srianand R., 2008, A&A, 481, 327
- Noterdaeme P. et al., 2012, A&A, 540, 63
- Osterbrock D. E., 1989, Astrophysics of Planetary Nebulae and Active Galactic Nuclei (University Science Books)
- Pei Y. C., 1992, ApJ, 395, 130
- Perley D. A. et al., 2008, ApJ, 672, 449
- Perley D. A. et al., 2013, ApJ, 778, 128
- Péroux C., Bouché N., Kulkarni V. P., York D. G., Vladilo G., 2012, MNRAS, 419, 3060
- Pettini M., Pagel B. E. J., 2004, MNRAS, 348, L59
- Pettini M., Smith L. J., Hunstead, R. W., King D. L., 1994, ApJ, 426, 79
- Pettini M., Rix S. A., Steidel C. C., Adelberger K. L., Hunt M. P., Shapley A. E., 2002, ApJ, 569, 742
- Prochaska J. X., Wolfe A. M., 1997, ApJ, 487, 73
- Prochaska J. X., Chen H. W., Bloom J. S., 2006, ApJ, 648, 95
- Prochaska J. X., Chen H. W., Dessauges-Zavadsky M., Bloom J. S., 2007, ApJ, 666, 267
- Prochaska J. X. et al., 2009, ApJ, 691, L27
- Reichart D., 2005, Nuovo Cimento C Geophysics Space Physics C, 28, 767
- Sari R., 1998, ApJ, 494, 49
- Savage B. D., Sembach K., R., 1996, ARA&A, 34, 279
- Savaglio S., 2001, IAUS, 204, 307
- Savaglio S., Fall S. M., Fiore F., 2003, ApJ, 585, 638
- Savaglio S., Fall S. M., 2004, ApJ, 614, 293
- Savaglio S., Glazebrook K., Le Borgne D., 2009, ApJ, 691, 182
- Savaglio S. et al., 2012, MNRAS, 449, 627
- Schady et al., 2012, ApJ, 537, 15
- Schlegel D. J., Finkbeiner D. P., Davis M., 1998, ApJ, 500, 525
- Schulze S. et al., 2014, A&A, 566, 102
- Sheffer Y., Prochaska J. X., Draine B. T., Perley D. A., Bloom J. S., 2009, ApJ, 701, 63
- Smette A., Sana H., Horst H., 2010, Highlights of Astronomy, 15, 533
- Smith et al., 2002, AJ, 123, 2121
- Sparre M. et al., 2011, ApJ, 735, L24
- Sparre M. et al., 2014, ApJ, 785, 105
- Stanek K. Z. et al., 2003, ApJ, 591, L17
- Starling R. L. C., Wijers R. A. M. J., Wiersema K., Rol E., Curran P. A., Kouveliotou C., van der Horst A. J.,
- Heemskerk M. H. M., 2007, ApJ, 661, 787
- Steeghs D., McClintock J. E., Parsons S. G., Reid M. J., Littlefair S., Dhillon V. S., 2013, ApJ, 768, 185
- Tody D., 1997, Astronomical Society of the Pacific Conference Series, 125, 451
- Treyer M. et al., 2007, ApJs, 173, 256
- Tumlinson J., Prochaska J. X., Chen H. W., Dessauges-Zavadsky M., Bloom J. S., 2007, ApJ, 668, 667
- Vernet J. et al., 2011, A&A, 536, A105
- Vladilo G., Centurión M., Levshakov S. A., Péroux C., Khare P., Kulkarni V. P., York D. G., 2006, A&A, 454, 151
- Vreeswijk P. M. et al., 2004, A&A, 419, 927
- Vreeswijk P. M. et al., 2007, A&A, 468, 83
- Vreeswijk P. M. et al., 2013, A&A, 549, 22
- Wakker B. P., York D. G., Wilhelm R., Barentine J. C., Richter P., Beers T. C., Ivezić Z, Howk J. C., 2008, ApJ, 672, 298
- Watson D., 2011, A&A, 533, 16
- Wiersema K. et al., 2007, A&A, 464, 529
- Wiersema K. et al., 2014, Nature, 509, 201
- Willingale R., Starling R. L. C., Beardmore A. P., Tanvir N. R., O'Brien P. T., 2013, MNRAS, 431, 394
- Wilms J., Allen A., McCray R., 2000, ApJ, 542, 914
- Wolfe A. M., Gawiser E., Prochaska J. X., 2005, ARAA, 43, 861
- Zafar T., Watson D., 2013, A&A, 560, 26
- Zafar T., Watson D., Fynbo J. P. U., Malesani D., Jakob-
- sson P., de Ugarte Postigo A., 2011, A&A, 532, 143

APPENDIX A: SKYNET MAGNITUDE TABLES

Tabels [A1,](#page-18-1) [A2,](#page-18-2) [A3,](#page-19-0) [A4,](#page-19-1) [A5](#page-19-2) and ?? give the magnitudes (not corrected for reddening) used for the optical light-curve input to model the distance between the excited gas (component a+b) and the burst itself.

Table A1. Skynet - Filter ${\cal R}$

Filter	Time (h)	Exposure time	S/N	Mag. (Vega)
\boldsymbol{R}	0.01656	1×10 s	27.7	15.01 ± 0.04
\boldsymbol{R}	0.02232	1×10 s	18.9	15.51 ± 0.06
\boldsymbol{R}	0.02760	1×10 s	12.1	15.88 ± 0.09
\boldsymbol{R}	0.03288	1×10 s	7.58	$16.0^{+0.2}_{-0.1}$
\boldsymbol{R}	0.04032	1×20 s	10.7	16.3 ± 0.1
R	0.04824	1×20 s	9.35	16.4 ± 0.1
R	0.06720	1×40 s	7.45	$16.5^{+0.2}_{-0.1}$
\boldsymbol{R}	0.08088	1×40 s	13.1	$16.68^{+0.09}_{-0.08}$
\boldsymbol{R}	0.10824	1×40 s	5.47	16.9 ± 0.2
R	0.18216	1×80 s	5.82	17.5 ± 0.2
\boldsymbol{R}	0.20880	1×80 s	3.58	17.0 ± 0.3
\boldsymbol{R}	0.24744	1×160 s	10.9	17.4 ± 0.1
\boldsymbol{R}	0.35520	1×160 s	5.37	17.8 ± 0.2
\boldsymbol{R}	0.40536	$1\times160\,\mathrm{s}$	6.08	17.6 ± 0.2
R	0.50928	1×160 s	2.86	$17.9^{+0.4}_{-0.3}$
\boldsymbol{R}	0.55584	1×160 s	6.08	17.5 ± 0.2
\boldsymbol{R}	0.66000	1×160 s	5.53	17.6 ± 0.2
\boldsymbol{R}	0.91992	3×160 s	5.79	18.4 ± 0.2
\boldsymbol{R}	1.28712	4×160 s	6.64	18.79 ± 0.2
R	1.84008	9×160 s	3.68	19.3 ± 0.3
\boldsymbol{R}	2.79768	7×160 s	6.72	$19.5^{+0.2}_{-0.1}$
\boldsymbol{R}	3.21240	7×160 s	8.31	19.8 ± 0.1
\boldsymbol{R}	3.59184	7×160 s	10.1	19.8 ± 0.1
\boldsymbol{R}	4.12824	12×160 s	10.7	19.8 ± 0.1
\boldsymbol{R}	4.9908	18×160 s	7.18	20.0 ± 0.1
\boldsymbol{R}	24.5124	40×160 s	1.94	$21.9^{+0.7}_{-0.4}$

Table A2. Skynet - Filter ${\cal B}$

Table A3. Skynet - Filter $\cal I$

Table A5. Skynet - Filter r'

Table A4. Skynet - Filter g'

