

## GENERALIZED HERMITE-HADAMARD TYPE INEQUALITIES INVOLVING LOCAL FRACTIONAL INTEGRALS

HUIXIA MO

School of Science, Beijing University of Posts and Telecommunications, Beijing, 100876, P. R. China  
Email: [huxmo@bupt.edu.cn](mailto:huxmo@bupt.edu.cn)

In the paper, two new identities involving the local fractional integrals have been established. Using these two identities, we obtain some generalized Hermite-Hadamard type integral inequalities for the local differentiable generalized convex functions.

**Key words:** fractal space, generalized Hadamard's inequality, local fractional derivative, local fractional integral

### 1. INTRODUCTION

The convex function is an important concept in the class mathematical analysis course. And many important inequalities are established for the class of convex functions. For example, the Hermite-Hadamard's inequality is one of the best known results in the literature, which can be stated as follows.

**Hermite-Hadamard's inequality[1]:** Let  $f$  be a convex function on  $[a,b]$  with  $a < b$ . If  $f$  is integral on  $[a,b]$ , then

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a)+f(b)}{2}. \quad (1.1)$$

Hadamard's inequality for convex functions has received renewed attention in recent years and a remarkable variety of refinements and generalizations have been found(see [2-6] and so on). In [2], Dragomir and Agarwal proved the following results which connected with the right part of (1.1).

**THEOREM A.** [2] Let  $I \subseteq R$  be an interval. Suppose that  $f : I^0 \subseteq R \rightarrow R$  ( $I^0$  is the interior of  $I$ ) is differentiable. If  $f' \in L[a,b]$  and  $|f'|$  is convex on  $[a,b]$  for  $a, b \in I^0$  with  $a < b$ , then

$$\left| \frac{f(a)+f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{b-a}{8} (|f'(a)| + |f'(b)|). \quad (1.2)$$

**THEOREM B.** [2] Let  $f : I^0 \subseteq R \rightarrow R$  be a differentiable mapping on  $I^0$ . If  $f' \in L[a,b]$  and  $|f'|^{p/(p-1)}$  is convex on  $[a,b]$  for  $p > 1$  and  $a, b \in I^0$  with  $a < b$ , then

$$\begin{aligned} & \left| \frac{f(a)+f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ & \leq \frac{b-a}{2(1+p)^{1/p}} \left[ \frac{|f'(a)|^{p/(p-1)} + |f'(b)|^{p/(p-1)}}{2} \right]^{(p-1)/p}. \end{aligned} \quad (1.3)$$

In [5], the above inequalities were generalized as follows.

**THEOREM C.** [5] Let  $f : I^0 \subseteq R \rightarrow R$  be a differentiable mapping on  $I^0$ . If  $f' \in L[a,b]$  and  $|f'|^q$  is convex on  $[a,b]$  for  $q \geq 1$  and  $a, b \in I^0$  with  $a < b$ , then the following equality holds:

$$\left| \frac{f(a)+f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{b-a}{4} \left[ \frac{|f'(a)|^q + |f'(b)|^q}{2} \right]^{1/q},$$

$$\left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{b-a}{4} \left[ \frac{|f'(a)|^q + |f'(b)|^q}{2} \right]^{1/q}.$$

In recent years, the fractal theory has received significantly remarkable attention [7-11]. In [9], the authors introduce the generalized convex function and establish the generalized Hermite-Hadamard's inequality on fractal space.

**THEOREM D. [9] (Generalized Hermite-Hadamard's inequality)** Let  $f(x) \in {}_a I_b^{(\alpha)}$  be a generalized convex function on  $[a, b]$  with  $a < b$ . Then

$$f\left(\frac{a+b}{2}\right) \leq \frac{\Gamma(1+\alpha)}{(b-a)^\alpha} {}_a I_b^{(\alpha)} f(x) \leq \frac{f(a)+f(b)}{2^\alpha}.$$

Inspired by these investigations, in the paper we will establish the generalized Hermite-Hadamard type integral inequalities for the local differentiable convex functions on fractal space.

## 2. PRELIMINARIES

Recall the set  $R^\alpha$  of real line numbers and use the Gao-Yang-Kang's idea to describe the definitions of the local fractional derivative and local fractional integral, see [10,11] and so on.

Recently, the theory of Yang's fractional sets [10] was introduced as follows.

For  $0 < \alpha \leq 1$ , we have the following  $\alpha$ -type set of element sets:

$Z^\alpha$  : The  $\alpha$ -type set of the integer is defined as the set  $\{0^\alpha, \pm 1^\alpha, \pm 2^\alpha, \dots, \pm n^\alpha, \dots\}$ .

$Q^\alpha$  : The  $\alpha$ -type set of the rational numbers is defined as the set  $\{m^\alpha = (p/q)^\alpha : p, q \in Z, q \neq 0\}$ .

$J^\alpha$  : The  $\alpha$ -type set of the irrational numbers is defined as the set  $\{m^\alpha \neq (p/q)^\alpha : p, q \in Z, q \neq 0\}$ .

$R^\alpha$  : The  $\alpha$ -type set of the real line numbers is defined as the set  $R^\alpha = Q^\alpha \cup J^\alpha$ .

If  $a^\alpha, b^\alpha$  and  $c^\alpha$  belong to the set  $R^\alpha$  of real line numbers, then

- |   |   |
|---|---|
| (1) $a^\alpha + b^\alpha$ and $a^\alpha b^\alpha$ belong to the set $R^\alpha$ ;                                    | (2) $a^\alpha + b^\alpha = b^\alpha + a^\alpha = (a+b)^\alpha = (b+a)^\alpha$ ; |
| (3) $a^\alpha + (b^\alpha + c^\alpha) = (a+b)^\alpha + c^\alpha$ ;  | (4) $a^\alpha b^\alpha = b^\alpha a^\alpha = (ab)^\alpha = (ba)^\alpha$ ;       |
| (5) $a^\alpha (b^\alpha c^\alpha) = (a^\alpha b^\alpha) c^\alpha$ ;   | (6) $a^\alpha (b^\alpha + c^\alpha) = a^\alpha b^\alpha + a^\alpha c^\alpha$ ;  |
| (7) $a^\alpha + 0^\alpha = 0^\alpha + a^\alpha = a^\alpha$ and $a^\alpha 1^\alpha = 1^\alpha a^\alpha = a^\alpha$ . |   |

The definitions of the local fractional derivative and local fractional can be given as follows.

**DEFINITION 2.1.[10]** A non-differentiable function  $f : R \rightarrow R^\alpha$ ,  $x \rightarrow f(x)$  is called to be local fractional continuous at  $x_0$ , if for any  $\varepsilon > 0$ , there exists  $\delta > 0$ , such that

$$|f(x) - f(x_0)| < \varepsilon^\alpha$$

holds for  $|x - x_0| < \delta$ , where  $\varepsilon, \delta \in R$ . If  $f(x)$  is local fractional continuous on the interval  $(a, b)$ , we denote  $f(x) \in C_\alpha(a, b)$ .

**DEFINITION 2.2.**[10] The local fractional derivative of  $f(x)$  of order  $\alpha$  at  $x = x_0$  is defined by

$$f^{(\alpha)}(x_0) = \frac{d^\alpha f(x)}{dx^\alpha} \Big|_{x=x_0} = \lim_{x \rightarrow x_0} \frac{\Delta^\alpha(f(x) - f(x_0))}{(x - x_0)^\alpha},$$

where  $\Delta^\alpha(f(x) - f(x_0)) \cong \Gamma(1+\alpha)(f(x) - f(x_0))$ .

If there exists  $f^{((k+1)\alpha)}(x) = \overbrace{D_x^\alpha \cdots D_x^\alpha}^{k+1 \text{ times}} f(x)$  for any  $x \in I \subseteq R$ , then we denoted  $f \in D_{(k+1)\alpha}(I)$ , where  $k = 0, 1, 2, \dots$ .

**DEFINITION 2.3.**[10] Let  $f \in C_\alpha[a, b]$ . Then the local fractional integral is defined by,

$${}_a I_b^{(\alpha)} f(x) = \frac{1}{\Gamma(1+\alpha)} \int_a^b f(t) (dt)^\alpha = \frac{1}{\Gamma(1+\alpha)} \lim_{\Delta t \rightarrow 0} \sum_{j=0}^{N-1} f(t_j) (\Delta t_j)^\alpha,$$

with  $\Delta t_j = t_{j+1} - t_j$  and  $\Delta t = \max\{\Delta t_1, \Delta t_2, \Delta t_3, \dots\}$ , where  $[t_j, t_{j+1}]$ ,  $j = 0, \dots, N-1$  and  $t_0 = a < t_1 < \dots < t_i < \dots < t_{N-1} < t_N = b$  is a partition of the interval  $[a, b]$ .

Here, it follows that  ${}_a I_a^{(\alpha)} f(x) = 0$  if  $a = b$  and  ${}_a I_b^{(\alpha)} f(x) = - {}_b I_a^{(\alpha)} f(x)$  if  $a < b$ . If for any  $x \in [a, b]$ , there exists  ${}_a I_x^{(\alpha)} f(x)$ , then we denoted by  $f(x) \in I_x^{(\alpha)}[a, b]$ .

**DEFINITION 2.4.** [9] Let  $f : I \subseteq R \rightarrow R^\alpha$ . For any  $x_1, x_2 \in I$  and  $\lambda \in [0, 1]$ , if the following inequality

$$f(\lambda x_1 + (1-\lambda)x_2) \leq \lambda^\alpha f(x_1) + (1-\lambda)^\alpha f(x_2)$$

holds, then  $f$  is called a generalized convex function on  $I$ .

Here are two basic examples of generalized convex functions:

$$(1) f(x) = x^{\alpha p}, x \geq 0, p > 1;$$

$$(2) f(x) = E_\alpha(x^\alpha), x \in R, \text{ where } E_\alpha(x^\alpha) = \sum_{k=0}^{\infty} \frac{x^{\alpha k}}{\Gamma(1+k\alpha)} \text{ is the Mittag-Leffler function.}$$

### LEMMA 2.1.[10]

(1) (**Local fractional integration is anti-differentiation**) Suppose that  $f(x) = g^{(\alpha)}(x) \in C_\alpha[a, b]$ , then we have

$${}_a I_b^\alpha f(x) = g(b) - g(a).$$

(2) (**Local fractional integration by parts**) Suppose that  $f(x), g(x) \in D_\alpha[a, b]$  and  $f^{(\alpha)}(x), g^{(\alpha)}(x) \in C_\alpha[a, b]$ , then we have

$${}_a I_b^\alpha f(x) g^{(\alpha)}(x) = f(x) g(x) \Big|_a^b - {}_a I_b^\alpha f^{(\alpha)}(x) g(x).$$

### LEMMA 2.2.[10]

$$\frac{d^\alpha x^{k\alpha}}{dx^\alpha} = \frac{\Gamma(1+k\alpha)}{\Gamma(1+(k-1)\alpha)} x^{(k-1)\alpha}; \quad \frac{1}{\Gamma(1+\alpha)} \int_a^b x^{k\alpha} (dx)^\alpha = \frac{\Gamma(1+k\alpha)}{\Gamma(1+(k+1)\alpha)} (b^{(k+1)\alpha} - a^{(k+1)\alpha}), \quad k \in R.$$

**LEMMA 2.3.[10] (Generalized Hölder's inequality)** Let  $f, g \in C_\alpha[a, b]$ ,  $p, q > 1$  with  $1/p + 1/q = 1$ , then

$$\frac{1}{\Gamma(1+\alpha)} \int_a^b |f(x)g(x)| (dx)^\alpha \leq \left( \frac{1}{\Gamma(1+\alpha)} \int_a^b |f(x)|^p (dx)^\alpha \right)^{1/p} \left( \frac{1}{\Gamma(1+\alpha)} \int_a^b |g(x)|^q (dx)^\alpha \right)^{1/q}.$$

### 3. MAIN RESULTS

**THEOREM 1.** Let  $I \subseteq R$  be an interval,  $f : I^0 \subseteq R \rightarrow R^\alpha$  ( $I^0$  is the interior of  $I$ ) such that  $f \in D_\alpha(I^\circ)$  and  $f^{(\alpha)} \in C_\alpha[a, b]$  for  $a, b \in I^0$  with  $a < b$ . Then the following equality holds:

$$\frac{f(a) + f(b)}{2^\alpha} - \frac{\Gamma(1+\alpha)}{(b-a)^\alpha} {}_a I_b^\alpha f(x) = \frac{(b-a)^\alpha}{2^\alpha} \frac{1}{\Gamma(1+\alpha)} \int_0^1 (1-2t)^\alpha f^{(\alpha)}(ta + (1-t)b) (dt)^\alpha.$$

**Proof:** From the local fractional integration by parts, it suffices to note that

$$\begin{aligned} & \frac{1}{\Gamma(1+\alpha)} \int_0^1 (1-2t)^\alpha f^{(\alpha)}(ta + (1-t)b) (dt)^\alpha \\ &= (1-2t)^\alpha \frac{f(ta + (1-t)b)}{(a-b)^\alpha} \Big|_0^1 - \frac{1}{(a-b)^\alpha} \frac{1}{\Gamma(1+\alpha)} \int_0^1 (-2)^\alpha \Gamma(1+\alpha) f(ta + (1-t)b) (dt)^\alpha. \\ &= \frac{f(a) + f(b)}{(b-a)^\alpha} - \frac{2^\alpha}{(b-a)^{2\alpha}} \frac{\Gamma(1+\alpha)}{\Gamma(1+\alpha)} \int_a^b f(x) (dx)^\alpha \\ &= \frac{f(a) + f(b)}{(b-a)^\alpha} - \frac{2^\alpha \Gamma(1+\alpha)}{(a-b)^{2\alpha}} {}_a I_b^\alpha f(x). \end{aligned}$$

Thus,  $\frac{(b-a)^\alpha}{2^\alpha} \frac{1}{\Gamma(1+\alpha)} \int_0^1 (1-2t)^\alpha f^{(\alpha)}(ta + (1-t)b) (dt)^\alpha = \frac{f(a) + f(b)}{2^\alpha} - \frac{\Gamma(1+\alpha)}{(b-a)^\alpha} {}_a I_b^\alpha f(x)$ .

**THEOREM 2.** Suppose that the assumptions of Theorem 1 are satisfied, then

$$f\left(\frac{a+b}{2}\right) - \frac{\Gamma(1+\alpha)}{(b-a)^\alpha} {}_a I_b^\alpha f(x) = \frac{1}{(b-a)^\alpha} \frac{1}{\Gamma(1+\alpha)} \int_a^b S(x) f^{(\alpha)}(x) (dx)^\alpha,$$

where

$$S(x) = \begin{cases} (x-a)^\alpha, & x \in [a, \frac{a+b}{2}], \\ (x-b)^\alpha, & x \in [\frac{a+b}{2}, b]. \end{cases}$$

**Proof.** From the definition of  $S(x)$ , we have

$$\frac{1}{(b-a)^\alpha} \frac{1}{\Gamma(1+\alpha)} \int_a^b S(x) f^{(\alpha)}(x) (dx)^\alpha$$

$= \frac{1}{(b-a)^\alpha} \frac{1}{\Gamma(1+\alpha)} \int_a^{\frac{a+b}{2}} (x-a)^\alpha f^{(\alpha)}(x)(dx)^\alpha + \frac{1}{(b-a)^\alpha} \frac{1}{\Gamma(1+\alpha)} \int_{\frac{a+b}{2}}^b (x-b)^\alpha f^{(\alpha)}(x)(dx)^\alpha$ . Firstly, let's estimate the first part.

In fact, using the local fractional integration by parts, we have

$$\frac{1}{\Gamma(1+\alpha)} \int_a^{\frac{a+b}{2}} (x-a)^\alpha f^{(\alpha)}(x)(dx)^\alpha = \left( \frac{b-a}{2} \right)^\alpha f\left( \frac{a+b}{2} \right) - \Gamma(1+\alpha) {}_a I_{\frac{a+b}{2}}^\alpha f(x).$$

Secondly, using the local fractional integration by parts, we have

$$\frac{1}{\Gamma(1+\alpha)} \int_{\frac{a+b}{2}}^b (x-b)^\alpha f^{(\alpha)}(x)(dx)^\alpha = \left( \frac{b-a}{2} \right)^\alpha f\left( \frac{a+b}{2} \right) - \Gamma(1+\alpha) {}_{\frac{a+b}{2}} I_b^\alpha f(x).$$

Combining the above formulas, we obtain

$$\frac{1}{(b-a)^\alpha} \frac{1}{\Gamma(1+\alpha)} \int_a^b S(x) f^{(\alpha)}(x)(dx)^\alpha = f\left( \frac{a+b}{2} \right) - \frac{\Gamma(1+\alpha)}{(b-a)^\alpha} {}_a I_b^\alpha f(x).$$

That is

$$f\left( \frac{a+b}{2} \right) - \frac{\Gamma(1+\alpha)}{(b-a)^\alpha} {}_a I_b^\alpha f(x) = \frac{1}{(b-a)^\alpha} \frac{1}{\Gamma(1+\alpha)} \int_a^b S(x) f^{(\alpha)}(x)(dx)^\alpha.$$

**THEOREM 3.** Let  $p, q \geq 1$  with  $1/p + 1/q = 1$ . Let  $f : I^0 \subseteq R \rightarrow R^\alpha$ , such that  $f \in D_\alpha(I^\circ)$  and  $|f^{(\alpha)}|^q \in C_\alpha[a, b]$  for  $a, b \in I^0$  with  $a < b$ . If  $|f^{(\alpha)}|^q$  is generalized convex on  $[a, b]$ , then

$$\begin{aligned} & \left| \frac{f(a)+f(b)}{2^\alpha} - \frac{\Gamma(1+\alpha)}{(b-a)^\alpha} {}_a I_b^\alpha f(x) \right| \\ & \leq \frac{(b-a)^\alpha}{2^\alpha} \left[ |f^{(\alpha)}(a)|^q + |f^{(\alpha)}(b)|^q \right]^{1/q} \left( \frac{\Gamma(1+\alpha)}{\Gamma(1+2\alpha)} \right)^{1/p} \left[ \left( \frac{3}{2} \right)^\alpha \frac{\Gamma(1+2\alpha)}{\Gamma(1+3\alpha)} - \left( \frac{1}{2} \right)^\alpha \frac{\Gamma(1+\alpha)}{\Gamma(1+2\alpha)} \right]^{1/q}. \end{aligned}$$

**Proof.** From Theorem 1 and the generalized Hölder's inequality (Lemma 3), it follows that

$$\begin{aligned} & \left| \frac{f(a)+f(b)}{2^\alpha} - \frac{\Gamma(1+\alpha)}{(b-a)^\alpha} {}_a I_b^\alpha f(x) \right| \\ & \leq \frac{(b-a)^\alpha}{2^\alpha} \frac{1}{\Gamma(1+\alpha)} \int_0^1 |(1-2t)^\alpha| \|f^{(\alpha)}(ta+(1-t)b)\| (dt)^\alpha \\ & \leq \frac{(b-a)^\alpha}{2^\alpha} \left( \frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^\alpha (dt)^\alpha \right)^{1/p} \left[ \frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^\alpha |f^{(\alpha)}(ta+(1-t)b)|^q (dt)^\alpha \right]^{1/q}. \quad (3.1) \end{aligned}$$

From Lemma 2, it's easy to see that

$$\frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^\alpha (dt)^\alpha = \frac{\Gamma(1+\alpha)}{\Gamma(1+2\alpha)}. \quad (3.2)$$

Since  $|f^{(\alpha)}|^q$  is generalized convex on  $[a, b]$ , thus

$$\begin{aligned} & \frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^\alpha |f^{(\alpha)}(ta+(1-t)b)|^q (dt)^\alpha \\ & \leq \frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^\alpha [t^\alpha |f^{(\alpha)}(a)|^q + (1-t)^\alpha |f^{(\alpha)}(b)|^q] (dt)^\alpha \\ & = \left[ |f^{(\alpha)}(a)|^q + |f^{(\alpha)}(b)|^q \right] \frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^\alpha t^\alpha (dt)^\alpha. \end{aligned} \quad (3.3)$$

It is obvious that

$$\frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^\alpha t^\alpha (dt)^\alpha = -\frac{\Gamma(1+\alpha)}{\Gamma(1+2\alpha)} \left(\frac{1}{2}\right)^\alpha + \frac{\Gamma(1+2\alpha)}{\Gamma(1+3\alpha)} \left(\frac{3}{2}\right)^\alpha. \quad (3.4)$$

Combining the estimates of (3.1)-(3.4), we have

$$\begin{aligned} & \left| \frac{f(a)+f(b)}{2^\alpha} - \frac{\Gamma(1+\alpha)}{(b-a)^\alpha} {}_a I_b^\alpha f(x) \right| \\ & \leq \frac{(b-a)^\alpha}{2^\alpha} \left[ |f^{(\alpha)}(a)|^q + |f^{(\alpha)}(b)|^q \right]^{1/q} \left( \frac{\Gamma(1+\alpha)}{\Gamma(1+2\alpha)} \right)^{1/p} \left[ \frac{\Gamma(1+2\alpha)}{\Gamma(1+3\alpha)} \left(\frac{3}{2}\right)^\alpha - \frac{\Gamma(1+\alpha)}{\Gamma(1+2\alpha)} \left(\frac{1}{2}\right)^\alpha \right]^{1/q}. \end{aligned}$$

**THEOREM 4.** Suppose that the assumptions of Theorem 3 are satisfied, then

$$\left| f\left(\frac{a+b}{2}\right) - \frac{\Gamma(1+\alpha)}{(b-a)^\alpha} {}_a I_b^\alpha f(x) \right| \leq \frac{(b-a)^\alpha}{4^\alpha} \left[ \frac{2^\alpha \Gamma(1+\alpha)}{\Gamma(1+2\alpha)} \right]^{1/p} \left( \frac{\Gamma(1+\alpha)[|f^{(\alpha)}(a)|^q + |f^{(\alpha)}(b)|^q]}{\Gamma(1+2\alpha)} \right)^{1/q}.$$

**Proof.** Similar to the estimate of Theorem 3, from Theorem 2 and Lemma 3, it follows that

$$\begin{aligned} & \left| f\left(\frac{a+b}{2}\right) - \frac{\Gamma(1+\alpha)}{(b-a)^\alpha} {}_a I_b^\alpha f(x) \right| \\ & \leq \frac{1}{(b-a)^\alpha} \frac{1}{\Gamma(1+\alpha)} \int_a^b |S(x)| |f^{(\alpha)}(x)| (dx)^\alpha \\ & \leq \frac{1}{(b-a)^\alpha} \left[ \frac{1}{\Gamma(1+\alpha)} \int_a^b |S(x)| (dx)^\alpha \right]^{1/p} \left[ \frac{1}{\Gamma(1+\alpha)} \int_a^b |S(x)| |f^{(\alpha)}(x)|^q (dx)^\alpha \right]^{1/q}. \end{aligned}$$

Let us estimate the first integral.

Using the expression of function  $S(x)$ , we can calculate that

$$\frac{1}{\Gamma(1+\alpha)} \int_a^b |S(x)| (dx)^\alpha = \frac{\Gamma(1+\alpha)}{\Gamma(1+2\alpha)} 2^\alpha \left(\frac{b-a}{2}\right)^{2\alpha}. \quad (3.5)$$

Secondly, for the another part, we have

$$\frac{1}{\Gamma(1+\alpha)} \int_a^b |S(x)| |f^{(\alpha)}(x)|^q (dx)^\alpha$$

$$= \frac{1}{\Gamma(1+\alpha)} \int_a^{\frac{a+b}{2}} (x-a)^\alpha |f^{(\alpha)}(x)|^q (dx)^\alpha + \frac{1}{\Gamma(1+\alpha)} \int_{\frac{a+b}{2}}^b (b-x)^\alpha |f^{(\alpha)}(x)|^q (dx)^\alpha.$$

Since  $|f^{(\alpha)}|^q$  is generalized convex on  $[a, b]$ , thus we have

$$\begin{aligned} & \frac{1}{\Gamma(1+\alpha)} \int_a^{\frac{a+b}{2}} (x-a)^\alpha |f^{(\alpha)}(x)|^q (dx)^\alpha \\ & \leq \frac{1}{\Gamma(1+\alpha)} \int_0^1 \left( \frac{b-a}{2} \right)^{2\alpha} t^\alpha \left| f^{(\alpha)} \left( \frac{a+b}{2} t + (1-t)a \right) \right|^q (dt)^\alpha \\ & \leq \frac{1}{\Gamma(1+\alpha)} \int_0^1 \left( \frac{b-a}{2} \right)^{2\alpha} t^\alpha \left[ t^\alpha \left| f^{(\alpha)} \left( \frac{a+b}{2} \right) \right|^q + (1-t)^\alpha |f^{(\alpha)}(a)|^q \right] (dt)^\alpha \\ & \leq \left( \frac{b-a}{2} \right)^{2\alpha} \left\{ \frac{|f^{(\alpha)}(a)|^q + |f^{(\alpha)}(b)|^q}{2^\alpha} \frac{\Gamma(1+2\alpha)}{\Gamma(1+3\alpha)} + |f^{(\alpha)}(a)|^q \left[ \frac{\Gamma(1+\alpha)}{\Gamma(1+2\alpha)} - \frac{\Gamma(1+2\alpha)}{\Gamma(1+3\alpha)} \right] \right\}. \end{aligned} \quad (3.6)$$

Similarly, we have

$$\begin{aligned} & \frac{1}{\Gamma(1+\alpha)} \int_{\frac{a+b}{2}}^b (b-x)^\alpha |f^{(\alpha)}(x)|^q (dx)^\alpha \\ & \leq \left( \frac{b-a}{2} \right)^{2\alpha} \left\{ \frac{|f^{(\alpha)}(a)|^q + |f^{(\alpha)}(b)|^q}{2^\alpha} \frac{\Gamma(1+2\alpha)}{\Gamma(1+3\alpha)} + |f^{(\alpha)}(b)|^q \left[ \frac{\Gamma(1+\alpha)}{\Gamma(1+2\alpha)} - \frac{\Gamma(1+2\alpha)}{\Gamma(1+3\alpha)} \right] \right\}. \end{aligned} \quad (3.7)$$

So, from (3.6) and (3.7), we get

$$\frac{1}{\Gamma(1+\alpha)} \int_a^b |S(x)| |f^{(\alpha)}(x)|^q (dx)^\alpha = \left( \frac{b-a}{2} \right)^{2\alpha} \left( |f^{(\alpha)}(a)|^q + |f^{(\alpha)}(b)|^q \right) \frac{\Gamma(1+\alpha)}{\Gamma(1+2\alpha)}. \quad (3.8)$$

Thus, combining (3.5) and (3.8), we obtain

$$\left| f \left( \frac{a+b}{2} \right) - \frac{\Gamma(1+\alpha)}{(b-a)^\alpha} {}_a I_b^\alpha f(x) \right| = \frac{(b-a)^\alpha}{4^\alpha} \left[ \frac{2^\alpha \Gamma(1+\alpha)}{\Gamma(1+2\alpha)} \right]^{1/p} \left( \frac{\Gamma(1+\alpha)[|f^{(\alpha)}(a)|^q + |f^{(\alpha)}(b)|^q]}{\Gamma(1+2\alpha)} \right)^{1/q}.$$

#### 4. APPLICATIONS TO SOME SPECIAL MEANS

As in [5], in the paper we consider some generalized means. We take

$$A(a, b) = \frac{a^\alpha + b^\alpha}{2^\alpha}; \quad L_n(a, b) = \left[ \frac{\Gamma(1+n\alpha)}{\Gamma(1+(n+1)\alpha)} (b^{(n+1)\alpha} - a^{(n+1)\alpha}) \right]^{1/n}, \quad n \in \mathbb{Z} \setminus \{-1, 0\}, \quad a, b \in \mathbb{R}, \quad a \neq b.$$

**Proposition 1.** Let  $a, b \in \mathbb{R}$ ,  $a < b$ ,  $0 \notin [a, b]$  and  $n \in \mathbb{Z}$ ,  $|n| \geq 2$ . Then for all  $p, q \geq 1$  with  $1/p + 1/q = 1$ ,

$$\begin{aligned} & \left| A(a^n, b^n) - \frac{\Gamma(1+\alpha)}{(b-a)^\alpha} (L_n(a, b))^n \right| \\ & \leq \frac{(b-a)^\alpha}{2^\alpha} \left[ \left( \frac{\Gamma(1+n\alpha)}{\Gamma(1+(n-1)\alpha)} \right)^q (a^{(n-1)q\alpha} + b^{(n-1)q\alpha}) \right]^{1/q} \left( \frac{\Gamma(1+\alpha)}{\Gamma(1+2\alpha)} \right)^{1/p} \left[ \frac{\Gamma(1+2\alpha)}{\Gamma(1+3\alpha)} \left( \frac{3}{2} \right)^\alpha - \frac{\Gamma(1+\alpha)}{\Gamma(1+2\alpha)} \left( \frac{1}{2} \right)^\alpha \right]^{1/q}; \end{aligned}$$

$$\left| \left( A(a,b) \right)^n - \frac{\Gamma(1+\alpha)}{(b-a)^\alpha} (L_n(a,b))^n \right| \leq \frac{(b-a)^\alpha}{4^\alpha} \left[ \frac{2^\alpha \Gamma(1+\alpha)}{\Gamma(1+2\alpha)} \right]^{1/p} \left[ \frac{\Gamma(1+\alpha)}{\Gamma(1+2\alpha)} \left( \frac{\Gamma(1+n\alpha)}{\Gamma(1+(n-1)\alpha)} \right)^q (a^{(n-1)q\alpha} + b^{(n-1)q\alpha}) \right]^{1/q}.$$

**Proof.** The proof is from Theorem 3 and Theorem 4 with  $f(x) = x^{n\alpha}$ ,  $x \in R$ ,  $n \in Z$ ,  $n \geq 2$ .

$$\begin{aligned} \left| A(a^n, b^n) - \frac{\Gamma(1+\alpha)}{(b-a)^\alpha} (L_n(a,b))^n \right| &= \left| \frac{f(a) + f(b)}{2^\alpha} - \frac{\Gamma(1+\alpha)}{(b-a)^\alpha} {}_a I_b^\alpha f(x) \right| \\ &\leq \frac{(b-a)^\alpha}{2^\alpha} \left[ \left( \frac{\Gamma(1+n\alpha)}{\Gamma(1+(n-1)\alpha)} \right)^q (a^{(n-1)q\alpha} + b^{(n-1)q\alpha}) \right]^{1/q} \left( \frac{\Gamma(1+2\alpha)}{\Gamma(1+3\alpha)} \left( \frac{3}{2} \right)^\alpha - \frac{\Gamma(1+\alpha)}{\Gamma(1+2\alpha)} \left( \frac{1}{2} \right)^\alpha \right)^{1/q}. \end{aligned}$$

And,

$$\begin{aligned} \left| \left( A(a,b) \right)^n - \frac{\Gamma(1+\alpha)}{(b-a)^\alpha} (L_n(a,b))^n \right| &= \left| f\left(\frac{a+b}{2}\right) - \frac{\Gamma(1+\alpha)}{(b-a)^\alpha} {}_a I_b^\alpha f(x) \right| \\ &\leq \frac{(b-a)^\alpha}{4^\alpha} \left[ \frac{2^\alpha \Gamma(1+\alpha)}{\Gamma(1+2\alpha)} \right]^{1/p} \left[ \frac{\Gamma(1+\alpha)}{\Gamma(1+2\alpha)} \left( \frac{\Gamma(1+n\alpha)}{\Gamma(1+(n-1)\alpha)} \right)^q (a^{(n-1)q\alpha} + b^{(n-1)q\alpha}) \right]^{1/q}. \end{aligned}$$

**Remark:** when  $\alpha = 1$ , this result reduce to Proposition 1 in [5].

## ACKNOWLEDGMENTS

This work has been supported by the National Natural Science Foundation of China (No. 11161042, 11471050).

## REFERENCES

- [1] D. S. MITRINOVIC AND I. B. LACKOVIC, *Hermite and convexity*, Aequationes Math., **28**, pp. 229-232 , 1985.
- [2] S. S. DRAGOMIR AND R. P. AGARWAL, *Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula*, Appl. Math. Lett., **11(5)**, pp. 91–95, 1998.
- [3] S. S. DRAGOMIR, *On some new inequalities of Hermite–Hadamard type for m-convex functions*, Tamk. J. Math., **3 (1)** , pp. 45-56, 2002.
- [4] Z. DAHMANI, L. TABHARIT AND S. TAF, *New generalizations of Gruss inequality usin Riemann–Liouville fractional integrals*, Bull. Math. Anal. Appl. **2 (3)**, pp. 93–99, 2010 .
- [5] C. E. M. PEARCE AND J. PECARIC, *Inequalities for differentiable mappings with application to specialmeans and quadrature formulae*, Appl. Math. Lett., **13(2)**, pp. 51-55, 2000.
- [6] M. Z. SARIKAYA, E. SET, H. YALDIZ AND N. BASAK, *Hermite–Hadamard’s inequalities for fractional integrals and related fractional inequalities*, Math. Comput. Model., **57 (9–10)**, pp. 2403–2407, 2013.
- [7] K. M. KOLWANKAR AND A. D. GANGAL, *Local Fractional Calculus: a Calculus for Fractal Space-time*, Fractals, Springer, London, PP.171-181, 1999.
- [8] K. GOLMANKHANEH AND D. BALEANU, *On a new measure on fractals*, J. Inequal. Appl., **522(1)**, pp. 1-9, 2013.
- [9] H. MO, X. SUI AND D. YU, *Generalized convex functions and some inequalities on fractal sets*, arXiv preprint arXiv: 1404.3964, 2014.
- [10] X. J. YANG, *Advanced Local Fractional Calculus and Its Applications*, World Science Publisher, New York, 2012.
- [11] J. YANG, D. BALEANU AND X. J. YANG, *Analysis of fractal wave equations by local fractional Fourier series method*, Adv. Math. Phys. , 2013 (2013), Article ID 632309