

Slope and curvature of the hadron vacuum polarization at vanishing virtuality from lattice QCD

Sz. Borsanyi,¹ Z. Fodor,^{1,2,3} T. Kawanai,³ S. Krieg,^{1,3} L. Lellouch,⁴
R. Malak,⁴ K. Miura,⁴ K.K. Szabo,^{1,3} C. Torrero,⁴ and B. Toth¹

¹*Department of Physics, Bergische Universität Wuppertal, Gausstr. 20, D-42119 Wuppertal*
²*Inst. for Theor. Physics, Eötvös University, Pázmány P. sét. 1/A, H-1117 Budapest, Hungary*
³*Jülich Supercomputing Centre, Forschungszentrum Jülich, D-52425 Jülich, Germany*
⁴*CNRS, Aix Marseille U., U. de Toulon, CPT, UMR 7332, F-13288 Marseille, France*
(Dated: September 17, 2018)

We compute the slope and curvature, at vanishing four-momentum transfer squared, of the leading order hadron vacuum polarization function, using lattice QCD. Calculations are performed with 2+1+1 flavors of staggered fermions directly at the physical values of the quark masses and in volumes of linear extent larger than 6 fm. The continuum limit is carried out using six different lattice spacings. All connected and disconnected contributions are calculated, up to and including those of the charm.

Introduction.— The vacuum expectation value of the product of two electromagnetic currents plays an important role in physics. It describes how virtual particle fluctuations polarize the vacuum as it is traversed by a propagating photon. While the contributions associated with virtual leptons and weak bosons can be computed in perturbation theory, those of quarks require nonperturbative methods for small photon virtuality, because of the confinement of quarks within hadrons. Here we focus on the latter, known as the hadron vacuum polarization or HVP.

The low energy behavior of the HVP is the limiting uncertainty in the standard model (SM) prediction of a number of quantities. It limits the precision with which many electroweak observables are determined [1]. It also represents the leading hadronic uncertainty in the SM prediction for the anomalous magnetic moments of leptons, a_ℓ with $\ell = e, \mu, \tau$ [2, 3]. In fact, it is the limiting factor in the SM prediction [2–6] of the much debated anomalous magnetic moment of the muon that is currently measured to 0.54 ppm [7].

Today the HVP is best determined using dispersion relations and the cross section of e^+e^- to hadrons or the rate of hadronic τ decays [4–6, 8]. However, since the pioneering work of [9], lattice QCD calculations of the leading order (LO) HVP contributions, $a_\mu^{\text{LO-HVP}}$, to a_μ have made significant progress [10–18]. Moreover, in the long run, this approach is likely to represent the most cost-effective way to increase the precision of the HVP to the levels that will soon be required by the new round of measurements of a_μ [19, 20] and, more generally, by particle physics phenomenology.

Here we present a full lattice QCD calculation of the first two derivatives of the HVP function at zero, euclidean virtuality. The calculation includes all contributions from u, d, s and c quarks, both in their quark-connected and quark-disconnected configurations, in the isospin limit. As shown in [21, 22], the slope of the polarization function provides an upper bound on the

HVP contribution to the anomalous magnetic moment of all three leptons. It also determines the whole of $a_\ell^{\text{LO-HVP}}/m_\ell^2$ in the limit that the lepton mass, m_ℓ , vanishes [21]. Moreover, together with the curvature, the slope gives $a_\mu^{\text{LO-HVP}}$ to within less than 2%. This fraction, which is indicative only, derives from comparing the value of $a_\mu^{\text{LO-HVP}}$ obtained from a simple phenomenological model to that obtained using a [1, 1] Padé approximant to describe the virtuality dependence of the HVP function. This approximant is constructed from the slope and curvature of the HVP function at vanishing virtuality, obtained in the same model. The model combines the e^+e^- experimental spectrum up to the ψ' that is compiled in [23], perturbative contributions above $s = 2.25 \text{ GeV}^2$ and dispersion relations. The use of Padé approximants for determining $a_\mu^{\text{LO-HVP}}$ was first proposed in [24] and was first used for a lattice QCD evaluation in [14].

The main challenge in calculating, on the lattice, derivatives of the HVP function at zero virtuality is the fact that it requires determining, with high precision, the dominant, and notoriously noisy, u and d quark contributions to the electromagnetic current correlation function at large euclidean distances. This is particularly clear when these derivatives are obtained by computing moments, in euclidean spacetime, of the quark, electromagnetic two-point function, as described below, around Eq. (1). In Fig. 1 we plot the kernels of two such time-moments. They correspond to those needed for the first two derivatives of the HVP function at vanishing virtuality (the higher derivatives are obtained from higher moments). As the figure shows, the distances that need to be reached to reliably determine the slope and curvature are above ~ 2 and ~ 4 fm, respectively. Higher derivatives require even larger distances. Here we address this challenge by performing a high-precision calculation on lattices of spatial extent $L \simeq 6$ fm and of time extent T , up to 11 fm. This calculation yields a prediction of QCD for these quantities that can be compared to present [25]




FIG. 1. Kernels of two moments of the spatial components of the zero three-momentum, electromagnetic current correlator as a function of euclidean time (see Eq. (1) with $\mu = i = 1, 2$ or 3 and $\nu = 0$). As described in the text, these moments can be used to determine the slope and curvature of the HVP function at vanishing virtuality. The electromagnetic current correlator is obtained using a phenomenological description of the e^+e^- data compiled in [23] and dispersion relations, as briefly described in the text.

and future phenomenological determinations. We leave to a forthcoming paper the computation of the anomalous magnetic moment of the muon.

Simulations.— We employ a tree-level improved Symanzik gauge action [26] and a fermion action for four flavors of stout-smear [27], staggered quarks. The up and down quark masses are treated as degenerate, their ratio to the strange quark mass is tuned to the vicinity of the physical point, which is defined from the Goldstone pion and kaon masses. The charm quark mass is fixed in units of the strange mass to $m_c/m_s = 11.85$ [28]. More information on the action together with simulation and algorithmic details can be found in [29].

To set the physical mass point we use the isospin corrected pion and kaon masses, $M_\pi = 134.8$ MeV and $M_K = 494.2$ MeV, from [30]. To convert the lattice results into physical units, we use the pion decay constant $f_\pi = 130.50(1)(3)(13)$ MeV [23] which is free of electromagnetic corrections and, to very good accuracy, equals to the decay constant in the $m_d = m_u$ limit [31]. This makes our definition of the physical point well defined in the isospin limit. In intermediate steps of the analysis we use the Wilson-flow-based [32] w_0 -scale [33]. For our finest lattice spacing the root mean squared pion mass is about 15% larger, than the Goldstone pion mass.

Table I lists the ensembles and the number of configurations used for quark-connected and disconnected measurements. A configuration corresponds to 10 unit length Rational Hybrid Monte Carlo (RHMC) [34] trajectories. The integration over the trajectory is improved with the gradient of the RHMC force [35, 36]. The topo-

β	a [fm]	$T \times L$	#conf-conn	#conf-disc
3.7000	0.134	64×48	1000	1000
3.7500	0.118	96×56	1500	1500
3.7753	0.111	84×56	1500	1500
3.8400	0.095	96×64	2500	1500
3.9200	0.078	128×80	3500	1000
4.0126	0.064	144×96	450	-

TABLE I. List of β , lattice spacings, sizes and number of configurations used for the connected and disconnected correlators.

logical charge undergoes sufficient number of tunnelings even on the finest lattices.

Observables.— The hadron vacuum polarization is derived from the electromagnetic current j_μ , which is defined as $j_\mu/e = \frac{2}{3}\bar{u}\gamma_\mu u - \frac{1}{3}\bar{d}\gamma_\mu d - \frac{1}{3}\bar{s}\gamma_\mu s + \frac{2}{3}\bar{c}\gamma_\mu c$, where e is the unit of electromagnetic charge. From this we build the current-current correlator $\langle j_\mu(x)j_\nu(0) \rangle$, in which we use the conserved lattice current at the source and sink. No renormalization is therefore necessary.

We split up the correlator in two different ways. First $\langle j_\mu j_\nu \rangle = \frac{e^2}{9}(5C_{\mu\nu}^l + C_{\mu\nu}^s + 4C_{\mu\nu}^c + C_{\mu\nu}^{\text{disc}})$, where the first three terms contain the connected contractions for the light, strange and charm flavors, and the last contains the disconnected contractions. Flavor mixing terms arise only in the latter. We can also separate the correlator according to isospin symmetry, which is exact in our simulations: $\langle j_\mu j_\nu \rangle = \langle j_\mu j_\nu \rangle_{I=0} + \langle j_\mu j_\nu \rangle_{I=1}$. Here the isospin singlet contribution is $\langle j_\mu j_\nu \rangle_{I=0} = \frac{e^2}{18}(C_{\mu\nu}^l + 2C_{\mu\nu}^s + 8C_{\mu\nu}^c + 2C_{\mu\nu}^{\text{disc}})$, whereas the isospin triplet one is $\langle j_\mu j_\nu \rangle_{I=1} = \frac{e^2}{2}C_{\mu\nu}^l$. The lowest energy state contains three/two pions in the isospin singlet/triplet channel. This fact determines the behavior of the correlator for large separations.

We calculate the connected contributions to the correlators using point sources. We use the all-mode-averaging technique (AMA) of [37] and 768 random source positions on each configuration for the light quarks, 64 sources for the strange (except at $\beta = 3.70$ where 128 were used) and 4 for the charm. To compute the quark-disconnected contributions, we apply AMA again, and exploit the approximate SU(3) flavor symmetry on around 6000 stochastic sources [15, 38]. We use random spatial wall sources, so only zero-momentum, time propagators are available. For the disconnected contribution of the charm we apply a hopping parameter expansion. The computer time required for this entire analysis is of the same order as the time needed for the generation of configurations.

The n -th coefficient of a Taylor expansion of the vacuum polarization scalar, $\Pi(Q^2)$, around $Q^2 = 0$ (i.e. $[\partial^n \Pi(Q^2)/(\partial Q^2)^n]_{Q^2=0}/n!$) can be written as $\Pi_n = \frac{1}{9}(5\Pi_n^l + \Pi_n^s + 4\Pi_n^c + \Pi_n^{\text{disc}})$, where each term is related to the respective, configuration-space correlator through




FIG. 2. Top: an upper/lower bound on Π_1^l is obtained by setting the timelike correlator to a two-pion decay/zero starting at a separation of t_c . Bottom: upper/lower bound on Π_1^{disc} is obtained by setting the correlator to zero/to the connected correlator with a two-pion decay. Results are for an ensemble at $\beta = 3.9200$.

moments [14, 39]

$$\Pi_{n,\mu\nu}^f = (-)^{n+1} \sum_x \frac{\hat{x}_\nu^{2n+2}}{(2n+2)!} C_{\mu\mu}^f(x) \quad (1)$$

for $f = \{l, s, c, \text{disc}\}$, with $\nu \neq \mu$ and \hat{x} defined as $\hat{x}_\nu = \min(x_\nu, L_\nu - x_\nu)$ and where L_μ is the size of the lattice in the μ -direction. In general the result depends on the choice of μ and ν . Three different averages, which are invariant under spatial cubic transformations, can be constructed: one which is an average over spatial moments, $\nu = 1, 2, 3$, of correlators of spatial currents, $\mu = 1, 2, 3 \neq \nu$; another, an average over spatial moments of correlators of timelike currents, $\mu = 4$; a third, an average over time moments, $\nu = 4$, of correlators of spatial currents. We call these averages $\Pi_{n,ss}$, $\Pi_{n,4s}$ and $\Pi_{n,s4}$, respectively. In the disconnected case we only have the $\Pi_{n,s4}$ average. The averages $\Pi_{n,ss}$, $\Pi_{n,4s}$ and $\Pi_{n,s4}$ can be different, which is a consequence of the finite lattice size and the asymmetry $T \neq L$.

In the case of the light and disconnected correlators the signal deteriorates quickly with increasing distance. In our analysis we consider the spatial moments, i.e. ss and $4s$, up to their full extent (ca. 3.1 fm). For the time-moment sums, $s4$, which can extend up to ca. 6 fm, we introduce a cut t_c in time. For times greater than t_c we replace the correlator by an upper and a lower

bound¹. The connected light correlator is proportional to the isospin triplet one, whose lowest-energy contribution comes from a two-pion state. Therefore up to exponentially suppressed corrections in T the correlator satisfies

$$0 \leq C^l(t) \leq C^l(t_c) \frac{\varphi(t)}{\varphi(t_c)}, \quad (2)$$

where $\varphi(t) = \cosh[E_{2\pi}(T/2 - t)]$ and $E_{2\pi}$ is the energy of two pions, each with the smallest nonvanishing lattice momentum, for which we use $2\pi/L$. Typically the two bounds agree for $t_c \gtrsim 3$ fm, as seen for example on the top plot of Fig. 2. A similar conclusion is reached for the second derivative $\Pi_{2,s4}^l$. In our analyses, we take $t_c = 3.1$ fm on the light connected timelike correlators and average the two bounds to get the final result. Pion-pion interactions can change the smallest two-pion momentum from $2\pi/L$ in that channel. Using the model of [40] and neglecting four-pion contributions, we determine the change in the momentum to be around 3%. We checked, that such a reduction of the momentum changes the result on Π^l by a small fraction of the statistical error.

The disconnected contribution alone can be constrained for large enough time separations, where the isospin singlet channel, dominated by three-pion states, can be neglected compared to the triplet, dominated by two-pion ones. Here we have

$$0 \geq [2C^s + 8C^c + 2C^{\text{disc}}](t) \geq -C^l(t_c) \frac{\varphi(t)}{\varphi(t_c)} \quad (3)$$

up to corrections exponentially suppressed in T . This gives an upper and a lower bound on $\Pi^s + 4\Pi^c + \Pi^{\text{disc}}$, which can be used to determine the time t_c after which the two bounds agree within errors. At large t , the connected strange and charm contributions in (3) are exponentially suppressed, and their presence does not make a difference when determining t_c so we neglect them. In the bottom plot of Fig. 2 we show the upper and lower bounds on Π^{disc} . In our analyses we average the two bounds for $t > t_c = 2.7$ fm.

Results. – To obtain our final results in the continuum limit and at the physical point, we fit the lattice results to a function which depends on the pion and kaon masses and on the lattice spacing squared a^2 . Since the simulations were done around the physical point, a linear pion/kaon mass dependence is always sufficient. For the light, strange and disconnected contributions reasonable fit qualities can be achieved with a linear a^2 dependence.

We begin with the light quark contribution Π^l , paying special attention to the difference between the averages Π_{ss}^l , Π_{4s}^l and Π_{s4}^l . Some ensemble show differences, but

¹ We are grateful to Christoph Lehner for a discussion of this issue.

	$\Pi_1[\text{GeV}^{-2}]$	$\Pi_2[\text{GeV}^{-4}]$
light	0.1653(17)(16)	-0.295(10)(7)
strange $\times 10^2$	6.57(1)(3)	-5.33(1)(4)
charm $\times 10^4$	40.3(2)(6)	-2.66(3)(11)
disconnected $\times 10^2$	-1.5(2)(1)	4.4(1.0)(0.4)
$I = 0$	0.0167(2)(2)	-0.018(1)(1)
$I = 1$	0.0827(8)(8)	-0.147(5)(4)
total	0.0993(10)(9)	-0.165(6)(4)
$I=1$ FV corr.	0.0006(23)	-0.016(10)
total + FV	0.0999(10)(9)(23)(13)	-0.181(6)(4)(10)(2)

TABLE II. Final results on the first and second derivative of the hadron vacuum polarization scalar at zero squared-momentum. The first four lines contain the continuum extrapolated lattice results. The next three lines are the isospin singlet, triplet and total contributions. The first error is statistical and the second is systematic. In the following line we give an estimate of the finite-volume effects of the $I = 1$ contribution using LO χ PT. In that approach, the finite-volume effects on Π^I/Π^{disc} are 2/-1 times those on the $I = 1$ contribution, whereas those on the $I = 0$ contribution vanish. The last line contains the total contribution corrected for finite-volume effects, the third uncertainty is the one associated with this correction. The fourth uncertainty corresponds to isospin breaking contributions (see text).




FIG. 3. Continuum extrapolations of the light, strange, charm and disconnected contributions to Π_1 . The lines are fits linear in a^2 , the different lines correspond to different cuts in the lattice spacing.

these are not significant statistically. If we assume no difference between the three averages and fit all ensembles and all three averages together in one fit, we get a reasonable fit quality $\chi^2/\text{dof} = 45/41$. Also, if we in-

clude additional fit parameters describing the difference between the three averages, they come out zero within error bars. This remains true after dropping the coarser lattices from the fit, for which the two-pion, finite-volume effects are reduced by taste violations. Therefore we average over the combinations ss , $4s$ and $s4$ in this work and leave an investigation of differences for the future, when statistical errors are reduced. The upper panel of Fig. 3 shows the continuum extrapolation of Π^I . The coarsest lattice gives a value about 15% smaller than the continuum limit. The final central value and systematic error on the continuum limit are the mean and standard error of the Akaike-Information-Criterion-weighted distribution obtained by imposing four different cuts on lattice spacing (no cut, $a \leq 0.118, 0.111, 0.095$ fm) in the extended frequentist approach of [41, 42]. The results for the first and the second moments are given in Table II.

The strange and charm quark contributions are plotted in the second and third panels of Fig. 3 and the respective continuum extrapolated values are given in Table II. The strange channel has much smaller lattice artefacts than the light, since it is much less affected by taste violations. The difference between the continuum and the coarsest lattice is about 2%. For the charm the fit qualities are much worse, because the precision of the data is orders of magnitude better than for the lighter flavors. Here we fit only a random subset of 10 configurations, which increases the statistical error and leads to good fit qualities. Results on the coarsest lattice deviate by about 20% from the continuum limit.

The lattice spacing dependence of the disconnected Π_1 is shown in the bottom panel of Fig. 3. We calculate the charm quark contribution to the disconnected term on the coarsest lattice: we find it to be 0.1% of the total disconnected result, i.e. much smaller than the total disconnected statistical error. We therefore discard the charm from the disconnected term at all lattice spac-

ings. Π_1^{disc} has the largest lattice artefacts among the four contributions. Since we have one less lattice spacing available than in the connected cases, we apply only three cuts in β . These extrapolations are also shown in Fig. 3.

The total result for Π_1 is the sum of the four contributions. It has a combined error of about 1.4%. All of these results apply to a box size of 6 fm and to the isospin symmetric case.

In the absence of a systematic study with simulations in a variety of volumes, only model estimates of finite-volume effects can be made. As argued in [43, 44], for large volumes those effects will be governed by pion contributions that can be computed in chiral perturbation theory (χ PT) [43]. Since the $I = 0$ channel is dominated by three-pion exchange, the finite-size effects are expected to be smaller than those of the $I = 1$ contribution, which are already small. Thus we consider only the latter. For all three index combinations, ss , $4s$ and $s4$, we calculate the difference between the infinite and finite volume moments at one loop in χ PT. We then take the average of the maximum and the minimum differences as our central value, with an uncertainty given by the half distance between the maximum and minimum. We record these corrections for Π_1 and Π_2 in Table II. For the first derivative the correction is on the level of 2%, whereas on the second derivative it is of order 10%. This correction increases rapidly with moment number, therefore we have chosen not to quote moments beyond the second one.

Concerning isospin breaking corrections, while little is known about how they modify the slope and curvature of the hadronic vacuum polarization function, more is known about their contribution to the anomalous magnetic moment of the muon. Compared to phenomenological determinations [4–6, 8], our $m_d = m_u$ calculation without QED is missing a number of effects². Noting their contributions to $a_\mu^{\text{LO-HVP}}$ in parentheses in units of 10^{-10} , these effects are $\rho\text{-}\omega$ (2.80 ± 0.19 from [45]) and $\rho\text{-}\gamma$ (-2.71 ± 0.27 from [46, 47]) mixing, final state radiation (3.86 ± 0.39 from [2] with a 10% error added), and the $\pi^0\gamma$ (4.42 ± 0.19 from [4]) and $\eta\gamma$ (0.64 ± 0.02 from [4]) contributions. This leads to a correction of $(9.0 \pm 0.5) \times 10^{-10}$, i.e. 1.3% of the result for $a_\mu^{\text{LO-HVP}}$ given in [4]. However, a competing effect enters. In our calculation without electromagnetism, the charged pion has a mass which is smaller than its physical value (see above). But a smaller charged-pion mass has the effect of enhancing the two-pion contribution to $a_\mu^{\text{LO-HVP}}$ and thus leads to a correction whose sign is opposite to the correction associated with the sum of effects discussed

above. Moreover, a phenomenological description based on e^+e^- data indicates that their magnitudes are very close [47]. Thus, we assume here that the total correction which has to be added to an isospin-limit determination of $a_\mu^{\text{LO-HVP}}$ is $(0.0 \pm 1.3)\%$, where we have taken the error to be of the typical size of the corrections themselves. Because of the dominant role which Π_1 plays in determining $a_\mu^{\text{LO-HVP}}$, one expects a tantamount correction on that coefficient. Inferring the correction on Π_2 is less direct, but we assume here that it is of the same size as for Π_1 . Thus, we add $(0.0 \pm 1.3)\%$ of Π_1 and Π_2 to our results for these quantities, after they have been corrected for finite-volume effects.

Putting everything together we quote our final results for the first two moments in the last row of Table II. Combining all four errors in quadrature, we obtain Π_1 with a total uncertainty of 2.9% and Π_2 of 7.2%.

Discussion.– It is interesting to compare these results with those in the literature. The only other lattice determination of Π_1 and Π_2 , near the physical mass point, comes from a single, $N_f = 2 + 1 + 1$ staggered simulation performed on a $(5.8^3 \times 7.7)$ fm⁴ lattice with spacing $a = 0.12$ fm [14, 48]. However, since the action used is different from ours and a comparison with our continuum limit results could be misleading, we choose not to exhibit such a comparison here.

The only phenomenological determination of the slope and derivative parameters, Π_1 and Π_2 , is the recent one of [25]. Taking their “data direct” results, which are obtained from an interpolation of $e^+e^- \rightarrow$ hadrons data, and converting them to our conventions, we get $\Pi_1 = 0.0990(7)$ GeV⁻² and $\Pi_2 = -0.2057(16)$ GeV⁻⁴. These numbers can be compared to our final results, i.e. those given in the last row of Table II. In absolute value, their result for Π_1 is 0.3 combined standard deviations smaller than ours and for Π_2 , 1.9 σ larger. The latter might be due to an underestimate of FV corrections in our determination of the second moment, or some problem with the experimental data used in the phenomenological analysis of [25].

Acknowledgments.– LL thanks M. Benayoun, C. Davies, F. Jegerlehner, C. Lehner, H. Leutwyler and E. de Rafael for very informative discussions. Computations were performed on JUQUEEN and JUROPA at Forschungszentrum Jülich, on Turing at the Institute for Development and Resources in Intensive Scientific Computing (IDRIS) in Orsay, on SuperMUC at Leibniz Supercomputing Centre in München, on Hermit at the High Performance Computing Center in Stuttgart. This project was supported in part by the OCEVU Laboratoire d’excellence (ANR-11-LABX-0060) and the A*MIDEX Project (ANR-11-IDEX-0001-02), which are funded by the “Investissements d’Avenir” French government program and managed by the “Agence nationale de la recherche” (ANR), by the Gauss Centre for Supercomputing e.V and by the GENCI-IDRIS

² We are grateful to Maurice Benayoun and Fred Jegerlehner for very informative discussions on this subject.

supercomputing grant No. 52275.

-
- [1] F. Jegerlehner, *PHIPSI08, proceedings of the International Workshop on $e+e-$ Collisions from ϕ to ψ , Frascati (Rome) Italy, 7-10 April 2008*, Nucl. Phys. Proc. Suppl. **181-182**, 135 (2008), arXiv:0807.4206 [hep-ph].
- [2] F. Jegerlehner and A. Nyffeler, Phys. Rept. **477**, 1 (2009), arXiv:0902.3360 [hep-ph].
- [3] J. P. Miller, E. de Rafael, B. L. Roberts, and D. Stöckinger, Ann. Rev. Nucl. Part. Sci. **62**, 237 (2012).
- [4] M. Davier, A. Hoecker, B. Malaescu, and Z. Zhang, Eur. Phys. J. **C71**, 1515 (2011), [Erratum: Eur. Phys. J. **C72**, 1874(2012)], arXiv:1010.4180 [hep-ph].
- [5] K. Hagiwara, R. Liao, A. D. Martin, D. Nomura, and T. Teubner, J. Phys. **G38**, 085003 (2011), arXiv:1105.3149 [hep-ph].
- [6] F. Jegerlehner, *Proceedings, Workshop on Flavour changing and conserving processes 2015 (FCCP2015)*, EPJ Web Conf. **118**, 01016 (2016), arXiv:1511.04473 [hep-ph].
- [7] G. W. Bennett *et al.* (Muon $g-2$), Phys. Rev. **D73**, 072003 (2006), arXiv:hep-ex/0602035 [hep-ex].
- [8] S. Eidelman and F. Jegerlehner, Z. Phys. **C67**, 585 (1995), arXiv:hep-ph/9502298 [hep-ph].
- [9] T. Blum, Phys. Rev. Lett. **91**, 052001 (2003), arXiv:hep-lat/0212018 [hep-lat].
- [10] C. Aubin and T. Blum, Phys. Rev. **D75**, 114502 (2007), arXiv:hep-lat/0608011 [hep-lat].
- [11] X. Feng, K. Jansen, M. Petschlies, and D. B. Renner, Phys. Rev. Lett. **107**, 081802 (2011), arXiv:1103.4818 [hep-lat].
- [12] M. Della Morte, B. Jager, A. Jüttner, and H. Wittig, JHEP **03**, 055 (2012), arXiv:1112.2894 [hep-lat].
- [13] F. Burger, X. Feng, G. Hotzel, K. Jansen, M. Petschlies, and D. B. Renner (ETM), JHEP **02**, 099 (2014), arXiv:1308.4327 [hep-lat].
- [14] B. Chakraborty, C. T. H. Davies, G. C. Donald, R. J. Dowdall, J. Koponen, G. P. Lepage, and T. Teubner (HPQCD), Phys. Rev. **D89**, 114501 (2014), arXiv:1403.1778 [hep-lat].
- [15] T. Blum, P. A. Boyle, T. Izubuchi, L. Jin, A. Jüttner, C. Lehner, K. Maltman, M. Marinkovic, A. Portelli, and M. Spraggs, Phys. Rev. Lett. **116**, 232002 (2016), arXiv:1512.09054 [hep-lat].
- [16] B. Chakraborty, C. T. H. Davies, J. Koponen, G. P. Lepage, M. J. Peardon, and S. M. Ryan, Phys. Rev. **D93**, 074509 (2016), arXiv:1512.03270 [hep-lat].
- [17] T. Blum *et al.* (RBC/UKQCD), JHEP **04**, 063 (2016), arXiv:1602.01767 [hep-lat].
- [18] T. Blum, M. Hayakawa, and T. Izubuchi, *Proceedings, 30th International Symposium on Lattice Field Theory (Lattice 2012)*, PoS **LATTICE2012**, 022 (2012), arXiv:1301.2607 [hep-lat].
- [19] F. Gray (New $(g-2)$), *Nuclear physics. Proceedings, 25th International Conference, INPC 2010, Vancouver, Canada, July 4-9, 2010*, J. Phys. Conf. Ser. **312**, 102006 (2011), arXiv:1009.0799 [nucl-ex].
- [20] M. Otani (E34), *Proceedings, 2nd International Symposium on Science at J-PARC: Unlocking the Mysteries of Life, Matter and the Universe (J-PARC 2014): Tsukuba, Japan, July 12-15, 2014*, JPS Conf. Proc. **8**, 025008 (2015).
- [21] J. S. Bell and E. de Rafael, Nucl. Phys. **B11**, 611 (1969).
- [22] E. de Rafael, Phys. Lett. **B736**, 522 (2014), arXiv:1406.4671 [hep-lat].
- [23] K. A. Olive *et al.* (Particle Data Group), Chin. Phys. **C38**, 090001 (2014).
- [24] C. Aubin, T. Blum, M. Golterman, and S. Peris, Phys. Rev. **D86**, 054509 (2012), arXiv:1205.3695 [hep-lat].
- [25] M. Benayoun, P. David, L. DelBuono, and F. Jegerlehner, (2016), arXiv:1605.04474 [hep-ph].
- [26] M. Luscher and P. Weisz, Commun. Math. Phys. **97**, 59 (1985), [Erratum: Commun. Math. Phys. **98**, 433(1985)].
- [27] C. Morningstar and M. J. Peardon, Phys. Rev. **D69**, 054501 (2004), arXiv:hep-lat/0311018 [hep-lat].
- [28] C. T. H. Davies, C. McNeile, K. Y. Wong, E. Follana, R. Horgan, K. Hornbostel, G. P. Lepage, J. Shigemitsu, and H. Trotter, Phys. Rev. Lett. **104**, 132003 (2010), arXiv:0910.3102 [hep-ph].
- [29] R. Bellwied, S. Borsanyi, Z. Fodor, S. D. Katz, A. Pasztor, C. Ratti, and K. K. Szabo, Phys. Rev. **D92**, 114505 (2015), arXiv:1507.04627 [hep-lat].
- [30] S. Aoki *et al.*, (2016), arXiv:1607.00299 [hep-lat].
- [31] J. Gasser and G. R. S. Zarnauskas, Phys. Lett. **B693**, 122 (2010), arXiv:1008.3479 [hep-ph].
- [32] M. Lüscher, JHEP **08**, 071 (2010), [Erratum: JHEP **03**, 092(2014)], arXiv:1006.4518 [hep-lat].
- [33] S. Borsanyi *et al.*, JHEP **09**, 010 (2012), arXiv:1203.4469 [hep-lat].
- [34] M. A. Clark and A. D. Kennedy, Phys. Rev. Lett. **98**, 051601 (2007), arXiv:hep-lat/0608015 [hep-lat].
- [35] M. A. Clark, B. Joo, A. D. Kennedy, and P. J. Silva, Phys. Rev. **D84**, 071502 (2011), arXiv:1108.1828 [hep-lat].
- [36] H. Yin and R. D. Mawhinney, *Proceedings, 29th International Symposium on Lattice field theory (Lattice 2011)*, PoS **LATTICE2011**, 051 (2011), arXiv:1111.5059 [hep-lat].
- [37] T. Blum, T. Izubuchi, and E. Shintani, Phys. Rev. **D88**, 094503 (2013), arXiv:1208.4349 [hep-lat].
- [38] V. Gülpers, A. Francis, B. Jäger, H. Meyer, G. von Hippel, and H. Wittig, *Proceedings, 32nd International Symposium on Lattice Field Theory (Lattice 2014): Brookhaven, NY, USA, June 23-28, 2014*, PoS **LATTICE2014**, 128 (2014), arXiv:1411.7592 [hep-lat].
- [39] X. Feng, S. Hashimoto, G. Hotzel, K. Jansen, M. Petschlies, and D. B. Renner, Phys. Rev. **D88**, 034505 (2013), arXiv:1305.5878 [hep-lat].
- [40] M. Luscher, Nucl. Phys. **B364**, 237 (1991).
- [41] S. Durr *et al.* (Budapest-Marseille-Wuppertal), Science **322**, 1224 (2008), arXiv:0906.3599 [hep-lat].

- [42] S. Borsanyi *et al.* (Budapest-Marseille-Wuppertal), *Science* **347**, 1452 (2015), arXiv:1406.4088 [hep-lat].
- [43] C. Aubin, T. Blum, P. Chau, M. Golterman, S. Peris, and C. Tu, *Phys. Rev.* **D93**, 054508 (2016), arXiv:1512.07555 [hep-lat].
- [44] A. Francis, B. Jaeger, H. B. Meyer, and H. Wittig, *Phys. Rev.* **D88**, 054502 (2013), arXiv:1306.2532 [hep-lat].
- [45] M. Davier, A. Hoecker, G. Lopez Castro, B. Malaescu, X. H. Mo, G. Toledo Sanchez, P. Wang, C. Z. Yuan, and Z. Zhang, *Eur. Phys. J.* **C66**, 127 (2010), arXiv:0906.5443 [hep-ph].
- [46] F. Jegerlehner and R. Szafron, *Eur. Phys. J.* **C71**, 1632 (2011), arXiv:1101.2872 [hep-ph].
- [47] M. Benayoun and F. Jegerlehner, *private communication, November 2016*.
- [48] B. Chakraborty, C. T. H. Davies, P. G. de Oliveira, J. Koponen, and G. P. Lepage, (2016), arXiv:1601.03071 [hep-lat].